

Molecular mechanisms in allergy and clinical immunology

Series editors: Joshua A. Boyce, MD, Fred Finkelman, MD, William T. Shearer, MD, PhD, and Donata Vercelli, MD

Asthma therapy and airway remodeling

Thais Mauad, MD, PhD,^a Elisabeth H. Bel, MD, PhD,^b and Peter J. Sterk, MD, PhD^b

São Paulo, Brazil, and Amsterdam, The Netherlands

This activity is available for CME credit. See page XXX for important information.

Asthma is characterized by variable degrees of chronic inflammation and structural alterations in the airways. The most prominent abnormalities include epithelial denudation, goblet cell metaplasia, subepithelial thickening, increased airway smooth muscle mass, bronchial gland enlargement, angiogenesis, and alterations in extracellular matrix components, involving large and small airways. Chronic inflammation is thought to initiate and perpetuate cycles of tissue injury and repair in asthma, although remodeling may also occur in parallel with inflammation. In the absence of definite evidence on how different remodeling features affect lung function in asthma, the working hypothesis should be that structural alterations can lead to the development of persistent airway hyperresponsiveness and fixed airway obstruction. It is still unanswered whether and when to begin treating patients with asthma to prevent or reverse deleterious remodeling, which components of remodeling to target, and how to monitor remodeling. Consequently, efforts are being made to understand better the effects of conventional anti-inflammatory therapies, such as glucocorticosteroids, on airway structural changes. Animal models, *in vitro* studies, and some clinical studies have advanced present knowledge on the cellular and molecular pathways involved in airway remodeling. This has encouraged the development of biologicals aimed to target various components of airway remodeling. Progress in this area requires the explicit linking of modern structure-function analysis with innovative biopharmaceutical approaches. (J Allergy Clin Immunol ■■■■;■■■:■■■-■■■.)

Key words: Asthma, therapy, remodeling, corticosteroids, leukotriene receptor antagonist, airway, basement membrane, smooth muscle, epithelium, extracellular matrix

Abbreviations used

ASM: Airway smooth muscle
BAL: Bronchoalveolar lavage
BM: Basement membrane
CTGF: Connective tissue growth factor
ECM: Extracellular matrix
FP: Fluticasone propionate
MMP: Matrix metalloproteinase
PDE: Phosphodiesterase
VEGF: Vascular endothelial growth factor

Asthma is a clinical diagnosis based on episodic symptoms and variable airways obstruction.¹ It has been generally recognized that the disease is also characterized by variable degrees of chronic inflammation and structural alterations in the airways.^{2,3} The structural alterations, collectively called *airway remodeling*, encompass complex changes in composition, content, and organization of the various cellular and molecular constituents of the airway wall.⁴ The most striking abnormalities are epithelial denudation, goblet cell metaplasia, subepithelial thickening, increased airway smooth muscle mass, bronchial gland enlargement, angiogenesis, and alterations in the extracellular matrix (ECM) components.⁵ These alterations involve large and small airways and the surrounding peribronchiolar areas³⁻⁶ (Fig 1).

Traditionally, it is thought that T_H2-mediated chronic inflammation triggers and perpetuates a vicious circle of tissue injury–tissue repair culminating in remodeling. Data on pediatric severe asthma showing the early presence of structural alterations in bronchial tissue raised the alternative hypothesis that inflammation and remodeling may occur in parallel, beginning at early stages of the disease.^{7,8} During normal lung development, there is a cross-talk between the bronchial epithelium and the underlying mesenchymal cells (the bronchial epithelial mesenchymal trophic unit), which is fundamental for adequate branching morphogenesis.⁹ Davies et al⁴ proposed that abnormal injury occurring in genetically susceptible bronchial epithelium because of components of the inhaled

From ^athe Department of Pathology, São Paulo University Medical School; and ^bthe Department of Respiratory Diseases, Academic Medical Center, University of Amsterdam.

Supported by Conselho Nacional de Desenvolvimento científico e Tecnológico (Brazilian National Research Council).

Disclosure of potential conflict of interest: The authors have declared that they have no conflict of interest.

Received for publication April 11, 2007; revised June 20, 2007; accepted for publication June 22, 2007.

Reprint requests: Peter J. Sterk, MD, PhD, Department of Respiratory Diseases, F5-259, Academic Medical Center, University of Amsterdam, PO Box 22700, NL-1100 DE Amsterdam, The Netherlands. E-mail: p.j.sterk@amc.nl.

0091-6749/\$32.00

© 2007 American Academy of Allergy, Asthma & Immunology

doi:10.1016/j.jaci.2007.06.031




FIG 1. A and C, Normal airways. B and D, Airways from a patient with fatal asthma. Mucus plug within the airway lumen, epithelial folding and thickened ASM layer in B. Mucus plugging and increased ASM thickness in D, with spreading of the inflammation to the surrounding peribronchiolar alveoli.* Hematoxylin and eosin. A and B, $\times 25$. C and D, $\times 100$. C, Cartilage; Ep, epithelium; M, mucus.

environment could reactivate the epithelial mesenchymal trophic unit in asthma, driving smooth muscle proliferation and remodeling. Therefore, myofibroblasts, mesenchymal cells that have the capacity to secrete ECM proteins and growth factors, are believed to play a key role in the remodeling process.

The current working hypothesis is that airway remodeling could explain at least some aspects of disease severity in animal models, such as airway hyperresponsiveness¹⁰ and fixed airway obstruction.¹¹ Southam et al¹² presented recent data on chronic allergen-sensitized mice, showing that a complex temporal interplay may exist between airway inflammation, structural changes, and the onset/maintenance of airway hyperresponsiveness. With regard to the human situation, we know that many patients exhibit persistent airway hyperresponsiveness, fixed airflow limitation, excessive airway narrowing, and/or enhanced airway closure¹³ despite maximal therapy.¹⁴ It is very likely that components of airway remodeling contribute to this, particularly in severe asthma. However, definite clinical evidence of a causal relationship is still missing. We do not know whether and how each component of airway remodeling, alone or in combination, affects lung function. Previous studies comparing fatal asthma and nonfatal asthma reported associations between increased airway thickness and asthma severity.¹⁵ Biopsy studies in patients with different asthma severities also demonstrated that increased collagen content and airway

smooth muscle (ASM) cell mass (among other remodeling features within the airway mucosa) were present in patients with severe asthma compared with patients with moderate and mild asthma.¹⁵⁻¹⁹ Other studies, however, had more difficulties in finding differences in collagen content between patients with asthma of different severities,²⁰ or associations between basement membrane thickening and progressive decline in lung function.²¹ Hence, the relevance of the different components of airways remodeling for the clinical severity and progression of asthma still remains to be established. As a consequence, the question is which components of airway remodeling should be treated.

Tissue turnover and restructuring is a physiological, homeostatic process.²² This may help to preserve optimal functional properties of the airways.²³ Airway wall thickening and loss of alveolar attachments²⁴ in asthma are likely to enhance airway narrowing.²⁵ However, the airways in patients with asthma are stiffer than in normal subjects,^{26,27} possibly because of fibrosis, mucosal folding, and/or smooth muscle stiffening,^{28,29} which may actually counteract airway narrowing.³⁰ Therefore, before considering any interventions aimed to prevent or reverse structural changes of the airways in asthma, there should be good evidence that the targeted tissue elements contribute to impaired airway function.

This raises a large challenge to the field. How should airway structure be sampled and monitored? Endobronchial

biopsies are sampling a very superficial and selected area of the airways,^{31,32} disregarding the small airways that are extremely difficult to assess.³³ Although high-resolution computed tomography scanning allows quantitative airway morphometry,³⁴ it cannot distinguish the histologic components of the airways. Hence, when targeting airway remodeling by therapeutic interventions, the outcome parameters should be carefully specified.

To assess the effects of different drugs on the cellular and molecular components of airway remodeling, animal experiments and *in vitro* models are indispensable despite their intrinsic limitations.³⁵ In such models, corticosteroids have been studied intensively, because these drugs are the most effective therapy for the control of airway inflammation in asthma.³⁶ However, less is known about how corticosteroids affect structural cells involved in remodeling, such as fibroblasts, myofibroblasts, ASM cells, and epithelial cells.³⁷ That is why the current review first focuses on data describing how corticosteroids and other currently available drugs affect each component of airway remodeling. Thereafter, we explore newly developed interventions³⁸ and their potential for influencing airway compartments involved in remodeling in asthma.

EFFECTS OF CURRENTLY AVAILABLE ASTHMA DRUGS ON AIRWAY STRUCTURE

Bronchial epithelium

Increased epithelial shedding caused by epithelial fragility is a feature frequently described in asthma,³⁹ leading to mucosal denudation and increased exposure of the mucosal nerve endings to irritant factors, enhanced penetration of allergens, and reduced mucociliary clearance (Fig 2). Although part of the observed shedding is probably artifactual because of bronchoscopy techniques,⁴⁰ the frequent sloughing of epithelial cells observed in bronchoalveolar lavage (BAL) fluid from patients with asthma suggests that some degree of epithelial shedding does occur in asthma.³¹ Goblet cell metaplasia is another important feature of asthma that, together with the bronchial gland enlargement, leads to hypersecretion and airway obstruction.

There is controversy about the role of corticosteroids on epithelial damage in asthma. Some *in vitro* studies suggested that corticosteroids increase apoptosis of epithelial cells, which could further contribute to the chronic epithelial damage already present in this disease.⁴¹ In contrast, glucocorticoids can inhibit cell death induced by cytokines.⁴² In mechanically denuded guinea pig tracheal epithelium, budesonide did not interfere with an efficient restitution of the epithelium,⁴³ whereas dexamethasone prolonged the repair potential after repeated episodes of epithelial injury in mucociliated human bronchial epithelial cells.⁴⁴

Defective mechanisms of epithelial repair are believed to occur in asthma. Persistent activation of the epithelial growth factor receptor with paradoxical increased expression of antiproliferative markers such as p21/waf have been described in adult and pediatric patients with severe

asthma,^{45,46} and both features seem to be unresponsive to steroid treatment. Hence, sustained epithelial activation associated with chronic inflammation is proposed to be the major trigger of the remodeling process.⁴ Partial restoration of the epithelium has been observed after the use of corticosteroids in a retrospective biopsy study⁴⁷ showing that treating patients with asthma with inhaled corticosteroids for 10 years decreased inflammation and partially improved epithelial damage. However, this was not associated with improvement of bronchial hyperresponsiveness. When examining this prospectively, compared with bronchodilator treatment alone, 3 months of budesonide therapy in patients with asthma increased the number of ciliated cells.⁴⁸

Goblet cell metaplasia and mucus hypersecretion are features of asthma, and airway mucus hypersecretion is recognized as an important contributor to morbidity and mortality in patients with lung diseases such as asthma and chronic obstructive pulmonary disease. Patients with asthma have altered mucin expression in the goblet cells, but the clinical or functional significance of this different composition is still unclear.⁴⁹ Corticosteroids and bronchodilators are not primarily targeted to act on goblet cell activity but seem to have direct and indirect suppressive effects on mucus production.⁴⁹ In addition, in animal models, corticosteroids appear to be effective in reducing goblet cell metaplasia.^{50,51} The effects of ciclesonide and fluticasone propionate (FP) on reducing goblet cell metaplasia in a rat model of allergic inflammation were comparable.⁵⁰ Apparently, this also occurs in human asthma, because De Kluijver et al⁵² observed a decrease in the number of goblet cells in patients with mild asthma after 2 weeks of inhaled corticosteroids.

Are there alternatives to corticosteroids available with regard to effects on bronchial epithelium? In mice, leukotriene receptor antagonists have a positive effect on reducing epithelial changes induced by allergic sensitization.^{53,54} In addition, pranlukast and zafirlukast greatly suppressed ovalbumin-induced secretion in the guinea pig trachea, suggesting a role of these drugs in the treatment of airway diseases with a hypersecretory component.⁵⁵

Basement membrane components and thickness

Because of its constant accessibility in bronchial biopsies, the basement membrane has been studied extensively in asthma⁶ (Fig 2). It has been considered “the remodeling marker” in several studies. It consists of the thickening of the subepithelial lamina reticularis that lies underneath the true basal lamina of the bronchial epithelium, sometimes referred to as *pseudo-thickening*. In patients without asthma, this structure measures approximately 5 to 6 μm , whereas it is consistently thickened in biopsies of patients with asthma with mean values of approximately 9 μm .⁵⁶ A study in symptomatic infants with reversible airway obstruction could not demonstrate basement membrane (BM) thickening,⁵⁷ whereas it was already present in older children with severe asthma,⁷ suggesting that BM thickness develops early in the disease course. Further, an association between BM thickness and disease duration and




FIG 2. A, Normal bronchial mucosa. The epithelium is intact and composed of ciliated columnar cells. **B,** Bronchial mucosa from a patient with fatal asthma. There is epithelial damage and basement membrane thickening (arrow). The lamina propria is thickened with inflammation and numerous capillaries. * The ASM layer is thickened. Hematoxylin and eosin. $\times 200$. Ep, Epithelium; SMG, submucosal glands.

disease duration or severity have not been consistently demonstrated.⁷

Accumulation of collagen III, I, and IV and fibronectin at the BM level occurs in patients with asthma.⁵⁸ More recent ultrastructural studies reported that accumulation of (thinner) reticulin fibers and not (thicker) interstitial collagen fibers accounts for the thickened BM in asthma, suggesting that this phenomenon in asthma is different from the collagen deposition observed in interstitial lung diseases.⁵⁹ It must be stressed that in sensitized animals, the typical characteristic hyaline thickening of the BM as seen in human asthma is not observed. What has been called *subepithelial thickening* in these models would correspond to the entire lamina propria in human airways.

Several studies assessed the effects of asthma treatment in the reversibility of BM thickness. Differences in

methodology, small samples, different corticosteroids and doses, and timing make results seemingly controversial and difficult to interpret. Some studies could not demonstrate reductions in BM thickness after the use of inhaled corticosteroids,^{60,61} whereas others could.^{62,63} The latter studies suggest that reducing BM thickness may require prolonged and high-dose inhaled steroid therapy. Reduction in collagen type III deposition, decrease of matrix metalloprotease (MMP)-9, and increase in tissue inhibitor of metalloprotease 1 levels could be involved in the mechanisms of the reversal of BM thickness.⁶⁴ It remains to be established whether such a decrease in BM thickness has any benefits on airway function.

Laminin is a glycoprotein found at the airway BM level, playing a major role in airway morphogenesis, particularly of the smooth muscle. Altraja et al⁶⁵ have previously

observed increased deposition of laminin chains at BM level in patients with chronic and occupational asthma. This could reflect an increased tissue turnover in the airway wall, possibly as a result of airway inflammation. Concomitant treatment with dexamethasone reduced laminin and receptor levels within the airways in an experimental model of asthma, but this was not accompanied by a reduction in airway hyperresponsiveness.⁶⁶ Furthermore, the expression of tenascin, one of the BM glycoproteins, was found to be decreased after the use of inhaled corticosteroids during the birch pollen season, but without any corresponding improvements in lung function.⁶⁷

All these studies point toward the capability of corticosteroids to change the thickness and constitution of the bronchial reticular basement membrane in asthma, but unless its functional benefit has been demonstrated, this should not be considered a therapeutic target.

Lamina propria

The lamina propria of the airways is composed of structural cells such as fibroblasts and myofibroblasts, vessels, ECM components, and inflammatory cells. In large airways, this wall layer accounts for a significant portion of the airway (Fig 2).

Fibroblasts and myofibroblasts. Fibroblasts and myofibroblasts play a crucial role in the mechanisms of altered airway structure in asthma because of their capacity to secrete growth factors and ECM elements.⁶⁸ It has been observed that the steroid triamcinolone acetonide and the combination of FP and salmeterol have the ability to downregulate fibroblast proliferation *in vitro*.^{69,70} However, studies using fibroblasts from patients with asthma show that dexamethasone can increase fibroblast proliferation and stimulate G1-S phase transition.^{71,72} This is an important observation, suggesting that corticosteroids may also promote elements of airway remodeling in asthma. FP displays anti-inflammatory effects on human lung fibroblasts during their myofibroblastic differentiation. At early stages of differentiation, FP inhibits the activation of Janus kinase/signal transducer and activator of transcription pathways induced by IL-13 in lung myofibroblasts.⁷³ FP also inhibits constitutive and TGF- β -induced expression of α -smooth muscle actin in fibroblasts, the main marker of myofibroblastic differentiation, both in very early and in mild differentiated myofibroblasts.⁷³ These *in vitro* results have been extended by *in vivo* studies. In a mouse model of chronic allergic sensitization, 3 months of systemic corticosteroids reduced myofibroblasts (defined as the cells that expressed α -smooth muscle actin and collagen I), TGF- β expression, and peribronchial fibrosis, but not ASM thickness.⁷⁴ However, to our knowledge, there are no clinical studies available on the effects of corticosteroids on fibroblast or myofibroblast activity and proliferation.

It is interesting to note that theophylline inhibits fibroblast proliferation and suppresses TGF- β -induced collagen I mRNA *in vitro*.⁷⁵ Furthermore, recent data suggest that the leukotriene receptor antagonist montelukast may lead to a decrease in airway wall myofibroblasts.⁷⁶

This indicates that it is worth examining other interventions than just corticosteroids in changing fibroblast or myofibroblast function in asthma.

Extracellular matrix elements. Collagens, elastin, proteoglycans, and glycoproteins compose the extracellular matrix of the airway wall and are mainly secreted by structural cells such as fibroblasts, myofibroblasts, and ASM cells.⁶⁸ Altered content and composition of extracellular matrix have been described in large and small airways of patients with asthma of different severities.^{52,77-81}

Aerosolized corticosteroids have shown to be effective in preventing and reversing enhanced fibronectin deposition during concomitant repeated allergen exposure.^{82,83} However, these effects appeared to be dependent on the timing^{82,83} and dose of the inhaled corticosteroids.⁸⁴ Treatment with low doses or treatment after allergen exposure had no effect on fibronectin deposition, which suggests that adequate therapy during a particular time window may be required to prevent and suppress components of airway remodeling.

There are few data on the ECM composition in the entire lamina propria in patients with asthma with or without antiasthma treatment. Wilson and Li⁷⁷ have shown that the bronchial submucosal region of patients with asthma has more type III and type V collagen than controls, but these results were not reproduced by Chu et al²⁰ when examining patients with asthma of different severities and controls. Patients with moderate-severe asthma have increased deposition of type III, type I collagens, and the profibrotic cytokines IL-11, IL-17, and TGF- β compared with healthy controls and patients with mild asthma.⁸⁵ When patients with asthma were treated with 2 weeks of oral corticosteroids, the levels of IL-11 and IL-17, but not of TGF- β or collagens, were reduced.⁸⁵ These results again suggest that some components of airway remodeling may not be responsive to steroid therapy.

Proteoglycans are important ECM components of the airway wall and are involved in mechanics, water balance, regulation of inflammation, cell migration, and proliferation.⁸⁶ Previous studies have demonstrated differential deposition of its components within the airways.^{52,79,81} The combination of budesonide and formoterol inhibits the serum-induced production of proteoglycans *in vitro*.⁸⁷ However, in asthmatic airways *in vivo*, we have observed that a 2-week course of inhaled corticosteroids resulted in increased density of the proteoglycans versican and biglycan, without associated changes in fibronectin or BM thickness.⁵² This study shows that treatment with corticosteroids differentially affects ECM components in asthma, but again it is unclear whether this has functional relevance. Long-term longitudinal studies are surely needed to determine whether the steroid induced changes are beneficial or detrimental to lung function decline in asthma.

Elastin is a major component of the lung ECM, presumably having a pivotal role in airway patency and lung elastic recoil. Changes in elastin have been described in patients with asthma,^{23,80,88} but to our knowledge, there are no intervention studies addressing elastin content in asthma.

Bronchial vessels

Engorgement of bronchial vessels at the large and small airway levels is a feature of asthma, and morphometric studies have revealed increases in both the number of vessels and the vessel area.⁸⁹⁻⁹² Furthermore, increased microvascular permeability is observed in asthmatic airways.⁹³ These features could indirectly amplify the inflammatory response and contribute to enhanced airway thickness.⁹⁴

Inhaled corticosteroids can decrease airway vascularity in the airways of patients with asthma, which is associated with a decrease in BM thickness, FEV₁, and airway responsiveness.⁹⁵ Different corticosteroids seem to have different potencies in the reduction of airway vasoconstriction, with FP and budesonide causing greater airway vasoconstriction than beclomethasone dipropionate.⁹⁶ Corticosteroids may also inhibit angiogenesis by regulating growth factors such as vascular endothelial growth factor (VEGF), an important regulator of angiogenesis. In patients with asthma treated with 800 µg/d beclomethasone, VEGF levels in induced sputum were decreased, and were associated with the degree of airway narrowing and vascular permeability.⁹⁷ Taken together, high doses of corticosteroids seem to be necessary to reduce structural changes in airway vessels that are accompanied by decreases in VEGF expression.⁹⁴ Feltis et al⁹⁸ recently extended these results, showing that reduction in subepithelial vascularity after 3 months of treatment with high-dose FP (750 µg twice daily) was associated with a concomitant reduction in VEGF-positive vessels, a reduction in angiogenic sprouts per vessel, and a decrease in VEGF receptors and in angiopoietin 1 in steroid-naïve patients with asthma.

There are few data about the effects of other antiasthma drugs on vessel remodeling. β₂-Agonists can have an inhibitory effect on plasma exudation and vascular permeability.⁹⁹ Indeed, adding salmeterol to low-dose inhaled steroid treatment (200–400 µg/d beclomethasone dipropionate [BDP] or 200–400 mg/d budesonide) results in decreased vessel density in the lamina propria of patients with asthma.¹⁰⁰ Leukotriene receptor antagonists may also have beneficial effects on vascular permeability and airway mucosa blood flow. In sensitized mice, the use of a leukotriene receptor antagonist led to a decrease in vascular permeability and a decrease in VEGF levels.¹⁰¹ In human beings, 2 weeks of daily treatment with 10 mg montelukast or 400 µg FP appeared to have equivalent effects on reducing airway mucosal blood flow in patients with asthma.¹⁰² So far, no human studies have addressed the effects on leukotriene receptor antagonists on the structural changes of the bronchial microvasculature.

ASM cells

Increase in ASM mass, caused by cell hyperplasia, hypertrophy, and/or increased extracellular matrix deposition, is an important component of the structurally altered airway in asthma (Figs 1 and 2) and is presumed to be a major determinant of enhanced bronchoconstriction and

airway hyperresponsiveness.¹⁰³ Increase in the ASM mass occurs along the entire airway tree, and it is the major contributor to the increased area of the inner airway wall in asthma.¹⁰³ Increases in ASM mass have been related to asthma severity,^{15,18} whereas smooth muscle amount and changes in its plasticity are likely to be key players in the dynamics of airway narrowing.^{29,104,105}

A shift toward a synthetic phenotype with increased proliferation rates of the ASM cells is believed to play a role in the mechanisms leading to the ASM thickening.¹⁰⁶ Several mediators in the asthmatic airways, such as contractile agonists, cytokines, growth factors, and ECM proteins, can induce ASM proliferation.¹⁰⁶ However, it must be emphasized that increased proliferation rates have not been confirmed in human beings *in vivo* yet. It is possible that ASM *in vivo* proliferation occurs very slowly, occurs only intermittently, or is already completed before or at the onset of asthma.¹⁰³

The mechanisms by which corticosteroids could affect ASM proliferation are of great interest and are currently being clarified.¹⁰⁷ Corticosteroids not only modulate the secretion of chemokines and cytokines that are involved in ASM function and proliferation,¹⁰⁸ but also can exert a direct effect on ASM cells. *In vitro* experiments have shown that the corticosteroids dexamethasone and FP arrest human ASM cells in the G1 phase of the cell cycle and inhibit some but not all growth factor–induced proliferation of ASM cells.¹⁰⁹ Corticosteroids have the capacity to inhibit the action of some, but not all, mitogens involved in ASM proliferation. Corticosteroids inhibit the ones signaling via G protein coupled receptors, but less efficiently the ones signaling via receptor tyrosine kinase pathway.¹¹⁰ A mechanism proposed to explain the absence of an effective antiproliferative action by corticosteroids on ASM cells of patients with asthma would be the absence of the CCAAT/enhancer binding protein α.¹¹¹ This protein forms a complex with the glucocorticoid receptor, leading to activation of the cell cycle inhibitor p21; its absence could explain the failure of glucocorticoids to inhibit ASM proliferation of asthmatic cell cultures *in vitro*.

It has been recently observed that corticosteroids have significant inhibitory effects on ASM contractile protein expression by reducing human ASM-actin protein abundance and incorporation into contractile filaments.¹¹² The authors suggest that these effects appear to be mediated via the attenuation of mRNA translation and enhancement of protein degradation.

Extracellular matrix components are known to affect ASM growth and its synthetic properties, and it has been hypothesized that altered composition of the ECM could contribute to ASM dysfunction in asthma.¹¹³ It was recently demonstrated that culturing these cells on collagen type I prevented the antimitogenic actions of glucocorticoids, but not of β₂-adrenoceptor agonists. In contrast, glucocorticoids were efficient in regulating ASM production of GM-CSF, whereas β₂-adrenoceptor agonists were not.¹¹⁴ These results suggest that combination therapy may have increased efficacy over glucocorticoids alone in controlling remodeling events.

When treating smooth muscle cells in culture after exposure to 10% serum from an individual with asthma with or without beclomethasone (0.01-100 nM), it appeared that there was a significant increase in the production of ECM components in the cells cultured with the asthmatic serum. However, beclomethasone did not reverse the increase in ECM protein.¹¹⁵ Burgess et al¹¹⁶ extended these results, showing that treating cells with a corticosteroid or long-acting β_2 -agonist did not inhibit TGF- β -stimulated release of connective tissue growth factor (CTGF), collagen I, fibronectin, and versican. Moreover, corticosteroid alone increased the release of CTGF, collagen I and fibronectin. These findings raise the point that corticosteroids may not be effective in reducing ECM deposition and that they may even increase matrix production in certain circumstances, as demonstrated by De Kluijver et al⁵² in bronchial biopsies.

To date, there are no longitudinal studies evaluating the effect of pharmacologic therapy on the structure of ASM in asthma. In a murine model of asthma, daily administration of intranasal FP for 3 months inhibited the thickening of the ASM layer, decreased TGF- β and its intracellular effectors pSmad2 and pSmad3, and increased pSmad7 levels.¹¹⁷ However, even though 3 months of systemic usage of dexamethasone prevented many structural alterations in mice submitted to allergic sensitization, it did not decrease ASM thickness.⁷⁴ Hence, the route of administration may be one of the determinants of steroid effects on ASM. Interestingly, the combination of allergen avoidance and treatment with dexamethasone for 1 month was capable of reverting airway remodeling in chronically sensitized mice, including the thickened ASM layer.¹¹⁸

How about other available antiasthma drugs? β -Agonists cause increases in the levels of 3',5' cAMP in ASM cells, and besides their bronchodilator effect could also inhibit mitogen-induced proliferation on ASM cells.¹¹⁹ Maintenance treatment with the long-acting anticholinergic, tiotropium bromide, appears to prevent ASM thickening and to decrease myosin expression and ASM contractility in a guinea pig model of chronic allergic sensitization.¹²⁰ In addition, montelukast reversed increased smooth muscle mass and subepithelial fibrosis in an animal model of chronic allergic sensitization,⁵³ although lower doses of leukotriene receptor antagonists were not able to decrease ASM mass in another experimental study.⁵⁴ Given the emerging importance of smooth muscle activity, proliferation, and plasticity in the pathogenesis of asthma,¹²¹ these findings indicate that it is worth revisiting the effects of widely used asthma therapy on these cells in asthma.

The issue has been raised that none of the physiological functions of the normal ASM are essential to lung physiology.¹²² Ablation of the ASM by thermoplasty has therefore been proposed as an alternative to treat/cure asthma.¹²³ Preliminary data indicate that bronchial thermoplasty reduces ASM¹²⁴ and increases airway compliance¹²⁵ without serious complications. The first uncontrolled study suggested that this intervention led to improvement in FEV₁ and in airway hyperresponsiveness.¹²⁶ However, a

recent randomized, controlled trial in patients with moderate to severe asthma reported improvements in the rate of mild exacerbations, morning expiratory peak flow, and symptom scores, but no effect on FEV₁ and airway responsiveness after 1 year of follow-up.¹²⁷ Even though it cannot be excluded that bronchial thermoplasty in a nonblind design may have introduced some level of placebo effect on clinical outcome, the current results certainly warrant more detailed studies on histologic and functional outcomes.

NEW DRUGS

Cytokine modulators

Anti-TNF- α . Evidence suggests that the TNF- α axis is upregulated in patients with refractory asthma, as indicated by increased levels of TNF- α , its receptors and converting enzyme in blood monocytes, bronchoalveolar lavage, and bronchial biopsies.^{128,129} Two (1 controlled, 1 uncontrolled) studies in patients with severe asthma revealed that the use for 10 to 12 weeks of etanercept, which binds specifically to both TNF- α and TNF- β , thereby preventing free cytokine binding to TNF receptors, resulted in decreases in asthma symptoms and bronchial hyperresponsiveness and an increase in lung function in absence of changes of cellular inflammation in induced sputum.^{128,129} This might point toward possible changes in airway remodeling, perhaps in ASM structure, which need to be addressed. Infliximab is a recombinant human-murine chimeric mAb that binds and neutralizes the soluble TNF- α homotrimer and its membrane-bound precursor. This drug was tested in patients with moderate asthma (3 infusions in 6 weeks) and resulted in the decrease of moderate exacerbations and sputum cytokine levels, but no effect was noted in lung function parameters. The drug was well tolerated.¹³⁰ The use of 2 dual inhibitors of TNF- α -converting enzyme and matrix metalloproteinases (PKF242-484, PKF241-466) in murine models of acute airway inflammation showed significant reductions on lung inflammation.¹³¹ If the adverse effects of this treatment remain to be medically acceptable, exploration of anti-TNF- α strategies in asthma require priority in the clinical research of severe asthma.

Blocking T_H2 cytokines. T_H2 cytokines play an important role in orchestrating the inflammatory pathways of allergic asthma. Attempts to develop drugs that block T_H2 cytokine action or inhibit their synthesis or release are therefore the subject of long-standing investigation.¹³² A nonspecific inhibitor, suplatast tosilate, suppresses selectively the synthesis of IL-4, IL-5 *in vitro*¹³³ and is available as an asthma controller in Japan. Beneficial effects on inflammation, symptoms, and lung function have been reported in steroid-dependent patients with asthma after 8 weeks of use.¹³⁴ Part of this effect may be a result of a decrease in PAS/alcian blue-positive bronchial goblet cells, as was measured by MUC5AC staining in bronchial epithelium by Hoshino et al.¹³⁵

TABLE I. Potential therapeutic targets with effect on airway structure in animal models

Target	Drug	Species	Effect on inflammatory parameters	Effect on airway structure	Reference
CCR3	Low-molecular-weight CCR3 antagonist	Mouse	Reduction in eosinophils, no changes in lymphocytes or macrophages	Prevention of goblet cell hyperplasia, subepithelial fibrosis, and accumulation of myofibroblasts	152
IL-13	Neutralizing IL-13 mAb	Mouse	Reduction in eosinophils, no changes in inflammatory cells	Reduction in mucus production and bronchiolar collagen deposition	153
TGF- β	1. Neutralizing anti-TGF- β antibody	1. Mouse	1. No effect on inflammation and T _H 2 cytokine production	1. Reduced peribronchiolar ECM deposition, ASM proliferation, and mucus production	1. 154
	2. Inhibitor of TGF- β receptor I kinase (SD-208)	2. Rat	2. Decrease in eosinophils, CD2 T-cell counts, and Smad2/3 intracellular expression	2. Decreased epithelial and ASM cell proliferation and goblet cell hyperplasia	2. 155
Tryptase	Tryptase inhibitor MOL 6131	Mouse	Reduction in total cells, eosinophils, and in the release of IL-4 and IL-13 in BAL fluid, reduction in airway tissue eosinophilia	Reduction in goblet cell hyperplasia, mucus secretion, and peribronchial edema	156
Immunomodulation	Immunostimulatory sequences—oligodeoxy-nucleotides (CpG-DNA)	1. Mouse	1. Decrease in VEGF, MMP-9 in bronchial epithelium, TGF- β levels, and CD4 T lymphocytes	1. Prevents collagen deposition and mucus production, reduction in angiogenesis, reversion of peribronchial collagen deposition.	1. 157-160
		2. Monkey	2. Fewer eosinophils and interstitial mast cells (no differences in glands or ASM mast cells)	2. Thinner basement membranes and fewer mucus cells	2. 161
Toll-like receptor 7/8	Toll-like receptor 7/8 ligand S28463	Rat	Reduction in airway inflammation and T _H 1/T _H 2 cytokine levels	Prevents goblet cell hyperplasia and ASM mass increase by reducing proliferation	162

IL-4 levels are important in regulating several steps of allergic inflammation, such as IgE isotype switching, upregulation of vascular adhesion molecule 1, and T_H2 cell commitment.¹³⁶ It has been previously shown that IL-4–deficient mice exhibit reduced collagen airway deposition after chronic allergen exposure.¹³⁷ An inhaled recombinant human soluble IL-4 receptor (altrakinecept) has been developed,¹³² but there are no data on possible effects on structural cells available in asthma yet. IL-13 may be a more attractive target with regard to airway remodeling. It is detectable in bronchial mucosa and sputum¹³⁸ and has been implicated in profibrotic activity.¹³⁹ Anti-IL-13 strategies, such as IL-13–deficient mice¹³⁷ or IL-4 receptor (R) α antisense oligonucleotides,¹⁴⁰ demonstrated reduced airway collagen and reduced goblet cell metaplasia and mucus staining, respectively. Hence, IL-13 seems to be a target worth exploring in the prevention (or reversal) of airway remodeling.

IL-5 is a major regulator of eosinophilopoiesis, and treatment with the anti-IL-5 humanized monoclonal antibody (hMoAB) (mepolizumab) almost abolished circulating and sputum eosinophils¹⁴¹ while reducing tissue and

bone marrow eosinophils by about 50%.¹⁴² Notably, such treatment resulted in a significant reduction in the expression of tenascin, lumican, and procollagen III in the bronchial mucosal reticular BM and in TGF- β –positive cells compared with placebo.¹⁴³ These data suggest a role for eosinophilic release of TGF- β in patients with asthma, which can have implications for the understanding of airway remodeling.

Anti-IgE

IgE is a key mediator of the inflammatory allergic reactions such as asthma and rhinitis. Omalizumab is a humanized anti-IgE mAb that binds free circulating IgE, thereby preventing the interaction between IgE and its receptors on inflammatory cells. Clinical trials have shown that omalizumab has beneficial effects in patients with severe persistent asthma by reducing the risk of exacerbations and hospitalization and improving symptom control, lung function, and quality of life in patients.¹⁴⁴ At the bronchial level, 16 weeks of treatment with omalizumab decreased IgE, associated with a marked reduction in airway eosinophilia and in expression of the

high-affinity IgE receptor and the T_H2 cytokine IL-4.¹⁴⁵ In epithelial cells, anti-IgE can reduce TNF- α and TGF- β expression, which may point toward possible indirect influence on airway structure.¹⁴⁶

Metalloprotease inhibitors

Matrix metalloproteases are a family of enzymes consisting of 24 zinc-dependent endopeptidases capable of degrading ECM components, having a role in cell migration, angiogenesis, and tissue remodeling.^{147,148} Expression of several MMPs has been associated with asthma. Increases in MMP-1, MMP-2, MMP-3, MMP-8, and MMP-9 have been described in sputum, BAL, or tissue from patients with asthma. MMP-9 BAL levels are significantly enhanced in patients with severe asthma and in patients experiencing acute severe attacks.¹⁴⁷ Targeting MMPs could be therefore an alternative to treat asthma. A broad spectrum synthetic inhibitor, marimastat, initially used in a clinical trial with patients with cancer, was tested in patients with mild asthma.¹⁴⁹ Compared with placebo, marimastat reduced bronchial hyperresponsiveness to inhaled allergen without significant changes in sputum inflammatory cells, exhaled nitric oxide, FEV₁, asthma symptoms, or albuterol use. In patients with cancer, serious musculoskeletal side effects were observed with this drug, and trials have stopped. Targeted therapy against specific MMPs will be probably necessary when treating lung diseases.

Phosphodiesterase inhibitors

The phosphodiesterases (PDEs) control the degradation of cAMP and cGMP to their inactive 5' monophosphates, having important roles in regulating signaling responses to intracellular gradients of cAMP and cGMP. In an animal model of chronic allergic sensitization, the PDE4 inhibitor roflumilast reduced airway inflammation, subepithelial collagen, and thickening of the airway epithelium, but not the accumulation of TGF- β .⁵¹ The same PDE4 inhibitor was able to reduce CTGF and ECM deposition from ASM cells *in vitro*, whereas corticosteroids and long-acting β_2 -agonists were not.¹¹⁶ This is indicative of the potential of PDE4 inhibitors to influence airway structure, which requires further testing in human *in vivo* studies.

Other drugs

Several new promising approaches and new targets for asthma are currently under investigation,^{150,151} but in many of them, there are no human data yet, or the effects on airway structure have not been investigated. Table I shows some new therapeutic targets which, in animals, have been shown to have effects on airway structure. Some of these approaches will surely deserve human proof of concept studies.

CONCLUSION

Airway remodeling has become a key concept in today's asthma research. However, the level of

uncertainties grows with the level of excitement. It appears to be an adequate working hypothesis that the observed changes in airway structure in asthma can have untoward effects on airway function, and thus should have therapeutic implications.

Progress in this area depends on linking disciplines: on explicit integration of innovative biopharmaceutics with modern structure-function analysis.^{163,164} Where can we find this in a single laboratory? Public and private institutes should collaborate to promote such synthesis. During the next 5 years, we must develop balanced answers to the following questions:

- At which stage is airway remodeling impairing lung function?
- What are the key components of airway structure that determine fixed airway obstruction and excessive airway narrowing?
- How are these components differentially regulated?
- How can such regulatory mechanisms selectively be manipulated?
- Can this adequately prevent and reverse the detrimental components of airway remodeling in asthma?

REFERENCES

1. Global Initiative for Asthma. Global strategy for asthma management and prevention. Revised 2006. Available at: <http://www.ginasthma.org/download.asp>. Accessed March 29, 2007.
2. Vignola AM, Kips J, Bousquet J. Tissue remodeling as a feature of persistent asthma. *J Allergy Clin Immunol* 2000;105:1041-53.
3. Pascual RM, Peters SP. Airway remodeling contributes to the progressive loss of lung function in asthma: an overview. *J Allergy Clin Immunol* 2005;116:477-86.
4. Davies DE, Wicks J, Powell RM, Puddicombe SM, Holgate ST. Airway remodeling in asthma: new insights. *J Allergy Clin Immunol* 2003;111:215-25.
5. Jeffery PK. Inflammation and remodeling in the adult and child with asthma. *Pediatr Pulmonol* 2001;21:3-16.
6. James AL, Maxwell PS, Pearce-Pinto G, Elliot JG, Carroll NG. The relationship of reticular basement membrane thickness to airway wall remodeling in asthma. *Am J Respir Crit Care Med* 2002;166:1590-6.
7. Payne DNR, Rogers AV, Adelroth E, Bandi V, Guntupalli KK, Bush A, et al. Early thickening of the reticular basement membrane in children with difficult asthma. *Am J Respir Crit Care Med* 2003;167:78-82.
8. Barbato A, Turato G, Baraldo S, Bazzan E, Calabrese F, Panizzolo C, et al. Epithelial damage and angiogenesis in the airways of children with asthma. *Am J Respir Crit Care Med* 2006;174:975-81.
9. Holgate ST, Davies DE, Powell RM, Howarth PH, Haitchi HM, Holloway JW. Local genetic and environmental factors in asthma disease pathogenesis: chronicity and persistence mechanisms. *Eur Respir J* 2007;29:793-803.
10. Locke NR, Royce SG, Wainwright JS, Samuel CS, Tang ML. Comparison of airway remodeling in acute, subacute, and chronic models of allergic airways disease. *Am J Respir Cell Mol Biol* 2007;36:625-32.
11. Kuhn Ch, Homer RJ, Zhu Z, Ward N, Flavell RA, Geba GP, et al. Airway hyperresponsiveness and airway obstruction in transgenic mice: morphologic correlates in mice overexpressing interleukin (IL)-11 and IL-6 in the lung. *Am J Respir Cell Mol Biol* 2000;22:289-95.
12. Southam DS, Ellis R, Wattie J, Inman MD. Components of airway hyperresponsiveness and their associations with inflammation and remodeling in mice. *J Allergy Clin Immunol* 2007;119:848-54.
13. in 't Veen JC, Beekman AJ, Bel EH, Sterk PJ. Recurrent exacerbations in severe asthma are associated with enhanced airway closure during stable episodes. *Am J Respir Crit Care Med* 2000;161:1902-6.

14. ten Brinke A, Zwinderman AH, Sterk PJ, Rabe KF, Bel EH. "Refractory" eosinophilic airway inflammation in severe asthma: effect of parenteral corticosteroids. *Am J Respir Crit Care Med* 2004;170:601-5.
15. Carroll N, Elliot J, Morton A, James A. The structure of large and small airway in nonfatal and fatal asthma. *Am Rev Respir Dis* 1993;147:405-10.
16. Milanese M, Crimi E, Scordamaglia A, Riccio A, Pellegrino R, Canonica GW, et al. On the functional consequences of bronchial basement membrane thickening. *J Appl Physiol* 2001;91:1035-40.
17. Shiba K, Kasahara K, Nakajima H, Adachi M. Structural changes of the airway wall impair respiratory function, even in mild asthma. *Chest* 2002;122:1622-6.
18. Benayoun L, Druilhe A, Dombret MC, Aubier M, Pretolani M. Airway structural alternations selectively associated with severe asthma. *Am J Respir Crit Care Med* 2003;167:1360-8.
19. Pepe C, Foley S, Shannon J, Lemiere C, Olivenstein R, Ernst P, et al. Differences in airway remodeling between subjects with severe and moderate asthma. *J Allergy Clin Immunol* 2005;116:544-9.
20. Chu HW, Halliday JL, Martin RJ, Leung DY, Szefer SJ, Wenzel SE. Collagen deposition in large airways may not differentiate severe asthma from milder forms of the disease. *Am J Respir Crit Care Med* 1998;158:1936-44.
21. van Rensen EL, Sont JK, Evertse CE, Willems LN, Mauad T, Hiemstra PS, et al. Bronchial CD8 cell infiltrate and lung function decline in asthma. *Am J Respir Crit Care Med* 2005;172:837-41.
22. Laurent GJ. Lung collagen: more than scaffolding. *Thorax* 1986;41:418-28.
23. McParland BE, Macklem PT, Pare PD. Airway wall remodeling: friend or foe? *J Appl Physiol* 2003;95:426-34.
24. Mauad T, Silva FF, Santos MA, Grinberg L, Bernardi FDC, Martins MA, et al. Abnormal alveolar attachments with decreased elastic fiber content in distal lung in fatal asthma. *Am J Respir Crit Care Med* 2004;170:857-62.
25. Macklem PT. A theoretical analysis of the effect of airway smooth muscle load on airway narrowing. *Am J Respir Crit Care Med* 1996;153:83-9.
26. Brackel HJL, Pedersen OF, Mulder PGH, Overbeek SE, Kerrebijn KF, Bogaard JM. Central airways behave more stiffly during forced expiration in patients with asthma. *Am J Respir Crit Care Med* 2000;162:896-904.
27. Ward C, Johns DP, Bish R, Pais M, Reid DW, Ingram C, et al. Reduced airway distensibility, fixed airflow limitation, and airway wall remodeling in asthma. *Am J Respir Crit Care Med* 2001;164:1718-21.
28. Seow CY, Wang L, Pare PD. Airway narrowing and internal structural constraints. *J Appl Physiol* 2000;88:527-33.
29. Fredberg J. Bronchospasm and its biophysical basis in smooth muscle. *Respir Res* 2004;5:2.
30. Niimi A, Matsumoto H, Takemura M, Ueda T, Chin K, Mishima M. Relationship of airway wall thickness to airway sensitivity and airway reactivity in asthma. *Am J Respir Crit Care Med* 2003;168:983-8.
31. Ward C, Reid DW, Orsida BE, Feltis B, Ryan VA, Johns DP, et al. Inter-relationships between airway inflammation, reticular basement membrane thickening and bronchial hyperreactivity to methacholine in asthma: a systematic bronchoalveolar lavage and airway biopsy analysis. *Clin Exp Allergy* 2005;35:1565-71.
32. Jeffery P, Holgate S, Wenzel S, on behalf of the Endobronchial Biopsy Workshop Authors. Methods for the assessment of endobronchial biopsies in clinical research: application to studies of pathogenesis and the effects of treatment. *Am J Respir Crit Care Med* 2003;168:S1-17.
33. Shaw RJ, Djukanovic R, Tashkin DP, Millar AB, Du Bois RM, Corris PA. The role of small airways in lung disease. *Respir Med* 2002;96:67-80.
34. De Jong PA, Muller NL, Pare PD, Coxson HO. Computed tomographic imaging of the airways: relationship to structure and function. *Eur Respir J* 2005;26:140-52.
35. Kips JC, Anderson GP, Fredberg JJ, Herz U, Inman MD, Jordana M, et al. Murine models of asthma. *Eur Respir J* 2003;22:347-82.
36. Barnes PJ. Corticosteroid effects on cell signalling. *Eur Respir J* 2006;27:413-26.
37. Panettieri RA. Effects of corticosteroids on structural cells in asthma and chronic obstructive pulmonary disease. *Proc Am Thorac Soc* 2004;1:231-4.
38. Barnes PJ. New drugs for asthma. *Nat Rev Drug Discov* 2004;3:831-44.
39. Djukanovic R, Roche WR, Wilson JW, Beasley CRW, Twentyman OP, Howarth PH, et al. Mucosal inflammation in asthma. *Am Rev Respir Dis* 1990;142:434-57.
40. Ordonez C, Ferrando R, Hyde DM, Wong HH, Fahy JV. Epithelial desquamation in asthma: artefact or pathology? *Am J Respir Crit Care Med* 2000;162:2324-9.
41. Dorscheid DR, Wojcik KR, Sun S, Marroquin B, White SR. Apoptosis of airway epithelial cells induced by corticosteroids. *Am J Respir Crit Care Med* 2001;164:1939-47.
42. Wen LP, Madani K, Fahrni JA, Duncan SR, Rosen GD. Dexamethasone inhibits lung epithelial cell apoptosis induced by IFN-gamma and Fas. *Am J Physiol* 1997;273:L921-9.
43. Erjefalt JS, Erjefalt I, Sundler F, Persson CG. Effects of topical budesonide on epithelial restitution in vivo in guinea pig trachea. *Thorax* 1995;50:785-92.
44. Wadsworth SJ, Nijmeh HS, Hall IP. Glucocorticoids increase repair potential in a novel in vitro human airway epithelial wounding model. *J Clin Immunol* 2006;26:376-87.
45. Puddicombe SM, Torres-Lozano C, Richter A, Buchieri F, Lordan JL, Howarth PH, et al. Increased expression of p21(waf) cyclin-dependent kinase inhibitor in asthmatic bronchial epithelium. *Am J Respir Cell Mol Biol* 2003;28:61-8.
46. Fedorov IA, Wilson SJ, Davies DE, Holgate ST. Epithelial stress and structural remodelling in childhood asthma. *Thorax* 2005;60:389-94.
47. Lundgren R, Soderberg M, Horstedt P, Stenling R. Morphological studies of bronchial mucosal biopsies from asthmatics before and after ten years of treatment with inhaled steroids. *Eur Respir J* 1998;11:883-9.
48. Laitinen LA, Laitinen A, Haahela T. A comparative study of the effects of an inhaled corticosteroid, budesonide, and a beta2-agonist, terbutaline, on airway inflammation in newly diagnosed asthma: a randomized, double-blind, parallel-group controlled trial. *J Allergy Clin Immunol* 1992;90:32-42.
49. Rogers DF, Barnes PJ. Treatment of airway mucus hypersecretion. *Ann Med* 2006;38:116-25.
50. Leung SY, Eynott P, Nath P, Chung KF. Effects of ciclesonide and fluticasone propionate on allergen-induced airway inflammation and remodeling features. *J Allergy Clin Immunol* 2005;115:989-96.
51. Kumar RK, Herbert C, Thomas PS, Wollin L, Beume R, Yang M, et al. Inhibition of inflammation and remodeling by roflumilast and dexamethasone in murine chronic asthma. *J Pharmacol Exp Ther* 2003;307:349-55.
52. De Kluijver J, Schrupf JA, Evertse CE, Sont JK, Roughley PJ, Rabe KF, et al. Bronchial matrix and inflammation respond to inhaled steroids despite ongoing allergen exposure in asthma. *Clin Exp Allergy* 2005;35:1361-9.
53. Henderson WR Jr, Chiang GK, Tien YT, Chi EY. Reversal of allergen-induced airway remodeling by CysLT1 receptor blockade. *Am J Respir Crit Care Med* 2006;173:718-28.
54. Muz MH, Deveci F, Bulut Y, Ilhan N, Yekeler H, Turgut T. The effects of low dose leukotriene receptor antagonist therapy on airway remodeling and cysteinyl leukotriene expression in a mouse asthma model. *Exp Mol Med* 2006;38:109-18.
55. Liu YC, Khawaja AM, Rogers DF. Effects of the cysteinyl leukotriene receptor antagonists pranlukast and zafirlukast on tracheal mucus secretion in ovalbumin-sensitized guinea-pigs in vitro. *Br J Pharmacol* 1998;124:563-71.
56. Jeffery PK, Wardlaw AJ, Nelson FC, Collins JV, Kay AB. Bronchial biopsies in asthma. An ultrastructural, quantitative study and correlation with hyperreactivity. *Am Rev Respir Dis* 1989;140:1745-53.
57. Saglani S, Malmstrom K, Pelkonen AS, Malmberg LP, Lindahl H, Kajosaari M, et al. Airway remodeling and inflammation in symptomatic infants with reversible airflow obstruction. *Am J Respir Crit Care Med* 2005;171:722-7.
58. Roche WR, Beasley R, Williams JH, Holgate ST. Subepithelial fibrosis in the bronchi of asthmatics. *Lancet* 1989;1:520-4.
59. Saglani S, Molyneux C, Gong H, Rogers A, Malmstrom K, Pelkonen A, et al. Ultrastructure of the reticular basement membrane in asthmatic adults, children and infants. *Eur Respir J* 2006;28:505-12.
60. Jeffery PK, Godfrey RW, Adelroth E, Nelson F, Rogers A, Johansson SA. Effects of treatment on airway inflammation and thickening of basement membrane reticular collagen in asthma: a quantitative light and electron microscopic study. *Am Rev Respir Dis* 1992;145:890-9.

61. Boulet LP, Turcotte H, Laviolette M, Naud F, Bernier MF, Martel S, et al. Airway hyperresponsiveness, inflammation, and subepithelial collagen-deposition in recently diagnosed versus long-standing asthma: influence of inhaled corticosteroids. *Am J Respir Crit Care Med* 2000; 165:1308-13.
62. Sont JK, Willems LN, Bel EH, van Krieken JH, Vandenbroucke JP, Sterk PJ. Clinical control and histopathologic outcome of asthma when using airway hyperresponsiveness as an additional guide to long-term treatment. The AMPUL Study Group. *Am J Respir Crit Care Med* 1999;159:1043-51.
63. Ward C, Pais M, Bish R, Reid D, Feltis B, Johns D, et al. Airway inflammation, basement membrane thickening and bronchial hyperresponsiveness in asthma. *Thorax* 2002;57:309-16.
64. Hoshino M, Takahashi M, Takai Y, Sim J. Inhaled corticosteroids decrease subepithelial collagen deposition by modulation of the balance between matrix metalloproteinase-9 and tissue inhibitor of metalloproteinase-1 expression in asthma. *J Allergy Clin Immunol* 1999;104: 356-63.
65. Altraja A, Laitinen A, Virtanen I, Kampe M, Simonsson BG, Karlsson SE, et al. Expression of laminins in the airways in various types of asthmatic patients: a morphometric study. *Am J Respir Cell Mol Biol* 1996; 15:482-8.
66. Christie PE, Jonas M, Tsai CH, Chi EY, Henderson WR. Increase in laminin expression in allergic airway remodelling and decrease by dexamethasone. *Eur Respir J* 2004;24:107-15.
67. Laitinen A, Altraja A, Kampe M, Linden M, Virtanen I, Laitinen LA. Tenascin is increased in airway basement membrane of asthmatics and decreased by an inhaled steroid. *Am J Respir Crit Care Med* 1997;156:951-8.
68. Fernandes DJ, Bonacci JV, Stewart AG. Extracellular matrix, integrins, and mesenchymal cell function in the airways. *Curr Drug Targets* 2006; 7:567-77.
69. Silvestri M, Fregonese L, Sabatini F, Dasic G, Rossi GA. Fluticasone and salmeterol downregulate in vitro, fibroblast proliferation and ICAM-1 or H-CAM expression. *Eur Respir J* 2001;18:139-45.
70. Oddera S, Cagnoni F, Mangraviti S, Giron-Michel J, Popova O, Canonica GW. Effects of triamcinolone acetonide on adult human lung fibroblasts: decrease in proliferation, surface molecule expression and mediator release. *Int Arch Allergy Immunol* 2002;129:152-9.
71. Kraft M, Lewis C, Pham D, Chu HW. IL-4, IL-13, and dexamethasone augment fibroblast proliferation in asthma. *J Allergy Clin Immunol* 2001;107:602-6.
72. Fouty B, Moss T, Solodushko V, Kraft M. Dexamethasone can stimulate G1-S phase transition in human airway fibroblasts in asthma. *Eur Respir J* 2006;27:1160-7.
73. Cazes E, Giron-Michel J, Baouz S, Doucet C, Cagnoni F, Oddera S, et al. Novel anti-inflammatory effects of the inhaled corticosteroid fluticasone propionate during lung myofibroblastic differentiation. *J Immunol* 2001;167:5329-37.
74. Miller M, Cho JY, McElwain K, McElwain S, Shim JY, Manni M, et al. Corticosteroids prevent myofibroblast accumulation and airway remodeling in mice. *Am J Physiol Lung Cell Mol Physiol* 2006;290: L162-9.
75. Yano Y, Yoshida M, Hoshino S, Inoue K, Kida H, Yanagita M, et al. Anti-fibrotic effects of theophylline on lung fibroblasts. *Biochem Biophys Res Commun* 2006;341:684-90.
76. Kelly MM, Chakir J, Vethanayagam D, Boulet LP, Laviolette M, Gaudie J, et al. Montelukast treatment attenuates the increase in myofibroblasts following low-dose allergen challenge. *Chest* 2006;130: 741-53.
77. Wilson JW, Li X. The measurement of reticular basement membrane and submucosal collagen in the asthmatic airway. *Clin Exp Allergy* 1997;27:363-71.
78. Pini L, Hamid Q, Shannon J, Lemelin L, Olivenstein R, Ernst P, et al. Differences in proteoglycan deposition in the airways of moderate and severe, asthmatics. *Eur Respir J* 2007;29:71-7.
79. Huang J, Olivenstein R, Taha R, Hamid Q, Ludwig M. Enhanced proteoglycan deposition in the airway wall of atopic asthmatics. *Am J Respir Crit Care Med* 1999;160:725-9.
80. Mauad T, Xavier ACG, Saldiva PHN, Dolhnikoff M. Elastosis and fragmentation of fibers of the elastic system in fatal asthma. *Am J Respir Crit Care Med* 1999;160:968-75.
81. de Medeiros Matsushita M, da Silva LF, dos Santos MA, Fernezlian S, Schrupf JA, Roughley P, et al. Airway proteoglycans are differentially altered in fatal asthma. *J Pathol* 2005;207:102-10.
82. Vanacker NJ, Palmans E, Kips JC, Pauwels RA. Fluticasone inhibits but does not reverse allergen-induced structural airway changes. *Am J Respir Crit Care Med* 2001;163:674-9.
83. Vanacker NJ, Palmans E, Pauwels RA, Kips JC. Fluticasone inhibits the progression of allergen-induced structural airway changes. *Clin Exp Allergy* 2002;32:914-20.
84. Vanacker NJ, Palmans E, Pauwels RA, Kips JC. Dose-related effects of inhaled fluticasone on allergen-induced airway changes in rats. *Eur Respir J* 2002;20:873-9.
85. Chakir J, Shannon J, Molet S, Fukakusa M, Elias J, Laviolette M, et al. Airway remodeling-associated mediators in moderate to severe asthma: effect of steroids on TGF- β , IL-11, IL-17, and type I and type III collagen expression. *J Allergy Clin Immunol* 2003;111:1293-8.
86. Roberts CR, Burke AK. Remodelling of the extracellular matrix in asthma: proteoglycan synthesis and degradation. *Can Respir J* 1998; 5:48-50.
87. Todorova L, Gurcan E, Miller-Larsson A, Westergren-Thorsson G. Lung fibroblast proteoglycan production induced by serum is inhibited by budesonide and formoterol. *Am J Respir Cell Mol Biol* 2006;34: 92-100.
88. Bousquet J, Lacoste JY, Chanez P, Vic P, Godard P, Michel FB. Bronchial elastic fibers in normal subjects and asthmatic patients. *Am J Respir Crit Care Med* 1996;153:1648-54.
89. Carroll NG, Cooke C, James AL. Bronchial blood vessel dimensions in asthma. *Am J Respir Crit Care Med* 1997;155:689-95.
90. Charan NB, Baile EM, Pare PD. Bronchial vascular congestion and angiogenesis. *Eur Respir J* 1997;10:1173-80.
91. Hoshino M, Takahashi M, Aoike N. Expression of vascular endothelial growth factor, basic fibroblast growth factor, and angiogenin immunoreactivity in asthmatic airways and its relationship to angiogenesis. *J Allergy Clin Immunol* 2001;107:295-301.
92. Hashimoto M, Tanaka H, Abe S. Quantitative analysis of bronchial wall vascularity in the medium and small airways of patients with asthma and COPD. *Chest* 2005;127:965-72.
93. Chung KF, Rogers DF, Barnes PJ, Evans TW. The role of increased airway microvascular permeability and plasma exudation in asthma. *Eur Respir J* 1990;3:329-37.
94. Chetta A, Zanini A, Olivieri D. Therapeutic approach to vascular remodelling in asthma. *Pulm Pharmacol Ther* 2007;20:1-8.
95. Hoshino M, Takahashi M, Takai Y, Sim J, Aoike N. Inhaled corticosteroids decrease vascularity of the bronchial mucosa in patients with asthma. *Clin Exp Allergy* 2001;31:722-30.
96. Mendes ES, Pereira A, Danta I, Duncan RC, Wanner A. Comparative bronchial vasoconstrictive efficacy of inhaled glucocorticosteroids. *Eur Respir J* 2003;21:989-93.
97. Asai K, Kanazawa H, Kamoi H, Shiraishi S, Hirata K, Yoshikawa J. Increased levels of vascular endothelial growth factor in induced sputum in asthmatic patients. *Clin Exp Allergy* 2003;33:595-9.
98. Feltis BN, Wignarajah D, Reid DW, Ward C, Harding R, Walters EH. Effects of inhaled fluticasone on angiogenesis and vascular endothelial growth factor in asthma. *Thorax* 2007;62:314-9.
99. Laitinen LA, Laitinen A, Widdicombe J. Effects of inflammatory and other mediators on airway vascular beds. *Am Rev Respir Dis* 1987; 135:S67-70.
100. Orsida BE, Ward C, Li X, Bish R, Wilson JW, Thien F, et al. Effect of a long-acting beta2-agonist over three months on airway wall vascular remodeling in asthma. *Am J Respir Crit Care Med* 2001;164:117-21.
101. Lee KS, Kim SR, Park HS, Jin GY, Lee YC. Cysteinyl leukotriene receptor antagonist regulates vascular permeability by reducing vascular endothelial growth factor expression. *J Allergy Clin Immunol* 2004; 114:1093-9.
102. Mendes ES, Campos MA, Hurtado A, Wanner A. Effect of montelukast and fluticasone propionate on airway mucosal blood flow in asthma. *Am J Respir Crit Care Med* 2004;169:1131-4.
103. James A. Remodelling of airway smooth muscle in asthma: what sort do you have? *Clin Exp Allergy* 2005;35:703-7.
104. Lambert RK, Wiggs BR, Kuwano K, Hogg JC, Pare PD. Functional significance of increased airway smooth muscle in asthma and COPD. *J Appl Physiol* 1993;74:2771-81.

105. Gunst SJ, Fredberg JJ. The first three minutes: smooth muscle contraction, cytoskeletal events, and soft glasses. *J Appl Physiol* 2003;95:413-25.
106. Hirst SJ, Walker TR, Chilvers ER. Phenotype diversity and molecular mechanisms of airway smooth muscle proliferatin in asthma. *Eur Respir J* 2000;16:159-77.
107. Halayko AJ, Tran T, Ji SY, Yamasaki A, Gosens R. Airway smooth muscle phenotype and function: interactions with current asthma therapies. *Curr Drug Targets* 2006;7:525-40.
108. Fernandes DJ, Mitchell RW, Lakser W, Dowell M, Stewart AG, Solway J. Do inflammatory mediators influence the contribution of airway smooth muscle contraction to airway hyperresponsiveness in asthma? *J Appl Physiol* 2003;95:844-53.
109. Fernandes D, Guida E, Koutsoubos V, Harris T, Vadeloo P, Wilson JW, et al. Glucocorticoids inhibit proliferation, cyclin D1 expression, and retinoblastoma protein phosphorylation, but not activity of the extracellular-regulated kinases in human cultured airway smooth muscle. *Am J Respir Cell Mol Biol* 1999;21:77-88.
110. Vlahos R, Lee KS, Guida E, Fernandes DJ, Wilson JW, Stewart AG. Differential inhibition of thrombin- and EGF-stimulated human cultured airway smooth muscle proliferation by glucocorticoids. *Pulm Pharmacol Ther* 2003;16:171-80.
111. Roth M, Johnson PRA, Borger P, Bihl MP, Rudiger JJ, King GC, et al. Dysfunctional interaction of C/EBP α and the glucocorticoid receptor in asthmatic bronchial smooth muscle cells. *N Engl J Med* 2004;351:560-74.
112. Goldsmith AM, Hershenson MB, Wolbert MP, Bentley JK. Regulation of airway smooth muscle α -actin expression by glucocorticoids. *Am J Physiol Lung Cell Mol Physiol* 2007;292:L99-106.
113. Black JL, Burgess JK, Johnson PR. Airway smooth muscle: its relationship to the extracellular matrix. *Respir Physiol Neurobiol* 2003;137:339-46.
114. Bonacci JV, Stewart AG. Regulation of human airway mesenchymal cell proliferation by glucocorticoids and β 2-adrenoceptor agonists. *Pulm Pharmacol Ther* 2006;19:32-8.
115. Johnson PR, Black JL, Carlin S, Ge Q, Underwood PA. The production of extracellular matrix proteins by human passively sensitized airway smooth-muscle cells in culture: the effect of beclomethasone. *Am J Respir Crit Care Med* 2000;162:2145-51.
116. Burgess JK, Oliver BG, Poniris MH, Ge Q, Boustany S, Cox N, et al. A phosphodiesterase 4 inhibitor inhibits matrix protein deposition in airways in vitro. *J Allergy Clin Immunol* 2006;118:649-57.
117. Lee SY, Kim JS, Lee JM, Kwon SS, Kim KH, Moon HS, et al. Inhaled corticosteroid prevents the thickening of airway smooth muscle in murine model of chronic asthma. *Pulm Pharmacol Ther* 2006 Oct 20; [Epub ahead of print].
118. Cho JY, Miller M, McElwain K, McElwain S, Broide DH. Combination of corticosteroid therapy and allergen avoidance reverses allergen-induced airway remodeling in mice. *J Allergy Clin Immunol* 2005;116:1116-22.
119. Ammit AJ, Panettieri RA Jr. Airway smooth muscle cell hyperplasia: a therapeutic target in airway remodeling in asthma? *Prog Cell Cycle Res* 2003;5:49-57.
120. Gosens R, Bos ST, Zaagsma J, Meurs H. Protective effects of tiotropium bromide in the progression of airway smooth muscle remodeling. *Am J Respir Crit Care Med* 2005;171:1096-102.
121. An SS, Bai TR, Bates JH, Black JL, Brown RH, Brusasco V, et al. Airway smooth muscle dynamics: a final common pathway of airway obstruction in asthma. *Eur Respir J* 2007;29:834-60.
122. Mitzner W. Airway smooth muscle: the appendix of the lung. *Am J Respir Crit Care Med* 2004;169:787-90.
123. Cox PG, Miller J, Mitzner W, Leff AR. Radiofrequency ablation of airway smooth muscle for sustained treatment of asthma: preliminary investigations. *Eur Respir J* 2004;24:659-63.
124. Miller JD, Cox G, Vincic L, Lombard CM, Loomas BE, Danek CJ. A prospective feasibility study of bronchial thermoplasty in the human airway. *Chest* 2005;127:1999-2006.
125. Brown RH, Wizeman W, Danek C, Mitzner W. Effect of bronchial thermoplasty on airway distensibility. *Eur Respir J* 2005;26:277-82.
126. Cox G, Miller JD, McWilliams A, FitzGerald JM, Lam S. Bronchial thermoplasty for asthma. *Am J Respir Crit Care Med* 2006;173:965-9.
127. Cox G, Thomson NC, Rubin AS, Niven RM, Corris PA, Siersted HC, et al. Asthma control during the year after bronchial thermoplasty. *N Engl J Med* 2007;356:1327-37.
128. Howarth PH, Babu KS, Arshad HS, Lau L, Buckley M, McConnell W, et al. Tumour necrosis factor (TNF- α) as a novel therapeutic target in symptomatic corticosteroid dependent asthma. *Thorax* 2005;60:1012-8.
129. Berry MA, Hargadon B, Shelley M, Parker D, Shaw DE, Green RH, et al. Evidence of role of tumor necrosis factor α in refractory asthma. *N Engl J Med* 2006;354:697-708.
130. Erin EM, Leaker BR, Nicholson GC, Tan AJ, Green LM, Neighbour H, et al. The effects of a monoclonal antibody directed against tumor necrosis factor- α in asthma. *Am J Respir Crit Care Med* 2006;174:753-62.
131. Trifilieff A, Walker C, Keller T, Kottirsch G, Neumann U. Pharmacological profile of PKF242-484 and PKF241-466, novel dual inhibitors of TNF- α converting enzyme and matrix metalloproteinases, in models of airway inflammation. *Br J Pharmacol* 2002;135:1655-64.
132. Hendeles L, Asmus M, Chesrown S. Evaluation of cytokine modulators for asthma. *Paediatr Respir Rev* 2004;5(suppl A):S107-12.
133. Oda N, Minoguchi K, Yokoe T, Hashimoto T, Wada K, Miyamoto M, et al. Effect of suplatat tosilate (IPD-1151T) on cytokine production by allergen-specific human Th1 and Th2 cell lines. *Life Sci* 1999;65:763-70.
134. Tamaoki J, Kondo M, Sakai N, Aoshiba K, Tagaya E, Nakata J, et al. Effect of suplatat tosilate, a Th2 cytokine inhibitor, on steroid-dependent asthma: a double-blind randomized study. *Lancet* 2000;356:273-8.
135. Hoshino M, Fujita Y, Saji J, Inoue T, Nakagawa T, Miyazawa T. Effect of suplatat tosilate on goblet cell metaplasia in patients with asthma. *Allergy* 2005;60:1394-400.
136. O'Byrne PM. Cytokines or their antagonists for the treatment of asthma. *Chest* 2006;130:244-50.
137. Leigh R, Ellis R, Wattie JN, Hirota JA, Matthaei KI, Foster PS, et al. Type 2 cytokines in the pathogenesis of sustained airway dysfunction and airway remodeling in mice. *Am J Respir Crit Care Med* 2004;169:860-7.
138. Berry MA, Parker D, Neale N, Woodman L, Morgan A, Monk P, et al. Sputum and bronchial submucosal IL-13 expression in asthma and eosinophilic bronchitis. *J Allergy Clin Immunol* 2004;114:1106-9.
139. Kaviratne M, Hesse M, Leusing M, Cheever AW, Davies SJ, McKerrrow JH, et al. IL-13 activates a mechanism of tissue fibrosis that is completely TGF- β independent. *J Immunol* 2004;173:4020-9.
140. Karras JG, Crosby JR, Guha M, Tung D, Miller DA, Gaarde WA, et al. Anti-inflammatory activity of inhaled IL-4 receptor- α antisense oligonucleotide in mice. *Am J Respir Cell Mol Biol* 2007;36:276-85.
141. Leckie MJ, ten Brinke A, Khan J, Diamant Z, O'Connor BJ, Walls CM, et al. Effects of an interleukin-5 blocking monoclonal antibody on eosinophils, airway hyperresponsiveness, and the late asthmatic response. *Lancet* 2000;356:2144-8.
142. Flood-Page PT, Menzies-Gow AN, Kay BA, Robinson DS. Eosinophil's role remains uncertain as anti-interleukin-5 only partially depletes numbers in asthmatic airways. *Am J Respir Crit Care Med* 2003;167:199-204.
143. Flood-Page P, Menzies-Gow A, Phipps S, Ying S, Wangoo A, Ludwig MS, et al. Anti-IL-5 treatment reduces deposition of ECM proteins in the bronchial subepithelial basement membrane of mild atopic asthmatics. *J Clin Invest* 2003;112:1029-36.
144. Strunk RC, Bloomberg GR. Omalizumab for asthma. *N Engl J Med* 2006;354:2689-95.
145. Djukanovic R, Wilson SJ, Kraft M, Jarjour NN, Steel M, Chung KF, et al. Effect of treatment with anti-immunoglobulin E antibody omalizumab on airway inflammation in allergic asthma. *Am J Respir Crit Care Med* 2004;170:583-93.
146. Huang YC, Leyko B, Frieri M. Effects of omalizumab and budesonide on markers of inflammation in bronchial epithelial cells. *Ann Allergy Asthma Immunol* 2005;95:443-51.
147. Demedts IK, Brusselle GG, Bracke KR, Vermaelen KY, Pauwels RA. Matrix metalloproteinases in asthma and COPD. *Curr Opin Pharmacol* 2005;5:257-63.
148. Greenlee KJ, Werb Z, Kheradmand F. Matrix metalloproteinases in lung: multiple, multifarious, and multifaceted. *Physiol Rev* 2007;87:69-98.

149. Bruce C, Thomas PS. The effect of marimastat, a metalloprotease inhibitor, on allergen-induced asthmatic hyper-reactivity. *Toxicol Appl Pharmacol* 2005;205:126-32.
150. Belvisi MG, Hele DJ, Birrell MA. New advances and potential therapies for the treatment of asthma. *BioDrugs* 2004;18:211-23.
151. Das AM, Griswold DE, Torphy TJ, Li L. Biopharmaceutical therapeutics for asthma remodeling. *Curr Pharm Des* 2006;12:3233-40.
152. Wegmann M, Goggel R, Sel S, Sel S, Erb KJ, Kalkbrenner F, et al. Effects of a low-molecular-weight CCR-3 antagonist on chronic experimental asthma. *Am J Respir Cell Mol Biol* 2007;36:61-7.
153. Yang G, Volk A, Petley T, Emmell E, Giles-Komar J, Shang X, et al. Anti-IL-13 monoclonal antibody inhibits airway hyperresponsiveness, inflammation and airway remodeling. *Cytokine* 2004;28:224-32.
154. McMillan SJ, Xanthou G, Lloyd CM. Manipulation of allergen-induced airway remodeling by treatment with anti-TGF-beta antibody: effect on the Smad signaling pathway. *J Immunol* 2005;174:5774-80.
155. Leung SY, Niimi A, Noble A, Oates T, Williams AS, Medicherla S, et al. Effect of transforming growth factor-beta receptor I kinase inhibitor 2,4-disubstituted pteridine (SD-208) in chronic allergic airway inflammation and remodeling. *J Pharmacol Exp Ther* 2006;319:586-94.
156. Oh SW, Pae CI, Lee DK, Jones F, Chiang GK, Kim HO, et al. Tryptase inhibition blocks airway inflammation in a mouse asthma model. *J Immunol* 2002;168:1992-2000.
157. Cho JY, Miller M, Baek KJ, Han JW, Nayar J, Rodriguez M, et al. Immunostimulatory DNA inhibits transforming growth factor-beta expression and airway remodeling. *Am J Respir Cell Mol Biol* 2004;30:651-61.
158. Youn CJ, Miller M, Baek KJ, Han JW, Nayar J, Lee SY, et al. Immunostimulatory DNA reverses established allergen-induced airway remodeling. *J Immunol* 2004;173:7556-64.
159. Cho JY, Miller M, McElwain K, McElwain S, Shim JY, Raz E, et al. Remodeling associated expression of matrix metalloproteinase 9 but not tissue inhibitor of metalloproteinase 1 in airway epithelium: modulation by immunostimulatory DNA. *Am J Respir Cell Mol Biol* 2004;30:651-61.
160. Lee SY, Cho JY, Miller M, McElwain K, McElwain S, Sriramaraio P, et al. Immunostimulatory DNA inhibits allergen-induced peribronchial angiogenesis in mice. *J Allergy Clin Immunol* 2006;117:597-603.
161. Fanucchi MV, Schelegle ES, Baker GL, Evans MJ, McDonald RJ, Gershwin LJ, et al. Immunostimulatory oligonucleotides attenuate airways remodeling in allergic monkeys. *Am J Respir Crit Care Med* 2004;170:1153-7.
162. Camateros P, Tamaoka M, Hassan M, Marino R, Moisan J, Marion D, et al. Chronic asthma-induced airway remodeling is prevented by Toll-like receptor-7/8 ligand S28463. *Am J Respir Crit Care Med* 2007;175:1241-9.
163. Bergeron C, Tullic MK, Hamid Q. Tools used to measure airway remodelling in research. *Eur Respir J* 2007;29:596-604.
164. Hasegawa M, Nasuhara Y, Onodera Y, Makita H, Nagai K, Fuke S, et al. Airflow limitation and airway dimensions in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2006;173:1309-15.