

IFN-stimulated Gene Expression, Type 2 Inflammation, and Endoplasmic Reticulum Stress in Asthma

Nirav R. Bhakta^{1*}, Stephanie A. Christenson^{1*}, Srilaxmi Nerella¹, Owen D. Solberg¹, Christine P. Nguyen¹, David F. Choy², Kyle L. Jung¹, Suresh Garudadri¹, Luke R. Bonser^{3,4}, Joshua L. Pollack⁴, Lorna T. Zlock⁵, David J. Erle^{3,4}, Charles Langelier⁶, Joseph L. Derisi⁷, Joseph R. Arron², John V. Fahy^{1,3}, and Prescott G. Woodruff^{1,3}

¹Division of Pulmonary, Critical Care, Sleep and Allergy, Department of Medicine, ³Cardiovascular Research Institute, ⁴Lung Biology Center, Department of Medicine, ⁵Department of Pathology, and ⁶Division of Infectious Diseases, Department of Medicine, University of California, San Francisco, San Francisco, California; ²Genentech, Inc., South San Francisco, California; and ⁷Department of Biochemistry and Biophysics, University of California at San Francisco and Howard Hughes Medical Institute, San Francisco, California

ORCID ID: 0000-0002-2135-7012 (N.R.B.).

Abstract

Rationale: Quantification of type 2 inflammation provided a molecular basis for heterogeneity in asthma. Non-type 2 pathways that contribute to asthma pathogenesis are not well understood.

Objectives: To identify dysregulated pathways beyond type 2 inflammation.

Methods: We applied RNA sequencing to airway epithelial brushings obtained from subjects with stable mild asthma not on corticosteroids ($n = 19$) and healthy control subjects ($n = 16$). Sequencing reads were mapped to human and viral genomes. In the same cohort, and in a separate group with severe asthma ($n = 301$), we profiled blood gene expression with microarrays.

Measurements and Main Results: In airway brushings from mild asthma on inhaled corticosteroids, RNA sequencing yielded 1,379 differentially expressed genes (false discovery rate < 0.01). Pathway analysis revealed increased expression of type 2 markers,

IFN-stimulated genes (ISGs), and endoplasmic reticulum (ER) stress-related genes. Airway epithelial ISG expression was not associated with type 2 inflammation in asthma or with viral transcripts but was associated with reduced lung function by FEV₁ ($\rho = -0.72$; $P = 0.0004$). ER stress was confirmed by an increase in *XBPI* (X-box binding protein 1) splicing in mild asthma and was associated with both type 2 inflammation and ISG expression. ISGs were also the most activated genes in blood cells in asthma and were correlated with airway ISG expression ($\rho = 0.55$; $P = 0.030$). High blood ISG expression in severe asthma was similarly unrelated to type 2 inflammation.

Conclusions: ISG activation is prominent in asthma, independent of viral transcripts, orthogonal to type 2 inflammation, and associated with distinct clinical features. ER stress is associated with both type 2 inflammation and ISG expression.

Keywords: asthma; type 2 inflammation; IFNs; endoplasmic reticulum stress; blood gene expression

(Received in original form June 1, 2017; accepted in final form October 20, 2017)

*These authors contributed equally to this work.

Supported by grants from the NIH: K23HL116657 (N.R.B.), K23HL123778 (S.A.C.), K12 HL11999 (S.A.C. and D.J.E.), P01HL107202 (J.V.F.), AI077439 (D.J.E. and P.G.W.), and HL137013, AI077439, and HL107202 (P.G.W.); from the Sandler Asthma Basic Research Center; and from Genentech.

Author Contributions: N.R.B., S.A.C., and P.G.W. conceived the hypothesis, interpreted data, and drafted the manuscript. N.R.B. and S.A.C. designed and performed analysis of epithelial brushing and blood gene expression and clinical data. D.J.E. gave guidance on gene expression analyses and revised the manuscript critically for intellectual content. S.N. and O.D.S. provided computational support. C.P.N. coordinated the bronchoscopy study and participated in collection of samples and clinical data. D.F.C. performed analysis of blood microarray data and exacerbations in the severe asthma study. K.L.J. performed qPCR and assisted in analysis. S.G. assessed RNA quality and quantity and contributed to library generation. L.R.B., L.T.Z., J.L.P., and D.J.E. designed, performed, and analyzed data from the epithelial cultures. C.L. and J.L.D. provided S.A.C. with assistance in the analysis of nonhuman reads. J.R.A. designed the study in mild asthma. J.V.F. and P.G.W. designed the study in mild asthma and with N.R.B. participated in the collection of samples and clinical data. All authors read and approved the final manuscript.

Correspondence and requests for reprints should be addressed to Prescott G. Woodruff, M.D., M.P.H., Division of Pulmonary and Critical Care Medicine, Department of Medicine and CVRI, University of California, San Francisco, Box 0130, 513 Parnassus Ave, San Francisco, CA 94143. E-mail: prescott.woodruff@ucsf.edu.

This article has an online supplement, which is accessible from this issue's table of contents at www.atsjournals.org.

Am J Respir Crit Care Med Vol 197, Iss 3, pp 313–324, Feb 1, 2018

Copyright © 2018 by the American Thoracic Society

Originally Published in Press as DOI: 10.1164/rccm.201706-1070OC on October 24, 2017

Internet address: www.atsjournals.org

At a Glance Commentary

Scientific Knowledge on the

Subject: Although a subgroup of patients with asthma benefits from blockade of the type 2 cytokines IL-4, IL-5, and IL-13, there is an unmet need to identify non-type 2 mechanisms in asthma. Dysregulation in non-type 2 pathways has been demonstrated; however, the relationship of these pathways to type 2 inflammation and lung function impairment, and the impact of their presence on treatment response, have not been reported.

What This Study Adds to the

Field: In this study, IFN-stimulated gene (ISG) expression was increased in mild asthma compared with health as assessed by RNA sequencing of airway epithelial brushings. ISG expression was orthogonal to type 2 inflammation, did not require the presence of virus, and was associated with reduced FEV₁ and enhanced endoplasmic reticulum stress. ISG expression was also increased in whole blood, where it was similarly orthogonal to type 2 inflammation in severe asthma. Thus, ISG expression may be a viable target for future therapy. Both type 2 inflammation and ISG expression are independently associated with endoplasmic reticulum stress, which therefore represents an additional potential therapeutic target.

Allergic, type 2 cytokine-driven inflammation is associated with airway obstruction, hyperresponsiveness, remodeling, and response to inhaled corticosteroids (ICS) in asthma (1, 2). Therapeutic targeting of type 2 inflammation has been successful in some but not all patients (3). The subset of patients who respond poorly to therapies targeting type 2 inflammation may have non-type 2 inflammatory mechanisms driving their disease. Furthermore, these non-type 2 pathways likely coexist with type 2 inflammation in some individuals and contribute to severe asthma (3).

The airway epithelium is central to the initiation and maintenance of allergic immune responses, and contributes to asthma pathogenesis, including mucus

hypersecretion and airflow obstruction (4). Microarray gene expression profiling of human airway epithelial cells was crucial in establishing the type 2 endotype (1). New technologies incorporating massively parallel RNA sequencing (RNA-seq) of airway samples overcome the sensitivity, specificity, and signal saturation limitations of microarrays, and can offer new insight into airway disease mechanisms (5, 6). In this study, we hypothesized that airway epithelial RNA-seq in asthma would identify non-type 2 pathways of inflammation and epithelial cell dysfunction that are associated with a more severe asthma phenotype. Some of the results of these studies have been previously reported in the form of abstracts (7, 8).

Methods

Additional details are reported in the METHODS section of the online supplement.

Subjects and Procedures

Mild asthma study. We recruited 22 healthy control subjects and 32 subjects with mild asthma not on ICS via community-based advertising (NCT00595153). Baseline characterization included medical history and examination, spirometry, measurement of provocative concentration causing a 20% fall in FEV₁ (PC₂₀) and fractional exhaled nitric oxide, complete blood count and differential, serum total IgE, and collection of PAXgene blood RNA tubes (Qiagen). A total of 1–2 weeks following the baseline visit 20 healthy subjects and 25 subjects with mild asthma underwent spirometry, bronchoscopy, and blood collection; reasons for dropout of subjects are described in the METHODS section of the online supplement. Subjects with asthma were thereafter treated with inhaled budesonide, 180 µg twice daily, for 8 weeks and returned for spirometry 4 weeks after ICS was started, and for repeat bronchoscopy and spirometry 8 weeks after ICS was started.

Severe asthma study. To validate findings in a severe asthma population, we used PAXgene whole-blood RNA samples from a previously reported prospective, 48-week double-blind randomized placebo-controlled trial of omalizumab (anti-IgE) (NCT00314574) (9). Subjects had poorly controlled asthma despite high-dose ICS

and long-acting β -agonists. A total of 427 subjects received omalizumab and 423 placebo, with PAXgene tubes collected in a subgroup (Figure 1).

Gene Expression Measurements

For RNA-seq of airway epithelial brushings, library preparation and multiplexing was done with the Illumina TruSeq Stranded Total RNA with Ribo-zero Human/Mouse/Rat Kit (Illumina Inc.) per manufacturer's protocol. We performed 100 base pair paired-end sequencing on multiplexed samples via an Illumina HiSeq 2500 at the University of California, San Francisco, Genomics Core. PAXgene RNA samples were run on Affymetrix HGU133A Plus 2.0 arrays as previously described (10).

Differential Expression

RNA-Seq. We performed read count normalization and differential expression analyses using the DESeq2 package (11), and corrected for multiple comparison using the false discovery rate (FDR) (12). Genes with an FDR less than 0.01 were called significant. We used regularized-logarithm transformed counts from DESeq2's rlog function for unsupervised clustering, heatmaps, and multigene metric generation.

Microarray. We performed moderated *t* tests with the limma package (13, 14) in R. Genes with an FDR less than 0.01 were called significant except in the case of whole-blood samples, which had weaker differential expression and for which an unadjusted *P* less than 0.05 was used before pathway analysis.

Comparison of Multigene Metrics with Clinical Variables

We performed unadjusted analyses using Spearman rank correlation and Welch's *t* test, as appropriate. In adjusted analyses, we developed linear models and sought interactions between multigene metrics in association with lung function. We computed annualized exacerbation rates using zero-intercept generalized linear models, and used a model with an intercept to compute exacerbation rate reduction and perform all statistical tests of significance for exacerbations.

Metatranscriptome Preprocessing and Differential Expression

We used a recently described custom bioinformatics pipeline to detect viral

transcripts (15). Briefly, sequencing reads underwent quality and complexity filtering (16) followed by subtractive alignment to the human genome (ENSEMBL GRCh38) using STAR (17). Remaining reads underwent additional subtractive alignment against phage and nonhuman tropic viruses using Bowtie2 (18). Identical reads were compressed (19) and remaining sequences were aligned to the National Center for Biotechnology Information nucleotide database using GSNAPL to identify viral transcripts (20).

Statistics and Plots

All analyses were done in the R statistical environment. Correlations were reported either as Spearman ρ or Pearson r as indicated. Heatmaps were created using the *gplots* package. All other plots were created with the *ggplot2* package (21). P less than 0.05 was taken as statistically significant without correction for multiple testing unless otherwise indicated. Hierarchical clustering was performed with Euclidean distance and complete linkage.

Study Approval

All study sites obtained institutional review board approval and written informed consent from study participants before enrollment.

Results

Differential Airway Epithelial Gene Expression by RNA-Seq

RNA-seq data from airway epithelial brushings passed quality control in 16 healthy control subjects and 19 subjects with mild asthma not on ICS (Figure 1). Subject demographics are described in Table 1. A total of 1,379 transcripts were differentially expressed between asthma and health ($FDR < 0.01$) (see Table E1 in the online supplement). The most highly differentially expressed genes included those known to be induced by type 2 cytokine exposure, such as *POSTN* (periostin), *SERPINE2* (serpin family B member 2), and *CLCA1* (chloride channel accessory 1) (10).

We used ingenuity pathway analysis (IPA) to predict upstream regulators of overrepresented genes (Table 2). As expected, regulators of type 2 inflammation including IL-5 and IL-13 scored well. In addition, type I (IFN- α , β) and III (IFN- λ) IFN and their downstream mediators (IRF




Figure 1. Cohorts and samples contributing to the results in this publication. (A) Sample numbers from the mild asthma cohort from University of California, San Francisco (UCSF) before treatment with inhaled corticosteroids. Dropout from low RNA quantity was caused by inadequate RNA remaining for sequencing given that microarray and qPCR studies had used RNA from this study population previously. After sequencing, four subjects with mild asthma and three healthy control samples were removed from further analysis because of high mitochondrial RNA and ribosomal RNA abundance leading to insufficient depth of sequencing of protein coding reads. (B) Sample numbers from the randomized placebo-controlled trial of omalizumab in severe asthma. PAXgene = whole blood RNA collection tube; RNA-seq = RNA sequencing.

[IFN regulatory factor]-1, IRF7, IFNAR2 [IFN α and β receptor subunit 2], STAT1 [signal transducer and activator of transcription 1]) were among the pathways most significantly associated with asthma (Table 2). Unlike the IFN-stimulated genes (ISGs), IFNs themselves and most other upstream regulators involved in IFN signaling as identified by IPA were not differentially expressed in brushings from subjects with asthma. For example, STAT1, an upstream regulator previously reported to be dysregulated in asthma (22), and secretory leukocyte protease inhibitor, previously reported to be repressed by and inversely related to IFN- γ in severe asthma (23), were not significantly altered in our asthma group (see Table E2).

Comparison of RNA-Seq and Microarrays

The presence of strong signals for non-type 2 inflammation in airway brushings analyzed by RNA-seq is a new finding in

comparison with our previous microarray performed with similar subjects (1, 10). Whereas our RNA-seq study found 1,379 differentially expressed genes at an FDR less than 0.01, the earlier microarray study reported only 22 genes as differentially expressed at P less than 0.05 using a Bonferroni correction, and this increased to only 36 genes on reanalysis with a FDR less than 0.01 cutoff (10). However, although the prior microarray study had fewer statistically significant genes, the overall pattern of expression differences between groups was quite similar between the two studies as assessed using the Camera competitive gene set test (24), with significant overlap of both upregulated genes ($n = 915$; $P = 3.46 \times 10^{-28}$) and downregulated genes ($n = 284$; $P = 4.12 \times 10^{-15}$) in subjects with asthma compared with healthy control subjects (see Figure E1A for enrichment plots). This current paper presents RNA-seq generated from samples collected in one stand-alone

Table 1. Baseline Subject Characteristics

	Healthy Control Subjects	Subjects with Asthma	P Value
Sample size of RNA-seq data	16	19	
Age, yr	34 ± 10	34 ± 13	0.94
BMI, kg/m ²	26 ± 4	29 ± 6	0.14
Female sex, %	38	58	0.39
Ethnicity*			
White	9	9	
Black or African American	1	1	
Hispanic	3	4	
Asian/Pacific Islander	2	5	
Native American	1	0	
FEV ₁ , % predicted	98 ± 13	84 ± 14	0.0042
Δ FEV ₁ with albuterol, % of baseline	5.2 ± 0.45	13.8 ± 10.2	0.0027
FEV ₁ /FVC	0.78 ± 0.061	0.74 ± 0.077	0.10
Methacholine PC ₂₀ , mg/ml	>10 (n = 15)	0.57 (0.34–1.55)	<0.0001
IgE, IU/ml	17 (11–29) (n = 15)	247 (129–361)	0.00040
Blood eosinophils, ×10 ⁹ /L	0.11 (0.06–0.19)	0.36 (0.13–0.50)	0.0011
F _{ENO} , ppb	16 (12–23)	66 (20–85) (n = 14)	0.010
Epithelial brushings			
Nonsquamous epithelial, %	96 (95–98)	98 (97–99)	0.051
Neutrophil	0.2 (0–0.4)	0 (0–0.2)	0.034
Lymphocyte	0.10 (0–0.45)	0 (0–0)	0.055
Eosinophil	0 (0–0)	0 (0–0.10)	0.29
Macrophage	3 (1.3–4.7)	1.4 (0.9–2.7)	0.066

Definition of abbreviations: BMI = body mass index; F_{ENO} = fractional exhaled nitric oxide; PC₂₀ = provocative concentration causing a 20% fall in FEV₁; RNA-seq = RNA sequencing. Values are reported as mean ± SD or median (25–75th percentile). All data are from the baseline visit except spirometry, which are reported from visit 2 (immediately before initiation of RNA sequencing). P values are based on *t* test (where means reported), Wilcoxon rank sum (where medians reported), or chi-square test for proportions.

*In cases of multiple ethnicities, the lowest-numbered ethnicity from this list is reported: 1) Hispanic, 2) black or African American, 3) Pacific Islander, 4) Native American, 5) Asian, and 6) white.

clinical study, whereas our prior report was derived from samples from four different banked clinical studies that used comparable sample collection methods. This design may have avoided several clinical and technical sources of variation and improved our statistical power to find differences.

To directly assess how the use of RNA-seq versus microarrays affected the ability to detect gene expression changes and the ISG signature, we compared our RNA-seq data with microarray data generated using the same RNA samples. These microarray data represent a subset of the data previously reported in our comparison of subjects with asthma and chronic obstructive pulmonary disease (25). At an FDR less than 0.01, a total of 1,734 genes were differentially expressed by microarray between subjects with asthma and healthy control subjects. Comparison of RNA-seq and microarray data from the same samples using the Camera competitive gene set test showed significant overlap of both upregulated

genes ($P = 1.11 \times 10^{-59}$) and downregulated genes ($P = 3.50 \times 10^{-42}$) (see Figure E1B). Across ISGs, microarray and sequencing showed concordant differences between asthma and health, but the magnitude of the fold differences were greater in the sequencing results (see Figure E2A). The finding that some ISGs were significantly increased in asthma by RNA-seq but not by microarrays could not be explained consistently by low microarray signal intensity (see Figure E2B).

Development of a Metric of Airway ISG Expression

We first focused on ISGs. To generate a gene expression signature specific to airway epithelial cells, we cultured human bronchial epithelial cells at an air-liquid interface and stimulated them with IFN- α (10 ng/ml for 24 h). We chose IFN- α given its significance in the IPA list and the largely overlapping gene expression responses to type I and type III IFNs (26). RNA-seq of IFN- α -stimulated epithelial

cells yielded 1,031 IFN- α -induced genes (FDR < 0.01; top 50 in Figure 2A). We restricted this top 50 list to those differentially expressed in asthmatic airway brushings (FDR < 0.01), yielding nine

Table 2. IPA Upstream Regulator Analysis of Epithelial Brushing RNA-Seq-based Gene Expression

Upstream Regulator	Activation z-Score	P Value of Overlap
<i>IL5*</i>	4.93	2.71×10^{-4}
<i>EIF4E[†]</i>	4.80	4.56×10^{-6}
Interferon- α group [‡]	4.79	1.23×10^{-3}
<i>CD38</i>	4.64	1.28×10^{-4}
<i>IFNA2[‡]</i>	4.41	9.97×10^{-3}
<i>IL13*</i>	4.31	1.40×10^{-2}
<i>XBP1[†]</i>	4.22	1.26×10^{-7}
<i>MYC</i>	4.16	7.30×10^{-3}
<i>STAT1[‡]</i>	4.15	3.12×10^{-2}
<i>IRF7[‡]</i>	4.14	7.77×10^{-3}
<i>IFNB1[‡]</i>	3.79	4.17×10^{-2}
<i>AGT</i>	3.74	2.44×10^{-2}
<i>IFNL1[†]</i>	3.45	1.24×10^{-3}
<i>TP53</i>	3.39	4.61×10^{-3}
<i>IRF1[‡]</i>	3.37	6.91×10^{-4}
<i>PRL</i>	2.89	1.21×10^{-4}
<i>VEGFA</i>	2.83	1.10×10^{-2}
<i>EIF2AK2[‡]</i>	2.71	4.97×10^{-3}
<i>HRAS</i>	2.71	1.03×10^{-2}
<i>PAF1</i>	2.65	1.95×10^{-2}
<i>IFNA1/IFNA13[‡]</i>	2.60	1.27×10^{-3}
<i>Irfn group[‡]</i>	2.49	4.59×10^{-2}
<i>IFNAR2[‡]</i>	2.45	1.37×10^{-3}
<i>MYCN</i>	2.28	5.21×10^{-4}
<i>THBS4</i>	2.20	1.93×10^{-2}
<i>APP</i>	2.02	4.28×10^{-3}

Definition of abbreviations: IPA = ingenuity pathway analysis; RNA-seq = RNA sequencing. Regulators are filtered by 1) genes, proteins, or RNA; 2) activated predicted state; 3) activation z-score ≥ 2 ; and 4) $P < 0.05$. The list is sorted by activation z-score. P value of overlap is the probability that the observed overlap between the list of genes differentially expressed between asthma and health, and the set of genes assigned by IPA as downstream of the listed upstream regulator, would occur by chance (Fisher's exact test). The strength of activity of the regulator in disease, as determined by the net observed direction of changes in the overlapping downstream genes, is represented by an activation score (a more positive/negative value indicates that a larger proportion of genes changed in a direction consistent with activation/inhibition of the regulator in asthma).

*Genes demonstrating presence of airway type 2 inflammation.

[†]Genes/groups demonstrating presence of epithelial endoplasmic reticulum stress.

[‡]Genes/groups demonstrating presence of airway IFNs leading to epithelial IFN-stimulated gene expression.




Figure 2. Development of an airway epithelial brushing IFN-stimulated gene expression metric in mild asthma. (A) Hierarchical clustering of primary human epithelial cells cultured at an air-liquid interface (columns) using the top 50 (by fold-difference) genes (rows) induced by IFN- α (yellow, $n = 6$) relative to unstimulated (black, $n = 6$). All gene expression measurements are from RNA sequencing, with blue indicating low relative expression, and red indicating high relative expression. (B) Visualization of intersection of genes upregulated in asthma relative to health (false discovery rate < 0.01) and IFN- α -stimulated genes from A by supervised hierarchical clustering. $n = 19$ asthma (red); $n = 16$ healthy (black).

genes: *IFI6* (IFN α -inducible protein 6), *OAS1* (2'-5'-oligoadenylate synthetase 1), *USP18* (ubiquitin specific peptidase 18), *OAS2*, *IFITM1* (IFN-induced transmembrane protein 1), *BST2* (bone marrow stromal cell antigen 2), *IFI27* (IFN α -inducible protein 27), and *IFI35* (IFN-induced protein 35). This degree of overlap is significant because the probability of finding nine or more genes overlapping with the set of 1,379 genes differentially expressed in human airway brushings, when drawing 50 genes randomly without replacement from 19,938 protein-coding genes in the epithelial culture sequencing data, is P equal to 0.0067 by a Fisher's exact test.

Grouping subjects by the expression of these nine IFN-induced genes shows that 12 of 19 subjects with asthma have coordinately elevated expression for almost all of these

genes (Figure 2B). The centered, regularized read counts for these nine genes were added to form an airway ISG metric. We found no association between the epithelial ISG metric and our previously published type 2 inflammatory metric ($\rho = 0.12$; $P = 0.62$) (2), nor other markers of allergic inflammation including blood eosinophils, serum IgE, and fractional exhaled nitric oxide (Figure 3).

Airway ISG Expression Is Associated with Reduced Lung Function

Higher airway ISG expression was associated with reduced FEV₁ ($\rho = -0.72$; $P = 0.0004$) and increased bronchodilator response among subjects with asthma ($\rho = 0.75$; $P = 0.0002$) (Figures 4A and 4B). To validate this on a genome-wide scale, we performed IPA of the pathways associated with lower FEV₁ in asthma using all the RNA-seq data and found that IFN-related

pathways represented 5 of the top 10 pathways (see Table E3). We found no relationship between the ISG metric and age, sex, body mass index, or PC₂₀. Weaker relationships were seen between type 2 inflammation and lung function ($\rho = -0.40$, $P = 0.050$ for FEV₁; $\rho = 0.44$, $P = 0.032$ for bronchodilator response). The association between ISG expression and FEV₁ remained significant when adjusted for the type 2 metric ($P = 0.0056$), and there was no significant interaction between the type 2 and ISG metrics ($P = 0.75$ for interaction term).

Effect of ICS on Airway ISG Expression and Lung Function

Airway epithelial brush ISG activity was significantly attenuated by 8 weeks of ICS (see Figure E3A; $P = 0.007$ before vs. after ICS). After ICS treatment, airway ISG activity was not associated with FEV₁ or




Figure 3. Airway IFN-stimulated gene expression is not associated with airway type 2 inflammation. The airway IFN-stimulated gene metric was calculated from the expression of the nine signature genes in airway epithelial brushings acquired at bronchoscopy. Covariates were measured at baseline. Scatterplots of covariates versus the airway IFN-stimulated gene metric are shown for the following variables. (A) Airway epithelial type 2 gene expression metric ($n = 16$ healthy [solid circles] and $n = 19$ mild asthma [open circles]; $\rho = 0.12$; $P = 0.62$ within asthma). (B) Exhaled nitric oxide on a $\log_{10}$ axis ($n = 16$ healthy [solid circles] and $n = 15$ mild asthma [open circles]; $\rho = 0.30$; $P = 0.28$ within asthma). (C) Blood eosinophils on a square root axis ($n = 16$ healthy [solid circles] and $n = 19$ mild asthma [open circles]; $\rho = 0.22$; $P = 0.36$ within asthma). (D) Total serum IgE ($n = 15$ healthy [solid circles] and $n = 19$ mild asthma [open circles]; $\rho = -0.023$; $P = 0.93$ within asthma). Spearman rank correlation used in all cases. $FeNO$ = fractional exhaled nitric oxide; ISG = IFN-stimulated gene.

bronchodilator reversibility (see Figures E3B and E3C). The relationship between baseline ISG expression and the change in FEV_1 was significant after 2 but not after 4 and 8 weeks of ICS treatment (see Table E4). Thus, ISG expression at baseline had a mixed effect on ICS response, unlike the more consistent relationship seen for type 2 inflammation (2) (see Table E4). Similarly, compared with type 2 inflammation, ISG expression had lower area under the curve by receiver operating characteristic analysis in predicting those with a positive ICS response (see Figure E3D).

Airway ISG Expression Is Not Associated with a Distinct Viral Signature

Given the well-established relationship between IFNs and viral infection, we screened the airway epithelial RNA-seq for viral transcripts using our metagenomics pipeline. Potential viral pathogens were identified in low abundance in a small subset of subjects. A virus had to be present at more than 15 reads to be considered present. We

identified 35 human rhinovirus A reads in one subject with asthma with high ISG levels (see Figure E4 for the alignment of viral reads from this subject against the human rhinovirus A genome). Human herpes virus was detected in four healthy control subjects and one asthma subject with low ISG levels. To confirm the overall negative findings, we assessed the presence of three common respiratory viruses (respiratory syncytial virus, human rhinovirus, and parainfluenza virus type 3) by validated qPCR assays. All three viruses were undetectable in the human airway brush RNA samples, but were detected in positive control samples. The discordance between qPCR and RNA-Seq for human rhinovirus in the one subject previously mentioned suggests that the limit of detection for the qPCR assay is near 35 RNA-Seq reads in this dataset.

Whole-Blood Gene Expression Reflects Airway ISG Expression

Whole-blood PAXgene profiling by microarrays in 23 subjects with asthma not on ICS and 20 healthy control subjects

identified 1,372 differentially expressed transcripts ($P < 0.05$, see Table E5). Eosinophil-associated genes (*SIGLEC8* [sialic acid binding Ig-like lectin 8], *CLC* [Charcot-Leyden crystal galectin], *IL5RA* [IL5 receptor subunit α], *PRSS33* [protease, serine 33], *P2RY14* [purinergic receptor P2Y14], *RNASE2* [ribonuclease A family member 2]) topped this list (Figure 5A) (27). The high degree of correlation between these genes and with the percentage of blood eosinophils, but not neutrophils (see Figure E5 and Table E6), suggests that the differential expression is caused by the differences in eosinophil numbers between subjects with asthma and healthy control subjects; other cell types were not significantly different between groups (see Figure E6).

We used IPA on the differentially expressed gene list to identify non-eosinophil-associated pathways and found




Figure 4. Relationship of airway IFN-stimulated gene expression to lung function in mild asthma. (A) Baseline FEV_1 as a percent of the predicted value, measured immediately before and on the day of bronchoscopy before bronchodilator (BD) ($\rho = -0.72$; $P = 0.0004$ within asthma). (B) Change in FEV_1 as a percentage of predicted after administration of BD ($\rho = 0.75$; $P = 0.0002$ within asthma). Spearman rank correlation used and $n = 16$ healthy (solid circles) and $n = 19$ asthma (open circles) in all cases. ISG = IFN-stimulated gene.




Figure 5. Whole-blood gene expression reflects airway IFN-stimulated gene expression in mild asthma. (A) Unsupervised hierarchical clustering of whole-blood gene expression by six eosinophil-associated genes differentially expressed in asthma ($P < 0.05$). Samples (columns) were collected on the day of and immediately preceding bronchoscopy. Gene expression measurements are from microarrays, with blue indicating low relative expression, and red indicating high relative expression. $n = 23$ asthma (red) and $n = 20$ healthy (black). (B) Unsupervised hierarchical clustering of whole-blood gene expression by seven IFN-stimulated genes previously identified in whole blood. (C) Whole-blood IFN-stimulated gene metric versus airway (epithelial brushing) IFN-stimulated gene metric ($n = 16$ asthma; $\rho = 0.55$; $P = 0.030$). (D) Whole-blood eosinophil-associated gene metric versus blood eosinophils ($n = 21$ asthma; $\rho = 0.68$; $P = 0.00094$). Whole-blood gene expression metrics were made in the same manner as they were for the airway samples and related across tissues. Spearman rank correlation used in all cases. ISG = IFN-stimulated gene.

that IFN-associated genes were the top-scoring upstream regulators in asthma (e.g., *IRF7* [IFN regulatory factor 7], *STAT1*, *IFNB1* [IFN β 1], *IFNA2* [IFN α 2]) (Table 3). We developed a blood ISG metric by combining the expression of seven intercorrelated genes known to be responsive to IFN signaling in peripheral blood mononuclear cells: *IFI27*, *IFI44*, *MX1* (MX dynamin like GTPase 1), *IFIT1* (IFN-induced protein with tetratricopeptide repeats 1), *OAS1*, *OAS2*, and *OAS3* (Figure 5B) (28). We found significant correlation between airway and blood ISG metrics across subjects with asthma ($\rho = 0.55$; $P = 0.03$) (Figure 5C). A similarly robust relationship was seen between a blood eosinophil gene expression metric and blood eosinophils ($\rho = 0.68$; $P = 0.00094$, within asthma) (Figure 5D). Unlike in the airway, blood ISG expression was not correlated with FEV₁ in asthma ($n = 23$; $\rho = -0.29$; $P = 0.19$) or affected by ICS (see Figure E3). There was a modest negative

correlation between blood ISG expression and basophils but no other cell types (see Table E5). The expression of a subset of differentially expressed genes was validated by qPCR, with good correlation between platforms and consistency of differential expression (see Figure E7 and Table E7).

ISG Expression Is Increased in Severe Asthma

Demographic data for the subjects from the severe asthma study, in which they were randomized to placebo or omalizumab, are available in Table E8 (9). Unsupervised clustering of the blood ISG genes at baseline using microarrays showed a substantial fraction of subjects with elevated expression (Figure 6A and Table 4). There was no relationship between the blood ISG metric and type 2 inflammation as assessed by blood eosinophils (Figure 6B) (2); whereas, as expected, the blood eosinophil gene metric was significantly associated with blood eosinophils (Figure 6C). Consistent

with the effects of ICS in mild asthma, blood ISG expression and FEV₁ were not correlated in severe asthma ($n = 301$; $\rho = 0.0030$; $P = 0.96$). The lack of associations agrees with a recent description of “T1”-high severe asthma (29). Blood ISG expression in severe asthma was also unaffected by oral corticosteroid use (Figure 6A; see Figure E3). High compared with low ISG expression was not associated with a different exacerbation rate in the placebo arm (Table 4). In this trial, treatment with omalizumab reduced the exacerbation rate over 48 weeks relative to placebo. Markers of type 2 inflammation identified individuals more likely to respond (30). In contrast, a high blood ISG metric was not associated with a reduction in exacerbations in response to omalizumab (exacerbation

Table 3. IPA Upstream Regulator Analysis of Whole-Blood Array-based Gene Expression

Upstream Regulator	Activation z-Score	P Value of Overlap
<i>IRF7</i> *	4.78	5.68×10^{-6}
<i>STAT1</i> *	3.72	1.32×10^{-3}
<i>Cg</i>	3.68	3.01×10^{-2}
Interferon- α group*	3.57	3.91×10^{-4}
<i>PRL</i>	3.31	2.90×10^{-2}
<i>IFNB1</i> *	3.29	7.80×10^{-3}
<i>IFNA2</i> *	3.18	5.06×10^{-6}
<i>IRF1</i> *	3.04	1.30×10^{-3}
IFN- β group*	3.01	8.66×10^{-3}
<i>TGM2</i>	3.00	3.15×10^{-2}
<i>Ifnar</i> group*	2.87	1.57×10^{-3}
<i>Ifn</i> group*	2.78	9.94×10^{-3}
<i>IFNG</i> *	2.71	2.26×10^{-4}
<i>GATA2</i>	2.58	1.14×10^{-2}
<i>SMARCA4</i>	2.58	1.28×10^{-3}
<i>JUN</i>	2.45	4.57×10^{-2}
<i>IL4</i> [†]	2.40	2.53×10^{-6}
<i>IL5</i> [†]	2.36	6.29×10^{-6}
<i>IL2</i>	2.30	2.00×10^{-4}
<i>STAT3</i> *	2.20	3.57×10^{-5}
<i>RNASE2</i>	2.17	4.03×10^{-2}
<i>IL27</i>	2.16	1.56×10^{-2}
<i>IFNL1</i> *	2.04	1.28×10^{-4}
<i>IFNAR2</i> *	2.00	2.70×10^{-2}

Definition of abbreviation: IPA = ingenuity pathway analysis.

Regulators are filtered by 1) genes, proteins, or RNA; 2) activated predicted state; 3) activation z-score ≥ 2 ; and 4) $P < 0.05$. The list is sorted by activation z-score.

*Genes/groups demonstrating presence of airway IFN leading to epithelial IFN-stimulated gene expression.

[†]Genes demonstrating presence of airway type 2 inflammation.

rate reduction = 1.27%; 95% confidence interval, -60.13 to 39.20; $P = 0.96$ for effect of treatment on number of exacerbations) (Table 4). Although there was a trend for a response to omalizumab for the group




Figure 6. IFN-stimulated gene (ISG) expression is increased in severe asthma and unrelated to type 2 inflammation. (A) Unsupervised hierarchical clustering of whole-blood gene expression by seven ISGs. Severe asthma samples (columns) were collected at baseline before change in treatment. Gene expression measurements are from microarrays, with blue indicating low relative expression, and red indicating high relative expression, $n = 301$ severe asthma (color-coded red if on oral corticosteroids). (B) Whole-blood ISG expression is not related to blood eosinophil count, a measure of type 2 inflammation ($n = 288$ severe asthma; $\rho = 0.057$; $P = 0.34$). (C) Whole-blood eosinophil-associated gene expression is associated with blood eosinophil count ($n = 288$ severe asthma; $\rho = 0.82$; $P < 2.2 \times 10^{-16}$). Spearman rank correlation was used in both B and C. eos = eosinophils; OCS = oral corticosteroids; sqrt = square root.

Table 4. Association of ISG Expression with Subject Characteristics and Exacerbation Rates in Response to Omalizumab in Severe Asthma

	ISG-High*	ISG-Low*	P Value
Age, yr [†]	45.5 ± 13.8	46.0 ± 13.8	0.72
Female, % [†]	77	57	0.0002
BMI, kg/m ^{2†}	33.2 ± 7.9	31.0 ± 6.9	0.01
Number of exacerbations in year before enrollment [†]	2 (0 to 11)	2 (0 to 8)	0.66
N, placebo/omalizumab	76/75	74/76	—
Annualized exacerbation rate			—
Placebo (95% CI)	1.0 (0.74 to 1.36)	0.80 (0.56 to 1.11)	
Omalizumab (95% CI)	1.0 (0.73 to 1.34)	0.55 (0.36 to 0.81)	
P value for effect of treatment [‡]	0.96	0.11	
Exacerbation rate reduction % (95% CI)	1.3 (-60.3 to 39.2)	31.1 (-7.9 to 56.0)	0.3 [§]

Definition of abbreviations: BMI = body mass index; CI = confidence interval; ISG = IFN-stimulated gene.

*ISG-high and -low are determined as ISG expression metrics above or below the median value for all subjects at baseline.

[†]Baseline traits are presented as mean ± SD or median (range), and statistical differences were assessed by a two-sided Welch two-sample *t* test for the former and a generalized linear model with quasi-Poisson error distribution for the latter, as described in the METHODS. For female percentage, a chi-square test for equality of proportions was performed.

[‡]P value for treatment term with ISG-low and -high modeled separately as described in the METHODS.

[§]P value for interaction term in a model of exacerbation rate including treatment and ISG status as predictors.

with a low blood ISG metric, the effect of ISG status on treatment response was not significant ($P = 0.3$ for interaction term) (Table 4).

Airway Epithelial Endoplasmic Reticulum Stress Is Common to ISG and Type 2 Inflammation in Asthma

In response to endoplasmic reticulum (ER) stress, cells activate an unfolded-protein response (UPR) initiated by one of three signal transduction pathways: 1) IRE1 α (inositol-requiring enzyme 1 α)-XBP1 (X-box binding protein 1), 2) PERK (protein kinase R [PKR]-like endoplasmic reticulum kinase), and 3) ATF6 (activation transcription factor 6). Our IPA analysis showed prominent activation of the IRE1 α -XBP1 (IPA regulator XBP1) pathway in mild asthmatic airway brushings (Table 2 and Figure 7A). Activation of IRE1 α leads to splicing of *XBP1* mRNA to an activated form, which we confirmed in epithelial brushings through genome-wide splicing analysis of the airway brushing RNA-seq data (FDR = 0.0001) (Figure 7B). Notably, IPA upstream regulator analysis did not reveal significant activation of the XBP1 pathway in whole-blood microarray gene expression (Table 2). There was significant attenuation of ER stress after 8 weeks of ICS (see Figure E3).

We examined the relationship between XBP1 activation and airway ISG expression and type 2 inflammation. Genes downstream of XBP1 were combined into a single metric of ER stress that was associated with heightened type 2 inflammation and ISG expression (Figures 7C and 7D). Individuals with asthma and low levels of type 2 inflammation and ISG expression did not have activation of the XBP1 arm of the UPR. Type 2 inflammation and ISG expression remained significantly associated with ER stress when added in a linear model ($P = 0.00076$ and 0.0059 , respectively), without a significant interaction ($P = 0.85$). Consistent with the association between XBP1 activation and both type 2 inflammation and ISG expression, stimulation of epithelial cell cultures by IL-13 or IFN- α at an air-liquid interface showed activation of all arms of the UPR (see Table E9).

Discussion

We found that a clinically distinct subset of asthma has increased ISG expression in the airway and in peripheral blood. In subjects with mild asthma not on ICS, this ISG-high subgroup had decreased lung function and increased bronchodilator




Figure 7. Epithelial endoplasmic reticulum (ER) stress markers are elevated in mild asthma and associated with both high type 2 inflammation and IFN-stimulated gene expression. (A) Hierarchical clustering of airway epithelial brushing gene expression samples (columns) using genes differentially expressed (false discovery rate < 0.01) in asthma versus health and downstream of XBP1 (X-box binding protein 1) in IPA. (B) XBP1 splicing assessed bioinformatically from RNA sequencing data through use of the rMATS package in R (false discovery rate = 0.0001 for genome-wide splicing analysis). (C) Relationship of an ER stress gene expression metric consisting of genes downstream of XBP1 per IPA, to airway type 2 inflammation ($n = 19$ mild asthma; $\rho = 0.62$; $P = 0.0058$). (D) ER stress gene expression metric versus the airway IFN-stimulated gene metric ($n = 19$ mild asthma; $\rho = 0.62$; $P = 0.0052$). Only asthma shown. IPA = ingenuity pathway analysis; ISG = IFN-stimulated gene.

reversibility. In both mild and severe asthma, ISG expression was independent of type 2 inflammation. These data extend prior reports that support the presence of airway IFNs in stable asthma (23, 29, 31–34) by showing that IFN-driven inflammation is orthogonal to type 2 inflammation and is not associated with the presence of viral infection. Furthermore, we found that epithelial ER stress is related to both ISG expression and type 2 inflammation, making it a potential common downstream therapeutic target.

We did not find an association between the presence of viral RNA as detected by

RNA-seq or qPCR and the IFN response in this immunocompetent and clinically stable group. One recent report found evidence for IFN responses associated with occult viral transcripts in nasal brushings in some asymptomatic children with asthma (35). Our experiments used similar PCR methods and did not detect viral transcripts in the lower airway in this subgroup, with the exception of one subject. Our findings may differ because of differences in location (upper vs. lower airway) and age. In contrast to stable asthma, a large percentage of asthma exacerbations are induced by

viral infection, particularly with rhinovirus, and subjects with asthma have increased lower respiratory tract symptoms after viral infection (36). Deficient type I and type III IFN production in response to viral infection has been observed in epithelial cells recovered from asthmatic airways (37, 38). These findings in an *ex vivo* infection model are nonetheless potentially compatible with our results of heightened responses during stable disease. In fact, baseline elevated IFN responses can lead to desensitization to type I IFN stimulation through the ISG *USP18*, which we found induced in airway brushings (39).

Although ISGs have been most studied in association with viral infection, host expression of ISGs can also occur in response to LPS-containing gram-negative bacteria and some intracellular bacteria and fungi (40, 41). Future studies will address the relationship between airway ISG expression and the microbiome, which is dysregulated in asthma (42–47).

In our study, type 2 inflammation and ISG expression were not correlated. Previous work found that T-helper cell type 1 and IFN responses have suppressive effects on type 2 responses (48), suggesting that the two should be inversely related. Our results do not exclude the possibility that ISG expression by epithelial cells represents a regulatory mechanism aimed at dampening excessive type 2 inflammation. If true, this could lead to dual responses in some individuals, whereas in others ISG expression may persist after type 2 responses recede.

Although type 2 and IFN-driven inflammation were unrelated in airway epithelial brushings, both were independently associated with markers of airway epithelial ER stress. Thus, IRE1-XBP1-regulated airway epithelial ER stress may be a final common pathway for both types of inflammation. ER stress is a pathway of established importance in many diseases and airway epithelial ER stress is emerging as potentially important in asthma. The inflammatory environment of the airway in asthma offers multiple ways in which ER stress, or the UPR, can be induced. In mice, house-dust mite allergen challenge is a potent activator of epithelial ER stress, and knockdown of two ER stress-associated proteins, ATF6 and ERp57, decreased airway inflammation, hyperresponsiveness, and fibrosis (49, 50). In another study, IRE1 β -initiated ER stress

was required for epithelial mucin production (51). These studies support the notion that type 2 inflammation, which is associated with increased epithelial production and secretion of mucus and innate cytokines, can lead to ER stress. How IFN-driven inflammation might induce ER stress is less well understood. In our data, there was evidence of heightened ER stress-related gene expression in airway brushings but not in blood samples in mild asthma, and therefore we were unable to assess this pathway in severe asthma. Further studies in severe asthma will be important because inhibition of ER stress may be a novel therapeutic strategy for asthma based on preclinical results from animal models (52).

The largely overlapping responses of type I (IFN- α/β), II (IFN- γ), and III (IFN- λ) IFNs preclude distinguishing these pathways using transcript profiling (26). The cellular source of airway IFNs in our study is also unknown. Although type I and type III IFNs are secreted by many cell types including epithelial and hematopoietic cells, type II IFN secretion is limited to specific immune cells including T-helper cell type 1, natural killer T, and natural killer cells. In our airway brushing RNA-seq data, IFNs were not differentially expressed, suggesting that epithelial cells or luminal hematopoietic cells are not a likely source. Uncollected airway cells, such as inflammatory cells in the lamina propria, may contribute to airway IFNs.

Recently, approximately 70% of humans with severe asthma were found to have levels of IFN- γ -producing BAL fluid CD4⁺ T cells higher than the upper limit seen in mild-moderate asthma (23). The relationships between these IFN- γ -producing T cells, type 2 inflammation, and lung function were not presented, which combined with the lack of healthy control subjects precludes direct comparison with our findings. Additional data from that study suggest that IFN- γ -induced downregulation of *SLPI* (secretory leukocyte protease inhibitor) in airway epithelial cells leads to increased

airway hyperresponsiveness in severe asthma. We did not find differential expression of the *SLPI* gene in mild asthma. Furthermore, the production of type I and type III IFNs was not reported in that study, leaving uncertain which IFNs drive airway ISG expression in asthma.

Other potential explanations for the increased ISG expression that we observed include impaired resolution of IFN-driven inflammation, a higher gain in the IFN response pathways (22), or expression of endogenous retroelements. An abnormally high gain has the most support based on the previously reported finding of constitutive activation of the STAT1 transcription factor, a key mediator of intracellular IFN responses, in the airway epithelium from subjects with mild asthma (22). In the current study, we were limited to studying gene expression and therefore were unable to address the activation of STAT1. Although STAT1 is known to be downstream of IFN stimulation and was induced by IFN- α in our epithelial cultures, it was not differentially upregulated in our airway brushings from asthma relative to health. Possible explanations for this difference are that there are stimuli acting on the epithelium beyond IFNs *in vivo* (e.g., type 2 inflammation, microbiome), and that the level and duration of IFN stimulation differs between *in vivo* and culture conditions.

Our paired whole-blood gene expression analysis showed that blood cell ISG expression is elevated in stable asthma, not just with asthma exacerbations and suboptimal control (53, 54). The ISG signature in blood is not as prominent as it is in the airway, and therefore we used a significance threshold of an unadjusted *P* less than 0.05 rather than a more stringent FDR cutoff to identify a sufficient number of genes for optimal pathway analysis. Our data further show that a subgroup of severe subjects with asthma has elevated whole-blood ISG expression and that as in mild asthma, ISG expression and type 2 inflammation are orthogonal. Our data also suggest that elevated ISG expression in severe asthma may be associated with decreased exacerbation reduction benefit from anti-IgE therapy. This is an intriguing

trend identified in an exploratory study and can be tested in adequately powered trials in the future. The finding raises the possibility that incorporation of blood gene expression biomarkers could complement type 2 inflammatory markers and support broader clinical detection of ISG expression in asthma.

Notably, the presence of subgroups with elevated ISG expression in both mild and severe asthma parallels what has been found for the established endotype of type 2 inflammation (1, 55). In addition, similar to characteristics of type 2 inflammation, we found that ICS treatment of subjects with mild asthma led to reduced ISG expression and loss of the relationship of ISG expression to lung function, and that ISG expression is unrelated to FEV₁ in severe asthma (2, 55). Thus, the presence of elevated ISG expression in mild asthma, and the effects of ICS on ISG expression and its relationship to FEV₁, does not diminish the potential importance of IFN-driven inflammation in severe asthma.

This study adds to an expanding literature on asthma phenotyping and demonstrates that differential ISG expression defines a clinically relevant subgroup of subjects with asthma. This work informs further studies in larger cohorts designed to delineate the diversity of ISG responses. Additional evaluation of ISG expression in severe asthma, including associations with responses to existing and emerging therapeutics, will be an important future direction. ER stress may be a common pathway for both type 2 and IFN-driven inflammation, and may offer a new target for the treatment of asthma. ■

Author disclosures are available with the text of this article at www.atsjournals.org.

Acknowledgment: The authors thank the University of California, San Francisco, Sandler Genomics Core and the Parnassus Genomics Core for providing high-quality RNA sequencing data. Walter E. Finkbeiner gave valuable support for the cell culture experiments and was funded by the Cystic Fibrosis Foundation (RDP-R613) and NIH DK072517.

References

1. Woodruff PG, Modrek B, Choy DF, Jia G, Abbas AR, Ellwanger A, *et al*. T-helper type 2-driven inflammation defines major subphenotypes of asthma. *Am J Respir Crit Care Med* 2009;180:388–395.
2. Bhakta NR, Solberg OD, Nguyen CP, Nguyen CN, Arron JR, Fahy JV, *et al*. A qPCR-based metric of Th2 airway inflammation in asthma. *Clin Transl Allergy* 2013;3:1–9.
3. Ray A, Raundhal M, Oriss TB, Ray P, Wenzel SE. Current concepts of severe asthma. *J Clin Invest* 2016;126:2394–2403.

4. Fahy JV. Type 2 inflammation in asthma: present in most, absent in many. *Nat Rev Immunol* 2015;15:57–65.
5. 't Hoen PA, Friedländer MR, Almlöf J, Sammeth M, Pulyakhina I, Anvar SY, et al.; GEUVADIS Consortium. Reproducibility of high-throughput mRNA and small RNA sequencing across laboratories. *Nat Biotechnol* 2013;31:1015–1022.
6. SEQC/MAQC-III Consortium. A comprehensive assessment of RNA-seq accuracy, reproducibility and information content by the Sequencing Quality Control Consortium. *Nat Biotechnol* 2014;32:903–914.
7. Christenson SA, Bhakta NR, Nerella S, Pollack JL, Barbeau R, Barczak AJ, et al. RNA-sequencing of airway epithelial brushings identifies interferon-driven gene expression changes in asthma [abstract]. *Am J Respir Crit Care Med* 2015;191:A3626.
8. Bonser LR, Christenson SA, Pollack JL, Barbeau R, Barczak AJ, Zlock L, et al. Investigating mRNA and miRNA alterations induced by asthma-associated cytokine stimulation in a human airway model using RNASeq [abstract]. *Am J Respir Crit Care Med* 2016;193:A4624.
9. Hanania NA, Alpan O, Hamilos DL, Condemi JJ, Reyes-Rivera I, Zhu J, et al. Omalizumab in severe allergic asthma inadequately controlled with standard therapy: a randomized trial. *Ann Intern Med* 2011;154:573–582.
10. Woodruff PG, Boushey HA, Dolganov GM, Barker CS, Yang YH, Donnelly S, et al. Genome-wide profiling identifies epithelial cell genes associated with asthma and with treatment response to corticosteroids. *Proc Natl Acad Sci USA* 2007;104:15858–15863.
11. Anders S, McCarthy DJ, Chen Y, Okoniewski M, Smyth GK, Huber W, et al. Count-based differential expression analysis of RNA sequencing data using R and Bioconductor. *Nat Protoc* 2013;8:1765–1786.
12. Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J R Stat Soc Series B Stat Methodol* 1995;57:289.
13. Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, et al. limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Res* 2015;43:e47.
14. Smyth GK. Linear models and empirical Bayes methods for assessing differential expression in microarray experiments. *Stat Appl Genet Mol Biol* 2004;3:1–25.
15. Doan T, Wilson MR, Crawford ED, Chow ED, Khan LM, Knopp KA, et al. Illuminating uveitis: metagenomic deep sequencing identifies common and rare pathogens. *Genome Med* 2016;8:1–9.
16. Ruby JG, Bellare P, Derisi JL. PRICE: software for the targeted assembly of components of (Meta) genomic sequence data. *G3 (Bethesda)* 2013;3:865–880.
17. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* 2013;29:15–21.
18. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. *Nat Methods* 2012;9:357–359.
19. Fu L, Niu B, Zhu Z, Wu S, Li W. CD-HIT: accelerated for clustering the next-generation sequencing data. *Bioinformatics* 2012;28:3150–3152.
20. Wu TD, Nacu S. Fast and SNP-tolerant detection of complex variants and splicing in short reads. *Bioinformatics* 2010;26:873–881.
21. Wickham H. ggplot2: elegant graphics for data analysis. New York: Springer-Verlag; 2009.
22. Sampath D, Castro M, Look DC, Holtzman MJ. Constitutive activation of an epithelial signal transducer and activator of transcription (STAT) pathway in asthma. *J Clin Invest* 1999;103:1353–1361.
23. Raundhal M, Morse C, Khare A, Oriss TB, Milosevic J, Trudeau J, et al. High IFN- γ and low SLP1 mark severe asthma in mice and humans. *J Clin Invest* 2015;125:3037–3050.
24. Wu D, Smyth GK. Camera: a competitive gene set test accounting for inter-gene correlation. *Nucleic Acids Res* 2012;40:e133.
25. Christenson SA, Steiling K, van den Berge M, Hijazi K, Hiemstra PS, Postma DS, et al. Asthma-COPD overlap. Clinical relevance of genomic signatures of type 2 inflammation in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2015;191:758–766.
26. Crotta S, Davidson S, Mhlakoiv T, Desmet CJ, Buckwalter MR, Albert ML, et al. Type I and type III interferons drive redundant amplification loops to induce a transcriptional signature in influenza-infected airway epithelia. *PLoS Pathog* 2013;9:e1003773.
27. Esnault S, Kelly EA, Schwantes EA, Liu LY, DeLain LP, Hauer JA, et al. Identification of genes expressed by human airway eosinophils after an in vivo allergen challenge. *PLoS One* 2013;8:e67560.
28. McBride JM, Jiang J, Abbas AR, Morimoto A, Li J, Maciuga R, et al. Safety and pharmacodynamics of rontalizumab in patients with systemic lupus erythematosus: results of a phase I, placebo-controlled, double-blind, dose-escalation study. *Arthritis Rheum* 2012;64:3666–3676.
29. Modena BD, Bleecker ER, Busse WW, Erzurum SC, Gaston BM, Jarjour NN, et al. Gene expression correlated with severe asthma characteristics reveals heterogeneous mechanisms of severe disease. *Am J Respir Crit Care Med* 2017;195:1449–1463.
30. Hanania NA, Wenzel S, Rosén K, Hsieh HJ, Mosesova S, Choy DF, et al. Exploring the effects of omalizumab in allergic asthma: an analysis of biomarkers in the EXTRA study. *Am J Respir Crit Care Med* 2013;187:804–811.
31. Cho SH, Stanciu LA, Holgate ST, Johnston SL. Increased interleukin-4, interleukin-5, and interferon-gamma in airway CD4+ and CD8+ T cells in atopic asthma. *Am J Respir Crit Care Med* 2005;171:224–230.
32. Krug N, Madden J, Redington AE, Lackie P, Djukanovic R, Schauer U, et al. T-cell cytokine profile evaluated at the single cell level in BAL and blood in allergic asthma. *Am J Respir Cell Mol Biol* 1996;14:319–326.
33. Bullens DM, Decraene A, Dilissen E, Meyts I, De Boeck K, Dupont LJ, et al. Type III IFN-lambda mRNA expression in sputum of adult and school-aged asthmatics. *Clin Exp Allergy* 2008;38:1459–1467.
34. da Silva J, Hilzendeger C, Moermans C, Schleich F, Henket M, Kebabdzic T, et al. Raised interferon beta, type 3 interferon and interferon stimulated genes: evidence of innate immune activation in neutrophilic asthma. *Clin Exp Allergy* 2016;47:313–323.
35. Wesolowska-Andersen A, Everman JL, Davidson R, Rios C, Herrin R, Eng C, et al. Dual RNA-seq reveals viral infections in asthmatic children without respiratory illness which are associated with changes in the airway transcriptome. *Genome Biol* 2017;18:1–17.
36. Corne JM, Marshall C, Smith S, Schreiber J, Sanderson G, Holgate ST, et al. Frequency, severity, and duration of rhinovirus infections in asthmatic and non-asthmatic individuals: a longitudinal cohort study. *Lancet* 2002;359:831–834.
37. Wark PA, Johnston SL, Bucchieri F, Powell R, Puddicombe S, Laza-Stanca V, et al. Asthmatic bronchial epithelial cells have a deficient innate immune response to infection with rhinovirus. *J Exp Med* 2005;201:937–947.
38. Contoli M, Message SD, Laza-Stanca V, Edwards MR, Wark PA, Bartlett NW, et al. Role of deficient type III interferon-lambda production in asthma exacerbations. *Nat Med* 2006;12:1023–1026.
39. Schneider WM, Chevillotte MD, Rice CM. Interferon-stimulated genes: a complex web of host defenses. *Annu Rev Immunol* 2014;32:513–545.
40. Monroe KM, McWhirter SM, Vance RE. Induction of type I interferons by bacteria. *Cell Microbiol* 2010;12:881–890.
41. Morton CO, Bouzani M, Loeffler J, Rogers TR. Direct interaction studies between *Aspergillus fumigatus* and human immune cells; what have we learned about pathogenicity and host immunity? *Front Microbiol* 2012;3:413.
42. Marri PR, Stern DA, Wright AL, Billheimer D, Martinez FD. Asthma-associated differences in microbial composition of induced sputum. *J Allergy Clin Immunol* 2013;131:346–352.
43. Huang YJ, Nelson CE, Brodie EL, Desantis TZ, Baek MS, Liu J, et al.; National Heart, Lung, and Blood Institute's Asthma Clinical Research Network. Airway microbiota and bronchial hyperresponsiveness in patients with suboptimally controlled asthma. *J Allergy Clin Immunol* 2011;127:372–381.
44. Bundy KW, Gent JF, Beckett W, Bracken MB, Belanger K, Triche E, et al. Household airborne *Penicillium* associated with peak expiratory flow variability in asthmatic children. *Ann Allergy Asthma Immunol* 2009;103:26–30.

45. Sharpe RA, Bearman N, Thornton CR, Husk K, Osborne NJ. Indoor fungal diversity and asthma: a meta-analysis and systematic review of risk factors. *J Allergy Clin Immunol* 2015;135:110–122.
46. Chou H, Wu KG, Yeh CC, Tai HY, Tam MF, Chen YS, *et al.* The transaldolase, a novel allergen of *Fusarium proliferatum*, demonstrates IgE cross-reactivity with its human analogue. *PLoS One* 2014;9:e103488.
47. Agbetile J, Fairs A, Desai D, Hargadon B, Bourne M, Mutalithas K, *et al.* Isolation of filamentous fungi from sputum in asthma is associated with reduced post-bronchodilator FEV1. *Clin Exp Allergy* 2012;42:782–791.
48. Gonzales-van Horn SR, Farrar JD. Interferon at the crossroads of allergy and viral infections. *J Leukoc Biol* 2015;98:185–194.
49. Hoffman SM, Chapman DG, Lahue KG, Cahoon JM, Rattu GK, Daphtary N, *et al.* Protein disulfide isomerase-endoplasmic reticulum resident protein 57 regulates allergen-induced airways inflammation, fibrosis, and hyperresponsiveness. *J Allergy Clin Immunol* 2016;137:822–832.e7.
50. Hoffman SM, Tully JE, Nolin JD, Lahue KG, Goldman DH, Daphtary N, *et al.* Endoplasmic reticulum stress mediates house dust mite-induced airway epithelial apoptosis and fibrosis. *Respir Res* 2013;14:1–12.
51. Martino MB, Jones L, Brighton B, Ehre C, Abdulah L, Davis CW, *et al.* The ER stress transducer IRE1 β is required for airway epithelial mucin production. *Mucosal Immunol* 2013;6:639–654.
52. Siddesha JM, Nakada EM, Mihavics BR, Hoffman SM, Rattu GK, Chamberlain N, *et al.* Effect of a chemical chaperone, tauroursodeoxycholic acid, on HDM-induced allergic airway disease. *Am J Physiol Lung Cell Mol Physiol* 2016;310:L1243–L1259.
53. Bjornsdottir US, Holgate ST, Reddy PS, Hill AA, McKee CM, Csimma CI, *et al.* Pathways activated during human asthma exacerbation as revealed by gene expression patterns in blood. *PLoS One* 2011;6:e21902.
54. Croteau-Chonka DC, Qiu W, Martinez FD, Strunk RC, Lemanske RF Jr, Liu AH, *et al.*; Asthma BioRepository for Integrative Genomic Exploration (Asthma BRIDGE) Consortium. Gene expression profiling in blood provides reproducible molecular insights into asthma control. *Am J Respir Crit Care Med* 2017;195:179–188.
55. Jia G, Erickson RW, Choy DF, Mosesova S, Wu LC, Solberg OD, *et al.*; Bronchoscopic Exploratory Research Study of Biomarkers in Corticosteroid-refractory Asthma (BOBCAT) Study Group. Periostin is a systemic biomarker of eosinophilic airway inflammation in asthmatic patients. *J Allergy Clin Immunol* 2012;130:647–654.e10.

Interferon-Stimulated Gene Expression, Type-2 Inflammation and Endoplasmic Reticulum Stress in Asthma

Nirav R. Bhakta, Stephanie A. Christenson, Srilaxmi Nerella, Owen D. Solberg, Christine P. Nguyen, David F. Choy, Kyle L. Jung, Suresh Garudadri, Luke R. Bonser, Joshua L. Pollack, Lorna T. Zlock, David J. Erle, Charles Langelier, Joseph L. Derisi, Joseph R. Arron, John V. Fahy, and Prescott G. Woodruff

ONLINE DATA SUPPLEMENT

Supplementary Methods, Tables and Figures:

Supplemental Methods:

Inclusion/Exclusion criteria for primary cohort of mild asthma and healthy controls

All participants were between 18 and 70 years of age. Healthy controls had no history of asthma or allergic rhinitis. Subjects with asthma had a participant-reported history of asthma and were required to not have used inhaled or oral steroids for at least 6 weeks prior to enrollment, to have airway hyper-responsiveness (AHR) defined by a 20% decrease in FEV₁ (forced expiratory volume in 1 second) in response to inhaled methacholine at a provocative concentration (PC₂₀) $\leq$ 8.0 mg/mL, and at least one of the following: asthma symptoms on at least two days per week, beta agonist use on at least two days per week, or FEV₁ < 85% of predicted. Inhaled corticosteroids were not stopped in order to meet enrollment criteria, and therefore most subjects had not been using ICS for much longer than 6 weeks, as part of their usual care. All subjects were non-smokers defined as never smoker or former smokers with no smoking for at least 1 year *and* total pack-years $\leq$ 15. Additional exclusion criteria were pregnancy, upper respiratory tract infection in the 4 weeks prior to enrollment, and the use of certain other medications. Three prior publications have described other results from this cohort (1-3).

Reasons for loss of subjects across the bronchoscopy study of mild asthma

Of 32 subjects with mild asthma that met inclusion/exclusion criteria and underwent baseline characterization, four did not undergo bronchoscopy or further visits because the subjects' schedule conflicts; two did not show up for a scheduled bronchoscopy; and in one

subject who underwent bronchoscopy, the procedure was not well-tolerated and therefore airway brushings were not obtained. Of the 22 healthy controls that met inclusion/exclusion criteria and underwent baseline characterization, two did not undergo bronchoscopy or further visits due to the subjects' schedule conflicts. Therefore, airway brushings were collected from 25 subjects with mild asthma and 20 healthy controls. The remaining reasons for sample loss are described in Figure 1 of the manuscript.

Epithelial brushing RNA sequencing

RNA was isolated using the Qiagen miRNeasy kit (Qiagen Inc., Valencia, CA) as per manufacturer's protocol. RNA quality and quantity were assessed using the Agilent 2100 Bioanalyzer (Agilent Technologies, Santa Clara, CA) and the NanoDrop Spectrophotometer (Thermo Fisher Scientific, Wilmington, DE). Sequencing libraries were prepared using 100ng of RNA input per subject at the UCSF Sandler Genomics Core Facility. Library preparation and multiplexing was done using the Illumina TruSeq Stranded Total RNA with Ribo Zero Human/Mouse/Rat kit (Illumina Inc, San Diego, CA) as per manufacturer's protocol. One subject was excluded due to low sample RNA input. 100 base pair paired end sequencing was done on multiplexed samples via an Illumina HiSeq 2000 at the UCSF Genomics Core. Airway brushing cell counts were performed on cytopsin preparations.

Host gene expression analyses

Raw sequencing data underwent adapter and quality trimming using the cutadapt and sickle programs, then alignment to the human genome using the STAR aligner and the ENSEMBL GRCh38 genome build (4-6). Transcript abundance was determined using HT-Seq (7). Quality assessment was conducted using FastQC and STAR (8). Twelve samples were excluded from analysis (3 healthy, 4 steroid-naïve asthma, 5 asthma after steroids) due to low alignment to the human genome (<75% of reads aligned). These twelve samples were also found to have high alignment rates to human ribosomal RNA (rRNA) and mitochondrial RNA (mtRNA) (average of 58% and 17%, respectively), and thus were thought to have high amounts of rRNA contamination accounting for the low alignment rates. Following exclusion of these samples, the average read count for the remaining 43 samples was 152 million reads (76 million aligned pairs). The average percent of reads uniquely mapping to the human genome was 91% with 6% aligning to human ribosomal RNA sequences and 2% aligning to human mitochondrial RNA sequences.

Airway epithelial cell cultures

Human bronchial epithelial cells obtained from the proximal airways of lungs rejected for transplant (5 originated from individuals with no history of airway disease, one had a history of asthma) were grown to confluence in an air-liquid interface culture for 28 days as described previously (9). Some cultures were stimulated with IFN- α (10 ng/mL) for the final 24 hours or IL-13 for the final 7 days. Cells cultured in the presence and absence of IFN- α were then harvested and RNA isolated and sequenced as above.

PAXgene whole blood and airway brushing gene expression microarrays

PAXgene tubes were collected and processed for RNA per manufacturer's recommendations. Cells spun down from brushings were resuspended in RLT buffer (Qiagen) and underwent RNA extraction with the RNeasy kit (Qiagen). Quantity and quality were assessed by NanoDrop and Agilent Bioanalyzer (Nano), respectively. Samples were prepared for and run on Affymetrix HGU133A Plus 2.0 arrays as previously described (10). Background adjustment (without the use of mismatch probes), quantile normalization, and probe summarization were done using the RMA algorithm with the hgu133plus2hsentrezg custom chip definition file for annotation (<http://brainarray.mbni.med.umich.edu>). Batch effect was minimized using Combat with both array batches and disease status specified (11). Low quality arrays were identified and removed with the ArrayQualityMetrics package (www.bioconductor.org), followed by reapplication of the RMA algorithm and Combat batch correction. The entire analysis pipeline was performed separately for different tissue types in the R statistical environment (12). Airway brushes were processed for microarray analysis as previously described (10) after which we followed the analysis steps described for the PAXgene whole blood samples above. The airway brush array data are available as GSE67472 for a slightly larger group of subjects, and were previously used for a gene set list derivation in (3).

To assess short-term stability of whole blood microarray gene expression profiles, PAXgene samples collected at the baseline visit were included in normalization, batch correction, and iterative application of quality assessment in a separate analysis. The PAXgene samples

from the severe asthma cohort was analyzed separately with the same steps described above; after removing microarrays with low quality scores, 150 samples on placebo and 151 on omalizumab remained. Figure 1 in the main manuscript outlines the steps taken to derive the final sample numbers.

qPCR and enzyme-linked immunosorbent assay

Two-step, nested-primer RT-qPCR of PAXgene RNA was performed as described previously (13). Data were normalized to the mean of GAPDH, PPIA, and RPL37A as previously described (1). Primer and probe sequences are listed in SupplementalTable

10. qPCR of epithelial brushing RNA for viruses was performed on 5 ng of RNA per sample and per virus using primers, probes and protocols previously described (14, 15).

Positive controls validated three assays in our laboratory: respiratory syncytial virus, human rhinovirus, and human parainfluenza virus type 3.

Statistical Analysis of Gene Expression

Read count normalization and differential expression analyses were performed using the DESeq2 package (16). DESeq2 uses generalized linear models following a negative binomial distribution to test for differences between groups. Results were trimmed to transcripts indexed in the HGNC database and with Ensembl gene biotype of “protein_coding”. For unsupervised clustering, heatmaps, and multi-gene metrics, regularized-logarithm transformed counts from DESeq2's rlog function were used to provide more stable estimates of expression at low read counts. For assessment of differential expression in whole blood microarray data, moderated t tests were performed with

the limma package in the R statistical environment without adjustment of the degrees of freedom for prior batch correction (17). Multiple comparisons corrections were done using FDR by the Benjamini-Hochberg method (18). Genome-wide quantification and differential expression results between asthma and health of alternatively spliced transcripts were obtained using the rMATS package (19).

Ingenuity Pathway Analysis

For analysis of differentially expressed genes (Tables 2 and 3 in the manuscript), Ingenuity Pathway Analysis was run with the log₂ fold-change between a two-group comparison as the quantitative input (asthma not on ICS vs healthy). For the epithelial brushing RNA sequencing data, the list of genes to input into IPA was generated by requiring an FDR<0.01 for the asthma vs healthy comparison, a HUGO gene name, and an Ensembl biotype of protein coding; 1379 genes met these requirements, with 1376 mapping to IPA entries and 1372 identified as analysis-ready by the software. For the PAXgene whole blood microarray data, the list of genes to input into IPA was generated by requiring $p < 0.05$; 1372 genes met this requirement, with 1287 mapping to IPA entries and identified as analysis-ready. After running a basic core analysis, upstream regulator analysis output was filtered within IPA for regulators that were either genes, RNA, or proteins, and for regulators that had a predicted activated state. For the analysis of epithelial brushing genes and pathways associated with FEV₁ (Supplemental Table E1), RNA sequencing-based regularized-logarithm transformed counts from DESeq2's rlog function were used in genome-wide Spearman rank correlations between gene expression and FEV₁ as a percent of predicted, within the group of 19 asthmatics not on ICS. 1,386 genes with absolute Spearman rho >

0.5 were run through a basic core analysis.

Generation of multi-gene metrics

Gene expression values in log₂-scale with centering only or with centering and scaling were summed with equal weighting to form multi-gene metrics. There is no consensus on whether individual genes require centering only, or centering and scaling prior to combination into multi-gene metrics. For consistency, we used the previously reported airway type-2 metric for this cohort, which was computed as the mean of mean-centered and standard-deviation-scaled, log₂ qPCR normalized relative expression values (1). Given the use of a larger number of genes in the new interferon-associated metrics presented here, we performed mean centering only so that fold changes in gene expression weigh the contribution of individual genes to the variance of the metric across the group. This mean-centering only approach was used in other studies by members of this research team (20, 21). Short and long-term stability of peripheral blood gene expression was assessed by comparing variability between type-2 and ISG gene metrics (Supplementary Figure E5). The standard deviation of between-visit differences as a fraction of the standard deviation for all subjects and visits was comparable between the ISG metric (1-2 week interval in primary cohort SD_{diff}/SD_{all}=0.69; 48 week interval in placebo arm of severe asthma trial SD_{diff}/SD_{all}=1.02) and the eosinophil gene metric (1-2 week SD_{diff}/SD_{all}=0.84; 48 week SD_{diff}/SD_{all}=1.04).

Analysis of exacerbation rate reduction

Unadjusted exacerbation rates and 95% CI were computed with a zero-intercept generalized

linear model with a 4 level predictor (on or off treatment with high or low gene metric), quasipoisson error distribution, log link function, and an offset for the number of days in the study. Exacerbation rate reduction percentages and 95% CI were computed using the same model, but with a 2 level treatment variable as the predictor, with an intercept, and separate models for high and low gene metric groups. Statistical significance of the interaction effect between treatment and gene metric status was assessed with the same model with an intercept but with treatment, gene metric status, and the interaction of both as predictors.

Metatranscriptome preprocessing

To identify viral transcripts, raw sequencing reads underwent subtractive alignment to the human genome using the STAR aligner and the ENSEMBL GRCh38 genome build

(6). Remaining reads underwent complexity and quality filtering using PRICESeqFilter

(22) followed by additional subtractive alignment against non-microbial sequences using Bowtie2 (23). Identical reads were compressed using the CD-HIT-DUP algorithm (24)

and remaining sequences were aligned to the National Center for Biotechnology

Information (NCBI) nucleotide (nt) database using GSNAPL to identify viral transcripts

(25). Reads were filtered to include only those reads in which read concordance between paired-end reads was greater than 80%.

Metatranscriptome differential expression

Raw taxon read counts were aggregated to each taxon level using the taxize package in R

(26). Taxon read counts were further filtered to include only those taxons with an average

read count of 10 across samples and those taxons with at least 2 reads present in at least 25%

of samples. Taxon read counts were concatenated with human gene counts and underwent normalization and differential expression analysis using the DESeq2 package. This normalization method avoids introducing the bias imposed by using the total viral read count for sequencing depth correction.. Given that the viral and human reads are present in the same sample, we were able to leverage the more complete human transcriptome data and use the less biased size factor normalization technique provided by DESeq2 to normalize the viral transcripts for sequencing depth.

Study approval

NCT00595153 was approved by the University of California at San Francisco Committee on Human Research, written informed consent was obtained from all subjects, and all studies were performed in accordance with the principles expressed in the Declaration of Helsinki. NCT00314574 was conducted according to U.S. Food and Drug Administration regulations and Unsupervised clustering the International Conference on Harmonisation E6 Guidelines for Good Clinical Practice; all study sites obtained institutional review board approval and written informed consent from study participants before enrollment.

References

1. Bhakta NR, Solberg OD, Nguyen CP, Nguyen CN, Arron JR, Fahy JV, Woodruff PG. A qPCR-based metric of Th2 airway inflammation in asthma. *Clin Transl Allergy* 2013;3:24-7022-3-24.
2. Solberg OD, Ostrin EJ, Love MI, Peng JC, Bhakta NR, Hou L, Nguyen C, Solon M, Nguyen C, Barczak AJ, Zlock LT, Blagev DP, Finkbeiner WE, Ansel KM, Arron JR, Erle DJ, Woodruff PG. Airway Epithelial miRNA Expression is Altered in Asthma. *Am J Respir Crit Care Med* 2012.
3. Christenson SA, Steiling K, van den Berge M, Hijazi K, Hiemstra PS, Postma DS, Lenburg ME, Spira A, Woodruff PG. Asthma-COPD overlap. Clinical relevance of genomic signatures of type 2 inflammation in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2015;191:758-766.
4. Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads. *Bioinformatics in Action* 2012;17:10.
5. Joshi NA, Fass JN. Sickle: A sliding-window, adaptive, quality-based trimming tool for FastQ files. 2011;Version 1.33.
6. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* 2013;29:15-21.
7. Anders S, Pyl PT, Huber W. HTSeq--a Python framework to work with high-throughput sequencing data. *Bioinformatics* 2015;31:166-169.
8. Andrews S. FastQC: a quality control tool for high throughput sequence data. 2010.
9. Bonser LR, Zlock L, Finkbeiner W, Erle DJ. Epithelial tethering of MUC5AC-rich mucus impairs mucociliary transport in asthma. *J Clin Invest* 2016.
10. Woodruff PG, Boushey HA, Dolganov GM, Barker CS, Yang YH, Donnelly S, Ellwanger A, Sidhu SS, Dao-Pick TP, Pantoja C, Erle DJ, Yamamoto KR, Fahy JV. Genome-wide profiling identifies epithelial cell genes associated with asthma and with treatment response to corticosteroids. *Proc Natl Acad Sci U S A* 2007;104:15858-15863.
11. Johnson WE, Li C, Rabinovic A. Adjusting batch effects in microarray expression data using empirical Bayes methods. *Biostatistics* 2007;8:118-127.
12. R Core Team. R: A Language and Environment for Statistical Computing. 2015;Version 3.2.1.
13. Dolganov GM, Woodruff PG, Novikov AA, Zhang Y, Ferrando RE, Szubin R, Fahy JV. A novel method of gene transcript profiling in airway biopsy homogenates reveals increased expression of a Na⁺-K⁺-Cl⁻ cotransporter (NKCC1) in asthmatic subjects. *Genome Res* 2001;11:1473-1483.
14. Ali SA, Gern JE, Hartert TV, Edwards KM, Griffin MR, Miller EK, Gebretsadik T, Pappas T, Lee WM, Williams JV. Real-world comparison of two molecular methods for detection of respiratory viruses. *Virol J* 2011;8:332-422X-8-332.
15. Wesolowska-Andersen A, Everman JL, Davidson R, Rios C, Herrin R, Eng C, Janssen WJ, Liu AH, Oh SS, Kumar R, Fingerlin TE, Rodriguez-Santana J, Burchard EG, Seibold MA. Dual RNA-seq reveals viral infections in asthmatic children without respiratory illness which are associated with changes in the airway transcriptome. *Genome Biol* 2017;18:12-016-1140-8.
16. Anders S, McCarthy DJ, Chen Y, Okoniewski M, Smyth GK, Huber W, Robinson MD. Count-based differential expression analysis of RNA sequencing data using R and Bioconductor. *Nat Protoc* 2013;8:1765-1786.
17. Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, Smyth GK. limma powers

- differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Res* 2015;43:e47.
18. Benjamini Y, Hochberg Y. Controlling the false discovery rate: A practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society. Series B* 1995;57:289.
 19. Shen S, Park JW, Lu ZX, Lin L, Henry MD, Wu YN, Zhou Q, Xing Y. rMATS: robust and flexible detection of differential alternative splicing from replicate RNA-Seq data. *Proc Natl Acad Sci U S A* 2014;111:E5593-601.
 20. McBride JM, Jiang J, Abbas AR, Morimoto A, Li J, Maciucă R, Townsend M, Wallace DJ, Kennedy WP, Drappa J. Safety and pharmacodynamics of rontalizumab in patients with systemic lupus erythematosus: results of a phase I, placebo-controlled, double-blind, dose-escalation study. *Arthritis Rheum* 2012;64:3666-3676.
 21. Choy DF, Hart KM, Borthwick LA, Shikotra A, Nagarkar DR, Siddiqui S, Jia G, Ohri CM, Doran E, Vannella KM, Butler CA, Hargadon B, Sciurba JC, Gieseck RL, Thompson RW, White S, Abbas AR, Jackman J, Wu LC, Egen JG, Heaney LG, Ramalingam TR, Arron JR, Wynn TA, Bradding P. TH2 and TH17 inflammatory pathways are reciprocally regulated in asthma. *Sci Transl Med* 2015;7:301ra129.
 22. Ruby JG, Bellare P, Derisi JL. PRICE: software for the targeted assembly of components of (Meta) genomic sequence data. *G3 (Bethesda)* 2013;3:865-880.
 23. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. *Nat Methods* 2012;9:357-359.
 24. Fu L, Niu B, Zhu Z, Wu S, Li W. CD-HIT: accelerated for clustering the next-generation sequencing data. *Bioinformatics* 2012;28:3150-3152.
 25. Wu TD, Nacu S. Fast and SNP-tolerant detection of complex variants and splicing in short reads. *Bioinformatics* 2010;26:873-881.
 26. Chamberlain SA, Szocs E. taxize: taxonomic search and retrieval in R. *F1000Res* 2013;2:191-191.v2. eCollection 2013.

NOTE: Supplementary Table E1 is available as a separate file due to its length.

Supplemental Table E2: Upstream regulators in interferon pathways in epithelial brushings assayed by RNA sequencing: all gene symbols beginning with "IFNA", "IFNB", "IFNL", "IFNG", "IRF" as well as STAT1, shown. NA indicates expression level too low for DESeq2 to give a result

Ensembl ID	hgnc_symbol	log2FoldDifference	pvalue	padj
ENSG00000197919	IFNA1	-0.10	8.02E-01	NA
ENSG00000186803	IFNA10	0.77	2.55E-01	NA
ENSG00000233816	IFNA13	0.57	3.24E-01	NA
ENSG00000228083	IFNA14	0.68	3.07E-01	NA
ENSG00000147885	IFNA16	-0.81	1.97E-01	NA
ENSG00000234829	IFNA17	-0.02	9.70E-01	NA
ENSG00000188379	IFNA2	-0.18	6.56E-01	NA
ENSG00000137080	IFNA21	0.19	6.15E-01	NA
ENSG00000236637	IFNA4	-0.85	1.92E-01	NA
ENSG00000147873	IFNA5	-0.41	5.38E-01	NA
ENSG00000120235	IFNA6	-0.13	5.83E-01	NA
ENSG00000214042	IFNA7	0.44	5.11E-01	NA
ENSG00000120242	IFNA8	-0.50	4.48E-01	NA
ENSG00000142166	IFNAR1	-0.06	4.08E-01	5.78E-01
ENSG00000159110	IFNAR2	-0.05	5.04E-01	6.59E-01
ENSG00000171855	IFNB1	-0.43	2.58E-01	NA
ENSG00000111537	IFNG	0.89	8.58E-02	2.06E-01
ENSG00000027697	IFNGR1	0.06	6.25E-01	7.54E-01
ENSG00000159128	IFNGR2	0.16	9.39E-02	2.19E-01
ENSG00000262795	IFNGR2	NA	NA	NA
ENSG00000182393	IFNL1	-0.25	7.11E-01	NA
ENSG00000183709	IFNL2	-0.01	9.69E-01	NA
ENSG00000197110	IFNL3	NA	NA	NA
ENSG00000185436	IFNLR1	0.06	3.91E-01	5.62E-01
ENSG00000125347	IRF1	0.22	6.13E-02	1.64E-01
ENSG00000168310	IRF2	0.20	6.54E-03	3.89E-02
ENSG00000170604	IRF2BP1	0.34	1.92E-02	7.71E-02
ENSG00000168264	IRF2BP2	-0.10	2.69E-01	4.40E-01
ENSG00000119669	IRF2BPL	0.05	5.12E-01	6.67E-01
ENSG00000126456	IRF3	0.24	4.33E-02	1.30E-01
ENSG00000137265	IRF4	0.25	2.58E-01	4.28E-01
ENSG00000128604	IRF5	-0.02	8.89E-01	9.35E-01
ENSG00000117595	IRF6	0.24	1.06E-01	2.37E-01
ENSG00000185507	IRF7	0.58	1.41E-03	1.53E-02
ENSG00000140968	IRF8	-0.03	9.05E-01	9.45E-01
ENSG00000213928	IRF9	-0.01	9.69E-01	9.83E-01
ENSG00000124107	SLPI	0.37	2.84E-02	9.91E-02
ENSG00000115415	STAT1	0.28	1.40E-01	2.86E-01

Supplementary Table E3: IPA upstream regulator analysis of epithelial brushing genes associated with FEV₁ as a % of predicted.

Upstream Regulator	Activation z-score	p-value of overlap
IFNG	-7.864	3.49E-09
TNF	-7.36	1.57E-07
Interferon alpha	-6.696	1.98E-12
PRL	-5.956	6.27E-09
IFNA2	-5.883	2.30E-11
IRF7	-5.27	1.79E-07
IL2	-5.015	3.03E-07
IL1B	-4.772	2.12E-05
NFkB (complex)	-4.716	1.04E-03
IFNL1	-4.679	8.20E-10
STAT1	-4.45	2.02E-05
TNFSF11	-4.426	3.73E-02
IFN Beta	-4.339	1.57E-05
TLR7	-4.263	1.72E-03
IL15	-4.255	6.23E-05
TLR3	-4.192	2.15E-04
STAT4	-4.16	4.65E-03
Ifnar	-4.131	7.92E-06
TLR9	-4.126	2.69E-03
SMARCA4	-4.033	3.58E-07
Ifn	-4.016	6.60E-05
EGF	-4.016	4.51E-02
IRF3	-3.993	1.55E-05
RAF1	-3.957	6.29E-04
CD40LG	-3.93	1.85E-03
NFE2L2	-3.927	3.99E-03
IL4	-3.891	1.24E-06
IFNB1	-3.886	5.89E-05
PAF1	-3.873	2.62E-07
NFATC2	-3.849	1.96E-04
F2	-3.781	2.42E-02
IL12 (complex)	-3.685	4.09E-05
IRF1	-3.657	2.37E-04
SASH1	-3.606	5.63E-04
CSF2	-3.585	1.65E-02
XBP1	-3.499	3.33E-03
IRF5	-3.498	1.13E-04

Upstream Regulator	Activation z-score	p-value of overlap
IL6	-3.495	1.97E-04
IL27	-3.476	1.50E-03
CXCL12	-3.393	1.16E-04
P38 MAPK	-3.39	3.88E-02
DOCK8	-3.317	4.54E-03
ARNT	-3.274	3.26E-03
IL18	-3.235	4.07E-03
HGF	-3.216	5.75E-04
IL21	-3.195	1.63E-05
OSM	-3.193	1.13E-03
MAVS	-3.132	7.98E-04
SAMSN1	-3.051	5.62E-03
EIF2AK2	-3.023	1.29E-04
PDGF BB	-3.016	5.04E-03
IL7	-2.995	1.22E-03
TGM2	-2.992	1.64E-07
IL12 (family)	-2.942	9.04E-03
TNFSF10	-2.933	7.94E-07
TGFB1	-2.891	1.88E-02
TCR	-2.868	1.04E-02
TLR4	-2.83	1.14E-02
ARHGAP21	-2.828	5.72E-03
IL13	-2.821	2.29E-04
MYC	-2.82	3.69E-03
AGT	-2.795	1.40E-03
MAPK9	-2.744	2.08E-03
HIF1A	-2.734	2.23E-03
PRKCD	-2.71	4.92E-03
IL1A	-2.698	3.07E-02
AKT1	-2.682	1.34E-02
APP	-2.671	1.94E-02
IFN alpha/beta	-2.655	1.25E-02
Vegf	-2.645	3.90E-06
CHUK	-2.645	1.43E-02
MMP2	-2.626	7.44E-03
HRAS	-2.588	1.56E-08
IFNA1/IFNA13	-2.577	2.18E-03
TMEM173	-2.55	2.71E-04
MET	-2.525	2.83E-02
LCN2	-2.512	5.45E-03
MYD88	-2.485	2.38E-03
PGR	-2.446	1.76E-03
EP300	-2.432	2.15E-03

Upstream Regulator	Activation z-score	p-value of overlap
FLT3LG	-2.425	1.98E-02
IFNE	-2.415	2.66E-02
IPMK	-2.414	2.20E-03
Gm-csf	-2.4	4.90E-02
ESRRG	-2.398	3.38E-02
IFN type 1	-2.393	3.27E-03
IFIH1	-2.39	1.32E-02
IFNA4	-2.389	3.55E-03
ITGA1	-2.383	3.55E-03
JAK1	-2.376	2.06E-03
LCK	-2.366	2.54E-03
NFKBIA	-2.353	1.54E-02
CCL5	-2.351	1.71E-02
STAT5a/b	-2.343	5.62E-03
IFNAR1	-2.329	2.10E-06
EGR2	-2.306	8.30E-03
NLRC5	-2.277	1.87E-05
IFNAR2	-2.236	2.57E-06
ANXA2	-2.219	1.12E-02
IFNL3	-2.213	7.69E-03
PLIN5	-2.213	3.18E-02
TSH	-2.209	3.86E-02
THBS4	-2.2	2.76E-02
MAPK3	-2.19	4.22E-04
CCL3	-2.177	2.02E-02
SMAD7	-2.171	1.96E-02
IL17A	-2.149	1.67E-02
PCGEM1	-2.141	1.76E-04
DDX58	-2.119	1.08E-03
TRB	-2.08	1.54E-02
KITLG	-2.044	1.22E-02
SPI1	-2.031	1.51E-02
IKBKG	-2.027	3.14E-02
CXCL1	-2	5.67E-03
Srebp	-2	1.03E-02
hemoglobin	-2	2.57E-02
PARP9	-2	3.09E-02

Regulators filtered by: 1) genes, proteins or RNA; 2) absolute value of activation z-score $\geq$ 2; 3) p<0.05. List sorted by activation z-score, most negative (i.e. most inversely associated with FEV₁ as a % of predicted) to most positive. See Supplemental Methods.

Supplemental Table E4

Predictors in a linear model	Response: % change in FEV ₁ at stated weeks of ICS treatment relative to the baseline value		
	2 weeks of ICS	4 weeks of ICS	8 weeks of ICS
Type-2 airway gene metric	N=22	N=23	N=22
Pearson correlation	r=0.68,p=0.00044	r=0.86,p=2.1x10 ⁻⁶	r=0.47,p=0.029
Spearman correlation	ρ=0.68,p=0.00068	ρ=0.79,p=7.1x10 ⁻⁶	ρ=0.37,p=0.089
ISG airway gene metric	N=18	N=18	N=17
Pearson correlation	r=0.50,p=0.036	r=0.23,p=0.35	r=0.37,p=0.15
Spearman correlation	ρ=0.55,p=0.019	ρ=0.030,p=0.91	ρ=0.31,p=0.23
Type-2 and ISG airway gene metrics both in model	N=18 with both	N=18 with both	N=17 with both
ISG	p=0.019	p=0.44	p=0.16
Type-2	p=0.0010	p=0.00017	p=0.069
Type-2 and ISG airway gene metrics both in a model with an interaction term	N=18 with both	N=18 with both	N=17 with both
Interaction term	p=0.087	p=0.22	p=0.36

Greyed boxes indicate non-significant (i.e. $p \geq 0.05$) results

NOTE: Supplementary Table E5 is available as a separate file due to its length.

Supplemental Table E6: Correlation between whole blood gene expression metrics in healthy controls and subjects with mild asthma vs blood cell frequencies

Gene expression metric	Group	n	Cell type	Spearman ρ	Spearman P value
ISG	Healthy	20	neutrophil	0.37	0.11
ISG	Healthy	20	lymphocyte	-0.37	0.11
ISG	Healthy	20	eosinophil	-0.10	0.67
ISG	Healthy	20	monocyte	-0.14	0.57
ISG	Healthy	20	basophil	-0.45	0.047
ISG	Asthma	23	neutrophil	-0.23	0.29
ISG	Asthma	23	lymphocyte	0.05	0.82
ISG	Asthma	23	eosinophil	0.38	0.071
ISG	Asthma	23	monocyte	0.22	0.31
ISG	Asthma	23	basophil	-0.50	0.017
Eosinophil-associated genes	Healthy	20	neutrophil	0.04	0.87
Eosinophil-associated genes	Healthy	20	lymphocyte	-0.21	0.36
Eosinophil-associated genes	Healthy	20	eosinophil	0.73	0.00039
Eosinophil-associated genes	Healthy	20	monocyte	0.10	0.66
Eosinophil-associated genes	Healthy	20	basophil	0.34	0.14
Eosinophil-associated genes	Asthma	23	neutrophil	-0.05	0.81
Eosinophil-associated genes	Asthma	23	lymphocyte	-0.25	0.24
Eosinophil-associated genes	Asthma	23	eosinophil	0.85	0.0000019
Eosinophil-associated genes	Asthma	23	monocyte	-0.15	0.49
Eosinophil-associated genes	Asthma	23	basophil	0.17	0.44

Supplementary Table E7: Correlation coefficients comparing PAXgene whole blood microarrays to qPCR measurements

Gene	Pearson's r	Pearson p value	Spearman ρ	Spearman P value
CEACAM1	0.69737	2.03E-07	7.11E-01	3.50E-07
IFI16	0.70597	1.24E-07	7.25E-01	2.11E-07
IFI30	0.53922	1.91E-04	5.15E-01	4.91E-04
IFI44	-0.01462	9.27E-01	4.28E-01	5.04E-03
IFI6	0.86869	4.40E-14	8.74E-01	0.00E+00
IFIH1	0.64338	3.26E-06	4.96E-01	8.44E-04
IFIT1	0.83904	2.12E-12	7.62E-01	5.00E-08
IFIT5	0.61153	1.32E-05	4.59E-01	2.16E-03
IFITM1	0.44442	2.83E-03	4.12E-01	6.42E-03
ISG15	0.88308	4.66E-15	7.17E-01	2.81E-07
MT2A	0.77595	1.58E-09	8.00E-01	3.88E-09
MX1	0.86966	3.82E-14	7.03E-01	4.78E-07
MX2	0.60827	2.46E-05	6.30E-01	1.60E-05
TDRD7	0.40546	6.99E-03	1.79E-01	2.50E-01
TNFSF10	0.75016	7.00E-09	6.70E-01	1.90E-06
P2RY14	0.57722	5.07E-05	5.85E-01	5.22E-05
RNASE2	0.90759	0.00E+00	8.67E-01	0.00E+00
CLC	0.91603	0.00E+00	9.46E-01	0.00E+00
IL5RA	0.83316	4.17E-12	8.72E-01	0.00E+00

Correlations across the combined asthma and healthy data presented in Supplemental Figure 5A.

Supplementary Table E8: Baseline demographics of subjects from a randomized, placebo-controlled trial of the anti-IgE monoclonal antibody omalizumab in severe asthma (Hanania et al. 2011)

	Placebo	Omalizumab
	150	151
Age	47 ± 14	45 ± 14
% Female	71	63
BMI	32.4 ± 7.0	31.7 ± 7.9
FEV ₁ , % of predicted	64.9 ± 13.6 (n=149)	65.0 14.5 (n=150)
Blood eosinophils (K/μl)	0.26 (0.02-1.99) (n=142)	0.26 (0.03-2.29) (n=146)

Data reported as mean±SD or median (range).

Supplementary Table E9: Unfolded Protein Response Activation in IL-13 and IFN- α Stimulated epithelial air-liquid interface cultures. Selected output from IPA upstream regulator analysis of RNA Seq-based gene expression.

	IREα/XBP1 Pathway	PERK Pathway	ATF6 Pathway
IL-13 stimulation	XBP1: activation z-score 5.40 overlap p-value 0.00040 IREα: activation z-score 2.60 overlap p-value 0.0011	PERK: activation z-score 1.58 overlap p-value 0.00083 CHOP: activation z-score 1.26 overlap p-value 0.012 ATF4: activation z-score 1.96 overlap p-value 0.0020	ATF6: activation z-score 2.55 overlap p-value 0.00033
IFN-α stimulation	XBP1: Not activated IREα: activation z-score 2.17 overlap p-value 0.00011	PERK: activation z-score 2.81 overlap p-value 1.42×10^{-6} CHOP: activation z-score 2.29 overlap p-value 0.00037 ATF4: activation z-score 3.40 overlap p-value 0.030	ATF6: Not activated

The overlap p-value is the probability that the observed overlap between the list of genes differentially expressed between cytokine and control, and the set of genes assigned by IPA as downstream of the listed upstream regulator, would occur by chance (Fisher's Exact Test). The strength of activity of the regulator under cytokine stimulation, as determined by the net observed direction of changes in the overlapping downstream genes, is represented by an activation score (a more positive (or negative) value indicates that a larger proportion of genes changed in a direction consistent with activation (inhibition) of the regulator with cytokine stimulation).

Supplementary Table E10: Primers and probes used in qPCR

Gene	Type	Sequence
CEACAM1	outer-forward	CATTCACTTTGTCAGGAAACCTTG
	outer-reverse	TGCAGCTTTCTAGCTGTAAACAATG
	inner-forward	CCTTGGCCTCTGCTAAGGTG
	inner-reverse	AACAATGCCACCAGCATGAG
	probe	CCCTGTAGGGTGCTCCCACTTCTCAA
CLC	outer-forward	GCTGCCTCTTTGTCTACTGGTTCT
	outer-reverse	CTGAAAGGGCATATTCTTGGATTC
	inner-forward	CTGAGATGAAGGAGGAATCAGACA
	inner-reverse	CCATACTCACGGCTGTTTCATGA
	probe	TCTTCCATTTCCAAGTGTGCTTTGGTCGT
GAPDH	outer-forward	CAATGACCCCTTCATTGACCTC
	outer-reverse	CTCGCTCCTGGAAGATGGTGAT
	inner-forward	GATTCCACCCATGGCAAATTC
	inner-reverse	GGGATTTCCATTGATGACAAGC
	probe	CGTTCTCAGCCTTGACGGTGCCA
IFI16	outer-forward	TGAGGATGCAGATACTGAAGGAAG
	outer-reverse	TCTTGGTTTCAACGTGGTTAAGAA
	inner-forward	AGCATAGGCCCAGCTGAGAG
	inner-reverse	TGGTTAAGAACTGCTGCTTGG
	probe	CCCACACTCCTCAGATGCCTCCATC
IFI30	outer-forward	TGTGACCCTCTACTATGAAGCACTG
	outer-reverse	TGTTGAATTTGCACTCCTCTTCTC
	inner-forward	AACATGGCTGTTGGTCATGG
	inner-reverse	TGTTCTGTGCGTTTCCGTA
	probe	CCTCAATGTCACGCTGGTGCCC
IFI44	outer-forward	TCTGCCTTGAGAACTTATGAACCA
	outer-reverse	CAGTTGTATTAGTGCCACCAAAG
	inner-forward	CCATATGGATCCCTGGTTCA
	inner-reverse	CTGGACTTCCCAGCTCCAAT
	probe	CAAATACGAATTCTGCTGCTGGGTCCA
IFI6	outer-forward	CATAGTACACTGCAGCCTCCAACT
	outer-reverse	GGGAGAGTGATAGACAAAGTTCTGG
	inner-forward	GCCTCCAACTCCTAGCCTCA


	inner-reverse probe	TTCTGGATTCTGGGCATCGT CCAAGTAGGATTACAAGCATGCGCCG
IFIH1	outer-forward outer-reverse inner-forward inner-reverse probe	CGGTAGACCCTGCTTCTCTAAGT AGAAAGGTCAGGTAGTCCAGCAC AATGGGTATTCCACAGACGAGAA CTCCACCTGGATGTACATTTTCAC TTCCGCTATCTCATCTCGTGCTTCAGG
IFIT1	outer-forward outer-reverse inner-forward inner-reverse probe	GTAATACCCTCCCTTCTCCAAGA AGGCTGCTCTCGAACTCCTG GGCTCACGCCTGTAATCTCA TCGAACTCCTGGGTTCAGGT ATCTGCCTGCCTCGTCCCATCG
IFIT5	outer-forward outer-reverse inner-forward inner-reverse probe	GTGCTGAGGAGAGAGCGATCC TGACATTCTAACTCCAACAGAATGG AGAGAGCGATCCGAGGGACT GAATGGCCTTCAAGGTGTCC CTCATGATGGCAGCGCTCTGCA
IFITM1	outer-forward outer-reverse inner-forward inner-reverse probe	AACCTTCACTCAACACTTCTTCC GATGTTGATCACGGTGGACCT CCAAAGCCAGAAGATGCACA ACCTTGGAAGGATGGTGCTG CCCAGCACAGCCACCTCATGTTCC
IL5RA	outer-forward outer-reverse inner-forward inner-reverse probe	ATGGCCACCGCATTTCTCA CCTCTTGTTCCGACCAGTACTCAA GCCAGGCACATTGAGCATT GTCTATGCTCGTGGCTGCAA TCCTGTGCCTGACGCTATGCTAGATGC
ISG15	outer-forward outer-reverse inner-forward inner-reverse probe	ATCTTTGCCAGTACAGGAGCTTG CGCCTTCAGCTCTGACACC ACCTGACGGTGAAGATGCTG GCTCTGACACCGACATGGAG ACACCTGGAATTCGTTGCCCGC
MT2A	outer-forward outer-reverse inner-forward	GTGCAACCTGTCCCGACTCTA TTTCTTGACAGGAGGTGCATTT TCTAGCCGCCTCTTCAGCTC

	inner-reverse probe	GGTGCATTTGCACTCTTTGC CCATGGATCCCAACTGCTCCTGC
MX1	outer-forward outer-reverse inner-forward inner-reverse probe	GGACAAAGGAACTGAAGACAAGGT TGAAATATGGGTGGTTCTCAAAGA CCTCGTGTTCCACCTGAAGA CTGGTCCTGGATCTCCTGCT CCCGGCACTTGACAATCATGTAACCC
MX2	outer-forward outer-reverse inner-forward inner-reverse probe	ACCAGAGCTCGGGCAAGA GTGTTCCGGTAGCTGATCCTTC CTCTGTGCTGGAGGCACTGT GCGGACACCTGGTTACGATT CCTCTGGGAAGCGCGACTCCTG
P2RY14	outer-forward outer-reverse inner-forward inner-reverse probe	GAGTCAGACCGAAGCTCATTACAG CCCTAAACGGCTGGCATAGA TCACTCTGCTACTATCTGCTGCAA CCCTAAACGGCTGGCATAGA TGCTTGGACCCTATTATTTATTCT
PPIA	outer-forward outer-reverse inner-forward inner-reverse probe	ATGAGAACTTCATCCTAAAGCATACG TTGGCAGTGCAGATGAAAACT ACGGGTCCTGGCATCTTGT GCAGATGAAAACTGGGAACCA ATGGCAAATGCTGGACCCAACACA
RNASE2	outer-forward outer-reverse inner-forward inner-reverse probe	GGAGCTACAGGCCGGAGACT AGCCCTCCACAGCCAGAAG CGGAGACTGGGAAACATGGT AAGCCCCAACAGAAGAAGCA CCAAAAGTGTCACTTCCCAAATTT
RPL37A	outer-forward outer-reverse inner-forward inner-reverse probe	TATGTGAAATAATCCGGACATGCT TGCAATTGTAAGGCGTCTCTCT CGGACATGCTTCTGCTATGG GGCGTCTCTCTCATCCAGGT CCCAGATTCGCAAACCTATGGCGACA
TDRD7	outer-forward outer-reverse inner-forward	AGAGATCACAAATGTTACAGCAG TTCTTGTAGAAACCGAGGTGGAAT CCGGGCTCTTGATGTTCACT


TNFSF10	inner-reverse	CGAGGTGGAATTTCCCTGAG
	probe	TCCTGGACTCTGGCACTGTGACATCTG
	outer-forward	GAAGCAACACATTGTCTTCTCCAA
	outer-reverse	GACCAGTTCACCATTCCTCAAGT
	inner-forward	GCTCTGGGCCGCAAAAT
	inner-reverse	AGTGCAAGTTGCTCAGGAATGA
	probe	TGCCCACTCCTTGATGATTCCCAGG

Probes used a 5' FAM fluorescent probe with a 3' BHQ quencher.

A.




B.




Supplemental Figure E1: Ranking of genes differentially expressed by RNA Seq in the current study within microarray data from both a prior study and the current study. The 1379 genes with $FDR < 0.01$ in the asthma (not on ICS) vs healthy differential expression analysis of the RNA Seq data were used as the test set. Barcode plots from limma package in R for enrichment of RNA Seq gene test set within differential expression results from A. previously published microarray data from epithelial brushings from asthmatics and healthy controls (Woodruff et al. PNAS 2007, available on GEO as GSE4302), and B. microarray data from epithelial brushings from the same asthmatics ($n=19$) and healthy controls ($n=16$) in the current study population (available as GSE67472 for a slightly larger group of subjects, and used for a gene set list derivation in Christenson et al. Asthma-COPD overlap. Clinical relevance of genomic signatures of type 2 inflammation in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med.* 2015 Apr 1;191(7):758-66).


A.




B.





Supplemental Figure E2: Comparison of gene expression profiles in airway epithelial brushings between RNA sequencing and microarrays for interferon-stimulated genes shows overall agreement. Microarray gene expression profiles were obtained on the same samples presented in this study (19 subjects with asthma, 16 healthy controls; available for a slightly larger group of subjects as GSE67472, and used for a gene set list derivation in Christenson et al. Asthma-COPD overlap. Clinical relevance of genomic signatures of type 2 inflammation in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med.* 2015 Apr 1;191(7):758-66). Genes in the interferon alpha group per IPA that were both differentially expressed by RNA Seq in asthma relative to health and with direction of difference consistent with activation of this pathway in asthma were analyzed. Of the 27 genes that met these criteria, 25 were annotated on the arrays and shown here. Of these 25 genes with FDR<0.01 by RNA Seq, 10 (40%) have FDR<0.01 on microarrays. A. Log₂ fold-difference between asthma and health is generally consistent between RNA Seq (black bars) and microarrays (grey bars). B. Average expression across all samples for RNA sequencing (top black bars) as assessed by baseMean from DESeq2, and for microarrays (bottom grey bars) as assessed by AveExpr from limma. Genes with discordant statistical significance between platforms do not necessarily all have low average expression. Red outlines indicate FDR<0.01, blue FDR>=0.01. FDR for microarray data computed using limma with Benjamini-Hochberg correction and no clinical covariates.



Supplemental Figure E3: Effect of corticosteroids on airway and blood ISG and ER stress metrics and lung function. A. Airway ISG metric for healthy controls (n=16) and subjects with asthma (n=19) for which there is paired data before and after 8 weeks of ICS (n=9); $p=0.012$ for the effect of ICS by a paired Mann-Whitney U (MWU) test). B. and C. Relationship of airway ISG metric from brushings acquired at 8 weeks to: B. FEV₁ as a percent of the predicted value after 8 weeks of ICS, measured immediately before and on the day of bronchoscopy and prior to bronchodilator (Spearman $\rho=0$, $p=1$, $n=13$ asthma after ICS) and C. change in FEV₁ as a percent of predicted after administration of bronchodilator (Spearman $\rho=0.039$, $p=0.90$, $n=13$ asthma after ICS). D. Receiver operating characteristic (ROC) curves for the ability of the airway ISG metric (n=17) and markers of type-2 inflammation (blood eosinophils (n=23) and airway type-2 gene expression metric (n=22)) to predict an improvement in FEV₁ after 8 weeks of ICS. Improvement in FEV₁ was defined as an increase of at least 200 mL and 12% relative to week 0. ROC curves for type-2 gene expression metric and airway ISG metric were not significantly different ($p=0.31$ by DeLong's test for two correlated ROC curves from the pROC package in R); however, the estimated power for detecting a significant difference between these two observed curves is only 30%. E. Effect of ICS on blood ISG metric in mild asthma ($p=0.012$ for $n=20$ healthy vs $n=21$ asthma pre-ICS by MWU test); $p=0.45$ paired MWU on $n=21$ asthma pre and post ICS). Effect of oral corticosteroid use on baseline blood ISG metric in severe asthma ($p=0.074$, $n=301$, MWU test) F. Airway ER stress (XBP1) metric for the same subjects in A ($p=0.00034$ for the effect of ICS in mild asthma by a paired t test).




Supplemental Figure E4: Mapping of viral reads from one asthma subject to genome of human rhinovirus A. Sequences were assembled against the human rhinovirus 22 strain ATCC VR-1132 (version FJ445122.1 GI:2173163) reference genome using the *Geneious* software package for the one asthma subject in which rhinovirus was detected by sequencing.




Supplemental Figure E5: Additional data for whole blood gene expression. Whole blood microarray expression levels of eosinophil-associated genes are significantly associated with **(A)** blood eosinophil percentages across subjects with asthma (red; n=23; all Pearson correlation coefficients with p<0.05 except for RNASE2) and healthy controls (black; n=20; all Pearson correlation coefficients with p<0.05 except RNASE2 and P2RY14), but unrelated to **(B)** blood neutrophil percentages (all Pearson correlation coefficients with p>0.05 for asthma and healthy).

A.




Supplemental Figure E6: Proportion of five cell types obtained from a standard automated clinical laboratory cell differential in healthy controls (H, n=20) vs asthmatics not on ICS (A, n=23) on the day of and immediately before bronchoscopy.

A.



B.



Supplemental Figure E7: qPCR validation of PAXgene whole blood microarrays. A. Cross-platform gene expression correlations, log-transformed, for 15 ISGs for which primers worked in multiplex (from genes marked as downstream of interferons per Ingenuity Pathway Analysis). Data for four eosinophil-associated genes for which primers worked is also shown (outlined by red box). Correlation coefficients are presented in Supplemental Table 4. **B.** Log₂ fold-change between asthma and healthy is generally consistent between microarrays (black bars) and qPCR (grey bars).