

Characterising the allergic profile of children with cystic fibrosis

Amy L. Faulkner¹  | Michael Grayling²  | Benjamin Shillitoe³  |
Malcolm Brodrie^{4,5}  | Louise J. Michaelis^{2,3} 

¹Faculty of Medical Sciences, Newcastle University, Newcastle upon Tyne, UK

²Population Health Sciences Institute, Faculty of Medical Sciences, Newcastle University, Newcastle upon Tyne, UK

³Department of Immunology, Allergy, and Infectious Diseases, Great North Children's Hospital, Newcastle upon Tyne Hospitals NHS Foundation Trust, Newcastle upon Tyne, UK

⁴Translational and Clinical Research Institute, Faculty of Medical Sciences, Newcastle University, Newcastle upon Tyne, UK

⁵Paediatric Respiratory Medicine, Great North Children's Hospital, Newcastle upon Tyne Hospitals NHS Foundation Trust, Newcastle upon Tyne, UK

Correspondence

Malcolm Brodrie, Paediatric Respiratory Medicine, Great North Children's Hospital, Newcastle upon Tyne Hospitals NHS Foundation Trust, Queen Victoria Road, Newcastle upon Tyne, NE1 4LP, UK.

Email: malcolm.brodrie@ncl.ac.uk

Abstract

Background: Cystic fibrosis (CF) is a genetic condition that affects multiple organ systems. Allergic bronchopulmonary aspergillosis (ABPA) is a well-recognised problem but other allergic conditions are less well documented in CF.

Objective: To characterise the allergic profile of a cohort of children with CF, with a focus on those with ABPA.

Methods: A cohort of children with CF were interviewed and retrospective data were collected regarding their allergic histories and other relevant clinical features.

Results: The cohort included 37 children with median age of 9 years (inter-quartile range: 6-12). There was a history of ≥ 1 allergic condition(s) in 28/37 children (76%). The most common allergic condition was allergic rhinitis (AR) in 21/37 (57%) and 16 of these 21 children (76%) had another allergic condition. All children with ABPA (8) had another allergic condition. In some children ABPA exacerbations appeared to be seasonal, suggesting possible cross-sensitisation between *Aspergillus fumigatus* and aeroallergens associated with seasonal AR. Allergic conditions were also common in children with *Pseudomonas aeruginosa* infection.

KEYWORDS

allergic rhinitis, *Aspergillus*, cystic fibrosis, hypersensitivity

1 | INTRODUCTION

Cystic fibrosis (CF) is an autosomal recessive genetic condition with a prevalence of around 1:2000 in the United Kingdom (UK).¹ CF affects many organs but most notably causes problems in the respiratory system, with disordered

airway surface liquid homeostasis occurring as a result of defective CF transmembrane conductance regulator function.² This causes impaired mucus clearance, susceptibility to infection and neutrophilic inflammation.^{1,3} *Pseudomonas aeruginosa* and *Aspergillus fumigatus* are amongst key pathogens that drive lung pathology.³⁻⁵

Malcolm Brodrie and Louise J Michaelis have equal contribution.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2021 The Authors. *Immunity, Inflammation and Disease* published by John Wiley & Sons Ltd.

Allergic bronchopulmonary aspergillosis (ABPA) is an allergic condition, which is rare in the general population, but has a prevalence of around 9% in people with CF.⁶ ABPA results from sensitisation to *A. fumigatus* and is characterised by peripheral eosinophilia, elevated serum immunoglobulin E (IgE), episodic bronchial obstruction and lung infiltrates with or without bronchiectasis.⁵ Severe episodes of ABPA can be life-threatening and require admission to intensive care.⁵ The reasons for the association between CF and ABPA are not well understood.

The pathogenesis of allergic disease is recognised to be multifactorial. According to several well accepted hypotheses, certain allergic conditions are on causative pathways for others, meaning that they tend to accumulate.^{7–10} The classical order of allergy progression during childhood is known as the “Atopic March”,^{11–13} and follows the sequence of: eczema, food allergy, allergic rhinitis (AR) and asthma. Estimated prevalence of these conditions in children in the United Kingdom, as self-reported in population level data, are eczema 11%–20%, food allergy 8%, AR 10%–15% and asthma 10%.^{14–17}

It is known that ABPA and drug allergy are common in CF,¹⁸ but besides this, literature regarding the profile of allergy in CF and possible mechanisms involved is limited. One area that has been researched is the role of the gut microbiome in the pathogenesis of ABPA. It is thought that gut dysbiosis (which is common in CF),^{19,20} might lead to dysregulation of the mycobiome, expansion of *A. fumigatus* and predispose individuals to airway sensitisation.²¹ Other possible factors include prolonged exposure to antibiotics, and a susceptibility to lung infection with *P. aeruginosa*, which are both associated with the development of allergy.^{22,23}

We undertook a hypothesis-generating single centre pilot study with the aim of increasing understanding of the allergic profile of children with CF and to explore factors that may contribute to the development of ABPA.

2 | METHODS

2.1 | Study design

A retrospective pilot study examining the allergic profile of a cohort of children with CF was carried out between February and July 2020. Participants were interviewed as part of routine clinic review. The sole inclusion criterion was that they had a clear diagnosis of CF.

2.2 | Data collection

A modified version of the validated “International Study of Asthma and Allergies in Childhood” questionnaire²⁴ was used to collect data regarding participants’ allergy-focused history and associated factors (Appendix 2). Results of allergic, immunological and nutritional investigations were collected from participants’ electronic medical records. Where data from the date of interview was not available the most recent results were recorded.

2.3 | Data and statistical analysis

Analyses were largely descriptive regarding patterns of allergy in the sample population. Allergic outcomes were defined as presence versus absence of allergic conditions.

Summary statistics were computed using the Statistical Package for the Social Sciences (SPSS).²⁵ Comparisons of proportions were performed in R,²⁶ using the Berger and Boos (1994) method via the Exact package.²⁷ Comparison of percentage predicted forced expiratory volume in 1 s (%FEV1) was performed using the Mann–Whitney *U* test in R.²⁶ In relation to possible associations between *P. aeruginosa* and allergy and median age at interview the Kruskal–Wallis and Fischer’s exact statistical tests were performed using SPSS.²⁵ A *p* value <.05 was considered statistically significant throughout. All data were handled in accordance with the General Data Protection Regulations.

3 | RESULTS

3.1 | Sample population demographics

In total, 37 participants were sampled as they attended outpatient clinics, (from the GNCH CF clinic of 176 children), with a median age of 9 years (interquartile range: 6–12). Other key demographics of the population and study cohort are provided in Table 1.

3.2 | Allergic profile of the cohort

In the cohort, 28/37 (75.7%) children reported history of ≥ 1 of the five main allergic conditions. Allergic comorbidity (≥ 2 of the main allergic conditions), was present in around half of the cohort, including all children with food allergy or asthma. Eight children had ≥ 3 allergic conditions (all including eczema). Furthermore, 5/6 with food allergy had a history of eczema. All with asthma had AR. None had all five allergic conditions.

TABLE 1 Demographics of the study cohort

	Age in years Median (IQR)	Male gender	White ethnicity	CFTR Genotype			%FEV1
				F508del/ F508del	F508del/ Other	Other/ Other	
Sample population [<i>n</i> = 176]	8.0 (5.0–12.0) ^a	92 (52.3%)	158 (89.8%)	86 (48.9%)	75 (42.6%)	15 (8.5%)	93.7 (83.4–103.8) [<i>n</i> = 110] <i>At most recent annual review</i>
Study cohort [<i>n</i> = 37]	9.0 (6.0–12.0)	22 (59.5%)	37 (100%)	23 (62.2%)	13 (35.1%)	1 (2.7%)	95.5 (85.2–103.5) [<i>n</i> = 33]
Eczema [<i>n</i> = 18]	8.5 (6.0–11.0)	12 (66.7%)	18 (100%)	11 (61.1%)	6 (33.3%)	1 (5.6%)	95.5 (78.2–98.8) [<i>n</i> = 15]
Seasonal Allergic Rhinitis [<i>n</i> = 13]	11.0 (10.0–13.0)	7 (53.8%)	13 (100%)	9 (69.2%)	3 (23.1%)	1 (7.7%)	85.2 (74.3–98.5) [<i>n</i> = 13]
Allergic Bronchopulmonary Aspergillosis [<i>n</i> = 8]	14.5 (11.8–16.0)	3 (37.5%)	8 (100.0%)	5 (62.5%)	2 (25.0%)	1 (12.5%)	81.4 (75.9–86.7)
Perennial Allergic Rhinitis [<i>n</i> = 6]	8.0 (6.3–9.0)	4 (66.7%)	6 (100%)	3 (50.0%)	3 (50.0%)	0 (0.0%)	102.3 (91.1–111.4) [<i>n</i> = 6]
Perennial Allergic Rhinitis with seasonal exacerbation [<i>n</i> = 2]	13.5 (12.3–14.8)	1 (50.0%)	2 (100%)	1 (50.0%)	1 (50.0%)	0 (0.0%)	86.0 (81.0–91.0) [<i>n</i> = 2]
Drug allergy [<i>n</i> = 7]	15.0 (12.0–16.0)	3 (42.9%)	7 (100%)	6 (85.7%)	1 (14.3%)	0 (0.0%)	86.2 (63.5–91.8) [<i>n</i> = 7]
Food allergy [<i>n</i> = 6]	9.0 (4.0–11.0)	4 (66.7%)	6 (100%)	2 (33.3%)	4 (66.7%)	0 (0.0%)	95.8 (94.0–97.1) [<i>n</i> = 4]
Asthma [<i>n</i> = 3]	11.0 (9.0–11.5)	2 (66.7%)	3 (100%)	1 (33.3%)	2 (66.7%)	0 (0.0%)	96.0 (90.6–102.7) [<i>n</i> = 3]
No history of allergic conditions [<i>n</i> = 9]	8.0 (5.0–9.0)	5 (55.6%)	9 (100%)	6 (66.7%)	3 (33.3%)	0 (0.0%)	104.2 (98.0–105.9) [<i>n</i> = 8]

Abbreviations: CFTR, cystic fibrosis transmembrane conductance regulator; %FEV1, percentage of predicted forced expiratory volume in 1 s.

^aAge as of 08/03/2021.

AR was the most commonly reported condition (Figure 1). All children with AR who had specific IgE testing had sensitivity to >1 aeroallergen. This included half being sensitised to *A. fumigatus*. One child had a history of anaphylaxis (related to food allergy) and two had experienced allergy-associated breathing difficulties (related to drug allergy). Of those with eczema or AR 88% used daily treatment regimens during symptomatic periods—91% with AR used oral antihistamines, 29% with AR used corticosteroid nasal sprays and two-thirds with eczema used topical steroid creams. None with eczema, asthma or AR used oral corticosteroids or other immunosuppressants.

Median %FEV₁ varied between allergic conditions. Those with seasonal allergic rhinitis (SAR) and perennial allergic rhinitis with seasonal exacerbation (PARwE) had the worst lung function and those with perennial allergic rhinitis (PAR) had the best (Table 1).

In terms of median age at time of interview, children with each of the allergic conditions, in ascending order, was: eczema 110 months ($n=18$), food allergy 113 months ($n=21$), AR 133 months ($n=21$), asthma 138 months ($n=3$) and drug allergy 184 months ($n=7$). The median ages of each subgroup were significantly different according to the Kruskal–Wallis test ($p=.004$).


3.3 | CF-specific features of the cohort

Based on the existing literature, three prominent features of CF were identified as being of interest in terms of possible association with atopic disease:

- Prolonged exposure to antibiotics: all were on daily flucloxacillin prophylaxis.
- Upper airway pathology: 16/21 (76.2%) of those with AR had allergic comorbidity. Upper airway pathology was also frequent in the ABPA subgroup: half had SAR, 12.5% had PAR and 12.5% had PARwE (Table 2).
- Respiratory infection with *P. aeruginosa*: present in 17/37 (49.4%) were and 14/17 of these children had allergy compared to 12/20 who were not infected. The difference in prevalence of AR between those with *P. aeruginosa* and those not (13/17 vs. 8/20) was statistically significant according to the Chi-square test ($p=.04$).

3.4 | Analysis of children with ABPA

Overall, 8/37 (21.6%) of the cohort had ABPA (diagnostic criteria in Appendix 1) and the median age of this



Abbreviations: IgE – immunoglobulin E, PAR – perennial allergic rhinitis, PARwE – perennial allergic rhinitis with seasonal exacerbation, SAR – seasonal allergic rhinitis.

FIGURE 1 Distribution of the main five allergic conditions in the study cohort. IgE, immunoglobulin E; PAR, perennial allergic rhinitis; PARwe, perennial allergic rhinitis with seasonal exacerbation; SAR, seasonal allergic rhinitis

TABLE 2 Demographics and allergic profile of the ABPA-positive versus ABPA-negative subgroups in the study cohort

	Whole cohort <i>n</i> = 37 (%)	ABPA subgroup <i>n</i> = 8 (%)	Non-ABPA subgroup <i>n</i> = 29 (%)	<i>p</i> value comparing subgroups
Demographics				
Age in years	9.0 (6.0–12.0)	14.5 (11.8–16.0)	8.0 (5.0–11.0)	<i>p</i> = .0005 ^a
Gender male	22 (59.5)	3 (8.1)	19 (51.4)	<i>p</i> = .140 ^b
Reported allergy	28 (75.7)	8 (21.6)	20 (54.1)	<i>p</i> = .084 ^b
Allergic rhinitis (any)	21 (56.8)	6 (16.2)	15 (40.5)	<i>p</i> = .288 ^b
Seasonal allergic rhinitis	13 (35.1)	4 (10.8)	9 (24.3)	<i>p</i> = .271 ^b
Perennial allergic rhinitis	6 (16.2)	1 (2.7)	5 (13.5)	<i>p</i> = .936 ^b
Perennial allergic rhinitis	2 (5.4)	1 (2.7)	1 (2.7)	<i>p</i> = .223 ^b
Drug allergy	7 (18.9)	4 (10.8)	3 (8.1)	<i>p</i> = .014 ^b
Eczema	18 (48.6)	2 (5.4)	16 (43.2)	<i>p</i> = .166 ^b
Food allergy	6 (16.2)	0 (0.0)	6 (16.2)	<i>p</i> = .263 ^b
Atopic asthma	3 (8.1)	1 (2.7)	2 (5.4)	<i>p</i> = .376 ^b
Results				
%FEV1 (at time of interview)	95.8 (84.4–103.8) [<i>n</i> = 34]	77.5 (75.7–85.7) [<i>n</i> = 7]	97.7 (89.6–104.8) [<i>n</i> = 27]	<i>p</i> = .032 ^a
Specific IgE (KAU/L)				
<i>Aspergillus fumigatus</i>	12 (42.9) [<i>n</i> = 28]	8 (28.6) [<i>n</i> = 8]	4 (14.3) [<i>n</i> = 20]	<i>p</i> = .00005 ^b
Grass Mix 1	4 (80.0) [<i>n</i> = 5]	2 (80.0) [<i>n</i> = 3]	2 (40.0) [<i>n</i> = 2]	<i>p</i> = .568 ^b
House Dust Mite	2 (66.7) [<i>n</i> = 3]	1 (66.7) [<i>n</i> = 2]	1 (33.3) [<i>n</i> = 1]	<i>p</i> = .750 ^b
Tree Mix 1	0 (0.0) [<i>n</i> = 4]	0 (0.0) [<i>n</i> = 3]	0 (0.0) [<i>n</i> = 1]	<i>p</i> = 1.000 ^b
Weed Mix 1	0 (0.0) [<i>n</i> = 3]	0 (0.0) [<i>n</i> = 2]	0 (0.0) [<i>n</i> = 1]	<i>p</i> = 1.000 ^b
Cat dander	2 (66.7) [<i>n</i> = 3]	0 (0.0) [<i>n</i> = 1]	2 (66.7) [<i>n</i> = 2]	<i>p</i> = .250 ^b
Dog dander	1 (50.0) [<i>n</i> = 2]	0 (0.0) [<i>n</i> = 1]	1 (50.0) [<i>n</i> = 1]	<i>p</i> = .500 ^b
Moulds Mix	1 (100.0) [<i>n</i> = 1]	1 (100.0) [<i>n</i> = 1]	0 (0.0) [<i>n</i> = 0]	<i>P</i> = N/A
Risk factors				
Prolonged antibiotic exposure	37 (100.0)	8 (21.6%)	29 (78.4)	<i>p</i> = 1.000 ^b
Chronic infection with <i>Pseudomonas aeruginosa</i>	17 (45.9)	6 (16.2)	11 (29.8)	<i>p</i> = .082 ^b

Abbreviations: ABPA, allergic bronchopulmonary aspergillosis; %FEV1, percentage of predicted forced expiratory volume in 1 s; IQR, interquartile range; SpIgE, specific immunoglobulin E.

^aStatistical test used: Mann–Whitney *U*.

^bStatistical test used: Berger and Boos. A *p* value of <.05 considered statistically significant.

subgroup was 9 years (Table 2). Median %FEV1 was 20.2% points lower in those with ABPA compared to those without. Other co-existent allergic conditions included AR (75%) and drug allergy (50%). The difference in prevalence of drug allergy between the whole cohort and the ABPA subgroup was statistically significant according to the Berger and Boos test (*p* = .014).

Having been identified as a subgroup of interest, retrospective allergic timelines for six children with ABPA from the cohort were compiled (Figure 2). These comprised serial total IgE (TIgE), eosinophil, basophil and lymphocyte counts and dates of hospital admissions due to ABPA exacerbations. Timeline length depended on availability of data and spanned between 6 and 10 years per child.


Peak silver birch pollen months: February to April⁴⁷

Peak tree pollen months: April to June⁴⁷

Peak grass pollen months: June to August⁴⁷


Peak House dust mite months: Winter with a peak in Autumn⁴⁸

Abbreviations: ABPA – allergic bronchopulmonary aspergillosis, HDM – House Dust Mite, IgE – immunoglobulin E, SpIgE – specific immunoglobulin E




Exacerbation 1: coincides with beginning of house dust mite season (SpIgE confirmed sensitivity)

Exacerbation 2: coincides with tree pollen season (allergy testing not carried out)



Exacerbation 1: coincides with house dust mite season (negative skin prick testing).



Query exacerbation: coincides with house dust mite season (allergy testing not carried out)

Exacerbation 1: coincides with grass pollen season (allergy testing not carried out)

Exacerbation 2: coincides with house dust mite season (allergy testing not carried out)

FIGURE 2 Allergic timelines for six children with allergic bronchopulmonary aspergillosis in the study cohort. In the United Kingdom peak months for silver birch pollen are February to April, tree pollen April to June, grass pollen June to August and house dust mite in Autumn and Winter. ABPA, allergic bronchopulmonary aspergillosis; HDM, house dust mite; IgE, immunoglobulin E, SpIgE, specific immunoglobulin E.




FIGURE 2 Continued

TiGE was raised during all ABPA exacerbations. Eosinophil, basophil and lymphocyte counts were measured during hospital admission for eight exacerbations. Eosinophil count was above normal in 7/8 exacerbations and basophil count in 6/8 (normal ranges in Appendix 1). ABPA exacerbations sometimes coincided with the peak months of certain aeroallergens (Figure 2).

4 | DISCUSSION

This pragmatic study of the allergic profile of a cohort of children with CF has produced several interesting but preliminary findings. Firstly, the prevalence of allergic conditions was more common in this cohort than has been estimated in the general UK paediatric population (albeit figures can not be directly compared due to differences in methodology).^{14–17,28} The second observation is that children with ABPA often have other allergic conditions.

There are several features in children with CF that may make allergic conditions more likely. Firstly, gut dysbiosis (which has long been implicated in allergy)^{29–32} may contribute to ABPA pathogenesis via the gut-lung axis.²¹ Furthermore, chronic infection and prolonged antibiotic exposure have been linked with allergy through Th2 skewing, and both were prolific in the cohort.²² Antibiotic use has also been identified as a factor for drug allergy,^{33–35} and it is probable that the widespread use in this cohort contributed to the frequency of drug reactions seen here.

Chronic respiratory infection with *P. aeruginosa* appeared to be associated with allergy. This is in accordance with recent literature suggesting that *P. aeruginosa* infection might contribute to allergy by stimulating histamine production²³ in addition to resulting in increased antibiotic exposure.²² There is also evidence to suggest that *P. aeruginosa* and *A. fumigatus* interact in the airway, though opinion is divided as to whether the relationship is mutually beneficial or antagonistic.³⁶ If mutually beneficial then *P. aeruginosa* may facilitate *A. fumigatus* to persist in the airways and predispose individuals to become sensitised to the fungus, resulting in ABPA.

AR was frequent in this cohort of children. Children with CF have defective mucosal clearance which may permit aeroallergens to persist in the upper airway. In combination with inflammatory changes to the airway epithelium this may create an environment highly conducive to sensitisation.³ This is analogous to concepts described in the “Dual Exposure” hypothesis,^{7–9} though in this context sensitisation might occur at damaged epithelia in the airway rather than through damaged

skin. Sensitisation to *A. fumigatus* was observed in ABPA and AR, suggesting a possible role for the fungus in producing symptoms in both the upper and lower airways. AR was seen to precede ABPA, which if reproducible on a larger scale, might suggest that *A. fumigatus*-associated allergy first arises in the upper airway in the form of AR before contributing to the pathogenesis of lower airway allergy (ABPA). This is in accordance with the “United Airways” hypothesis.¹⁰ Furthermore, some with ABPA had apparent seasonality of exacerbations suggestive of a possible interaction between aeroallergens classically associated with SAR and ABPA. This concept is not previously described in the literature, but is comparable to the relationship between AR and atopic asthma observed in phenomena such as “thunderstorm asthma”.^{10,37,38}

Comparing the median ages of each allergic subgroup in the study cohort provides some suggestions as to the chronology of allergy in CF, which appears similar to that in the general population^{11–13} but with additional features (chronic *P. aeruginosa* infection, ABPA and drug allergy). It is important to note that this was based on age at time of interview as age of onset of the conditions was not elicited.

It may seem counter-intuitive that reported asthma was uncommon in what appears to be a highly allergic group of children. However, respiratory symptoms are common in children with CF and there is often overlap, making it challenging to distinguish between CF-related symptoms and asthma.

4.1 | Limitations

Limitations to this pragmatic study include the restricted cohort size, possible recruitment bias and missing data. The COVID-19 pandemic limited the scale of the study. However, the demographics of the study cohort are broadly similar to the whole clinic and these preliminary observations are useful to trigger further research.

4.2 | Future directions

No firm conclusions can be drawn about the allergic profile of children with CF from a retrospective study of this size and nature, although it has produced findings that should prompt further research to characterise the allergic profile of children with CF. This would involve multi-centre allergy testing and prospective monitoring of allergic disease and risk factors of children with CF. We propose that further understanding of Th2 allergic inflammation in children with CF is required to elucidate

the immune mechanisms involved and to potentially facilitate the use of targeted therapies to improve quality of life in this chronic disease.

ACKNOWLEDGEMENTS

We are thankful to all who supported this work including the multidisciplinary cystic fibrosis and allergy teams at the Great North Children's Hospital, Newcastle upon Tyne Hospitals NHS Foundation Trust. Additional gratitude goes to Professor Andrew Gennery for comments on the manuscript.

CONFLICT OF INTERESTS

Malcolm Brodlie reports investigator-led research grants from Pfizer and Roche Diagnostics; speaker fees paid to Newcastle University from Novartis, TEVA and Roche Diagnostics and travel expenses to educational meetings from Boehringer Ingelheim and Vertex Pharmaceuticals. Louise J. Michaelis reports investigator-led research grants from Danone and Sanofi; Speaker fees paid to Newcastle University from Danone and Sanofi, advisory board and travel expenses to educational meetings from Danone, Novartis, Allergy Therapeutics and Sanofi Pharmaceuticals. The other authors have no conflict of interests.

AUTHOR CONTRIBUTIONS

Amy L. Faulkner: investigation, methodology, formal analysis, writing—original draft. **Michael Grayling:** formal analysis, writing—review and editing. **Benjamin Shillitoe:** formal analysis, writing—review and editing. **Malcolm Brodlie:** conceptualization, methodology, formal analysis, supervision, writing—review and editing. **Louise J. Michaelis:** conceptualization, methodology, formal analysis, supervision, writing—review and editing.

ETHICAL STATEMENT

The research was registered (audit reference 10146) and initiated as part of a service evaluation of allergy care provided to children with CF at the Great North Children's Hospital (GNCH) in accordance with national clinical guidelines, both with regards to CF and allergy^{39–44}

ORCID

Amy L. Faulkner  <http://orcid.org/0000-0001-5390-3869>

Michael Grayling  <http://orcid.org/0000-0002-0680-6668>

Benjamin Shillitoe  <http://orcid.org/0000-0003-4407-5998>

Malcolm Brodlie  <http://orcid.org/0000-0003-4591-8299>

Louise J. Michaelis  <http://orcid.org/0000-0001-8229-154X>

REFERENCES

1. Wilkinson I, Raine T, Wiles K, Goodhart A, Hall C, O'Neill H. *Oxford Handbook of clinical medicine*. 10th ed Oxford: Oxford University Press; 2017.
2. Haq IJ, Gray MA, Garnett JP, Ward C, Brodlie M. Airway surface liquid homeostasis in cystic fibrosis: pathophysiology and therapeutic targets. *Thorax*. 2016;71(3):284–287.
3. Elborn J. Cystic fibrosis. *The Lancet*. 2016;388(10059):2519–2531.
4. Pitcher-Wilmott RL, Turner RJ, Matthew MW, Pseudomonas DJ. infection, allergy, and cystic fibrosis. *Arch Dis Child*. 1982;57:582–586.
5. Tracy MC, Okorie CUA, Foley EA, Moss RB. Allergic bronchopulmonary aspergillosis. *J Fungi*. 2016;2(2):17.
6. Knutsen A. Allergic bronchopulmonary aspergillosis in asthma. *Expert Rev Clin Immunol*. 2017;13(1):11–14.
7. Du Toit G, Sampson HA, Plaut M, Burks AW, Akdis CA, Lack G. Food allergy: update on prevention and tolerance. *J Allergy Clin Immunol*. 2018;141(1):30–40.
8. Lack G. Epidemiologic risks for food allergy. *JACI*. 2008;121(6):1331–1336.
9. Perkin MR, Logan K, Marrs T, et al. Enquiring About Tolerance (EAT) study: feasibility of an early allergenic food introduction regimen. *J Allergy Clin Immunol*. 2016;137(5):1477–1486.e1478.
10. Scadding G, Scadding G. United airways paradigm in guidelines and clinical practice. In: Bachert CBA, Chenez P, eds. *The Nose and Sinuses in Respiratory Disorders: ERS Monograph*. Lausanne: European Respiratory Society, 2017.
11. Cantani A. The growing genetic links and the early onset of atopic diseases in children stress the unique role of the atopic March: a meta-analysis. *J Invest Allergol Clin Immunol*. 1999;9(5):314–320.
12. Spergel J, Paller A. Atopic dermatitis and the atopic march. *J Allergy Clin Immunol*. 2003;112(6):S118–S127.
13. Hill D, Spergel J. The atopic march Critical evidence and clinical relevance. *Ann Allergy Asthma Immunol*. 2018;120:131–137.
14. Cork M, Danby S, Ogg G. Atopic dermatitis epidemiology and unmet need in the United Kingdom. *J Dermatolog Treat*. 2019;21:1–9.
15. Loh W, Tang M. The epidemiology of food allergy in the global context. *Int J Environ Res Public Health*. 2018;15(9):2043.
16. Pols D, Wartna J, Moed H, van Alphen E, Bohnen A, Bindels P. Atopic dermatitis, asthma and allergic rhinitis in general practice and the open population: a systematic review. *Scand J Prim Health Care*. 2016;34(2):143–150.
17. National Statistics. Health Survey for England 2018—Asthma. Health Survey for England. 2019.
18. Parmar JS, Nasser S. Antibiotic allergy in cystic fibrosis. *Thorax*. 2005;60(6):517–520.
19. Vernocchi P, Del Chierico F, Russo A, et al. Gut microbiota signatures in cystic fibrosis: Loss of host CFTR function drives the microbiota enterophenotype. *PLoS One*. 2018;13(12):e0208171.
20. Dorsey J, Gonska T. Bacterial overgrowth, dysbiosis, inflammation, and dysmotility in the Cystic Fibrosis intestine. *J Cystic Fibrosis*. 2017;16:S14–S23.
21. Mac Aogáin M, Vidaillac C, Chotirmall SH. Fungal Infections and ABPA. In: Davis SD, Rosenfeld M, Chmiel J, eds.,

- Cystic Fibrosis: A Multi-Organ System Approach*. Cham: Springer International Publishing; 2020:93-126.
22. Wypych T, Marsland B. Antibiotics as instigators of microbial dysbiosis: implications for asthma and allergy. *Trends Immunol*. 2018;39(9):697-711.
 23. Potera C. Common bacterium induces histamine production in neutrophils. *Environ Health Perspect*. 2012;120(5):A190.
 24. The International Study of Asthma and Allergies in Childhood (ISAAC). Worldwide variations in the prevalence of asthma symptoms: the International Study of Asthma and Allergies in Childhood (ISAAC). *Eur Respir J*. 1998;12(2):315-335.
 25. IBM SPSS Statistics for Mac [computer program]. Version 26.0. Armonk, NY: IBM Corp.; 2019.
 26. R: A language and environment for statistical computing. [computer program]. Vienna, Austria, 2020.
 27. *Exact: Unconditional exact test. R package version 2.1*. [computer program]. 2020.
 28. Gomes ER, Brockow K, Kuyucu S, et al. Drug hypersensitivity in children: report from the pediatric task force of the EAACI Drug Allergy Interest Group. *Allergy*. 2016;71:149-161.
 29. West CD, Prescott SL, Jenmalm MC. Bugging allergy; role of pre-, pro- and synbiotics in allergy prevention. *Allergol Int*. 2017;66(4):529-538.
 30. Sampson HA, O'Mahony L, Burks AW, Plaut M, Lack G, Akdis CA. Mechanisms of food allergy. *JACI*. 2018;141(1):11-19.
 31. Arrieta MC, Stiemsma LT, Dimitriu PA, et al. Early infancy microbial and metabolic alterations affect risk of childhood asthma. *Sci Transl Med*. 2015;7(307): 307ra152.
 32. Strachan D. Hay fever, hygiene, and household size. *BMJ*. 1989;299(6710):1259-1260.
 33. Braun C, Reix P, Durieu I, et al. The diagnosis of hypersensitivity to antibiotics is rarely confirmed by allergy work-up in cystic fibrosis patients. *Pediatr Allergy Immunol*. 2020;31(4):396-404.
 34. Anstey KM, Choi L, Dawson D, Kleinhenn ME, Otani IM. Enabling antibiotic allergy evaluations and reintroduction of first-line antibiotics for patients with cystic fibrosis. *Ann Allergy Asthma Immunol*. 2021. <https://pubmed.ncbi.nlm.nih.gov/34048884/>
 35. Wright MFA, Bush A, Carr SB. Hypersensitivity reactions to intravenous antibiotics in cystic fibrosis. *Paediatr Respir Rev*. 2018;27:9-12.
 36. Zhao J, CW, He X, Liu Y. The co-colonization prevalence of *Pseudomonas aeruginosa* and *Aspergillus fumigatus* in cystic fibrosis: a systematic review and meta-analysis. *Microb Pathog*. 2018;125:122-128.
 37. Hew M, Lee J, Susanto NH, et al. The 2016 Melbourne thunderstorm asthma epidemic: riskfactors for severe attacks requiring hospital admission. *EAACI*. 2018;74(1): 122-130.
 38. D'Amato G, Liccardi G, Frenguelli G. Thunderstorm-asthma and pollen allergy. *Allergy*. 2007;62:11-16.
 39. National Institute for Health and Care Excellence. Cystic fibrosis: diagnosis and management. In: Excellence NifHaC, ed 2017.
 40. National Institute for Health and Care Excellence. *Atopic eczema in under 12s: diagnosis and management*. In. London: NICE; 2007.
 41. National Institute for Health and Care Excellence. *Asthma: diagnosis, monitoring and chronic asthma management > Recommendations*. In. Vol 2020. London: NICE; 2020.
 42. National Institute for Health and Care Excellence. Food allergy in under 19s: assessment and diagnosis. In. London: NICE; 2011.
 43. Scadding GK, Kariyawasam HH, Scadding G, et al. BSACI guideline for the diagnosis and management of allergic and non-allergic rhinitis (Revised Edition 2017; First edition 2007). *Clinical & Experimental Allergy*. 2017;47(7):856-889.
 44. National Institute for Health and Care Excellence. *Drug allergy: diagnosis and management*. In. London: NICE; 2014.

SUPPORTING INFORMATION

Additional supporting information may be found in the online version of the article at the publisher's website.

How to cite this article: Faulkner AL, Grayling M, Shillitoe B, Brodlie M, Michaelis LJ. Characterising the allergic profile of children with cystic fibrosis. *Immun Inflamm Dis*. 2021;1-10. <https://doi.org/10.1002/iid3.540>

Appendices

Appendix 1: Definitions and diagnostic criteria

Allergic Conditions

Allergic Bronchopulmonary Aspergillosis¹	Isolation of <i>Aspergillus fumigatus</i> from a respiratory sample or raised IgE in combination with a) Reduced lung function b) Respiratory symptoms in the form of wheeze c) Shortness of breath and/or cough. d) Symptomatic treatment with oral prednisolone +/- antifungals.
Allergic Rhinitis Perennial Allergic Rhinitis (PAR) Seasonal Allergic Rhinitis (SAR) Perennial Allergic Rhinitis with seasonal exacerbation (PARwe)	Participant positive for condition if: a) It had been diagnosed by medical professional - OR b) They had self-reported symptoms consistent with the condition as per the ISAAC questionnaire², e.g. answered yes when asked: i. <i>"Have you ever had a problem with sneezing, or a runny, or blocked nose when you DID NOT have a cold or the flu"</i> PAR: symptoms every month of the year. SAR: symptoms only in some months of the year. PARwe: symptoms every month of the year but worse in certain months.
Atopic Asthma	Participant positive for condition if: a) It had been diagnosed by medical professional - OR b) They had self-reported symptoms consistent with the condition as per the ISAAC questionnaire², e.g. answered yes when asked: i. <i>"Has your child ever had wheezing or whistling in the chest at any time in the past?"</i>
Drug Allergy	Participant positive for condition if: a) It had been diagnosed by medical professional - OR b) They had self-reported symptoms consistent with the condition as per the RCPCH care pathway³, e.g. answered yes when asked: i. <i>"Has your child ever experienced an allergic reaction to a prescribed drug?"</i>
Eczema	Participant positive for condition if: a) It had been diagnosed by medical professional - OR b) They had self-reported symptoms consistent with the condition as per the ISAAC questionnaire², e.g. answered yes when asked: i. <i>"Has your child ever had an itchy rash which was coming and going for at least six months?"</i>
Food Allergy	Participant positive for condition if: a) It had been diagnosed by medical professional - OR b) They had self-reported symptoms consistent with the condition as per the RCPCH care pathway⁴, e.g. answered yes when asked: i. <i>"Has your child ever had an allergic reaction to something they have eaten?"</i>

Other definitions

Aeroallergens included in allergy testing	House dust mite, moulds mix, grass mix, tree mix, weed mix, silver birch, dog, cat.
Basophil Count	Normal range as per the Newcastle upon Tyne Hospitals NHS Foundation Trust ⁵ : 0.00 - 0.10 x 10 ⁹ /L
Cystic Fibrosis	Children with a clear diagnosis of cystic fibrosis under the care of the Great North Children's Hospital's regional multi-disciplinary team.
Eosinophil Count	Age-specific normal ranges as per the Newcastle upon Tyne Hospitals NHS Foundation Trust ⁵ : between 0.3-0.8 and 0.04-0.4 x 10 ⁹ /L
Total Immunoglobulin E	Normal ranges as per the Newcastle upon Tyne Hospitals NHS Foundation Trust ⁵ : Neonates 0-1.5 IU/L, infants 0-15 IU/L, children aged 1-5 years 0-60 IU/L, children aged 6-9 years 0-90 IU/L, children aged 10-15 years 0-200 IU/L, 16 years and over 0-100 IU/L.

¹Sisodia J, Bajaj T. Allergic Bronchopulmonary Aspergillosis. [Updated 2021 Mar 2]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2021 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK542329/>

² The International Study of Asthma and Allergies in Childhood (ISAAC). Worldwide variations in the prevalence of asthma symptoms: the International Study of Asthma and Allergies in Childhood (ISAAC). Eur Respir J. 1998;12(2):315-335

³ Royal College of Paediatrics and Child Health, 2011. Allergy care pathway for drug allergy. Available at: <https://www.rcpch.ac.uk/resources/allergy-care-pathway-drug-allergy>. [Accessed 07/04/2021]

⁴ Royal College of Paediatrics and Child Health, 2011. Allergy care pathway for food allergy. Available at: <https://www.rcpch.ac.uk/resources/allergy-care-pathway-food-allergy>. [Accessed 07/04/2021]

⁵ The Newcastle upon Tyne Hospitals Trust. Home - Newcastle Laboratories. <https://www.newcastlelaboratories.com/>. Published 2020. Accessed 14/11/2020, 2020.

Appendix 2: Modified ISAAC questionnaire used

Questions regarding drug and food allergy, not ordinarily included in the International Study of Asthma and Allergies in Childhood (ISAAC) questionnaire, were additional as per the Royal College of Paediatrics and Child Health Allergy Pathways^{1,2}. Appendix 1 outlines the definitions and diagnostic criteria used for conditions examined in the study.

¹Royal College of Paediatrics and Child Health, 2011. Allergy care pathway for drug allergy. Available at: <https://www.rcpch.ac.uk/resources/allergy-care-pathway-drug-allergy>. [Accessed 07/04/2021]

² Royal College of Paediatrics and Child Health, 2011. Allergy care pathway for food allergy. Available at: <https://www.rcpch.ac.uk/resources/allergy-care-pathway-food-allergy>. [Accessed 07/04/2021]

FOOD ALLERGY

1.

- a) Has your child ever had an allergic reaction to something they have eaten?
Yes ☐ No ☐ Don't Know ☐

IF THE ANSWER TO a) IS YES, COMPLETE QUESTION. IF NO, SKIP TO NEXT QUESTION

- b) Which food has caused your child their MOST SEVERE allergic reaction? This may have been the ONLY reaction)
- c) Thinking about the symptoms of your child's allergy/allergies please consider the symptoms associated with his or her MOST SEVERE reaction (this may have been the ONLY reaction) and tick all that apply.

- | | |
|---|---|
| Vomiting ☐ | Abdominal pain ☐ |
| Irritability ☐ | Rash ☐ |
| Face Swelling ☐ | Tingling or sore mouth ☐ |
| Swelling of the lips or tongue ☐ | Throat tightening or difficulty swallowing ☐ |
| Breathing difficulties ☐ | Wheeze ☐ |
| Blue around the lips ☐ | Collapse or faint ☐ |
| Other (please specify): | |

- d) Thinking about the MOST SEVERE allergic reaction that your child has experienced:

- i. What treatment did your child take?

- | | | |
|--|---|-------------------------------------|
| Anti-Histamines ☐ | Adrenaline Injection (Epi-Pen) ☐ | None ☐ |
| Oral Steroid ☐ | Salbutamol Inhaler ☐ | Don't know ☐ |

- ii. Did your child go to hospital for the reaction?

- Yes ☐ No ☐ Don't Know ☐

- iii. If 4d)ii was yes, did your child stay overnight in hospital?

- Yes ☐ No ☐ Don't Know ☐

- iv. Was your child admitted to:

- High Dependency Ward ☐ Other Ward ☐ Don't know ☐

- e) Apart from the most severe reaction, has your child had any allergic reactions to other foods? If so state what he or she had a reaction to.

- f) Has your child ever had any of the following?

- | | | |
|--|---|---|
| Difficulty swallowing ☐ | Pain during swallowing ☐ | Inability to swallow ☐ |
| None ☐ | Don't know ☐ | |

NON-FOOD ALLERGY

2.

- a) Has your child ever experienced an allergic reaction to something other than a food or a drug?
Yes ☐ No ☐ Don't Know ☐

IF THE ANSWER TO a) IS YES, COMPLETE QUESTION. IF NO, SKIP TO NEXT QUESTION

- b) Thinking about your child's MOST SEVERE non-food reaction, what were they allergic to?
- c) Thinking about the symptoms of your child's non-food allergy/allergies please consider the symptoms associated with his or her MOST SEVERE reaction (this may have been the ONLY reaction) and tick all that apply.

- | | |
|---|---|
| Vomiting ☐ | Abdominal pain ☐ |
| Irritability ☐ | Rash ☐ |
| Face Swelling ☐ | Tingling or sore mouth ☐ |
| Swelling of the lips or tongue ☐ | Throat tightening or difficulty swallowing ☐ |
| Breathing difficulties ☐ | Wheeze ☐ |
| Blue around the lips ☐ | Collapse or faint ☐ |
| Other (please specify): | |

- d) Thinking about the most severe allergic reaction that your child has experienced:
- i. What treatment did your child take?
- | | | |
|--|---|-------------------------------------|
| Anti-Histamines ☐ | Adrenaline Injection (Epi-Pen) ☐ | None ☐ |
| Oral Steroid ☐ | Salbutamol Inhaler ☐ | Don't know ☐ |
- ii. Did your child go to hospital for the reaction?
- Yes ☐ No ☐ Don't Know ☐
- iii. If d)ii. was yes, did your child stay overnight in hospital?
- Yes ☐ No ☐ Don't Know ☐
- iv. Was your child admitted to:
- | | | |
|---|-------------------------------------|-------------------------------------|
| High Dependency Ward ☐ | Other Ward ☐ | Don't know ☐ |
|---|-------------------------------------|-------------------------------------|
- e) Have any of your child's allergies (food, drug or other) been diagnosed by a medical professional?
- Yes ☐ No ☐ Don't Know ☐
- f) If e) was Yes, which allergy or allergies have been diagnosed?

IF THE ANSWER TO e) IS YES, COMPLETE QUESTION. IF NO, SKIP TO NEXT QUESTION

- g) Has your child been referred to an allergy specialist? If so, for which allergy?
- | | |
|------------------------------|-------------------------------------|
| Yes ☐ | Allergy: _____ |
| No ☐ | Don't Know ☐ |
- h) Does your child have an Adrenaline Auto-injector (EpiPen/JEXT)
- Yes ☐ No ☐ Don't Know ☐

DRUG REACTIONS

3.

- a) Has your child ever experienced an allergic reaction to a prescribed drug?
- Yes ☐ No ☐ Don't Know ☐

IF THE ANSWER TO a) IS YES, COMPLETE QUESTION. IF NO, SKIP TO NEXT QUESTION

b) Thinking about your child's MOST SEVERE drug reaction, which drug did they react to?

c) Thinking about the symptoms of your child's non-food allergy/allergies please consider the symptoms associated with his or her MOST SEVERE reaction (this may have been the ONLY reaction) and tick all that apply.

- | | |
|---|---|
| Vomiting ☐ | Abdominal pain ☐ |
| Irritability ☐ | Rash ☐ |
| Face Swelling ☐ | Tingling or sore mouth ☐ |
| Swelling of the lips or tongue ☐ | Throat tightening or difficulty swallowing ☐ |
| Breathing difficulties ☐ | Wheeze ☐ |
| Blue around the lips ☐ | Collapse or faint ☐ |
| Headache ☐ | Blood in Stool ☐ |
| Diarrhoea ☐ | Localised reaction to infusion site (If IV drug) ☐ |
| Other (please specify): | |

d) Thinking about the most severe allergic reaction that your child has experienced:

i. What treatment did your child take?

- | | | |
|--|---|-------------------------------------|
| Anti-Histamines ☐ | Adrenaline Injection (Epi-Pen) ☐ | |
| Oral Steroid ☐ | Salbutamol Inhaler ☐ | None ☐ |
| IV Steroids ☐ | IV Antihistamine ☐ | Don't know ☐ |

ii. Did the reaction occur in hospital or elsewhere? (e.g: at home)

- Hospital ☐ Elsewhere ☐

iii. If the answer to ii. was elsewhere, was your child taken to hospital because of their reaction?

- Yes ☐ No ☐

iv. If iii. was yes, was your child admitted to:

- High Dependency Ward ☐ Other Ward ☐ Don't know ☐

e) Apart from the most severe reaction, has your child had any allergic reactions to other prescribed drugs? If so state which drug/drugs he or she has had a reaction to

ECZEMA

4.

a) Has your child ever been diagnosed with **Eczema/Atopic Dermatitis**?

- Yes ☐ No ☐ Don't Know ☐

b) Has your child ever had an itchy rash which was coming and going for at least six months?

- Yes ☐ No ☐ Don't Know ☐

IF THE ANSWER TO a) OR b) IS YES, COMPLETE QUESTION. IF NO, SKIP TO NEXT QUESTION

c) Has your child had this itchy rash at any time within the last 12 months?

- Yes ☐ No ☐ Don't Know ☐

d) Has this itchy rash affected any of the following places at any time? (i.e. not just within the last 12 months)
Folds of elbows, behind the knees, in front of ankles, under the buttocks, or around the neck, ears or eyes?

- Yes ☐ No ☐ Don't Know ☐

e) At what age did this rash first occur?

- Under 2 years old ☐ 2-4 years old ☐ 5 or older ☐ Don't know ☐

- f) In the past 12 months, how often on average has your child been kept awake at night by this itchy rash?
 Never in the past 12 months ☐
 Less than one night per week ☐
 One or more nights a week ☐
- g) Has your child ever been prescribed topical steroid cream for this rash?
 Yes ☐ No ☐ Don't Know ☐
- h) Tick any of the following treatments that your child requires most days for their eczema when it is present:
 Emollients ☐ Topical Steroid creams ☐ Soap Substitute ☐
 Bath Additives ☐ Oral Steroids ☐ IV Steroids ☐
 Skinny Garments ☐ Icthopaste Bandages ☐ Immune Suppressive ☐
 Biologics ☐ None ☐

ATOPIC ASTHMA

5.

- a) Has your child been diagnosed with Atopic Asthma?
 Yes ☐ No ☐ Don't Know ☐
- b) Has your child's chest ever sounded wheezy?
 Yes ☐ No ☐ Don't Know ☐

IF THE ANSWER TO a) OR b) IS YES, COMPLETE QUESTION. IF NO, SKIP TO NEXT QUESTION

- c) How many attacks of wheezing has your child had in the last 12 months?
 None ☐ 1 to 3 ☐ 4 to 12 ☐ More than 12 ☐
- d) In the past 12 months, how often has your child's sleep been disturbed due to wheezing?
 Never woken with wheezing ☐
 Less than one night per week ☐
 One or more nights per week ☐
 Don't know ☐
- e) In the past 12 months, has wheezing ever been severe enough to limit your child's speech to only one or two words at a time between breaths?
 Yes ☐ No ☐ Don't Know ☐
- f) Has your child ever sounded wheezy during or after exercise?
 Yes ☐ No ☐ Don't Know ☐
- g) Has your child sounded wheezy during or after exercise in the last 12 months?
 Yes ☐ No ☐ Don't Know ☐
- h) Has your child been prescribed any of the following inhalers for asthma or wheeze?
 Reliever inhaler (blue) ☐ Preventor inhaler (brown) ☐
 Neither ☐ Don't Know ☐

- i) Has your child ever been prescribed oral steroids for asthma or wheeze?
Yes ☐ No ☐ Don't Know ☐
- j) In the last 12 months, has your child had a dry cough at night that wasn't associated with a cold or a chest infection?
Yes ☐ No ☐ Don't Know ☐

ALLERGIC RHINITIS

6.

- a) Has your child been diagnosed with **Hayfever / Allergic Rhinitis**?
Yes ☐ No ☐ Don't Know ☐
- b) Does your child ever experience an itchy and/or runny or blocked nose when they do not have a cold?
Yes ☐ No ☐ Don't Know ☐

IF THE ANSWER TO a) OR b) IS YES, COMPLETE QUESTION. IF NO, SKIP TO NEXT QUESTION

- c) In which of the past 12 months did this nose problem occur?
January ☐ April ☐ July ☐ October ☐
February ☐ May ☐ August ☐ November ☐
March ☐ June ☐ September ☐ December ☐
None ☐ Don't Know ☐
- d) Does your child require any of the following treatments MOST DAYS during the problematic months?
Antihistamines ☐ Inhaled Nasal Steroids ☐
Eye Drops ☐ Anti-Leukotriene Antagonist ☐
Sublingual Immunotherapy ☐ Subcutaneous Immunotherapy ☐
Oral Steroids ☐
None ☐ Don't know ☐
- e) In the last 12 months, how much did this nose problem interfere with your child's daily activities?
Not at all ☐ A Little ☐ A Moderate amount ☐ A lot ☐

7.

- a) Has your child ever been diagnosed with Allergic Bronchopulmonary Aspergillosis
Yes ☐ No ☐ Don't Know ☐
- b) If 7a) was yes, has your child received oral steroids for the condition?
Yes ☐ No ☐ Don't Know ☐
- c) If 7a) was yes, has your child received other medication for the condition apart from oral steroids?
Yes ☐ No ☐ Don't Know ☐

GENERAL MEDICAL HISTORY

8. Thinking about your child's current or past feeding (depending on their age)
- a) Does / Did your child have
Breast Milk only ☐ Formula Milk only ☐ Breast Milk and Formula milk ☐
- b) ~~Yes~~ ☐ If your child has / had formula milk, what is the name or brand?
No ☐
9. Does your child have any other medical conditions for which he/she sees the doctor regularly about? If yes please state condition(s)
Don't Know ☐
10. Has your child been admitted to Hospital for any reason other than stated previously in the last two years? If yes state what for
Yes ☐
No ☐
Don't Know ☐

DRUG HISTORY

11. *Only answer if answered yes to having hayfever symptoms*

a) Has your child ever been prescribed any of the following for their hayfever?

Anti-Histamines ☐

Nasal Spray ☐

Eye Drops ☐

Sublingual Immunotherapy ☐

Subcutaneous Immunotherapy ☐

None ☐

Don't know ☐

12. Is your child on any other regular prescribed medications that aren't for Cystic Fibrosis? If so state what

SOCIAL HISTORY

Thinking about the house which your child spends the most time in:

a) Do any smokers live at the house?

Yes – they smoke outside ☐

Yes – they smoke inside ☐

No ☐

b) Do any pets live inside the house?

Dog ☐

Cat ☐

Bird ☐

Other: