



Contents lists available at ScienceDirect

Journal of Biotechnology

journal homepage: www.elsevier.com/locate/jbiotec



Effects of orally applied Fes p1-displaying *L. plantarum* WCFS1 on Fes p1 induced allergy in mice

Rajna Minic^{a,*}, Marija Gavrovic-Jankulovic^c, Vladimir Petrusic^a, Irena Zivkovic^a, Vincent G.H. Eijsink^b, Ljiljana Dimitrijevic^a, Geir Mathiesen^b

^a Institute of Virology, Vaccines and Sera, Torlak, Belgrade, Serbia

^b Department of Chemistry, Biotechnology and Food Science, Norwegian University of Life Sciences, Aas, Norway

^c Department of Biochemistry, Faculty of Chemistry, University of Belgrade, Serbia

ARTICLE INFO

Article history:

Received 16 October 2014

Received in revised form 8 December 2014

Accepted 14 January 2015

Available online xxx

Keywords:

Allergy

Oral therapy

Recombinant *L. plantarum* WCFS1

Festuca pratensis

ABSTRACT

Group I grass pollen allergens are major contributors to grass pollen-related seasonal allergic rhinitis, and as such a primary target for allergen specific immunotherapy. In this study the potential therapeutic role of oral application of *Lactobacillus plantarum* WCFS1, directing cell wall attachment of the recombinant Fes p1 allergen, from *Festuca pratensis* was tested in a mouse model of Fes p1 allergy. For surface expression of Fes p1 allergen in *L. plantarum* WCFS1 pSIP system with inducible expression was used. Balb/c mice were sensitized with Fes p1 protein in alum and subsequently received live recombinant *L. plantarum* orally. Antibody levels (IgE, total IgG, IgG1, IgG2a, and IgA) were determined by ELISA. Differential eosinophil count in peripheral blood was performed. Reduced peripheral blood eosinophilia and increased serum IgG2A levels was detected in both groups which received live *L. plantarum* orally. Specific serum IgA levels were increased only in mice treated with the recombinant bacteria. Oral application of *L. plantarum* WCFS1 has a beneficial therapeutic effect in a mouse model of Fes p1 allergy. Cell surface expression of Fes p1 allergen potentiates this phenomenon in an allergen specific way.

© 2015 Published by Elsevier B.V.

1. Introduction

Type I allergy is one of the leading disorders of the immune system affecting about 25% of the population worldwide. Subcutaneous allergen specific immunotherapy has been used for almost a century and numerous studies have shown the benefits of this strategy (Bousquet et al., 1998). However, the efficacy of this strategy varies and it is associated with a risk of anaphylaxis and lengthy therapeutic regimes. Mucosal tissues contain antigen-presenting cells with a tolerogenic phenotype and their immunostimulation is less likely to give rise to anaphylactic shock (Calderon et al., 2012). Therefore, it is of interest to target mucosal tissues when developing strategies to combat allergy.

In recent years probiotic lactic acid bacteria have received massive attention because of their potential beneficial effects on human health (Isolauri et al., 2002). There is increasing experimental evidence for claims that lactic acid bacteria exert beneficial effects on gut health (Kalliomaki et al., 2003; Sazawal et al., 2006) and that

some of these effects relate to favorable immunomodulating properties, such as tolerance induction (van Baarlen et al., 2009). To date, several clinical trials have shown that probiotic strains may reduce allergy symptoms (Kalliomäki et al., 2010; Snel et al., 2011). In order to achieve immunological memory towards the specific allergen it is of interest to use the relevant allergen together with lactobacilli, to prevent the possibility of future sensitization to the same allergen.

L. plantarum WCFS1 is one of the best characterized lactic acid bacteria both in terms of genome analysis, behavior in the gastrointestinal tract and immunomodulatory properties (Kleerebezem et al., 2003; Marco et al., 2009; van Hemert et al., 2010). Considering the tolerance-inducing immunomodulatory effects of *L. plantarum* (van Baarlen et al., 2009), it is of interest to explore the possibility to use allergen expressing lactobacilli as delivery vehicles of allergy vaccines or in immunotherapy (Kruisselbrink et al., 2001; Daniel et al., 2006; Rigaux et al., 2009). Indeed, it has been shown that intranasal prophylactic application of allergen-expressing *L. plantarum* in different mouse allergy models reduces allergic sensitization by shifting the immune response to a non-allergic Th1 immune response as evidenced by production of specific IgG2a antibodies and increased secretion of IFN- γ (Repa et al., 2003; Daniel et al., 2006; Rigaux et al., 2009; Schwarzer et al., 2011).

* Corresponding author at: Institute of Virology, Vaccines and Sera, Torlak, Vojvode stepa 458, Belgrade, Serbia. Tel.: +381 648852431; fax: +381 113975501. E-mail address: minicrajna@gmail.com (R. Minic).

It has also been shown that therapeutic oral administration of *L. acidophilus* ATCC 4356 expressing the Der p5 allergen results in drastically decreased allergen-specific serum IgE levels and lowered eosinophilia in bronchoalveolar lavage samples (Charng et al., 2006).

Based on these promising previous results, the goal of the present study was to explore the possibility to use an engineered *L. plantarum* with a surface-displayed respiratory allergen Fes p1 in immunotherapy for pollen allergy. Covalent surface-anchoring of the antigen has the advantage that it minimizes the risk of the antigen entering the blood stream. Fes p1, a group I allergen from *Festuca pratensis* (Poeae grasses) was chosen because of its global distribution across cool, temperate areas (Borrill et al., 1976). Furthermore, sensitization to Fes p1 usually implies sensitization to several grass species due to group I grass pollen allergens being highly conserved among these species (Laffer et al., 1994). One closely related allergen is Phl p1 from *Phleum pratense*, which was recently found to be the initiator of IgE sensitization in 75% of children with grass pollen-related seasonal allergic rhinitis in Germany (Hatzler et al., 2012). Group I allergens are usually glycoproteins, with a carbohydrate content comprising up to 5% of the molecular mass (Petersen et al., 1995).

To engineer a suitable recombinant *L. plantarum* strain, we used recently developed genetic tools (Sørvig et al., 2003, 2005; Mathiesen et al., 2009; Fredriksen et al., 2010) for expression and cell-wall anchoring of heterologous proteins in lactobacilli. Thus, a *L. plantarum* strain was generated in which Fes p1 is covalently anchored to the bacterial cell wall and displayed on the bacterial surface, as documented by immunofluorescence microscopy. The effect of this engineered strain on Fes p1-induced allergy was studied in a mouse model.

2. Methods

2.1. Bacterial strains and culture conditions

E. coli Top 10 cells (Life Technologies Ltd., Paisley, UK) were grown in BHI broth (Oxoid Ltd., Basingstoke, UK) at 37 °C with shaking. *Lactobacillus plantarum* WCSF1 (Kleerebezem et al., 2003) was propagated in MRS (Oxoid) medium at 37 °C without agitation. For plates, the medium was solidified with 1.5% agar (w/v). The antibiotic concentrations were 200 µg ml⁻¹ and 10 µg ml⁻¹ erythromycin for *E. coli* and *L. plantarum*, respectively.

2.2. Plasmid construction and transformation

All vectors used in the present study are derivatives of the modular pSIP vectors developed for inducible expression in *Lactobacillus* (Sørvig et al., 2003, 2005) and further developed for secretion and anchoring of target proteins (Mathiesen et al., 2008, 2009). The gene for Fes p1 (GenBank: CCD28290) with codon usage optimized for *L. plantarum* and inserted in pUC57 was purchased from GenScript (Piscataway, NJ, USA). The sequence was N-terminally truncated, and started with amino acid 43. The coding sequence was PCR amplified using Phusion® High-Fidelity DNA Polymerase (New England Biolabs, Ipswich, MA) with the primers: Forward, 5'-TCAGAAGAGGATCTGGATGCTAAAAGTACATGGT-3', and Reverse 5'-AAAACGCGTCTTTGCAGCATAACTCGT-3' (MluI site italics), both from Operon Biotechnologies GmbH (Cologne, Germany). A Myc-tag (N-EQKLISEEDL-C) was fused N-terminally to the coding sequence by reamplification of the PCR product with an extended forward primer 5'-AAAGTCGACGAACAAAACTCATCTCAGAGGATCTG-3' (Sall site italics). The ~700bp PCR fragment was digested with Sall/MluI and ligated into Sall/MluI digested ~6.4bp

fragment of pLp_0373sOFACwa2 (Fredriksen et al., 2010), yielding pLpFesp1CWA. All restriction enzymes were purchased from New England Biolabs. In the constructed plasmid, Fes p1 containing N-terminal Myc-tag is N-terminally fused to the *L. plantarum* signal peptide of lp_0373 (Mathiesen et al., 2008), and C-terminally fused to the 194 amino acid long LPxTG anchor sequence of the lp_2578 gene (Fredriksen et al., 2012). *E. coli* Top 10 cells made competent by calcium chloride were used for subcloning, and the final plasmid, pLpFesp1CWA, was transformed to electrocompetent *L. plantarum* WCSF1 (Aukrust et al., 1995). The expression of Fes p1 in *L. plantarum* WCSF1 was checked by Western blot analysis using an Anti-Myc antibody (Life Technologies). *L. plantarum* harboring a pSIP plasmid without the Fes p1 gene (named pEV) (Fredriksen et al., 2012) was used as a negative control.

2.3. Flow cytometry and indirect immunofluorescence microscopy of Fes p1 expressing *L. plantarum*

Surface display of Fes p1 was analyzed by both flow cytometry and indirect immunofluorescence microscopy of *L. plantarum*. The staining procedure was the same for both analyses. For Fes p1 expression the bacteria were induced at optical density (610 nm) of 0.25 by the addition of the inducing peptide (IP-673) (CASLO Laboratory ApS, Denmark) at 10 ng ml⁻¹ final concentration, essentially as described previously (Halbmayer et al., 2008; Mathiesen et al., 2008). Two hours post-induction, the bacteria were harvested by centrifugation at 3000 × g for 5 min, and washed once with PBS containing 2% bovine serum albumin (PBS/2% BSA). Approximately 2 × 10⁷ of harvested bacteria were resuspended in 0.5 ml PBS/2% BSA and incubated with 5 µg Anti-Myc antibody, for 15 min at 4 °C. After incubation the cells were washed twice with 1 ml 2% B-PBS, followed by incubation in 0.1 ml PBS/2% BSA containing 5 µg Anti-Mouse IgG (Fc specific)–FITC antibody produced in goat (Sigma–Aldrich, St. Louis, MO, USA). After incubation for 15 min at 4 °C, the bacteria were collected by centrifugation at 3000 × g for 2 min and washed twice in PBS. Flow cytometry analysis was done on MACSQuant analyzer (Miltenyi Biotec GmbH, Bergisch Gladbach, Germany), following the manufacturer's instructions. The immunofluorescence microscopy signal was visualized using a Leica SP5 confocal scanning laser microscope using a 488-nm argon laser (FITC photomultiplier tube [PMT]) and a bright field (BF) PMT for transmitted light (Leica Microsystems, GmbH, Wetzlar, Germany). *L. plantarum* harboring a pSIP plasmid without the Fes p1 gene (named pEV) (Fredriksen et al., 2012) was used as a negative control.

2.4. Purification of Fes p1

Festuca pratensis pollen was obtained from the Institute of Virology, Vaccines and Sera, Torlak, Serbia. The pollens were lysed by incubation of 1 g of pollen in 10 ml 20 mM acetate buffer (pH 4.5) over night at 4 °C, after which debris was removed by centrifugation at 10,000 × g for 10 min. The supernatant was passed through a 5 cm × 0.7 cm CM-cellulose column (Sigma–Aldrich, St. Louis, MO, USA), equilibrated with 20 mM acetate buffer (pH 4.5). Stepwise elution was performed with 100 mM, 200 mM, 500 mM, and 1000 mM NaCl in 20 mM acetate buffer (pH 4.5) and the protein was eluted from the column in the 1 M NaCl fraction. Protein concentrations were determined by measuring absorbance at 280 nm, using the theoretical extinction coefficient for Fes p1.

2.5. Experimental protocol for animal tests

The therapeutic potential of the engineered *L. plantarum* strains was tested *in vivo* using a mouse model of Fes p1 induced allergy. Four week old female Balb/c mice were purchased from Military




Fig. 1. Overview of the experimental design. On days 0 and 7 Balb/c mice received intraperitoneal injections of 7 μ g Fes p1 allergen in alum. Oral administration of lactobacilli was performed on 12 days, divided into four periods: 18–20, 25–27, 55–57, and 60–62. At day 68 the mice were sacrificed and blood samples were taken for analysis.

Medicine Academy (VMA, Belgrade, Serbia), divided into groups of ten animals, kept in individual ventilated cages, and fed with standard food and water *ad libitum* in a pathogen-free part of the animal house. Humidity, temperature and light/dark cycles were maintained at $55\% \pm 5\%$, $21 \pm 2^\circ\text{C}$ and 12/12 h, respectively. All experiments were approved by the Ethics Committee for the welfare of experimental animals, at the Institute of Virology, Vaccines and Sera in Torlak, Serbia, and conformed to Serbian laws and European regulations on animal welfare (Approval No. 011-00-00510/2011-05/4). At day 0 and 7 the mice were sensitized by intra peritoneal (i. p.) injection with 7 μ g Fes p1 allergen in 100 μ l PBS mixed with an equal amount of aluminum hydroxide (alum) suspension (Fig. 1). At days 18, 19, 20, 25, 26, 27, 55, 56, 57, 60, 61 and 62 (Fig. 1) the mice were given 1×10^9 recombinant *L. plantarum* in 20 μ l fresh MRS medium without antibiotics by oral gavage. The bacteria were grown, induced and harvested as described for immunofluorescence microscopy in Section 2.3. Untreated mice were subjected to the same experimental protocol, but were given sterile MRS medium. At day 68 all mice were euthanized by cervical dislocation and samples of peripheral blood were collected, PBMC were immediately isolated, and serum samples were collected and stored at -80°C .

2.6. Specific antibody detection

MaxiSorp plates (Nunc A/S, Denmark) were coated with 0.5 μ g purified Fes p1 allergen in 50 μ l 0.1 M sodium carbonate buffer (pH 9.6) overnight at 4°C . After blocking with PBS/2% BSA (1 h, 37°C), the wells were incubated with 50 μ l diluted mice sera (100 fold dilution for IgG, IgG1, IgG2a, IgG3 and IgA and 6 fold dilution for IgE) for 1 h at 37°C . Dilutions were made in PBS containing 1% BSA. Biotinylated anti-mouse antibodies (Biolegend, San Diego, CA) were used for specific antibody class/subclass determination. Streptavidin peroxidase (Sigma–Aldrich, St. Louis, MO, USA) was used for detection with *ortho*-phenyldiamin as the substrate.

2.7. Eosinophil counts

To count eosinophils among leukocytes, peripheral blood smears from each mouse were prepared and after fixation in methanol for 10 min the leukocytes were stained with May–Grünwald Giemsa. Eosinophils were counted in a double blind fashion, using a light microscope at 400 fold magnification, with the total count being 300 leukocytes per slide.

2.8. Statistical analysis

Statistical analysis of antibody levels measurement was done with use of One-Way ANOVA test; Tukey test for means comparison. Statistical analysis of eosinophil number was done with use of Student's *t*-test for independent samples. Results are presented as mean values $\pm$ standard error of the mean.




Fig. 2. Expression of Fes p1 in *L. plantarum*, and purification of Fes p1 from pollen. (A) Western blot analysis of the whole cell lysates. Lane 1, *L. plantarum* WCFS1 harboring pLpFesp1CWA; lane 2, *L. plantarum* WCFS1 harboring pEV; lane mm, molecular marker (kDa) (Life Technologies). The predicted mass of the N- and C-terminally tagged Fes p1 protein is 53 kDa. (B) Coomassie-Brilliant blue stained 12% SDS-PAGE gel showing purification of Fes p1 allergen. mm, molecular marker (kDa); lanes 1–4, step elution of the CM-cellulose column with 100 mM NaCl, 250 mM NaCl, 500 mM NaCl and 1 M NaCl, respectively. The predicted mass of the non-glycosylated Fes p1 protein is 28 kDa.

3. Results

3.1. Protein expression analysis and protein purification

The Fes p1 allergen was cloned, with N-terminal Myc tag, into a previously developed pSIP vector (Fredriksen et al., 2010) where it becomes linked to N-terminal signal sequence for secretion and a C-terminal cell wall anchor for sortase-mediated covalent attachment of the allergen to the surface of *L. plantarum* WCFS1. Western blot analysis of cell lysates harvested 2 h after induction clearly showed that *L. plantarum* harboring pLpFesp1CWA produces Fes p1 (Fig. 2A). A slightly larger molecular weight of Fes p1, than the predicted 53 kDa is observed, which might be a consequence of covalent binding to the bacterial cell wall. Fes p1 allergen was isolated from *Festuca pratensis* pollen, and the fraction used was eluted with 1 M NaCl (Fig. 2B). The localization of the allergen was analyzed using flow cytometry and indirect immunofluorescence microscopy, which confirmed that Fes p1 was displayed at the surface of the recombinant *L. plantarum* cells (Fig. 3A and B).

3.2. Specific antibody production

Peripheral blood samples were taken at day 68 and used for analysis of Fes p1-specific immunoglobulins using purified Fes p1 (the purity was estimated to be in around 90% (Fig. 2B)) in the ELISA. As expected, IgE levels increased in all groups injected with purified Fes p1 allergen, with slightly lower values in pLp.pEV group (Fig. 4A). This shows that the allergy induction protocol was successful in terms of IgE production.

In order to see whether oral administration of *L. plantarum* resulted in changes in allergen specific levels of protective IgG and IgA antibodies the levels of these immunoglobulin classes was determined. All immunized groups showed elevated levels of Fes p1-specific total IgG, with no significant difference between the




Fig. 3. Flow cytometry and fluorescent microscopy analysis confirming the surface localization of the recombinant protein. (A) Flow cytometry analysis using mouse Anti-Myc antibody and FITC conjugated anti mouse IgG. *L. plantarum* WCFS1 harboring pEV shown in gray; *L. plantarum* WCFS1 harboring pLpFesp1CWA no color. (B) Immuno-fluorescence microscopy using mouse Anti-Myc antibody and FITC conjugated anti mouse IgG. 1, *L. plantarum* WCFS1 harboring pLpFesp1CWA; 2, *L. plantarum* WCFS1 harboring pEV.




Fig. 4. (A–E) Analysis of Fes p1 specific immunoglobulins in mice sera sampled at day 68. (A) Fes p1 specific IgE; (B) Fes p1 specific total IgG; (C) Fes p1 specific IgG1; (D) Fes p1 specific IgG2a; (E) Fes p1 specific IgA. (F) Peripheral blood eosinophil counts, expressed as percentage of eosinophils in the total number of leukocytes. C+, mice i.p. injected with Fes p1 at day 0 and 7; pLp.pEV, Fes p1 immunized mice treated by oral application of *L. plantarum* WCFS1 carrying pEV (control vector); pLp.Fesp1CWA, Fes p1 immunized mice treated by oral application of *L. plantarum* carrying pLpFesp1CWA. Bars with different superscripts are significantly different (* $p < 0.05$).

treated and the non-treated mice (Fig. 4B). ELISA showed that the majority of IgG was of the IgG1 subclass; also with respect to IgG1 levels there were no statistically significant difference between the immunized groups (Fig. 4C). The level of subclass IgG2a was also analyzed, and the production was increased in both groups administered with bacteria (Fig. 4D).

Fes 1p-specific IgA levels are of particular interest because production of these secreted antibodies is indicative of a mucosal immune response, i.e. of the type of response one hopes to achieve when using antigen-displaying lactobacilli. The measurements showed that IgA production was significantly higher in mice treated with *L. plantarum* harboring pLpFesp1CWA compared to any of the other groups (Fig. 4E).

3.3. Peripheral eosinophil counts

Increased eosinophil levels (i.e. higher than 5% of the total number of leukocytes) in peripheral blood are generally considered as evidence of an increased IL-5 production (Fulkerson and Rothenberg, 2013), which is indicative of a Th2 type of response.

Fig. 4F shows eosinophil levels in the four different mice groups of this study. In the positive control group (c+) the eosinophil level was higher than 5% of the total leukocytes, confirming that these mice have developed eosinophilia as part of the allergic reaction towards the Fes p1 protein. Treatment with *L. plantarum* cells with or without the allergen resulted in reduced numbers of eosinophils in the peripheral blood. Similar level of eosinophils as in the negative control mice was found in both mouse groups which received the bacteria.

4. Discussion

In the present study a major group I allergen from *F. pratensis* pollen, Fes p1, was successfully expressed and displayed at the surface of *L. plantarum* WCFS1 (Fig. 2A and 3). Surface-display of heterologous proteins in lactobacilli is still not too common and the results obtained with Fes p1 are encouraging. The idea behind the construction of the Fes p1-displaying strain was that surface-exposure will facilitate the interaction with the mucosal immune system. In addition, covalent attachment to the bacterial cell wall

prevents entering the blood stream by the allergen, which in principle could cause anaphylactic shock.

Oral application of both pEV and pLpFesp1CWA harboring *Lactobacilli* resulted in amelioration of several allergy parameters: reduction of peripheral blood eosinophilia and increase in production of anti Fes p1 IgG2A production antibodies all being indicative of a reduced Th2 response. These effects were further increased in mice treated with the Fes p1 producing *Lactobacillus* which shows accumulation of serum Fes p1 specific IgA. Previous studies have shown that *L. plantarum* acts in a Th1 adjuvant manner (Repa et al., 2003; Rigaux et al., 2009), we found similar effects in Th1-mediated responses of *L. plantarum* bearing recombinant allergen.

Importantly, we detected significant increase in allergen specific IgA levels in mice treated with *L. plantarum* carrying the surface-exposed allergen (pLpFesp1CWA). Serum levels of allergen-specific IgA are indicative of secreted IgA levels, which again are indicative of a mucosal immune response (Pabst, 2012). Interestingly, a previous study on the use *L. plantarum* engineered to intracellularly produce the major birch pollen allergen Bet v1 also showed allergen-specific secretory IgA responses after treatment with the allergen-producing strain only (Daniel et al., 2006).

Eosinophilia is an important effect of eosinophil-associated disorders, such as asthma and allergy (Fulkerson and Rothenberg, 2013). Oral application of *L. plantarum*/pEV and *L. plantarum*/pLpFesp1CWA led to a clear reduction of eosinophilia in the peripheral blood of the experimental mice (Fig. 4F), thus providing clear evidence of the beneficial effect of the bacterial treatment. As eosinophilia is a consequence of increases in IL-5 production, a marker of a Th2 response, a decrease in eosinophilia shows amelioration of the pathological Th2 response.

In conclusion, our data show that the use of recombinant *L. plantarum* WCFS1 as a delivery vehicle for surface-displayed, covalently anchored antigens represents a promising strategy for antigen delivery in general and allergy treatment. The *L. plantarum* WCFS1 does not colonize the gastrointestinal tract but it can be found in the human intestines for approximately 1 week after the application of a ten times smaller dose than the dose used in this study (Vesa et al., 2000). Several parameters indicate that in the current experimental setting, oral application of *L. plantarum* cells alone, even without the specific antigen, has a beneficial therapeutic effect, which is not unexpected considering the results of previous studies (Kalliomäki et al., 2010; Snel et al., 2011). However, the data showed that surface display of the Fes p1 antigen potentiates this phenomenon as evidenced by an increase in specific IgA production. As therapeutic oral administration of lactic acid bacteria is cost effective and safe, and since humans consume probiotic *Lactobacilli* as part of their normal nutrition, it would be interesting to carry out human trials with the type of *Lactobacillus* strains engineered in this study.

Author contributions

RM, MGJ, VGHE, GM and LD conceived and designed the experiments. RM, IZ and GM performed the experiments. RM, VP, LD and GM analyzed the data. MGJ, GM, VGHE and LD contributed reagents/materials/analysis tools. RM, GM and VE wrote the paper.

Conflict of interest

The authors have declared no conflict of interest.

Uncited references

Chacko and Appukuttan (2003).

Acknowledgements

This work was supported by The Research Council of Norway, project numbers 196363 & 202845, and the Ministry of Education, Science and Technological Development of the Republic of Serbia, project number 172049. *Festuca pratensis* pollen was a kind gift from Dr. Lidija Burazer from the Institute of Virology, Vaccines and Sera, Torlak, Belgrade, Serbia.

References

- Aukrust, T.W., Brurberg, M.B., Nes, I.F., 1995. Transformation of *Lactobacillus* by electroporation. *Methods Mol. Biol.* 47, 201–208.
- Borrell, M., Tyler, B.F., Morgan, W.G., 1976. Studies in *Festuca*. 7. Chromosome atlas. 2. Appraisal of chromosome race distribution and ecology, including *Festuca pratensis* Var *apennina* (De Not) Hack - tetraploid. *Cytologia (Tokyo)* 41, 219–236.
- Bousquet, J., Lockey, R., Malling, H.J., 1998. Allergen immunotherapy: therapeutic vaccines for allergic diseases. A WHO position paper. *J. Allergy Clin. Immunol.* 102, 558–562.
- Calderon, M.A., Simons, F.E., Malling, H.J., Lockey, R.F., Moingeon, P., Demoly, P., 2012. Sublingual allergen immunotherapy: mode of action and its relationship with the safety profile. *Allergy* 67, 302–311.
- Chacko, B.K., Appukuttan, P.S., 2003. Dextran-binding human plasma antibody recognizes bacterial and yeast antigens and is inhibited by glucose concentrations reached in diabetic sera. *Mol. Immunol.* 39, 933–939.
- Chang, Y.C., Lin, C.C., Hsu, C.H., 2006. Inhibition of allergen-induced airway inflammation and hyperactivity by recombinant lactic-acid bacteria. *Vaccine* 24, 5931–5936.
- Daniel, C., Repa, A., Wild, C., Pollak, A., Pot, B., Breiteneder, H., Wiedermann, U., Mercenier, A., 2006. Modulation of allergic immune responses by mucosal application of recombinant lactic acid bacteria producing the major birch pollen allergen Bet v 1. *Allergy* 61, 812–819.
- Fredriksen, L., Kleiveland, C.R., Olsen Hult, L.T., Lea, T., Nygaard, C.S., Eijsink, V.G., Mathiesen, G., 2012. Surface display of N-terminally anchored invasins by *Lactobacillus plantarum* activates NF- κ B in monocytes. *Appl. Environ. Microbiol.* 78, 5864–5871.
- Fredriksen, L., Mathiesen, G., Sioud, M., Eijsink, V.G.H., 2010. Cell wall anchoring of the 37-Kilodalton oncofetal antigen by *Lactobacillus plantarum* for mucosal cancer vaccine delivery. *Appl. Environ. Microbiol.* 76, 7359–7362.
- Fulkerson, P.C., Rothenberg, M.E., 2013. Targeting eosinophils in allergy, inflammation and beyond. *Nat. Rev. Drug Discov.* 12, 117–129.
- Halbmayer, E., Mathiesen, G., Nguyen, T.H., Maischberger, T., Peterbauer, C.K., Eijsink, V.G., Haltrich, D., 2008. High-level expression of recombinant β -galactosidases in *Lactobacillus plantarum* and *Lactobacillus sakei* using a Sakacin P-based expression system. *J. Agric. Food Chem.* 56, 4710–4719.
- Hatzler, L., Panetta, V., Lau, S., Wagner, P., Bergmann, R.L., Illi, S., Bergmann, K.E., Keil, T., Hofmaier, S., Rohrbach, A., Bauer, C.P., Hoffman, U., Forster, J., Zepp, F., Schuster, A., Wahn, U., Matricardi, P.M., 2012. Molecular spreading and predictive value of preclinical IgE response to *Phleum pratense* in children with hay fever. *J. Allergy Clin. Immunol.* 130, 894–901, e895.
- Isolauri, E., Kirjavainen, P.V., Salminen, S., 2002. Probiotics: a role in the treatment of intestinal infection and inflammation? *Gut* 50 III54–59.
- Kalliomäki, M., Antoine, J.-M., Herz, U., Rijkers, G.T., Wells, J.M., Mercenier, A., 2010. Guidance for substantiating the evidence for beneficial effects of probiotics: prevention and management of allergic diseases by probiotics. *J. Nutr.* 140, 713S–721S.
- Kalliomäki, M., Salminen, S., Poussa, T., Arvilommi, H., Isolauri, E., 2003. Probiotics and prevention of atopic disease: 4-year follow-up of a randomised placebo-controlled trial. *Lancet* 361, 1869–1871.
- Kleerebezem, M., Boekhorst, J., van Kranenburg, R., Molenaar, D., Kuipers, O.P., Leer, R., Turchini, R., Peters, S.A., Sandbrink, H.M., Fiers, M.W.E.J., Stiekema, W., Lankhorst, R.M.K., Bron, P.A., Hoffer, S.M., Groot, M.N.N., Kerkhoven, R., de Vries, M., Ursing, B., de Vos, W.M., Siezen, R.J., 2003. Complete genome sequence of *Lactobacillus plantarum* WCFS1. *Proc. Natl. Acad. Sci. U.S.A.* 100, 1990–1995.
- Kruisselbrink, A., Den Bak-Glashouwer, M.J.H., Havenith, C.E.G., Thole, J.E.R., Janssen, R., 2001. Recombinant *Lactobacillus plantarum* inhibits house dust mite-specific T-cell responses. *Clin. Exp. Immunol.* 126, 2–8.
- Laffer, S., Valenta, R., Vrtala, S., Susani, M., van Ree, R., Kraft, D., Scheiner, O., Duchene, M., 1994. Complementary DNA cloning of the major allergen Phl p I from timothy grass (*Phleum pratense*); recombinant Phl p I inhibits IgE binding to group I allergens from eight different grass species. *J. Allergy Clin. Immunol.* 94, 689–698.
- Marco, M.L., Peters, T.H., Bongers, R.S., Molenaar, D., van Hemert, S., Sonnenburg, J.L., Gordon, J.L., Kleerebezem, M., 2009. Lifestyle of *Lactobacillus plantarum* in the mouse caecum. *Environ. Microbiol.* 11, 2747–2757.
- Mathiesen, G., Sveen, A., Brurberg, M.B., Fredriksen, L., Axelsson, L., Eijsink, V.G., 2009. Genome-wide analysis of signal peptide functionality in *Lactobacillus plantarum* WCFS1. *BMC Genomics* 10, 425.
- Mathiesen, G., Sveen, A., Piard, J.C., Axelsson, L., Eijsink, V.G.H., 2008. Heterologous protein secretion by *Lactobacillus plantarum* using homologous signal peptides. *J. Appl. Microbiol.* 105, 215–226.

- 432 Pabst, O., 2012. New concepts in the generation and functions of IgA. Nat. Rev.
433 Immunol. 12, 821–832.
- 434 Petersen, A., Becker, W.M., Moll, H., Blumke, M., Schlaak, M., 1995. Studies on the car-
435 bohydrate moieties of the timothy grass pollen allergen Phl p 1. Electrophoresis
436 16, 869–875.
- 437 Repa, A., Grangette, C., Daniel, C., Hochreiter, R., Hoffmann-Sommergruber, K., Thal-
438 hamer, J., Kraft, D., Breiteneder, H., Mercenier, A., Wiedermann, U., 2003. Mucosal
439 co-application of lactic acid bacteria and allergen induces counter-regulatory
440 immune responses in a murine model of birch pollen allergy. Vaccine 22, 87–95.
- 441 Rigaux, P., Daniel, C., Hisbergues, M., Muraille, E., Hols, P., Pot, B., Pestel, J., Jacquet,
442 A., 2009. Immunomodulatory properties of *Lactobacillus plantarum* and its use
443 as a recombinant vaccine against mite allergy. Allergy 64, 406–414.
- 444 Sazawal, S., Hiremath, G., Dhingra, U., Malik, P., Deb, S., Black, R.E., 2006. Efficacy
445 of probiotics in prevention of acute diarrhoea: a meta-analysis of masked, ran-
446 domised, placebo-controlled trials. Lancet Infect. Dis. 6, 374–382.
- 447 Schwarzer, M., Repa, A., Daniel, C., Schabussova, I., Hrnčir, T., Pot, B., Stepankova,
448 R., Hudcovic, T., Pollak, A., Tlaskalova-Hogenova, H., Wiedermann, U., Kozakova,
449 H., 2011. Neonatal colonization of mice with *Lactobacillus plantarum* producing
450 the aeroallergen Bet v 1 biases towards Th1 and T-regulatory responses upon
451 systemic sensitization. Allergy 66, 368–375.
- 452 Snel, J., Vissers, Y.M., Smit, B.A., Jongen, J.M., van der Meulen, E.T., Zwijsen, R.,
Ruinemans-Koerts, J., Jansen, A.P., Kleerebezem, M., Savelkoul, H.F., 2011.
Strain-specific immunomodulatory effects of *Lactobacillus plantarum* strains
on birch-pollen-allergic subjects out of season. Clin. Exp. Allergy 41, 232–
242.
- Sørvig, E., Grönqvist, S., Naterstad, K., Mathiesen, G., Eijsink, V.G.H., Axelsson, L.,
2003. Construction of vectors for inducible gene expression in *Lactobacillus sakei*
and *L. plantarum*. FEMS Microbiol. Lett. 229, 119–126.
- Sørvig, E., Mathiesen, G., Naterstad, K., Eijsink, V.G.H., Axelsson, L., 2005. High-
level, inducible gene expression in *Lactobacillus sakei* and *Lactobacillus plantarum*
using versatile expression vectors. Microbiology 151, 2439–2449.
- van Baarlen, P., Troost, F.J., van Hemert, S., van der Meer, C., de Vos, W.M., de Groot,
P.J., Hooiveld, G.J.E.J., Brummer, R.-J.M., Kleerebezem, M., 2009. Differential NF-
kB pathways induction by *Lactobacillus plantarum* in the duodenum of healthy
humans correlating with immune tolerance. Proc. Natl. Acad. Sci. U.S.A. 106,
2371–2376.
- van Hemert, S., Meijerink, M., Molenaar, D., Bron, P., de Vos, P., Kleerebezem, M.,
Wells, J., Marco, M., 2010. Identification of *Lactobacillus plantarum* genes modu-
lating the cytokine response of human peripheral blood mononuclear cells. BMC
Microbiol. 10, 293.
- Vesa, T., Pochart, P., Marteau, P., 2000. Pharmacokinetics of *Lactobacillus plantarum*
NCIMB 8826, *Lactobacillus fermentum* KLD, and *Lactococcus lactis* MG 1363 in the
human gastrointestinal tract. Aliment Pharmacol. Ther. 14, 823–828.