

Aeroallergen-induced IL-33 predisposes to respiratory virus-induced asthma by dampening antiviral immunity

Jason P. Lynch, PhD,^{a,*} Rhiannon B. Werder, B Biomed Sci (Hons),^{a,*} Jennifer Simpson, B Biomed Sci (Hons),^a Zhixuan Loh, B Biomed Sci (Hons),^a Vivian Zhang, PhD,^a Ashraf Haque, PhD,^b Kirsten Spann, PhD,^{c,d} Peter D. Sly, MD, PhD,^{d,e} Stuart B. Mazzone, PhD,^a John W. Upham, MD, PhD,^{d,f} and Simon Phipps, PhD^{a,d} Brisbane, Australia

Background: Frequent viral lower respiratory infections in early life are an independent risk factor for asthma onset. This risk and the development of persistent asthma are significantly greater in children who later become sensitized.

Objective: We sought to elucidate the pathogenic processes that underlie the synergistic interplay between allergen exposures and viral infections.

Methods: Mice were inoculated with a murine-specific *Pneumovirus* species (pneumonia virus of mice [PVM]) and exposed to low-dose cockroach extract (CRE) in early and later life, and airway inflammation, remodeling, and hyperreactivity assessed. Mice were treated with anti-IL-33 or apyrase to neutralize or block IL-33 release.

Results: PVM infection or CRE exposure alone did not induce disease, whereas PVM/CRE coexposure acted synergistically to induce the hallmark features of asthma. CRE exposure during viral infection in early life induced a biphasic IL-33 response and impaired IFN- α and IFN- λ production, which in turn increased epithelial viral burden, airway smooth muscle growth, and type 2 inflammation. These features were ameliorated when CRE-induced IL-33 release was blocked or neutralized, whereas substitution of CRE with exogenous IL-33 recapitulated the phenotype observed in PVM/CRE-coexposed mice.

Mechanistically, IL-33 downregulated viperin and interferon regulatory factor 7 gene expression and rapidly degraded IL-1 receptor-associated kinase 1 expression in plasmacytoid dendritic cells both *in vivo* and *in vitro*, leading to Toll-like receptor 7 hyporesponsiveness and impaired IFN- α production.

Conclusion: We identified a hitherto unrecognized function of IL-33 as a potent suppressor of innate antiviral immunity and demonstrate that IL-33 contributes significantly to the synergistic interplay between respiratory virus and allergen exposures in the onset and progression of asthma. (J Allergy Clin Immunol 2016;■■■:■■■-■■■.)

Key words: *Pneumonia virus of mice, IL-33, type 2 innate lymphoid cell, antiviral, respiratory syncytial virus, interferon, plasmacytoid dendritic cell*

Pathologically, asthma is characterized by airway inflammation and airway remodeling; structural changes including goblet cell metaplasia, increased deposition of extracellular matrix proteins, and increased airway smooth muscle (ASM) mass. These features collectively contribute to airway narrowing, loss in lung function, and airway hyperreactivity (AHR) and are poorly responsive to conventional therapies.^{1,2} The inflammatory response is typically of a type 2 cytokine profile, which promotes the recruitment and survival of key effector cells, such as eosinophils and mast cells, and can induce features of airway remodeling through direct activation of airway epithelial and mesenchymal cells.^{3,4} Both CD4⁺ T_H2 cells and type 2 innate lymphoid cells (ILCs) produce the type 2 cytokines IL-5 and IL-13⁵ in response to the tissue alarmin IL-33 and work collaboratively to affect their teleological role, namely antihelminthic immunity.⁶ However, why type 2 immunity is increased during acute exacerbations of asthma, most commonly triggered by a viral respiratory tract infection, remains less clear.

Epidemiologic studies have reproducibly demonstrated that severe/frequent viral lower respiratory tract infections (vLRTIs) in early life are a major independent risk factor for asthma inception.⁷⁻⁹ Notably, this association is markedly increased in children who later become sensitized to food allergens or aeroallergens and, significantly, confers the greatest risk for progression to asthma that is severe, persistent, or both.¹⁰⁻¹² The underlying mechanisms remain poorly defined, although it is likely that the synergistic interaction between allergen exposures and viral respiratory tract infection is underpinned by genetic defects, functional defects, or both in a shared antiviral immune pathway necessary for viral control in early life.¹¹ Phenotypically, this can manifest as a deficiency in production of the antiviral cytokines IFN- α , IFN- β , and IFN- λ by leukocytes and airway epithelial cells (AECs),¹³⁻¹⁵ although whether such defects are virus specific, genetic, or acquired (eg, as a consequence of a type 2 inflammatory milieu) remains highly contentious.¹⁶⁻¹⁸

From ^athe School of Biomedical Sciences, ^dthe Australian Infectious Diseases Research Centre, ^eCentre for Children's Health Research, University of Queensland; ^bQIMR Berghofer Medical Institute; ^cthe School of Biomedical Sciences, Queensland University of Technology; and the ^fLung and Allergy Research Centre, School of Medicine, University of Queensland, Princess Alexandra Hospital.

*These authors contributed equally to this work.

Supported by an equipment grant (Rebecca L. Cooper Medical Research Foundation), an Australian Infectious Disease Research Excellence Award awarded (to S.P., J.P.L., and J.W.U.) and an Australian Research Council Future Fellowship (to S.P.).

Disclosure of potential conflict of interest: A. Haque has received a grant from the National Health and Medical Research Council (1028641). J. W. Upham is on advisory boards for Menarini Pharmaceuticals, Novartis, AstraZeneca, and GlaxoSmithKline. S. Phipps has received a grant from the National Health and Medical Research Council Australia and Pfizer. The rest of the authors declare that they have no relevant conflicts of interest.

Received for publication July 10, 2015; revised January 25, 2016; accepted for publication February 9, 2016.

Corresponding author: Simon Phipps, PhD, School of Biomedical Sciences, University of Queensland, St Lucia, Queensland 4072, Australia. E-mail: s.phipps@uq.edu.au. 0091-6749/\$36.00

© 2016 American Academy of Allergy, Asthma & Immunology
<http://dx.doi.org/10.1016/j.jaci.2016.02.039>

Abbreviations used

AEC: Airway epithelial cell
 AHR: Airway hyperreactivity
 ASM: Airway smooth muscle
 BALF: Bronchoalveolar lavage fluid
 CRE: Cockroach extract
 HDM: House dust mite
 ILC: Innate lymphoid cell
 IL1RL1: IL-1 receptor-like 1
 IRAK: IL-1 receptor-associated kinase 1
 ISG: Interferon-stimulated gene
 pDC: Plasmacytoid dendritic cell
 pfu: Plaque-forming unit
 PVM: Pneumonia virus of mice
 RSV: Respiratory syncytial virus
 TLR: Toll-like receptor
 vLRI: Viral lower respiratory tract infection

Plasmacytoid dendritic cells (pDCs) are among the first cells to respond to a viral infection, sensing viral RNA and producing vast amounts of IFN- α/β downstream of Toll-like receptor (TLR) 7 activation.^{19,20} Defects in pDC and TLR7 responsiveness are associated with bronchiolitis, infant wheeze, and childhood asthma.^{21,22} In recent studies, we have used the natural mouse pathogen pneumonia virus of mice (PVM), which, unlike human respiratory syncytial virus (RSV), replicates more readily in mice, to induce the more severe symptoms of infantile RSV bronchiolitis.^{20,23} *Tlr7* gene deletion delayed the antiviral response, causing severe viral bronchiolitis and release of the alarmin IL-33 in early life, and consequently, an asthma-like pathology developed on viral challenge in later life.^{20,24} Of note, the transfer of *Tlr7*-sufficient, but not deficient, pDCs ameliorated bronchiolitis, suggesting that perturbations to the pDC compartment in early life predispose to AEC damage and IL-33 release and hence might underlie the association between severe vLRI and asthma onset.²¹ Intriguingly, IL-33 and its receptor, IL-1 receptor-like 1 (IL1RL1, also known as ST2), are susceptibility loci for wheezing in infancy (and asthma),^{25,26} although the molecular basis for this association is not known. This led us to question whether allergen-induced release of IL-33 increases the severity of virus-induced bronchiolitis by dampening pDC-orchestrated innate antiviral immunity and to assess whether this early-life interaction causes persistent changes to host immunity that underlie disease progression in response to viral challenge, allergen challenge, or both in later life.

METHODS**Experimental procedures**

A detailed description of the materials and procedures used is provided in the [Methods](#) section in this article's Online Repository at www.jacionline.org.

Induction of virus- plus allergen-coinduced asthma and associated perturbations

Specific pathogen-free BALB/c mice or 4C13R mice²⁷ were inoculated with PVM (strain J3666; 1 plaque-forming unit [pfu] in early life and 20 pfu in later life) or cockroach extract (CRE; 1 μ g), as previously described^{20,28} and as outlined in the study design ([Fig 1, A](#)). In some experiments mice were exposed to house dust mite (HDM) extract (5 or 100 μ g), LPS (186 pg), recombinant IL-33 (10 ng), anti-IL-33/isotype control

antibody (200 μ g), apyrase (4 U/mL), or pyridoxalphosphate-6-azophenyl-2',4'-disulfonic acid (100 μ M in 10 μ L). All studies were approved by the University of Queensland Animal Ethics Committee. Sample processing is described in the [Methods](#) section in this article's Online Repository.

Flow cytometry

Lungs were digested to single-cell suspensions, as previously described,²⁴ and then incubated with anti-Fc γ RIII/II before incubation with fluorochrome-conjugated antibodies at 4°C for 30 minutes, as described in the [Methods](#) section in this article's Online Repository.

Histology and immunohistochemistry

Paraffin-embedded lung sections were prepared, as previously described,²⁰ and then stained with Chromotrope 2R, periodic acid-Schiff, Masson trichrome, anti-IL-33, anti-PVM G protein, anti- α -smooth muscle actin, or anti-periostin, as described in the [Methods](#) section in this article's Online Repository.

Measurement of protein expression

IL-33, IFN- λ 2/3, IL-12p40, IFN- γ , IL-13, and IL-5 were quantified by means of ELISA. IFN- α and IL-13 were quantified by using a cytometric bead array. IgG_{1a} was detected by using an in-house ELISA system.²⁸

Quantitative real-time PCR

Total RNA was isolated with TRI Reagent solution, followed by phenol-chloroform extraction. Reverse transcription was performed with M-MLV reverse transcriptase and random primers. Quantitative RT-PCR was performed with SYBR Green by using the primers described in [Table E1](#) in this article's Online Repository at www.jacionline.org.

Airway function assessment

Airways resistance was determined by using a forced oscillation technique in response to methacholine, as previously described.²⁹

Type 2 ILC culture

Type 2 ILCs were FACS sorted from PVM/CRE-coexposed mice at 10 dpi. Cells were cultured with IL-2 (30 ng/mL) and preincubated with IFN- α (5000 U/mL) before stimulation with IL-33 (30 ng/mL).

pDC culture


Bone marrow-derived pDCs were generated, as described previously.²⁰ pDCs were preincubated with IL-33 (3 ng/mL) and then stimulated with imiquimod (3 μ g/mL).

Statistical analyses

Data were analyzed with a Student *t* test, 1-way ANOVA with a Tukey *post hoc* test, or 2-way ANOVA with a Sidak *post hoc* test, as appropriate, by using GraphPad Prism software (version 5.0; GraphPad Software, La Jolla, Calif). A *P* value of less than .05 was considered statistically significant.

RESULTS***Pneumovirus* and allergen coexposure synergize to promote asthma onset and progression**

To interrogate the synergistic interaction between a viral respiratory tract infection and allergen exposures, we first identified a dose of CRE that by itself would not induce allergic inflammation. Because we developed a previous mouse model of



to use 1 μg of CRE for all further studies. To assess for a synergistic effect between PVM and CRE, we superimposed a low-dose PVM infection, which by itself did not induce features of type 2 inflammation (Fig 1, B-D, and see Fig E2 in this article's Online Repository at www.jacionline.org) onto the low-dose CRE exposure model (Fig 1, A). Eosinophilic inflammation, mucus hypersecretion, CRE-specific IgG_{1a} levels (Fig 1, B-D), and mucosal mast cell numbers (data not shown) were greatest in mice exposed to both PVM and CRE. By contrast, these features of type 2 inflammation were absent in mice exposed to PVM




FIG 2. Early-life CRE exposure increases viral load and dampens antiviral cytokine production. **A**, Viral load in AECs detected by using immunohistochemistry and enumerated as a percentage of total AECs. **B**, Edema in lung parenchyma. **C**, Weight gain. **D**, IFN- α , IFN- λ , and IFN- γ levels in BALF and IL-12p40 levels in lung homogenate. **E**, mRNA expression of interferon-regulatory factor 7 (*IRF7*), viperin, and signal transducer and activator of transcription 1 (*STAT1*) in lungs relative to those in vehicle-treated mice. Data are means $\pm$ SEMs and representative of 2 to 3 experiments ($n = 4$ -6 mice per group). * $P < .05$, ** $P < .01$, and *** $P < .001$ compared with vehicle-treated mice. # $P < .05$, ## $P < .01$, and ### $P < .001$ compared with PVM/CRE mice.

alone or low-dose CRE alone in both early and later life or low-dose CRE alone in later life (Fig 1, B-D, and see Fig E2). Similarly, features of airway remodeling, including collagen and periostin deposition, and ASM remodeling (Fig 1, E-G) were only evident in PVM/CRE-coexposed mice.

To compare our findings to contemporary acute models of allergen-induced disease,³⁰ we exposed a group of mice to high-dose (100 μ g) CRE in later life alone. Strikingly, this regimen elicited type 2 inflammation but was not sufficient to elicit features of airway remodeling (see Fig E2). Thus low-dose virus and low-dose allergen coexposures acted synergistically to induce all of the cardinal features of asthma.

Virus and allergen exposure in early and later life are necessary for maximal asthma severity

We next addressed the relative contribution of the virus and allergen exposures in early and later life. PVM/CRE coexposure in early life alone was not sufficient to induce type 2 inflammation or airway remodeling compared with the vehicle-treated group (see Fig E2). The importance of both the virus and allergen challenge for asthma progression was confirmed when the omission of either exposure significantly diminished airway inflammation, airway remodeling, and AHR when assessed at 76 days of age (see Fig E2). These findings led us to question the importance of the CRE exposure during primary viral infection. Remarkably, when this exposure was omitted, type 2 inflammation, airway remodeling, and AHR were completely

absent (Fig 1, final row), even though the mice had been infected with PVM in early and later life and then challenged repeatedly with CRE. Taken together, these data suggested that the CRE exposure in early life fundamentally altered the nature of the host response to secondary virus and/or allergen exposure in later life and that this effect was necessary for disease progression.

Allergen exposure increases viral load and lung tissue damage and dampens antiviral cytokine production

Aeroallergens are known to rapidly induce innate inflammation in the airways.^{28,31,32} We hypothesized that allergen-induced inflammation might perturb the host's antiviral response to PVM, thus increasing the severity of viral bronchiolitis. Viral load in the airway epithelium was significantly greater and persisted for longer in PVM/CRE-coexposed mice compared with that seen in mice inoculated with virus alone (Fig 2, A, and see Fig E3, A, in this article's Online Repository at www.jacionline.org). In children with severe RSV bronchiolitis, the virus can spread to the parenchyma³³; however, few PVM-immunoreactive cells were observed in the alveoli (data not shown). Strikingly, PVM/CRE coexposure stunted weight gain and heightened tissue edema, indicative of severe disease (Fig 2, B and C). The increased viral illness in PVM/CRE-coexposed mice was associated with significantly lower IFN- α , IFN- λ , and IL-12p40, but not IFN- γ , production in bronchoalveolar lavage fluid (BALF; Fig 2, D) and attenuated transcription of interferon-stimulated genes (ISGs) involved in TLR7 and

type I interferon receptor signaling, including *Irf7*, *Viperin*, and *Stat1* (Fig 2, E). Intriguingly, CRE exposure alone significantly downregulated the expression of *Viperin*, a component of the TLR7 signaling cascade in pDCs.³⁴ Although TLR7-mediated activation of pDCs is critical for host defense against acute PVM infection,²⁰ CRE exposure did not affect pDC numbers in the lung (see Fig E3, B).

Type 2 inflammation and ASM remodeling are increased in PVM/CRE-coexposed mice in early life

Allergen exposure is well known to induce the release of IL-33 to promote type 2 inflammation.³⁵ Analyzing IL-33 protein expression across the time course of infection revealed a significant increase in lung homogenates at 3 dpi (sampling 2 hours after CRE exposure) and 7 dpi in the coexposed mice but not in those exposed to PVM alone (Fig 3, A, upper panel). Although CRE exposure did not increase lung IL-33 expression at 3 dpi, it did increase airway luminal IL-33 levels measured in the BALF at this time, both in PVM-infected mice and noninfected mice (Fig 3, B). Consistent with other reports,³⁶ under homeostatic conditions, IL-33 was expressed predominantly in resident myeloid cell populations and alveolar (but not airway) epithelial cells (Fig 3, C, top left panel). However, by 7 dpi, IL-33-immunoreactive AECs were evident in PVM alone and PVM/CRE-exposed mice (Fig 3, C, and see Fig E3, C), which was notable as a second phase of IL-33 release occurred at 10 dpi (Fig 3, B). In contrast to IL-33, expression of IL-25 and thymic stromal lymphopoietin was unaffected by PVM exposure, CRE exposure, or both (data not shown). Critically, the IL-33 response was associated with increased numbers of type 2 ILCs and eosinophils ($CD4^+$ and $CD8^+$ T-cell numbers were unaffected, data not shown), and type 2 cytokine expression (Fig 3, D-F, and see Fig E4, A, in this article's Online Repository at www.jacionline.org). Using IL-4/IL-13 reporter mice,²⁷ we revealed that $CD8^+$, but not $CD4^+$, T cells contributed to IL-4 and IL-13 production. However, the majority of type 2 ILCs were IL-13⁺, and the proportion of IL-13-expressing type 2 ILCs, but not $CD4^+$ or $CD8^+$ T cells, increased in PVM/CRE-coexposed mice (see Fig E4, B). Similar to our observations of ASM remodeling in later life (Fig 1, G), PVM infection or CRE exposure alone had no effect on ASM growth, whereas coexposure induced a 3-fold increase in ASM area (Fig 3, G).

Increased viral load is not sufficient to induce type 2 inflammation and ASM remodeling

To address whether increased viral load was causally related to the development of ASM growth, we first inoculated mice with a 10× higher dose of PVM. Although this increased the viral load and induced severe illness and mortality (Fig 4, A and B), it was not sufficient to increase IL-33 release at 3 dpi (even when the dose was increased 100-fold [data not shown]) or induce ASM growth (Fig 4, C and D). We then assessed whether HDM was able to affect viral load and ASM growth at 7 and 10 dpi, respectively (when these pathologies peaked). Substitution of CRE with HDM significantly increased viral load similar to CRE coexposure but had no effect on ASM growth (Fig 4, E and F), which was consistent with a lack of effect on antiviral cytokine expression, type 2 ILCs in the lungs, or IL-33 release

(Fig 4, G-I). Notably, HDM induced IL-33 release was detectable at 7 dpi, which is consistent with the findings of others (data not shown).^{28,31,37-39} Because the level of endotoxin in the CRE was high (395 EU per 100 µg of protein), we coexposed mice to PVM and an equivalent dose of LPS. LPS induced a massive IFN-α response and decreased viral load (Fig 4, H and I), suggesting LPS contamination of CRE was not the cause of impaired antiviral immunity. Taken together, our findings suggested that ASM growth and the onset of type 2 inflammation was associated with an IFN-α/λ^{low}IL-33^{high} cytokine microenvironment rather than viral load.

Anti-IL-33 prevents type 2 inflammation and remodeling in response to virus and allergen coexposure in early and later life

Mice were treated with a neutralizing antibody in both early and later life to assess the contribution of IL-33 (see Fig E5, A, in this article's Online Repository at www.jacionline.org). Although mice treated with an isotype-matched control had all of the hallmark features of asthma, treatment with anti-IL-33 abolished these features (see Fig E5, B-E). Similarly, anti-IL-33 treatment in early life ablated ASM growth and the onset of type 2 inflammation (Fig 5 and see Fig E5, F). As expected, anti-IL-33 had no effect in mice infected with PVM alone.

IL-33 is a negative regulator of innate antiviral immunity

We hypothesized that CRE might dampen innate antiviral immunity through IL-33 release. Strikingly, anti-IL-33 reversed the increased viral burden, dampened IFN-α and IFN-λ production, and attenuated ISG expression caused by CRE exposure but did not alter pDC or CD8 T-cell numbers in the lung (Fig 6, A-C, and see Fig E5, G and H). In contrast, anti-IL-33 had no effect on the antiviral response in mice infected with PVM alone. CRE-induced IL-33 release occurs downstream of ATP-mediated purinergic receptor activation.³⁵ In our model treatment with apyrase (to catalyze the hydrolysis of ATP) significantly decreased IL-33 levels in BALF (2 hours after CRE exposure; see Fig E6, A, in this article's Online Repository at www.jacionline.org) and significantly decreased viral load at 7 dpi in the airway epithelium of PVM/CRE-coexposed mice (Fig 6, D). Treatment with the broad-spectrum P2R antagonist pyridoxalphosphate-6-azophenyl-2',4'-disulfonic acid had the same effect (see Fig E6, A and B), further implicating nucleoside/purinergic receptor signaling in mediating IL-33 release. As with anti-IL-33, blocking IL-33 release recapitulated IFN-α production and ISG expression in the lung (Fig 6, E and F), decreased IL-13 expression (data not shown), and ablated ASM growth (Fig 6, G), again highlighting the protective nature of an IFN-α/λ^{high}IL-33^{low} cytokine environment. This led us to question whether IFN-α suppresses IL-33-induced cytokine production by type 2 ILCs. Notably, FACS-purified type 2 ILCs expressed the type I interferon receptor (see Fig E4, A) and produced less IL-5 and IL-13 in response to IL-2/IL-33 stimulation when pretreated with IFN-α (Fig 6, H). This response was unrelated to cell death (see Fig E6, C and D); however, we observed that IFN-α attenuated IL-2/IL-33-induced upregulation of ST2 (Fig 6, I).




FIG 3. Type 2 inflammation in early life is increased in PVM/CRE-coexposed mice. **A** and **B**, IL-33 in lung homogenate (Fig 3, A) and BALF (Fig 3, B). **C**, Representative micrographs of IL-33 immunostaining plus 4',6-diamidino-2-phenylindole (DAPI) counterstaining. Scale bar = 50 μm . **D**, Type 2 ILCs in lung (Lineage⁻CD45⁺CD90.2⁺CD25⁺ST2⁺). **E**, Peribronchial eosinophils. **F**, IL-5 and IL-13 in BALF. **G**, ASM area. BM, Basement membrane. Data are means $\pm$ SEMs and representative of 2 independent experiments ($n = 6$ -8 mice per group). * $P < .05$, ** $P < .01$, and *** $P < .001$ compared with vehicle-treated mice. ## $P < .05$, ### $P < .01$, and #### $P < .001$ compared with PVM/CRE mice.




FIG 4. High viral load alone does not promote type 2 inflammation and airway remodeling. **A** and **E**, Viral load in AECs. **B**, Survival curve after intranasal infection with 1 or 10 pfu of PVM. **C** and **G**, IL-33 in BALF. **D** and **F**, ASM area. **H**, IFN- α , IFN- λ , and IFN- γ levels in BALF and IL-12p40 levels in lung homogenate. **I**, Type 2 ILCs in lung. **BM**, Basement membrane. Data are means $\pm$ SEMs and representative of 2 independent experiments ($n = 6-8$ mice per group). * $P < .05$, ** $P < .01$, and *** $P < .001$ compared with PVM-treated mice. # $P < .05$, ## $P < .01$, and ### $P < .001$ compared with PVM/CRE mice.

Exogenous IL-33 increases viral load and decreases IL-1 receptor–associated kinase 1 expression and antiviral cytokine production by pDCs

We next examined the effect of low-dose exogenous IL-33 during acute PVM infection. Similar to CRE, exposure to exogenous IL-33 (10 ng at 3, 4, and 5 dpi; intranasal route) during acute PVM infection did not affect the infiltration of pDCs (see Fig E7, A and B, in this article's Online Repository at www.jacionline.org); however, it significantly diminished IFN- α and IFN- λ levels in the BALF, increased epithelial viral load, and ablated ISG expression in the lung (Fig 7, A-D). Moreover, IL-33/PVM coexposure significantly increased ASM growth in contrast to IL-33 or PVM alone (Fig 7, E). Lastly, we questioned whether IL-33 affects TLR7-mediated activation of pDCs by rapidly

depleting the intracellular adaptor molecules IL-1 receptor–associated kinase 1 (IRAK1) and viperin.^{40,41} Consistent with this possibility, ST2 was expressed on pDCs in bone marrow, lung, and mediastinal lymph nodes (Fig 7, F). Notably, in PVM/CRE-coexposed mice, intracellular IRAK1 and viperin expression in pDCs was significantly diminished within 2 hours of CRE administration (Fig 7, G). To directly assess the role of IL-33, we stimulated pDCs *in vitro* with IL-33. As shown *in vivo*, IRAK1 expression by pDCs was rapidly decreased (Fig 7, H). Additionally, IL-33 treatment significantly diminished the production of IFN- α (Fig 7, I) and IL-12p40 (data not shown) in response to TLR7 stimulation. Collectively, these data suggest that IL-33 negatively regulates the early antiviral response by inducing a state of TLR7 hyporesponsiveness in pDCs.




FIG 5. Anti-IL-33 prevents type 2 inflammation and remodeling in response to PVM and CRE coexposure in early life. **A**, Type 2 ILCs in lung. **B**, IL-13 in BALF. **C**, Peribronchial eosinophils. **D**, ASM area. Data are means $\pm$ SEMs and representative of 2 experiments ($n = 6$ -7 mice per group). *BM*, Basement membrane. * $P < .05$, ** $P < .01$, and *** $P < .001$ compared with isotype-treated PVM/CRE-coexposed mice.

DISCUSSION

IL-33 and its receptor, ST2, have been reproducibly identified in genetic studies as asthma susceptibility loci, and clinical and experimental studies have revealed a preeminent role for IL-33 in the development and expansion of T_H2 immunity through activation of $CD4^+$ T_H2 cells, type 2 ILCs, and various type 2 effector cells. In the present study we extend this paradigm to show that IL-33 dampens antiviral cytokine production, thus removing an inhibitory tonic and establishing a cytokine microenvironment that is conducive for the expansion of type 2 inflammation and ASM growth. Critically, using a novel preclinical model of asthma, we demonstrate that this dual activity of IL-33 underlies the synergistic effect of respiratory virus and allergen exposures on the development and progression of asthma.

We repeatedly inoculated mice with low-dose virus and allergen to better simulate the natural course of events and demonstrated that each of these environmental exposures in both early and later life was necessary to maximally induce all of the cardinal features of asthma. Several investigators have developed experimental mouse models of virus/allergen coexposure to show that a viral respiratory tract infection facilitates sensitization to an otherwise innocuous allergen; however, in these models, as in ours, allergic sensitization was not apparent until after allergen challenge. We note that analysis from one birth cohort study found that allergic sensitization precedes rhinovirus-induced wheezing.⁴² However, it is important to note that although rhinovirus-associated illness occurs in older children, severe RSV bronchiolitis often occurs in the first year of life, when aeroallergen-specific IgE titers remain low.¹⁰ Because an antibody-mediated response did not account for the predisposing effect of CRE in early life, we questioned whether the primary allergen exposure altered the course of the viral respiratory tract infection. Unexpectedly, we found that CRE exposure attenuated antiviral cytokine production and doubled the viral load in the airway epithelium. Treatment of PVM/CRE-coexposed mice with anti-IL-33 or apyrase restored IFN- α production and decreased the viral burden to levels observed with PVM alone, whereas substitution of CRE with exogenous IL-33 attenuated IFN- α production and increased viral load, implicating IL-33 as a potent suppressor of antiviral immunity. Mechanistically, we found that IRAK1 and viperin expression by pDCs was lower *in vivo* in PVM/CRE-coexposed mice and IRAK1 expression

was lower *in vitro* after IL-33 treatment. Significantly, IL-33 impaired IFN- α production by pDCs, which is consistent with our earlier report that TLR7 hyporesponsiveness is a feature of asthma.²² Additionally, CRE or IL-33 exposure decreased *Viperin* and *Irf7* gene expression *in vivo*, whereas anti-IL-33 or apyrase treatment attenuated this downregulation. Because viperin interacts with IRAK1 to induce the nuclear translocation of IRF7 and downstream interferon production by pDCs,³⁴ our findings illustrate that IL-33 negatively regulates TLR7 signaling by at least 3 key points: viperin, IRAK1, and IRF7.

Presently, the role of type I and III interferons in asthma pathogenesis is highly contentious. The first reports of impaired IFN- β and IFN- λ production by *ex vivo* virus-stimulated AECs from asthmatic patients^{13,14} have not been universally replicated by other investigators,¹⁶⁻¹⁸ leading to suggestions that the phenotype is only present in a subpopulation of patients or might stem from the cytokine milieu.⁴³ In light of our findings, these concepts might not be mutually exclusive because the IL-33 receptor IL1RL1, a common susceptibility locus for asthma,²⁵ is expressed on AECs⁴⁴ and rhinovirus-infected AECs secrete IL-33.⁴⁵ Accordingly, it will be important to test whether the increased viral load in nasal lavage fluid, as reported by Jackson et al,⁴⁵ is related to the suppressive effects of IL-33 on antiviral immunity.

The phenotype of our early-life model shows a number of striking similarities to pediatric patients with severe therapy-resistant asthma, including impaired type I and III interferon production, increased collagen deposition, and eosinophilic inflammation.⁴⁶⁻⁴⁸ It remains to be established whether the cause of disease in this patient group relates to severe/frequent vLRIs; however, it is noteworthy that they present with increased IL-33 expression⁴⁷ and therefore have an IFN- $\alpha/\lambda^{\text{low}}$ IL-33 $^{\text{high}}$ cytokine microenvironment, as observed in our model. We and others have shown that type I and III interferons suppress $CD4^+$ T_H2 cytokine production,⁴⁹ and here we show that IFN- α suppressed IL-33-induced type 2 cytokine production by type 2 ILCs, in part by preventing ST2 upregulation. Our findings highlight an important counterregulatory process whereby IL-33 suppresses antiviral cytokine production. Such systems are critically dependent on the timing of expression, emphasizing the need to tightly regulate the release of prestored IL-33. Evidently, if IL-33 release is triggered inadvertently, as we show here in response to CRE exposure, this can have profound short- and long-term effects, which




FIG 6. IL-33 blockade in PVM/CRE-coexposed mice prevents excessive viral load and reverses dampened antiviral immunity induced by CRE. **A** and **D**, AEC viral load. **B** and **E**, IFN- α and IFN- λ levels in BALF and IL-12p40 levels in lung homogenate. **C** and **F**, mRNA expression of ISGs relative to vehicle-treated mice. **G**, ASM area. **H**, Type 2 ILCs were treated with IL-2 with or without IFN- α for 30 minutes, followed by culture with IL-33 and IL-2 for 72 hours, before the supernatant was probed for IL-5 and IL-13. **I**, ST2 expression on cultured type 2 ILCs. *MFI*, Mean fluorescence intensity. Data are means $\pm$ SEMs and representative of 2 to 3 independent experiments ($n = 6$ -10 mice per group). *BM*, Basement membrane. * $P < .05$, ** $P < .01$, and *** $P < .001$ compared with PVM/CRE/isotype mice or as indicated.




FIG 7. Exogenous IL-33 dampens IFN- α production and increases viral load by decreasing IRAK1 expression and antiviral cytokine production by pDCs. **A** and **B**, IFN- α (Fig 7, **A**) and IFN- λ (Fig 7, **B**) in BALF. **C**, Viral load in AECs. **D**, mRNA expression relative to vehicle-treated mice. **E**, ASM area. **F**, ST2 staining of bone marrow (BM), lung, and mediastinal lymph node (LN) pDCs (solid line). Fluorescence minus one for ST2 staining is shown in gray. **G**, IRAK1 and viperin intracellular expression in pDC *in vivo* at 2 hours after CRE administration. **H**, Mean fluorescence intensity. **H**, Intracellular IRAK1 staining of bone marrow-derived pDCs preincubated with vehicle (solid line) or IL-33 (3 ng/mL, dotted line) for 0.5 hours. Fluorescence minus one for IRAK staining is shown in gray. **I**, IFN- α production in bone marrow-derived pDC cell-culture supernatants. BM, Basement membrane. Data are means $\pm$ SEMs and representative of 2 independent experiments ($n = 7$ mice per group). n.d., Not detected. * $P < .05$, ** $P < .01$, and *** $P < .001$ compared with mice exposed to PVM alone or vehicle-treated bone marrow-derived pDCs.

can be exacerbated in subjects with a gain-of-function single nucleotide polymorphism in the IL-33 or IL1RL1 gene.⁵⁰ The importance of timing was further emphasized when we switched CRE with HDM; although HDM did induce IL-33, as others have shown, the response was too slow to affect the antiviral response. Thus we predict that exposure to HDM at the time of inoculation would predispose to viral bronchiolitis and subsequent asthma.

In the naive mouse lung IL-33 is primarily expressed by alveolar epithelial cells; however, in response to ovalbumin challenge, several different types of hematopoietic cells also produce IL-33.³⁶ In our model PVM/CRE coexposure of neonatal mice induced IL-33 release in a biphasic manner, peaking at 3 dpi (ie, 2 hours after CRE exposure) and then at 10 dpi. Notably, CRE

alone induced IL-33 release at 3 dpi but not 10 dpi, PVM infection alone induced IL-33 release at 10 dpi only, while in coexposed mice, expression was significantly increased at both time points. PVM infection increased IL-33 expression in AECs, although not until 7 dpi, implicating infiltrating inflammatory cells or alveolar epithelial cells as the source of IL-33 at 3 dpi. This release of IL-33 was ATP dependent³⁵; however, the late release, which was associated with PVM infection, might involve a separate mechanism. Indeed, recent reports have shown that the release of active IL-33 can also occur downstream of necroptosis, a programmed form of necrotic cell death, that can be initiated by viral activation of TLR3,⁵¹ which is consistent with the apparent association between late IL-33 release and viral load in our model at 10 dpi.

A striking feature of our model was the development of ASM growth, which to our knowledge has not been observed previously in an experimental model of virus-induced asthma. Furthermore, we demonstrated that this hallmark pathology of asthma began to develop in early life, which is consistent with a recent clinical report in which ASM changes were evident in preschool-aged children with wheeze who are later given a diagnosis of asthma.⁵² In fact, evidence of ASM proliferation is infrequent in patients with established asthma,⁵³ emphasizing the need for new therapies to target prevention in early life, rather than resolution of ASM mass in later life. Neutralization or blockade of IL-33 release significantly decreased the development of ASM remodeling, together with other features of asthma, in both early and later life. IL-33 has been shown to directly induce collagen synthesis from fibroblasts⁴⁷; however, ASM cells do not express ST2,⁴⁴ and hence it is likely that the effects of IL-33 on ASM growth in our model were indirect, perhaps being mediated through activation of type 2 ILCs, eosinophils, or both.

In summary, we have developed a novel model to study the molecular processes that underlie the synergistic relationship between vLRI and allergen exposure and the onset of asthma. CRE exposure rapidly induces the release of IL-33, which downregulates components of the TLR7 signaling pathway causing TLR7 hyporesponsiveness in pDCs. The ensuing IFN- α/γ ^{low}IL-33^{high} cytokine microenvironment allows for expansion of type 2 inflammation and increased ASM growth in early life. This in turn leads to persistent alterations to resident airway cells, immune cells, or both, which are necessary for disease progression after challenge with virus and allergen in later life. Thus emerging therapies aimed at targeting IL-33 will not only decrease T_H2 inflammation but also likely boost innate antiviral immunity.

We thank Drs Nocka, Kasaian, and Bloom (Pfizer) for supplying anti-IL-33. We also thank Dr Ashik Ullah (Queensland Institute of Medical Research) and Dr Ben Roediger (Centenary Institute) for technical assistance.

Key messages

- Aeroallergen-induced IL-33 release suppresses antiviral immunity, increasing the severity of viral bronchiolitis.
- IL-33 impairs antiviral immunity by degrading IRAK1 and viperin in pDCs, leading to a state of TLR7 hyporesponsiveness.
- Coexposure to virus and allergen in early life predisposes to asthma progression in later life.

REFERENCES

- James AL, Pare PD, Hogg JC. The mechanics of airway narrowing in asthma. *Am Rev Respir Dis* 1989;139:242-6.
- Lambert RK, Wiggs BR, Kuwano K, Hogg JC, Pare PD. Functional significance of increased airway smooth muscle in asthma and COPD. *J Appl Physiol* 1993; 74:2771-81.
- Holgate ST, Davies DE, Lackie PM, Wilson SJ, Puddicombe SM, Lordan JL. Epithelial-mesenchymal interactions in the pathogenesis of asthma. *J Allergy Clin Immunol* 2000;105:193-204.
- Fichtner-Feigl S, Strober W, Kawakami K, Puri RK, Kitani A. IL-13 signaling through the IL-13 α 2 receptor is involved in induction of TGF- β 1 production and fibrosis. *Nat Med* 2006;12:99-106.
- Neill DR, Wong SH, Bellosi A, Flynn RJ, Daly M, Langford TK, et al. Nuocytes represent a new innate effector leukocyte that mediates type-2 immunity. *Nature* 2010;464:1367-70.
- McKenzie Andrew NJ, Spits H, Eberl G. Innate lymphoid cells in inflammation and immunity. *Immunity* 2014;41:366-74.
- Stein RT, Sherrill D, Morgan WJ, Holberg CJ, Halonen M, Taussig LM, et al. Respiratory syncytial virus in early life and risk of wheeze and allergy by age 13 years. *Lancet* 1999;354:541-5.
- James KM, Gebretsadik T, Escobar GJ, Wu P, Carroll KN, Li SX, et al. Risk of childhood asthma following infant bronchiolitis during the respiratory syncytial virus season. *J Allergy Clin Immunol* 2013;132:227-9.
- Sigurs N, Bjarnason R, Sigurbergsson F, Kjellman B. Respiratory syncytial virus bronchiolitis in infancy is an important risk factor for asthma and allergy at age 7. *Am J Respir Crit Care Med* 2000;161:1501-7.
- Holt PG, Rowe J, Kusel M, Parsons F, Hollams EM, Bosco A, et al. Toward improved prediction of risk for atopy and asthma among preschoolers: a prospective cohort study. *J Allergy Clin Immunol* 2010;125:653-9.
- Holt PG, Sly PD. Viral infections and atopy in asthma pathogenesis: new rationales for asthma prevention and treatment. *Nat Med* 2012;18:726-35.
- Oddy WH, de Klerk NH, Sly PD, Holt PG. The effects of respiratory infections, atopy, and breastfeeding on childhood asthma. *Eur Respir J* 2002; 19:899-905.
- Contoli M, Message SD, Laza-Stanca V, Edwards MR, Wark PA, Bartlett NW, et al. Role of deficient type III interferon-lambda production in asthma exacerbations. *Nat Med* 2006;12:1023-6.
- Wark PA, Johnston SL, Bucchieri F, Powell R, Puddicombe S, Laza-Stanca V, et al. Asthmatic bronchial epithelial cells have a deficient innate immune response to infection with rhinovirus. *J Exp Med* 2005;201:937-47.
- Gill MA, Bajwa G, George TA, Dong CC, Dougherty II, Jiang N, et al. Counterregulation between the TLR7 pathway and antiviral responses in human plasmacytoid dendritic cells. *J Immunol* 2010;184:5999-6006.
- Spann KM, Baturcam E, Schagen J, Jones C, Straub CP, Preston FM, et al. Viral and host factors determine innate immune responses in airway epithelial cells from children with wheeze and atopy. *Thorax* 2014;69:918-25.
- Sykes A, Macintyre J, Edwards MR, Del Rosario A, Haas J, Gielen V, et al. Rhinovirus-induced interferon production is not deficient in well controlled asthma. *Thorax* 2014;69:240-6.
- Bochkov YA, Hanson KM, Keles S, Brockman-Schneider RA, Jarjour NN, Gern JE. Rhinovirus-induced modulation of gene expression in bronchial epithelial cells from subjects with asthma. *Mucosal Immunol* 2010;3:69-80.
- Swiecki M, Gilfillan S, Vermi W, Wang Y, Colonna M. Plasmacytoid dendritic cell ablation impacts early interferon responses and antiviral NK and CD8(+) T cell accrual. *Immunity* 2010;33:955-66.
- Davidson S, Kaiko G, Loh Z, Lalwani A, Zhang V, Spann K, et al. Plasmacytoid dendritic cells promote host defense against acute pneumovirus infection via the TLR7-MyD88-dependent signaling pathway. *J Immunol* 2011; 186:5938-48.
- Lynch JP, Mazzone SB, Rogers MJ, Arikatt JJ, Loh Z, Pritchard AL, et al. The plasmacytoid dendritic cell: a cell at the cross-roads in asthma. *Eur Respir J* 2014; 43:264-75.
- Roponen M, Yerkovich ST, Hollams E, Sly PD, Holt PG, Upham JW. Toll-like receptor 7 function is reduced in adolescents with asthma. *Eur Respir J* 2010; 35:64-71.
- Bem RA, Domachowski JB, Rosenberg HF. Animal models of human respiratory syncytial virus disease. *Am J Physiol Lung Cell Mol Physiol* 2011; 301:L148-56.
- Kaiko GE, Loh Z, Spann K, Lynch JP, Lalwani A, Davidson S, et al. TLR7 gene deficiency and early-life *Pneumovirus* infection interact to predispose toward the development of asthma-like pathology in mice. *J Allergy Clin Immunol* 2013; 131:1331-9.
- Traister RS, Uvalle CE, Hawkins GA, Meyers DA, Bleecker ER, Wenzel SE. Phenotypic and genotypic association of epithelial IL1RL1 to human TH2-like asthma. *J Allergy Clin Immunol* 2015;135:92-9.
- Savenije OE, Mahachie John JM, Granel R, Kerkhof M, Dijk FN, de Jongste JC, et al. Association of IL33-IL-1 receptor-like 1 (IL1RL1) pathway polymorphisms with wheezing phenotypes and asthma in childhood. *J Allergy Clin Immunol* 2014;134:170-7.
- Huang Y, Guo L, Qiu J, Chen X, Hu-Li J, Siebenlist U, et al. IL-25-responsive, lineage-negative KLRG1(hi) cells are multipotential 'inflammatory' type 2 innate lymphoid cells. *Nat Immunol* 2015;16:161-9.
- Ullah MA, Loh Z, Gan WJ, Zhang V, Yang H, Li JH, et al. Receptor for advanced glycation end products and its ligand high-mobility group box-1 mediate allergic airway sensitization and airway inflammation. *J Allergy Clin Immunol* 2014;134: 440-50.

29. Phipps S, Hansbro N, Lam CE, Foo SY, Matthaei KI, Foster PS. Allergic sensitization is enhanced in early life through toll-like receptor 7 activation. *Clin Exp Allergy* 2009;39:1920-8.
30. Hammad H, Chieppa M, Perros F, Willart MA, Germain RN, Lambrecht BN. House dust mite allergen induces asthma via Toll-like receptor 4 triggering of airway structural cells. *Nat Med* 2009;15:410-6.
31. Willart MA, Deswarte K, Pouliot P, Braun H, Beyaert R, Lambrecht BN, et al. Interleukin-1 α controls allergic sensitization to inhaled house dust mite via the epithelial release of GM-CSF and IL-33. *J Exp Med* 2012;209:1505-17.
32. Guo L, Huang Y, Chen X, Hu-Li J, Urban JF Jr, Paul WE. Innate immunological function of TH2 cells in vivo. *Nat Immunol* 2015;16:1051-9.
33. Aherne W, Bird T, Court SD, Gardner PS, McQuillin J. Pathological changes in virus infections of the lower respiratory tract in children. *J Clin Pathol* 1970;23:7-18.
34. Saitoh T, Satoh T, Yamamoto N, Uematsu S, Takeuchi O, Kawai T, et al. Antiviral protein Viperin promotes Toll-like receptor 7- and Toll-like receptor 9-mediated type I interferon production in plasmacytoid dendritic cells. *Immunity* 2011;34:352-63.
35. Kouzaki H, Iijima K, Kobayashi T, O'Grady SM, Kita H. The danger signal, extracellular ATP, is a sensor for an airborne allergen and triggers IL-33 release and innate Th2-type responses. *J Immunol* 2011;186:4375-87.
36. Hardman CS, Panova V, McKenzie ANJ. IL-33 citrine reporter mice reveal the temporal and spatial expression of IL-33 during allergic lung inflammation. *Eur J Immunol* 2013;43:488-98.
37. Tjota MY, Williams JW, Lu T, Clay BS, Byrd T, Hrusch CL, et al. IL-33-dependent induction of allergic lung inflammation by Fc γ RIII signaling. *J Clin Invest* 2013;123:2287-97.
38. Chu DK, Llop-Guevara A, Walker TD, Flader K, Goncharova S, Boudreau JE, et al. IL-33, but not thymic stromal lymphopoietin or IL-25, is central to mite and peanut allergic sensitization. *J Allergy Clin Immunol* 2013;131:187-200, e1-8.
39. Llop-Guevara A, Chu DK, Walker TD, Goncharova S, Fattouh R, Silver JS, et al. A GM-CSF/IL-33 pathway facilitates allergic airway responses to sub-threshold house dust mite exposure. *PLoS One* 2014;9:e88714.
40. Liu YC, Simmons DP, Li XL, Abbott DW, Boom WH, Harding CV. TLR2 signaling depletes IRAK1 and inhibits induction of type I IFN by TLR7/9. *J Immunol* 2012;188:1019-26.
41. Sandig H, Jobbings CE, Roldan NG, Whittingham-Dowd JK, Orinska Z, Takeuchi O, et al. IL-33 causes selective mast cell tolerance to bacterial cell wall products by inducing IRAK1 degradation. *Eur J Immunol* 2013;43:979-88.
42. Jackson DJ, Evans MD, Gangnon RE, Tisler CJ, Pappas TE, Lee WM, et al. Evidence for a causal relationship between allergic sensitization and rhinovirus wheezing in early life. *Am J Respir Crit Care Med* 2012;185:281-5.
43. Baraldo S, Saetta M, Barbato A, Contoli M, Papi A. Rhinovirus-induced interferon production in asthma. *Thorax* 2014;69:772.
44. Yagami A, Orihara K, Morita H, Futamura K, Hashimoto N, Matsumoto K, et al. IL-33 mediates inflammatory responses in human lung tissue cells. *J Immunol* 2010;185:5743-50.
45. Jackson DJ, Makrinioti H, Rana BMJ, Shamji BWH, Trujillo-Torralbo M-B, Footitt J, et al. IL-33-dependent type 2 inflammation during rhinovirus-induced asthma exacerbations in vivo. *Am J Respir Crit Care Med* 2014;15:1373-82.
46. Edwards MR, Regamey N, Vareille M, Kieninger E, Gupta A, Shoemark A, et al. Impaired innate interferon induction in severe therapy resistant atopic asthmatic children. *Mucosal Immunol* 2013;6:797-806.
47. Saglani S, Lui S, Ullmann N, Campbell GA, Sherburn RT, Mathie SA, et al. IL-33 promotes airway remodeling in pediatric patients with severe steroid-resistant asthma. *J Allergy Clin Immunol* 2013;132:676-85.e13.
48. Bossley CJ, Fleming L, Gupta A, Regamey N, Frith J, Oates T, et al. Pediatric severe asthma is characterized by eosinophilia and remodeling without T(H)2 cytokines. *J Allergy Clin Immunol* 2012;129:974-82.e13.
49. Pritchard AL, Carroll ML, Burel JG, White OJ, Phipps S, Upham JW. Innate IFNs and plasmacytoid dendritic cells constrain Th2 cytokine responses to rhinovirus: a regulatory mechanism with relevance to asthma. *J Immunol* 2012;188:5898-905.
50. Grotenboer NS, Ketelaar ME, Koppelman GH, Nawijn MC. Decoding asthma: translating genetic variation in IL33 and IL1RL1 into disease pathophysiology. *J Allergy Clin Immunol* 2013;131:856-65.
51. Pasparakis M, Vandenabeele P. Necroptosis and its role in inflammation. *Nature* 2015;517:311-20.
52. O'Reilly R, Ullmann N, Irving S, Bossley CJ, Sonnappa S, Zhu J, et al. Increased airway smooth muscle in preschool wheezers who have asthma at school age. *J Allergy Clin Immunol* 2012;131:1024-32.
53. Stewart A. More muscle in asthma, but where did it come from? *Am J Respir Crit Care Med* 2012;185:1035-7.

Online Repository

Aeroallergen-induced IL-33 predisposes to respiratory virus-induced asthma by dampening anti-viral immunity

Jason P. Lynch, PhD,^{1*}, Rhiannon B. Werder, BSc (Hons),^{1*}, Jennifer Simpson, B Biomed Sci (Hons),¹, Zhixuan Loh, B Biomed Sci (Hons),¹, Vivian Zhang, PhD,¹ Ashraful Haque, PhD,², Kirsten Spann, PhD,^{2,3}, Peter D. Sly, MD, PhD,^{3,4}, Stuart Mazzone, PhD,¹, John W. Upham, MD, PhD,^{3,5}, and Simon Phipps, PhD,^{1,3}

Online Methods

Induction of co-virus and allergen-induced asthma and associated perturbations.

Specific pathogen-free BALB/c mice or 4C13R mice¹ were inoculated with PVM (strain J3666; 1 pfu ‘early life’; 20 pfu ‘later life’) or cockroach allergen extract (CRE, 1 µg, Greer Laboratories) as previously described^{2, 3} and as outlined in the study design (Figure 1A). In some experiments, mice were exposed to HDM extract (5 µg or 100ug, Greer Laboratories) or LPS (186 pg, Sigma-Aldrich), recombinant IL-33 (10 ng, eBioscience), anti-IL-33/isotype control antibody (200 µg, Pfizer, Inc), apyrase (4 U/mL, Sigma-Aldrich) or pyridoxalphosphate-6-azophenyl-2',4'-disulfonic acid (PPADS, 100 µm, Sigma-Aldrich). All studies were approved by The University of Queensland Animal Ethics Committee.

Sample extraction and processing

Following euthanasia by pentobarbitone overdose, blood was obtained by cardiac puncture, centrifuged twice (13,000 rpm, 4°C) and the serum stored at -80°C. A bronchoalveolar lavage

(BAL) was performed with 400 μ L (neonate/‘early life’) or 600 μ L (adult/‘later life’) of PBS. The BAL fluid was centrifuged at 5,000 rpm, 4°C for 5 min and the supernatant stored at -80°C until analysis by cytokine bead array (CBA) or ELISA. Lung lobes were excised and processed as previously described ⁴. Briefly, the left lung lobe was processed immediately for flow cytometry and the superior right lobe fixed in 10% formalin neutral buffer overnight before storage in 70% ethanol. The post-caval and inferior lobes were pooled and snap frozen before mechanical digestion and clarification, followed by analysis by ELISA. The inferior right lobe was snap frozen before RNA extraction. All snap frozen lungs were stored at -80 °C.

Flow cytometry

Flow cytometry was performed on lung tissue digest cells as previously described ⁴. Briefly, single cell suspensions were incubated with anti-Fc γ RIII/II (Fc block) for 15 min at 4°C then incubated with the following fluorochrome-conjugated antibodies at 4°C for 30 minutes: anti-mouse CD2-FITC (RM2-5), CD4-AF488 (RM4-5), Gr-1-AF488 (RB6-8C5), CD11c-AF488 (HL3), CD11b-AF488 (M1/70), B220-AF488 (RA36B2), CD3-AF488 (145-2C11) CD45RA-PE (14.8), B220-V500 (RA36B2), CD8a-PerCP (53-6.7), Sca1-PE (E13-161.7) (all BD Bioscience), CD19-AF488 (6D5), CD45-BV421 (30-F11), CD11b-BV421 (M1/70), ST-2-APC (DIH9), IFNAR (MAR1-5A3), ICOS-PE (7E.17G9), IL-7Ra-PE (SB/199) (all Biolegend), IL-17RB-PE-Cy7 (eBio17B7) (eBioscience) and Siglec-H-APC (511.3D3) (Miltenyi Biotec). 7-AAD (eBioscience) was used to exclude dead cells. For intracellular staining, cells were fixed and permeabilised using the BD Cytofix/Cytoperm kit as per the manufacturer’s instructions, followed by staining with rabbit anti-mouse IRAK1 (D51G7, Cell Signalling) or anti-mouse Viperin (HM1016, Hycult) for 30 minutes. Cells were then washed followed by incubation with

goat anti-rabbit AF647 (Invitrogen) for 30 minutes. Samples were collected with an LSR Fortessa X-20 (BD Biosciences) and the data analysed with FACSDiva v8 (BD Biosciences) and FlowJo v8.8 (Treestar). Type 2 ILCs were identified as Lineage⁻ (CD2, Gr-1, CD3, CD11b, B220, CD4, CD19), CD45⁺, CD25⁺, CD90.2⁺, ST-2⁺, ICOS⁺, IL-7Ra⁺, IL-17RB⁺ (Fig E4A). pDCs were identified as CD11b⁻, CD11c⁺, CD45RA⁺, B220⁺, Siglec-H⁺.

Histology and Immunohistochemistry

Paraffin-embedded lung sections were prepared as previously described ³. Lung tissue sections were stained with Chromotrope 2R, Periodic acid-Schiff or Masson's Trichrome, to enumerate eosinophils, mucus-secreting cells and collagen deposition respectively. For immunohistochemistry, lung sections were pretreated with 10% normal goat serum for 30 min. Sections were probed with anti-IL-33 (AF3626, R&D), anti-PVM G protein (kindly provided by Dr Ulla Buchholz), anti- α -SM actin and anti-periostin (both Sigma-Aldrich) overnight at 4°C. Following incubation with appropriate secondary antibodies, immunoreactivity was developed with Fast Red (Sigma-Aldrich) and counterstained with Mayer's hematoxylin (bright field) or with 4',6-diamidino-2-phenylindole (DAPI, Sigma-Aldrich) (fluorescence). The percentage of PVM positive or mucous AECs was quantified in 5 airways per mouse (Scanscope XT, Aperio). Oedema was assessed by point counting of fluid-filled airspaces. Eosinophils were enumerated around the airways and expressed as cells per 100 μ m of epithelial basement membrane. Airway smooth muscle mass and collagen deposition around the small airways (defined as a circumference <500 μ m for neonates and <800 μ m for mice aged >7 week) was measured using Scanscope XT software and expressed as area per μ m of basement membrane. Periostin expression was quantified as a % of the airway circumference. Photomicrographs were taken at

400x and 1000x magnification using an Olympus BX-51 microscope with an Olympus DP-72 camera at room temperature and acquired using Olympus Image Analysis Software. IL-33 images were false colored using Adobe Photoshop CS6 software. Immunofluorescent images were taken using Discovery Spinning disk confocal using Nikon Viewer software. Images were processed using Image J and Imaris software.

Measurement of protein expression

IL-33 and IFN- λ 2/3 (R&D Systems), IL-12p40 and IFN- γ (Biolegend) and IL-5 (BD Biosciences) expression was quantified by ELISA. IFN- α (eBioscience) and IL-13 (Quantikine kit, R&D or Enhanced Sensitivity Flex Set, BD Biosciences) expression was quantified by CBA. IgG1a was detected in the serum using an in-house ELISA system ².

Quantitative real time PCR

Total RNA was isolated from the inferior right lung lobe with TriReagent solution (Ambion) followed by phenol-chloroform extraction. DNase digestion was performed with Turbo DNase (Ambion), according to the manufacturer's instructions. Reverse transcription was performed using M-MLV reverse transcriptase and random primers (Invitrogen). qRT-PCR was performed with SYBR Green (Life Technologies) with the primers described in Table S1. Expression values were normalized to *Hprt* and expressed as fold change over vehicle mice, as described ^{3,4}.

Airway function assessment

AHR was measured as described previously ⁵. Briefly, airways resistance was determined by forced oscillation technique (Flexivent, Scireq) in response to nebulized methacholine (0.3 to 10 mg/mL; Sigma-Aldrich).

Type-2 innate lymphoid cell culture and activation

Lungs were excised from PVM/CRE co-exposed mice at 10 dpi and digested by gentleMACS dissociation (Miltenyi Biotech). Type 2 ILCs (7-AAD⁻ Lineage⁻ CD45⁺, CD25⁺, CD90.2⁺) were FACS-sorted to 96% purity (Fig E6C) using a BD FACS-Aria, and cultured in the presence of IL-2 (30 ng/mL, eBioscience). Cells (10,000/well) were pre-incubated with IFN- α (5000U/mL, Hycult Biotech) for 30 minutes, before stimulation with IL-33 (30 ng/mL, eBioscience) for 72 hours. Supernatant was collected and probed for cytokine production.

Plasmacytoid dendritic cell culture

Bone marrow-derived pDC were generated as described previously ³. On day 8 of culture, pDC were pre-incubated with IL-33 (3 ng/mL, eBioscience) then stimulated with imiquimod (3 μ g/mL, Sigma-Aldrich).

Statistical analyses

GraphPad Prism version 5.0 software (La Jolla, California) was used for all statistical analyses. A Student's t-test, one-way ANOVA with a Tukey post-hoc test or two-way ANOVA with a Sidak post-hoc test were applied as appropriate. A *P* value <0.05 was considered statistically significant.

Supplemental Figure Legends

Figure E1. Determination of CRE allergen dose (A) Study design. (B) Peribronchial eosinophils. BM, basement membrane. (C) Mucous-secreting cells. AEC, airway epithelial cells. Data are mean $\pm$ SEM, representative of 2 experiments, n=5-9 mice per group.

Figure E2. Both virus and allergen challenge in later life are necessary for asthma progression. (A) Peribronchial eosinophils. Basement membrane (BM). (B) Mucous-secreting airway epithelial cells (AECs). (C) Cockroach-specific IgG1a in serum. (D) Collagen area. (E) ASM area. (F) Airway resistance (Rn) in response to increasing doses of methacholine (MCh), response at 10mg/mL shown. (G) Representative micrograph (x400 magnification) of peribronchial eosinophils (chromotrope 2R), scale bar = 50 μ m, white arrows indicate eosinophils. Data are mean $\pm$ SEM, representative of three independent experiments, n=6-9 mice per group. * compared with vehicle mice. # compared with PVM/CRE/PVM/CRE mice. Dashed line denotes vehicle treated mice. Related to Figure 1.

Figure E3. Expression of virus and IL-33 in airway epithelial cells. (A) Representative micrographs of PVM immunostaining, scale bar = 50 μ m, arrows indicate PVM+ airway epithelial cells. (B) pDC 7 days post infection, enumerated as CD11b- CD11c+ Siglec-H+ B220+ CD45RA+. (C) Quantification of airway epithelial cells (AECs) expressing IL-33 as % of AECs. (D) Bronchoalveolar lavage total and differential counts.

Figure E4. Type 2 innate lymphoid cell and T cell responses. (A) Representative scatter plots showing gating strategy for enumeration of type 2 ILCs and their immunophenotype in the lung. Grey filled histogram = fluorescence of minus one control for staining; solid line = expression of ILC2 markers in lung. Values are % of total lung cells. (B) ILC2, CD4- and CD8-T cell expression of IL-13 and IL-4 at 10 DPI in 4C13R mice. Values are % parent population.

Figure E5. Anti-IL-33 prevents type 2 inflammation and remodeling in response to virus and allergen co-exposure in later-life. (A) Study plan. (B) Airway resistance (Rn) in response to increasing doses of methacholine (MCh). (C) Mucous-secreting airway epithelial cells (AECs). (D) Peribronchial eosinophils. Basement membrane (BM). (E) ASM area. Data are mean $\pm$ SEM, representative of 2 experiments, n=6-7 mice per group. Dashed line denotes vehicle treated mice, solid line denotes PVM treated mice. (F) Bronchoalveolar lavage total and differential counts. (G) pDC number and (H) Sca-1+ CD8 T cells in the lung.

Figure E6. Blockade of IL-33 release and type-2 ILC sort purity and viability in culture following stimulation. (A) IL-33 in bronchoalveolar lavage (BALF), following treatment with apyrase or pyridoxalphosphate-6-azophenyl-2',4'-disulfonic acid (PPADS). (B) Viral load in airway epithelial cells and IL-12p40 in lung. (C) Representative scatter plots showing gating strategy for FACS sorting of lung type-2 ILCs. Values are % of parent population. (D) Viable (7-AAD-) type-2 ILCs following 3 day culture.

Figure E7. Exogenous IL-33 dampens IFN- α production and increases viral load (A) Study design. Mice were inoculated with PVM at 7 dpi, then exposed (i.n.) to 10 ng IL-33 3, 4 and 5 days later. (B) Plasmacytoid dendritic cell (pDC), (C) Type-2 ILC and (D) Eosinophil number in

the lung. (E-I) Bronchoalveolar lavage total and differential counts. (E) Representative scatter plots showing gating strategy for cultured pDCs.

Supplemental Table Legends

Table E1. Oligonucleotide sequences used in this study are shown.

Supplemental References

1. Huang Y, Guo L, Qiu J, Chen X, Hu-Li J, Siebenlist U, et al. IL-25-responsive, lineage-negative KLRG1(hi) cells are multipotential 'inflammatory' type 2 innate lymphoid cells. *Nat Immunol* 2015; 16:161-9.
2. Ullah MA, Loh Z, Gan WJ, Zhang V, Yang H, Li JH, et al. Receptor for advanced glycation end products and its ligand high-mobility group box-1 mediate allergic airway sensitization and airway inflammation. *J Allergy Clin Immunol* 2014; 134:440-50.
3. Davidson S, Kaiko G, Loh Z, Lalwani A, Zhang V, Spann K, et al. Plasmacytoid dendritic cells promote host defense against acute pneumovirus infection via the TLR7-MyD88-dependent signaling pathway. *J Immunol* 2011; 186:5938-48.
4. Kaiko GE, Loh Z, Spann K, Lynch JP, Lalwani A, Davidson S, et al. TLR7 gene deficiency and early-life Pneumovirus infection interact to predispose toward the development of asthma-like pathology in mice. *J Allergy Clin Immunol* 2013; 131:1331-39.
5. Phipps S, Lam CE, Kaiko GE, Foo SY, Collison A, Mattes J, et al. Toll/IL-1 signaling is critical for house dust mite-specific helper T cell type 2 and type 17 [corrected] responses. *Am J Respir Crit Care Med* 2009; 179:883-93.

Table S1

Name	Oligonucleotide Primer
<i>Irf7</i>	Forward: 5'-CTTAGCCGGGAGCTTGGATCTACT-3' Reverse: 5'-CCCTTGTACATGATGGTCACATCC-3'
<i>Stat1</i>	Forward: 5'-ACAGTGGTTCGAGCTTCAG-3' Reverse: 5'-GGCCAGGTACTGTCTGATTT-3'
<i>Viperin</i>	Forward: 5'- CGAAGACATGAATGAACACATCAA-3' Reverse: 5'-AATTAGGAGGCACTGGAAAACCT-3'
<i>Hprt</i>	Forward: 5'- AGGCCAGACTTTGTTGGATTTGAA-3' Reverse: 5'-CAACTTGCGCTCATCTTAGGCTTT-3'




Fig E1.

Jason P. Lynch, Rhiannon B. Werder, Jennifer Simpson, Zhixuan Loh, Vivian Zhang, Ashraful Haque, Kirsten Spann, Peter D. Sly, Stuart B. Mazzone, John W. Upham, Simon Phipps

Aeroallergen-induced IL-33 predisposes to respiratory virus-induced asthma by dampening antiviral immunity

Journal of Allergy and Clinical Immunology, 2016, Available online 25 April 2016

<http://dx.doi.org/10.1016/j.jaci.2016.02.039>




Fig E2.

Jason P. Lynch, Rhiannon B. Werder, Jennifer Simpson, Zhixuan Loh, Vivian Zhang, Ashraful Haque, Kirsten Spann, Peter D. Sly, Stuart B. Mazzone, John W. Upham, Simon Phipps

Aeroallergen-induced IL-33 predisposes to respiratory virus-induced asthma by dampening antiviral immunity

Journal of Allergy and Clinical Immunology, 2016, Available online 25 April 2016

<http://dx.doi.org/10.1016/j.jaci.2016.02.039>




Fig E3.

Jason P. Lynch, Rhiannon B. Werder, Jennifer Simpson, Zhixuan Loh, Vivian Zhang, Ashraful Haque, Kirsten Spann, Peter D. Sly, Stuart B. Mazzone, John W. Upham, Simon Phipps

Aeroallergen-induced IL-33 predisposes to respiratory virus-induced asthma by dampening antiviral immunity

Journal of Allergy and Clinical Immunology, 2016, Available online 25 April 2016

<http://dx.doi.org/10.1016/j.jaci.2016.02.039>




Fig E4.

Jason P. Lynch, Rhiannon B. Werder, Jennifer Simpson, Zhixuan Loh, Vivian Zhang, Ashraf Haque, Kirsten Spann, Peter D. Sly, Stuart B. Mazzone, John W. Upham, Simon Phipps

Aeroallergen-induced IL-33 predisposes to respiratory virus-induced asthma by dampening antiviral immunity

Journal of Allergy and Clinical Immunology, 2016, Available online 25 April 2016

<http://dx.doi.org/10.1016/j.jaci.2016.02.039>




Fig E5.

Jason P. Lynch, Rhiannon B. Werder, Jennifer Simpson, Zhixuan Loh, Vivian Zhang, Ashraf Haque, Kirsten Spann, Peter D. Sly, Stuart B. Mazzone, John W. Upham, Simon Phipps

Aeroallergen-induced IL-33 predisposes to respiratory virus-induced asthma by dampening antiviral immunity

Journal of Allergy and Clinical Immunology, 2016, Available online 25 April 2016

<http://dx.doi.org/10.1016/j.jaci.2016.02.039>




Fig E6.

Jason P. Lynch, Rhiannon B. Werder, Jennifer Simpson, Zhixuan Loh, Vivian Zhang, Ashraf Haque, Kirsten Spann, Peter D. Sly, Stuart B. Mazzone, John W. Upham, Simon Phipps

Aeroallergen-induced IL-33 predisposes to respiratory virus-induced asthma by dampening antiviral immunity

Journal of Allergy and Clinical Immunology, 2016, Available online 25 April 2016

<http://dx.doi.org/10.1016/j.jaci.2016.02.039>




Fig E7.

Jason P. Lynch, Rhiannon B. Werder, Jennifer Simpson, Zhixuan Loh, Vivian Zhang, Ashraful Haque, Kirsten Spann, Peter D. Sly, Stuart B. Mazzone, John W. Upham, Simon Phipps

Aeroallergen-induced IL-33 predisposes to respiratory virus-induced asthma by dampening antiviral immunity

Journal of Allergy and Clinical Immunology, 2016, Available online 25 April 2016

<http://dx.doi.org/10.1016/j.jaci.2016.02.039>