

Response to self antigen imprints regulatory memory in tissues

Michael D. Rosenblum^{1*}, Iris K. Gratz^{2*}, Jonathan S. Paw^{1,2}, Karen Lee³, Ann Marshak-Rothstein⁴ & Abul K. Abbas²

Immune homeostasis in tissues is achieved through a delicate balance between pathogenic T-cell responses directed at tissue-specific antigens and the ability of the tissue to inhibit these responses. The mechanisms by which tissues and the immune system communicate to establish and maintain immune homeostasis are currently unknown. Clinical evidence suggests that chronic or repeated exposure to self antigen within tissues leads to an attenuation of pathological autoimmune responses, possibly as a means to mitigate inflammatory damage and preserve function. Many human organ-specific autoimmune diseases are characterized by the initial presentation of the disease being the most severe, with subsequent flares being of lesser severity and duration¹. In fact, these diseases often spontaneously resolve, despite persistent tissue autoantigen expression². In the practice of antigen-specific immunotherapy, allergens or self antigens are repeatedly injected in the skin, with a diminution of the inflammatory response occurring after each successive exposure³. Although these findings indicate that tissues acquire the ability to attenuate autoimmune reactions upon repeated responses to antigens, the mechanism by which this occurs is unknown. Here we show that upon expression of self antigen in a peripheral tissue, thymus-derived regulatory T cells (T_{reg} cells) become activated, proliferate and differentiate into more potent suppressors, which mediate resolution of organ-specific autoimmunity in mice. After resolution of the inflammatory response, activated T_{reg} cells are maintained in the target tissue and are primed to attenuate subsequent autoimmune reactions when antigen is re-expressed. Thus, T_{reg} cells function to confer 'regulatory memory' to the target tissue. These findings provide a framework for understanding how T_{reg} cells respond when exposed to self antigen in peripheral tissues and offer mechanistic insight into how tissues regulate autoimmunity.

We hypothesized that exposure of immune cells to self antigen in peripheral tissues induces stable regulatory mechanisms that limit autoimmune injury. To test this and to define the nature of these control mechanisms, we created a novel mouse model of inducible tissue-specific self-antigen expression. We crossed transgenic mice expressing a membrane-bound form of ovalbumin (Ova) under the control of a tetracycline response element to transgenic mice expressing the tetracycline transactivator protein under the control of the keratin 5 (K5, also called Krt5) promoter (Fig. 1a)⁴. In the resultant K5/TGO double transgenic mice (see Methods for definition of TGO), Ova expression in the skin is tightly controlled, as adoptively transferred Ova-specific DO11.10 (DO11) $CD4^{+}$ T cells become activated and proliferate in skin-draining lymph nodes (SDLNs) of recipient mice only after treatment with doxycycline (Fig. 1b). In addition, Ova messenger RNA is detected in epidermal cell suspensions only after induction with doxycycline (Supplementary Fig. 1).

To define functional Ova expression in the thymus, K5/TGO mice were crossed with the DO11.10 T-cell receptor (TCR)-transgenic

strain⁵. Deletion of $CD4^{+}$ DO11 T cells or increased production of $Foxp3^{+}CD4^{+}$ DO11 T_{reg} cells in K5/TGO/DO11 mice is a sensitive indicator of thymic Ova expression. K5/TGO/DO11 triple transgenic mice as well as TGO/DO11 double transgenic mice have modest deletion of $CD4^{+}$ DO11 cells (Fig. 1c) and a pronounced increase in DO11 T_{reg} cells in both the thymus and SDLNs (Fig. 1c, d). Between 30–40% of antigen-specific $CD4^{+}$ DO11 T cells in the SDLNs of K5/TGO/DO11 (and TGO/DO11) mice co-express CD25 and $Foxp3$ (Fig. 1d). These results demonstrate that Ova is constitutively expressed in the thymus, independent of doxycycline treatment and dependent only on the presence of the TGO transgene. Thus, K5/TGO/DO11 mice represent a unique model in which antigen is continuously expressed in the thymus and tightly controlled in the periphery, mimicking the pattern of tissue-specific self-antigen expression in mice and humans^{6,7}.

Despite the presence of a large percentage of Ova-specific T_{reg} cells, induction of cutaneous Ova expression in K5/TGO/DO11 mice results in a pronounced inflammatory dermatitis (Fig. 2a). Disease peaks at





Figure 1 | Characterization of K5/TGO/DO11 mice. **a**, Construct for double transgenic mice expressing ovalbumin (Ova) driven by the cytokerin 5 (K5) promoter in a tetracycline-inducible fashion. **b**, Lymph node cells from DO11 TCR-transgenic mice were labelled with CFSE and injected into K5/TGO mice. Recipient mice were fed doxycycline chow and DO11 cell proliferation (CFSE dilution) and CD44 expression was measured 3 days later by flow cytometry. **c**, SDLN cell numbers from DO11 and TGO/DO11 mice in the absence of doxycycline treatment. **d**, Thymus and SDLN cells from DO11, TGO/DO11 and K5/TGO/DO11 mice in the absence of doxycycline treatment. Thymocytes are gated on $CD4^{+}KJ^{+}CD8^{-}$ cells. KJ, KJ1-26 monoclonal antibody. Lymph node cells are gated on $CD4^{+}KJ^{+}$ cells. * $P < 0.05$ (t -test). Error bars represent s.d. Results are representative of 3 replicate experiments with $n = 3$ –4 mice per group.

¹Department of Dermatology, University of California San Francisco, San Francisco, California 94115, USA. ²Department of Pathology, University of California San Francisco, San Francisco, California 94143, USA. ³Department of Pediatrics, Columbia University Medical Center, New York, New York 10032, USA. ⁴Department of Medicine, Rheumatology Division, University of Massachusetts, Worcester, Massachusetts 01655, USA.

*These authors contributed equally to this work.

10–14 days after antigen induction and is characterized by marked erythema, scaling and alopecia (Fig. 2b). The skin infiltrate at the height of disease is composed primarily of antigen-specific DO11 T cells and GR-1⁺CD11b⁺ myeloid cells (Fig. 2c). Ova-specific skin-infiltrating CD4⁺ T cells produce interferon (IFN)- γ and interleukin (IL)-17 (Fig. 2j).

Interestingly, skin inflammation spontaneously resolves. Twenty to 30 days after antigen induction, mice begin to show signs of clinical and histological improvement, and by 40 to 60 days, there is complete resolution of cutaneous inflammation (Fig. 2d–f). Disease resolves despite continued doxycycline treatment, with ongoing Ova expression in the skin and continuous thymic output of Ova-specific T cells (Supplementary Fig. 2).



Given the large percentage of antigen-specific T_{reg} cells present in the SDLNs of K5/TGO/DO11 mice, it is surprising that cutaneous inflammation results when antigen is induced. We were further intrigued by the spontaneous resolving nature of the autoimmune response, as it suggests that an immune regulatory mechanism(s) is initiated in an attempt to minimize tissue damage. To determine if T_{reg} cells have a role in resolving skin inflammation, we depleted these cells before antigen induction and at the height of disease. Treatment of K5/TGO/DO11 mice with CD25-depleting antibody (PC61) before antigen induction resulted in pronounced disease exacerbation with delayed resolution of skin inflammation (Fig. 2g, h). In some experiments, depletion of T_{reg} cells resulted in dehydration, weight loss and death, most likely from severely disrupted skin barrier function (Supplementary Fig. 3). Depletion of T_{reg} cells at the height of disease (that is, 10 days after antigen induction) resulted in delayed kinetics of resolution of skin inflammation, with some mice having to be euthanized secondary to non-resolving skin disease (Fig. 2i). Higher percentages of both IFN- γ and IL-17-secreting DO11 cells were observed in the skin and SDLN after T_{reg} cell depletion (Fig. 2j). Anti-CD25 antibody was used in these studies because CD25 is constitutively expressed on T_{reg} cells and antibody treatment can be accurately titrated to achieve optimal deletion of these cells (Supplementary Fig. 4). A caveat of this approach is that other CD25-expressing cells (which may be either tolerance-promoting or pathogenic) might also be deleted. The finding that anti-CD25 antibody treatment inhibits or prevents disease resolution strongly suggests that T_{reg} cells are the major targets of the antibody and have an essential role in resolving skin disease in our model; however, depletion of other CD25-expressing regulatory populations cannot be definitively ruled out.

The emergence of disease despite the presence of T_{reg} cells and the T_{reg} cell-dependent suppression of disease after antigen induction suggested that expression of tissue antigen activates T_{reg} cells and enhances their suppressive function. To test this, we induced Ova expression in K5/TGO/DO11 mice and characterized T_{reg} cells in both the skin and SDLNs. Foxp3⁺ DO11 cells robustly expanded in SDLNs upon antigen induction (Fig. 3a). Expansion peaked at 12–14 days, when these cells were increased ~15-fold over baseline numbers. In contrast, the peak expansion of Foxp3⁻ DO11 effector cells occurred earlier and these cells only expanded ~4-fold over baseline (Fig. 3a). At the peak of expansion, more than 50% of Foxp3⁺ DO11 cells had markedly reduced CD25 on the cell surface (Fig. 3b). Loss of CD25 was transient, associated only with the proliferative burst, as Foxp3⁺ DO11 cells in SDLNs during the resolution phase of disease (>20 days) express CD25 (Fig. 3b). Loss of CD25 expression on *in vivo* proliferating T_{reg} cells (with preservation of suppressive capacity)

Figure 2 | K5/TGO/DO11 mice develop autoimmune skin disease that resolves spontaneously. **a**, Mean clinical scores of K5/TGO/DO11 mice left untreated or fed doxycycline chow. **b**, Skin lesions in a representative K5/TGO/DO11 mouse at the height of clinical disease. **c**, Flow cytometry of skin-infiltrating cells in K5/TGO/DO11 mice treated with doxycycline for 12 days. **d**, Representative mice from the height of disease (day 12) and after disease resolution (day 86). **e**, Skin histology of K5/TGO/DO11 mice after beginning doxycycline. **f**, Flow cytometry of skin-infiltrating cells in K5/TGO/DO11 mice at 11 and 43 days after beginning doxycycline. **g**, Disease scores of K5/TGO/DO11 mice treated with anti-CD25 monoclonal antibody (PC61) or control (PBS or isotype control antibody) at 10 days and 3 days before beginning doxycycline. **h**, Flow cytometry of skin-infiltrating cells in K5/TGO/DO11 mice untreated or treated with PC61 at 23 days after beginning doxycycline. **i**, Disease scores of K5/TGO/DO11 mice treated with PC61 or isotype control antibody at the height of disease. Results are pooled data from two replicate experiments with $n = 3$ –4 mice per group. **j**, Intracellular cytokine stains of SDLN cells and skin-infiltrating cells from PC61- or isotype control-treated (before antigen induction) K5/TGO/DO11 mice at 11 days after beginning doxycycline treatment. Gated on CD4⁺KJ⁺ cells. All error bars represent s.d. Results are representative of 3 replicate experiments with $n = 3$ –4 mice per group. Euthanization of two PC61-treated mice secondary to non-resolving skin disease is indicated by the ‡ symbol.




Figure 3 | **T_{reg} cells are activated upon induction of peripheral antigen.**

a, Quantification of CD4⁺KJ⁺Fcpx3⁺ and CD4⁺KJ⁺Fcpx3⁻ cell numbers from SDLNs of K5/TGO/DO11 mice after beginning doxycycline treatment. **b**, Flow cytometry of SDLN cells from K5/TGO/DO11 mice after beginning doxycycline treatment. Gated on CD4⁺KJ⁺ cells. **c**, Flow cytometry of SDLN cells from K5/TGO/DO11 mice left untreated (naive CD25⁺) or treated for 6 days with doxycycline (activated CD25⁺ or CD25⁻ populations). **d**, Flow cytometry of skin-infiltrating cells from K5/TGO/DO11 mice after beginning doxycycline. Gated on live CD4⁺KJ⁺ cells. **e**, *In vitro* suppression assay using sorted CD4⁺KJ⁺Fcpx3⁺CD25⁺ cells from non-induced naive mice (nCD25⁺), CD4⁺KJ⁺Fcpx3⁺CD25⁺ cells from doxycycline-treated mice

has been reported by others⁸. It is possible that proliferating T_{reg} cells reduce expression of CD25 in a negative feedback loop to limit proliferation and regulate T_{reg} cell numbers. SDLN T_{reg} cells in the resolution phase of disease have increased expression of Fcpx3 and CD25 when compared to T_{reg} cells from non-induced mice (Fig. 3b and Supplementary Fig. 5). These results indicate that T_{reg} cells are activated upon exposure to peripheral antigen. To test this further, we analysed expression levels of various markers shown to be involved in T_{reg} cell function early after antigen induction. Six days after inducing Ova expression in the skin, >80% of T_{reg} cells have entered the cell cycle, as shown by Ki67 expression, and proliferating CD25⁺ and CD25⁻ T_{reg} cells express higher levels of CTLA-4 compared to T_{reg} cells from non-induced mice (Fig. 3c).

Expansion of T_{reg} cells in SDLNs upon antigen induction is associated with an accumulation of these cells in the skin (Fig. 3d). Before antigen induction, CD4⁺ DO11 cells are barely detectable in the skin. At the height of disease, most skin-infiltrating DO11 cells are Fcpx3⁻. However, resolution of inflammation is associated with a preferential accumulation of Fcpx3⁺ cells in the skin, with >60% of DO11 cells in the skin expressing Fcpx3 in mice maintained on doxycycline that have resolved disease (that is, clinically and histologically normal appearing skin) (Fig. 3d).

We next tested whether T_{reg} cells acquire a more suppressive phenotype after exposure to tissue antigen. To do so, we crossed K5/TGO/

(activated CD25⁺, aCD25⁺), or CD4⁺KJ⁺Fcpx3⁺CD25⁻ cells from doxycycline-treated mice (aCD25⁻) as suppressors and CFSE-labelled DO11 lymph node cells as responders. **f**, *In vivo* suppression assay using suppressors (as in part **e**) mixed 1:1 with naive CFSE-labelled Thy1.1⁺ DO11 lymph node cells and adoptively transferred into K5/TGO mice. Recipient mice were treated with doxycycline and DO11 cell proliferation and intracellular cytokine expression (**g**) were assayed 3 days later. Gated on Thy1.1⁺CD4⁺KJ⁺ cells. **P* < 0.05 (*t*-test). All error bars represent s.d. Results are representative of 3 replicate experiments with *n* = 3–4 mice per group except for *in vitro* and *in vivo* suppression assays, which are representative of 2 replicate experiments with *n* = 2–3 mice per group.

DO11 mice with Fcpx3^{GFP} reporter mice⁹. K5/TGO/DO11/Fcpx3^{GFP} mice were treated with doxycycline to induce antigen expression in the skin, and 6 days later, CD4⁺Fcpx3⁺CD25⁺ and CD4⁺Fcpx3⁺CD25⁻ DO11 cells were isolated from SDLNs and their suppressive capacity tested *in vitro* and *in vivo*. Both CD25⁺ and CD25⁻ T_{reg} cells from antigen-induced mice were more potent suppressors of effector T-cell proliferation *in vitro* when compared to T_{reg} cells isolated from non-induced mice (that is, peripheral antigen naive) at all effector to suppressor ratios examined (Fig. 3e). In addition, T_{reg} cells from antigen-induced mice were more potent suppressors of cutaneous autoimmune responses *in vivo*. Upon co-transfer with naive DO11 responder cells into K5/TGO double transgenic mice (and treatment of recipient mice with doxycycline to induce Ova expression in the skin), both CD25⁺ and CD25⁻ Fcpx3⁺ T_{reg} cells isolated from antigen-induced K5/TGO/DO11/Fcpx3^{GFP} mice were more potent at inhibiting T-cell proliferation when compared to Fcpx3⁺ cells isolated from non-induced mice (Fig. 3f). These cells were also more potent inhibitors of IL-17 and IFN-γ production from effector T cells (Fig. 3g).

The mechanism of enhanced T_{reg} cell suppression after peripheral antigen induction may involve CTLA-4, which is expressed at markedly higher levels after activation of T_{reg} cells (Fig. 3c) and has been shown to be a critical mediator of T_{reg} function in suppressing tissue inflammation¹⁰. In addition, CD25⁺ and CD25⁻ T_{reg} cells had similarly high

levels of CTLA-4 expression and are equally suppressive *in vitro* and *in vivo*. We were unable to detect cytokine production (including IL-10, IL-17 and IFN- γ) from T_{reg} cells before or after antigen induction, suggesting that a cytokine-mediated mechanism of suppression is unlikely (Fig. 2j and data not shown).

Our results indicate that early after exposure to tissue autoantigen, T_{reg} cells are activated, proliferate, differentiate into more potent suppressors, and progressively accumulate in the target tissue. These results are consistent with recent findings that two major populations of Foxp3⁺ T_{reg} cells exist in human peripheral blood, which have been called resting T_{reg} cells and activated T_{reg} cells¹¹. Resting T_{reg} cells are recently derived from the thymus, express low levels of CTLA-4, are not actively cycling, and are inferior suppressors when compared to activated T_{reg} cells. In contrast, activated T_{reg} cells are constitutively cycling, express higher levels of Foxp3 and CTLA-4, and are more potent suppressors. Longitudinal studies showed ongoing conversion of resting T_{reg} cells to activated T_{reg} cells in a healthy human volunteer, with T-cell-receptor clonotypes initially detected in the resting T_{reg} cell subset found 18 months later in the activated T_{reg} cell subset. These results suggest that in normal healthy individuals, resting T_{reg} cells are continuously activated and proliferate into more potent activated T_{reg} cells. Taken together with results presented here, we speculate that exposure to tissue self antigen may drive this process as a mechanism to inhibit and/or suppress autoimmunity.

We hypothesized that once T_{reg} cells have been activated and accumulate in the skin, these cells are relatively stable and persist in the tissue. They would then be capable of suppressing subsequent autoimmune reactions when antigen is re-expressed. To test this, we induced antigen expression in K5/TGO/DO11 mice and upon resolution of skin disease, removed doxycycline for >30 days to effectively 'turn antigen off' in the skin. We then re-induced antigen expression and followed mice clinically. To confirm the absence of antigen expression upon cessation of doxycycline, we assessed activation and proliferation of adoptively transferred DO11 cells and activation of endogenous DO11 cells in K5/TGO/DO11 mice (as a surrogate marker of antigen expression) at various times after discontinuing doxycycline. In addition, we measured Ova mRNA levels in the skin. At 20 days after discontinuing doxycycline treatment there was no proliferation of adoptively transferred DO11 cells and by 29 days, CD44 and CD69 expression had returned to baseline levels on adoptively transferred and endogenous DO11 cells, respectively (Fig. 4a and Supplementary Fig. 6a). Expression of Ova mRNA was undetectable in all mice at 32 days after stopping doxycycline treatment (Supplementary Fig. 6b). Taken together, these data confirm that antigen expression is effectively 'turned off' in all mice by 30 days after cessation of doxycycline treatment. Analysis of DO11 T cells in the skin of K5/TGO/DO11 mice that had been off doxycycline for >30 days revealed a persistent Foxp3⁺ population comprising >50% of DO11 cells in the skin (Fig. 4b). The proportion of DO11 cells in the skin that were Foxp3⁺ was enriched relative to SDLNs, indicating a preferential accumulation of T_{reg} cells in the target tissue upon cessation of antigen expression (Fig. 4b). Furthermore, T_{reg} cells in the skin continued to express high levels of CTLA-4 (Fig. 4c). When compared to T_{reg} cells in the SDLN, cells that persisted in the skin (in the absence of antigen expression) expressed lower levels of CD25 and higher levels of CTLA-4, CD127 and KLRG1 (Fig. 4d). The persistence of activated T_{reg} cells in the skin indicated that there may be an attenuation of cutaneous inflammation upon antigen re-expression, as T_{reg} cells must be present in the skin to suppress autoimmunity¹². To test this, we re-started doxycycline treatment and followed mice clinically. Upon antigen re-expression, K5/TGO/DO11 mice developed skin disease phenotypically similar to the disease that developed upon initial antigen expression; however, the severity of skin disease was significantly reduced and it resolved with accelerated kinetics (Fig. 4e). Attenuated disease upon antigen re-exposure was not secondary to reduced numbers of DO11 cells, as these cells returned to baseline




Figure 4 | Memory T_{reg} cells attenuate skin disease upon re-expression of tissue antigen. **a**, Flow cytometry of CFSE-labelled Thy1.1⁺DO11⁺ SDLN cells 3 days after adoptive transfer into K5/TGO/DO11 mice that had been on doxycycline for >30 days and off for 20 days. **b**, Flow cytometry and T_{reg} cell percentages of SDLN and skin-infiltrating CD4⁺ DO11 cells isolated from K5/TGO/DO11 mice that were on doxycycline for >30 days and off for >30 days. Gated on CD4⁺KJ⁺ cells. **c**, Flow cytometry of CD4⁺DO11 cells from the skin of K5/TGO/DO11 mice that have not been treated with doxycycline (naive) or on for >30 days and off doxycycline for >30 days. **d**, Phenotype of CD4⁺Foxp3⁺ DO11 cells from skin (shaded) and SDLN (unshaded) from K5/TGO/DO11 mice that were on doxycycline for >30 days and off for >40 days. **e**, Clinical skin disease upon re-starting doxycycline in K5/TGO/DO11 mice that had been on doxycycline for >30 days and off for >30 days. **f**, Clinical skin disease of K5/TGO/DO11 mice treated with either PC61 or isotype control antibody before re-starting doxycycline. In all mice, antigen was induced, skin disease developed, and disease had resolved before antibody treatment. All error bars represent standard error of samples within each group. Mean clinical scores from individual mice are shown. Results are representative of 3 replicate experiments with 3–4 mice per group except for **f**, which is combined data from 2 replicate experiments with 2–4 mice per group. **P* < 0.05 (*t*-test).

levels relatively early after cessation of doxycycline treatment (Supplementary Fig. 7). In addition, antigen-experienced DO11 T cells from K5/TGO/DO11 mice that had resolved disease and were either maintained on doxycycline or were taken off doxycycline after resolution readily induced skin disease when adoptively transferred into new antigen-expressing hosts (Supplementary Fig. 8). This suggests that cell-intrinsic anergy in responding T cells does not seem to have a major role in attenuating skin inflammation. To test whether reduced disease severity upon second antigen exposure was mediated by T_{reg} cells, we depleted these cells before re-inducing antigen expression. Depletion of T_{reg} cells resulted in a complete abrogation of the attenuated response observed upon re-expression of Ova in the skin (Fig. 4f).

Our results indicate that exposure to tissue autoantigens leads to the activation of self-reactive T_{reg} cells that were generated by self-antigen expression in the thymus. Activated T_{reg} cells persist in the target tissue and suppress autoimmune responses upon repeated or chronic encounters with tissue autoantigen. Thus, tissues that have undergone autoimmune inflammatory reactions develop a property we have termed regulatory memory, which serves to limit the severity of future such reactions. Memory T_{reg} cells have several characteristics of effector memory cells^{13,14}, including survival without antigen, residence in non-lymphoid tissues, phenotypic markers suggestive of prior activation, and enhanced functional activity. The life history of

T_{reg} cells is fundamentally similar to that of conventional T cells, passing through defined phases that include generation in the thymus, activation in the periphery leading to proliferation and differentiation into functionally more active cells, and survival as memory populations. Although our data support an obligatory role for T_{reg} cells in suppressing primary and memory responses in the skin, it is possible that these cells work together with other T_{reg} -cell-independent mechanisms to suppress tissue-specific autoimmunity. Identifying and harnessing the network of regulatory pathways that serve to limit tissue inflammation will undoubtedly offer new strategies for controlling autoimmune reactions and preserving end-organ function.

METHODS SUMMARY

Mouse model. TRE-TGO mice were crossed with K5-rtTA mice to generate mice with inducible epidermal expression of Ova. The TGO construct encodes a fusion protein linking the transferrin receptor transmembrane domain, green fluorescent protein (GFP) and amino acids 230–359 of chicken ovalbumin. To create K5/TGO/DO11 mice, K5-rtTA/TRE-TGO mice were crossed onto the DO11.10 TCR-transgenic background. In all experiments, mice were gender-matched and between 5 and 24 weeks of age. All mice were bred and maintained in a specific pathogen-free facility in accordance with the guidelines of the Laboratory Animal Resource Center of the University of California San Francisco.

Inflammatory skin disease model. To induce expression of the TGO transgene, K5/TGO/DO11 mice were maintained on 1 g kg⁻¹ doxycycline chow. A 12-point clinical scoring scale was used to quantify skin disease. Scaling, alopecia, erythema and level of activity were each given a score from 0 to 3. Individual scores were summed and mean scores per group are displayed.

T_{reg} cell depletion. PC61 (anti-CD25 monoclonal antibody) or isotype control were injected intraperitoneally at 10 and 3 days before doxycycline treatment. For depletion at the height of disease, PC61 monoclonal antibody was injected on days 10 and 11 after starting K5/TGO/DO11 mice on doxycycline. For experiments in which CD25⁺ cells were deleted before antigen re-expression, K5/TGO/DO11 mice that had been off doxycycline for >30 days were treated with PC61 at 7 and 3 days before restarting doxycycline.

In vitro and in vivo suppression assays. K5/TGO/DO11 mice were crossed to Foxp3-GFP transgenic mice. K5/TGO/DO11/Foxp3-GFP mice were started on doxycycline and 6 days later, T_{reg} cells were isolated. For *in vitro* suppression assays, T_{reg} cells were cultured with CFSE-labelled SDLN cells isolated from DO11.10⁺/Rag2^{-/-}/Thy1.1⁺ mice and Ova peptide-pulsed bone-marrow-derived dendritic cells. For *in vivo* suppression assays, purified T_{reg} cells were mixed 1:1 with CFSE-labelled DO11 effector cells from DO11.10⁺/Rag2^{-/-}/Thy1.1⁺ mice. Cell mixtures were injected intravenously into K5/TGO mice and recipient mice were started on doxycycline.

Full Methods and any associated references are available in the online version of the paper at www.nature.com/nature.

Received 9 April; accepted 21 October 2011.

Published online 27 November 2011.

1. James, W. D. *Andrews' Diseases of the Skin: Clinical Dermatology* (Saunders Elsevier, 2006).

2. Lara-Corrales, I. & Pope, E. Autoimmune blistering diseases in children. *Semin. Cutan. Med. Surg.* **29**, 85–91 (2010).
3. Sabatos-Peyton, C. A., Verhagen, J. & Wraith, D. C. Antigen-specific immunotherapy of autoimmune and allergic diseases. *Curr. Opin. Immunol.* **22**, 609–615 (2010).
4. Diamond, I., Owolabi, T., Marco, M., Lam, C. & Glick, A. Conditional gene expression in the epidermis of transgenic mice using the tetracycline-regulated transactivators tTA and rTA linked to the keratin 5 promoter. *J. Invest. Dermatol.* **115**, 788–794 (2000).
5. Murphy, K. M., Heimberger, A. B. & Loh, D. Y. Induction by antigen of intrathymic apoptosis of CD4⁺CD8⁺TCR^{lo} thymocytes *in vivo*. *Science* **250**, 1720–1723 (1990).
6. Wada, N. *et al.* Aire-dependent thymic expression of desmoglein 3, the autoantigen in pemphigus vulgaris, and its role in T-cell tolerance. *J. Invest. Dermatol.* **131**, 410–417 (2011).
7. Mouquet, H. *et al.* Expression of pemphigus-autoantigen desmoglein 1 in human thymus. *Tissue Antigens* **71**, 464–470 (2008).
8. Gavin, M. A., Clarke, S. R., Negrou, E., Gallegos, A. & Rudensky, A. Homeostasis and anergy of CD4⁺CD25⁺ suppressor T cells *in vivo*. *Nature Immunol.* **3**, 33–41 (2002).
9. Fontenot, J. D. *et al.* Regulatory T cell lineage specification by the forkhead transcription factor Foxp3. *Immunity* **22**, 329–341 (2005).
10. Wing, K. *et al.* CTLA-4 control over Foxp3⁺ regulatory T cell function. *Science* **322**, 271–275 (2008).
11. Miyara, M. *et al.* Functional delineation and differentiation dynamics of human CD4⁺ T cells expressing the FoxP3 transcription factor. *Immunity* **30**, 899–911 (2009).
12. Dudda, J. C., Perdue, N., Bachtanian, E. & Campbell, D. J. Foxp3⁺ regulatory T cells maintain immune homeostasis in the skin. *J. Exp. Med.* **205**, 1559–1565 (2008).
13. Kurtulus, S., Tripathi, P., Opferman, J. T. & Hildeman, D. A. Contracting the “mus cells”—does down-sizing suit us for diving into the memory pool? *Immunol. Rev.* **236**, 54–67 (2010).
14. Akbar, A. N., Vukmanovic-Stejic, M., Taams, L. S. & Macallan, D. C. The dynamic evolution of memory and regulatory CD4⁺ T cells in the periphery. *Nature Rev. Immunol.* **7**, 231–237 (2007).

Supplementary Information is linked to the online version of the paper at www.nature.com/nature.

Acknowledgements We thank C. Benetiz for assistance with animal husbandry, S. Isakson for genotyping, S.-w. Jiang and M. Lee for cell sorting, and K. Ravid and G. Martin for derivation of TRE-TGO transgenic mice. We thank S. Ziegler, Benaroya Research Institute, for transgenic mice. M.D.R. is supported by a Dermatology Foundation Career Development Award and the UCSF Department of Dermatology. This work was partially funded through NIH grants P01 AI35297, R01 AI73656 and U19 AI56388 (to A.K.A.); NIH grant AR055634 to (A.M.-R.); and the Scleroderma Research Foundation (A.M.-R.). I.K.G. is supported by an Erwin Schrodinger Fellowship from the Austrian Science Fund (FWF), J2997-B13.

Author Contributions M.D.R. and I.K.G. contributed equally to this work and designed the studies, performed the experiments and analysed the data. M.D.R. and A.K.A. wrote the manuscript. J.S.P. collected and analysed data as well as helped with mouse husbandry. K.L. engineered and derived the TRE-TGO mice in the laboratory of A.M.-R. A.K.A. oversaw all study design and data analysis. A.M.-R. was involved in study design and data analysis. All authors discussed results and commented on the manuscript.

Author Information Reprints and permissions information is available at www.nature.com/reprints. The authors declare no competing financial interests. Readers are welcome to comment on the online version of this article at www.nature.com/nature. Correspondence and requests for materials should be addressed to A.K.A. (Abul.Abbas@ucsf.edu).

METHODS

Mouse model. K5/rTA mice were generated as described⁴. The TGO construct encodes for a fusion protein linking the transferrin receptor transmembrane domain (Tfr-tm), GFP and amino acids 230–359 of chicken ovalbumin (Ova)¹⁵. TGO was cloned upstream of the tetracycline response element (TRE2) and transgenic mice with stable incorporation of the TRE-TGO construct were generated. GFP is expressed at levels too low to detect in these mice. TRE-TGO mice were crossed with K5-rTA mice to generate mice with inducible epidermal expression of Ova. To create K5/TGO/DO11 mice, K5-rTA/TRE-TGO mice were crossed onto the DO11.10 TCR transgenic background, which expresses Ova-specific CD4⁺ T cells⁵. KJ1-26 monoclonal antibody (KJ) specifically recognizes the DO11 TCR. All mice were bred and maintained in a specific pathogen-free facility in accordance with the guidelines of the Laboratory Animal Resource Center of the University of California San Francisco.

T_{reg} cell depletion. Anti-CD25 monoclonal antibody (PC61) preferentially depletes T_{reg} cells¹⁶, especially in the BALB/c strain¹⁷. PC61 or isotype control (UCSF Monoclonal Antibody Core) were injected (0.5 mg per mouse) intraperitoneally at 10 and 3 days before doxycycline treatment. PC61 treatment of TGO/DO11 mice results in a > 90% reduction of CD4⁺CD25⁺Foxp3⁺ DO11 cells at 7 days after injection (Supplementary Fig. 4). For depletion of CD25⁺ cells at the height of disease, PC61 monoclonal antibody (0.5 mg per mouse) was injected on days 10 and 11 after starting K5/TGO/DO11 mice on doxycycline chow. For experiments in which CD25⁺ cells were deleted before antigen re-expression, K5/TGO/DO11 mice that had been off doxycycline for >30 days were treated with PC61 at 7 and 3 days before restarting doxycycline.

Inflammatory skin disease model. To induce expression of the TGO transgene in the skin, K5/TGO/DO11 mice were maintained on 1 g kg⁻¹ doxycycline chow

(Bio-Serv). A 12-point clinical scoring scale was used to quantify skin disease. The clinical parameters of scaling, alopecia, erythema and level of activity were each given a score from 0 to 3. Scores for individual parameters were summed, and each mouse was assigned a clinical severity score out of 12. Individual mouse scores were averaged to get mean clinical scores per group.

In vitro and in vivo suppression assays. K5/TGO/DO11 mice were crossed to Foxp3-GFP transgenic mice⁹, provided by A. Rudensky. K5/TGO/DO11/Foxp3-GFP mice were started on doxycycline chow or left untreated and 6 days later, SDLN cells were isolated and purified by FACS. For *in vitro* suppression assays, cells were cultured in varying ratios with freshly isolated CFSE-labelled SDLN cells isolated from DO11.10⁺/Rag2^{-/-}/Thy1.1⁺ mice and Ova peptide-pulsed bone-marrow-derived dendritic cells. For *in vivo* suppression assays, FACS-purified T_{reg} cells were mixed 1:1 with CFSE-labelled DO11 effector cells from DO11.10⁺/Rag2^{-/-}/Thy1.1⁺ mice. Cell mixtures were injected intravenously into K5/TGO mice and recipient mice were started on doxycycline chow. Three days later, SDLN cells were isolated and CFSE dilution and intracellular cytokine expression by DO11.10⁺/Thy1.1⁺ cells was quantified by flow cytometry.

15. Saff, R. R., Spanjaard, E. S., Hohlbaum, A. M. & Marshak-Rothstein, A. Activation-induced cell death limits effector function of CD4 tumor-specific T cells. *J. Immunol.* **172**, 6598–6606 (2004).
16. Setiady, Y. Y., Coccia, J. A. & Park, P. U. *In vivo* depletion of CD4⁺FOXP3⁺ Treg cells by the PC61 anti-CD25 monoclonal antibody is mediated by FcγRIII⁺ phagocytes. *Eur. J. Immunol.* **40**, 780–786 (2010).
17. Tenorio, E. P., Fernández, J., Olguín, J. E. & Saavedra, R. Depletion with PC61 mAb before *Toxoplasma gondii* infection eliminates mainly Tregs in BALB/c mice, but activated cells in C57BL/6J mice. *FEMS Immunol. Med. Microbiol.* **62**, 362–367 (2011).