

DR LINDA P. SIZIBA (Orcid ID : 0000-0002-4773-252X)

DR SUSANNE HARNER (Orcid ID : 0000-0001-6733-5823)

PROFESSOR CHRISTIAN APFELBACHER (Orcid ID : 0000-0003-3805-8219)

DR JON GENUNEIT (Orcid ID : 0000-0001-5764-1528)

PROFESSOR MICHAEL KABESCH (Orcid ID : 0000-0003-0697-1871)

Article type : Original Article

Food allergy in infants assessed in two German birth cohorts 10 years after the EuroPrevall Study

Daniela V. Röhl¹, Susanne Brandstetter^{2,3}, Linda P. Siziba⁴, Dietrich Rothenbacher⁵ Vincent D. Gaertner¹, Susanne Harner¹, Angela Köninger^{2,6}, Christian Apfelbacher^{2,7}, Michael Melter^{2,3}, Jon Genuneit^{* 4,5}, Michael Kabesch^{*1,2} and the KUNO Kids study group

¹Department of Pediatric Pneumology and Allergy, University Children`s Hospital Regensburg (KUNO), of the University of Regensburg and the Order of St. John at the St. Hedwig Hospital, Regensburg, Germany

²Science and Innovation Campus Regensburg (WECARE) of the Order of St. John at the St. Hedwig Hospital, Regensburg, Germany

³University Children`s Hospital Regensburg (KUNO) of the University of Regensburg and the Order of St. John at the St. Hedwig Hospital, Regensburg, Germany

⁴ Pediatric Epidemiology, Department of Pediatrics, Medical Faculty, Leipzig University, Leipzig, Germany

⁵Institute of Epidemiology and Medical Biometry, Ulm University, Ulm, Germany

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/PAI.13689](#)

This article is protected by copyright. All rights reserved

⁶Department of Obstetrics and Gynecology of the University of Regensburg and the Order of St. John
at the St. Hedwig Hospital

⁷Institute of Social Medicine and Health Systems Research, Otto von Guericke University Magdeburg, Germany

* These authors contributed equally

Running title: Food allergy in two German birth cohorts

Correspondence:

Michael Kabesch, M.D.

University Children's Hospital Regensburg (KUNO) at the St. Hedwig Hospital of the Order of St. John

Department of Pediatric Pneumology and Allergy

Steinmetzstr. 1-3

D-93049 Regensburg, Germany

Phone: +49-941-369-5900

Fax: +49-941-369-5802

E-mail: Michael.kabesch@barmherzige-regensburg.de

Word count: max 2500 words (PAI)

Number of tables and figures: 6

Supplementary material: 2 tables

Conflict of interest statement

D. Rothenbacher received research grants from Danone Nutritia Research and J. Genuneit is the project manager of research grants from Danone to both Ulm University

and Leipzig University in relation to studies of the composition of breast milk including data of the Ulm SPATZ Health Study. All authors declare that they have no competing financial or personal interests.

Funding

The KUNO Kids study is funded by research grants of the EU (HEALS: 603946) and the German Federal Ministry for Education and Research (SYSINFLAME: 01ZX1306E and NAMIBIO: 01EA2108B). Further financial support was provided by the University Children's Hospital of the University of Regensburg and the Order of St. John. The funding bodies had neither influence on the design of the study, on the collection, analysis and interpretation of data, nor in the writing of the manuscript. The Ulm SPATZ Health Study was funded through an unrestricted grant by the Medical Faculty of Ulm University and by the German Federal Ministry for Education and Research (NAMIBIO: 01EA2108B).

Abstract:

Background: The prevalence of food allergies (FA) in children increased rapidly at the turn of the century. The EuroPrevall study identified Germany as a country with very high prevalence of FA at that time. Using two large German birth cohorts we provide an update of the status quo ten years later.

Methods: KUNO Kids and Ulm SPATZ Health studies are two ongoing prospective birth cohorts. Information on FA was obtained by questionnaires at birth and after 6, 12 and 24 months. Univariable and multivariable logistic regression analyses were performed to investigate risk factors during pregnancy, birth and early childhood.

Results: In 1139 and 1006 children from KUNO Kids and SPATZ the point prevalence of parent-reported FA symptoms at the ages of 1 and 2 years was 13.2% (95% CI: 11.2-15.2) and 13.9% (95% CI: 11.5-17.2) in KUNO Kids. Doctor's diagnosed FA at 1 and 2 years was 2.4% (95% CI: 1.6-3.4) and 2.7% (95% CI: 1.2-4.3) in KUNO Kids and 2.3% (95% CI: 1.3-3.6) and 3% (95% CI: 2.0-4.5) in SPATZ. Cow's milk and citrus fruits were most frequently suspected by parents to cause FA symptoms. Atopy in the child was associated with a higher frequency of FA at any time, whereas atopy in first degree relatives was only associated with FA at year 1. Smoke exposure during pregnancy was a risk for FA at age 2.

Conclusion: The prevalence of food allergy seems to have plateaued in the last 10 years in Germany. FA is often suspected by parents but only rarely diagnosed by oral food challenge. Risk factor analysis may help to establish personalized health approaches.

Keywords: Food hypersensitivity, infant, birth cohort, epidemiology

Key messages:

In two ongoing prospective birth cohorts in Germany the prevalence of parent-reported FA in Germany was assessed and did not increase further in comparison to the EuroPrevall study ten years ago. Parents suspect a multitude of different foods to cause

FA symptoms and oral food challenge is seldom used to confirm the diagnosis. Risk factors for FA include atopy and smoke exposure, seem to be age specific and could provide targets for personalized medicine approach.

Abbreviations used

FA	Food allergy
OR	Odds ratio
BMI	Body mass index
CI	Confidence interval
DBPCFC	Double-blind placebo-controlled food challenge
GDM	Maternal gestational diabetes
SD	Standard Deviation

Introduction

Within the last five decades of the 20th century, the prevalence of atopic diseases like asthma, atopic dermatitis and allergic rhinitis increased dramatically in children across the world¹⁻³ and remained high^{4,5}. Food allergy (FA) seems to be the epidemic of the new century with rapidly increasing prevalence rates³, becoming a major health burden in many countries, especially in children^{6,7}. The EuroPrevall birth cohort is the first and largest study to investigate FA in Europe⁸. When recruitment was done in 2005, Germany was identified as one of the European countries with the highest prevalence of FA^{9,10}. However, it remains unclear whether FA further increased since then and if risk factors for the disease changed over time.

Therefore, we investigated the prevalence of FA in infants and young children approximately 10 years after the EuroPrevall study in two large and currently ongoing German birth cohorts, the KUNO Kids Health study (KUNO Kids)¹¹ and the Ulm SPATZ Health Study (SPATZ)^{12,13}. Furthermore, we explored the role of factors potentially associated with FA at ages 1 and 2 years, such as medical features of pregnancy and birth, nutrition of the child and environmental exposures during pregnancy and infancy¹⁴⁻¹⁶ in KUNO Kids and sought for replication in SPATZ. Finally, we compared results to extractable EuroPrevall data.

Methods

Description of populations

The KUNO Kids Health study

KUNO Kids is an ongoing prospective birth cohort started in 2015 which aims to explore the determinants of child health in a holistic approach. Recruitment is still ongoing. For the purpose of this study, we used data from the first and second phase of the study, which comprises 3100 newborns and their families recruited between June 2015 and March 2019. Data captured by follow-up questionnaires received by March 2020 were included in statistical analysis.

The precise methodology and design of the study have been described elsewhere¹¹. Briefly, written informed consent was obtained from all eligible mothers who gave birth at the children's and women's hospital St. Hedwig in Regensburg. Exclusion criteria were unavailability for informed consent (i.e., insufficient German language skills) and mother's age below the legal age of 18. The study design and procedures were approved by the ethics committee of the University of Regensburg (file number: 14-101-0347). An interview was conducted within days after birth and follow-up questionnaires were sent to the families after four weeks, six months and on every birthday of the child to collect additional information on the index child and the family. Socio-demographic information, medical history, feeding practices, lifestyle behaviors (nutrition, physical activity, smoking, alcohol consumption), and psychosocial constructs were collected from parents and siblings. Where available, validated questions and measurement instruments were used to allow for joint analyses of data with pre-existing studies, including SPATZ.

In KUNO Kids, parent-reported FA was defined in concordance with the EuroPrevall (using the same questionnaire) as a positive history of gastrointestinal, dermatological or respiratory symptoms, which appeared in timed correlation to the food intake without other explanations for the symptoms. Gastrointestinal symptoms were nausea, vomiting, diarrhea, abdominal pain, constipation and soft stools. Dermatological reactions comprised urticaria, angioedema, itching and the appearance or worsening of eczema. Respiratory symptoms were defined as wheezing, dyspnoea, coughing, stridor and allergic rhinitis symptoms. Information on signs, symptoms, point of time and clinical

diagnostics of FA was collected using a validated questionnaire at 6 months, 1 and 2 years postpartum. Additional information on lifestyle-related factors like tobacco exposure during pregnancy, birth mode, infant feeding practices, environmental exposures during infancy, maternal and infant antibiotic intake as well as atopic diseases within the family was obtained from the KUNO Kids database. Parent-reported FA in KUNO Kids was defined as a parent report of FA related symptoms in the past 12 months, while the doctor reported FA was defined as a positive report of a doctor's diagnosis of FA at 1 and 2 years.

The Ulm SPATZ Health Study

The Ulm SPATZ Health Study is an ongoing birth cohort study. A total of 1006 newborns and their 970 mothers were enrolled into the study from the general population soon after delivery at the University Medical Centre Ulm, between April 2012 and May 2013. A detailed description of the study can be found elsewhere^{12,13}. Mothers were excluded from the study if (i) their German language skills were insufficient to understand the study procedure and consent forms, (ii) they had outpatient child birth, (iii) they were <18 years of age, (iv) there was postpartum transfer of mother or child to an intensive care unit, or stillbirth. Participation in the study was entirely voluntary after informed consent was given. The study was approved by the Ethics board of Ulm University (No. 311/11).

Demographic, lifestyle and birth-related data including child gender, delivery mode, birth weight, maternal age, education, maternal body mass index (BMI), smoking status (within one year prior to delivery), maternal, paternal and child history of allergic disease was collected using self-administered questionnaires, hospital electronic charts and paper-documented routine screening during pregnancy. Smoking status and alcohol consumption were assessed and defined as previously explained¹².

A reported doctor's diagnosis of FA in the past 12 months was assessed at 1 year, 2 years and at each further yearly follow-up by separate self-administered questionnaires from parents and the children's primary care pediatricians. In this case, the reports were distinguished between "parent-reported" FA and "pediatrician-reported" FA, however,

both were reports of a doctor's diagnosis. The parent-reported FA in SPATZ equals the doctor's diagnosed FA in the KUNO Kids study. FA symptoms were not assessed in SPATZ.

Statistical analysis

The statistical analysis was conducted using Statistical Package for Social Sciences® (SPSS.24) and SAS 9.4 (The SAS® Institute, Cary, NC, USA). P-values <0.05 were considered statistically significant. Descriptive statistics are displayed as median and interquartile range and mean and standard deviation for non-parametric and parametric data, respectively. First, univariable logistic regression analyses were used to explore the association of the different previously suggested risk factors with FA at age 1 and at age 2, in the KUNO Kids and in the Ulm SPATZ cohort, respectively. Then, multivariable regression modeling was performed, for the respective FA outcome at age 1 and at age 2 in both cohorts. Risk factors were included in the multivariable models if they were associated with the outcome ($p < 0.2$) at age 1 or at age 2 in the respective cohort. Odds ratios (OR) and 95% confidence intervals (CI) are displayed.

Results

Of the 3100 newborns and their families enrolled in the KUNO Kids study until March 2020, 2679 were at least one year old when the dataset for this analysis was closed. FA data at one or more time points (6 months, 1 year, 2 years) were available from 1139 infants. Compared to the total study population, caesarean section, having older siblings, exposure to household smoking and a migration background were less frequent in children with completely available FA questionnaires until 1 year of age, while older mothers and a higher educational level were more frequent (Table 1 and supplementary Table 1).

Cumulatively, 23.9% of parents in the KUNO Kids study reported that their child had an adverse reaction to food in the first 2 years of life, not distinguishing between IgE and non-IgE mediated self-reported food allergy. The point prevalence was slightly higher at age 6 months and gastrointestinal and dermatological symptoms were most frequent (Figure 1). Cow's milk, citrus fruits and wheat were most frequently described by parents as allegedly causative, while eggs were mentioned less frequently (Figure 2). In infants younger than 4 months, parents mentioned cow's milk most often to be responsible for the symptoms and fruits in 4 to 6 months old infants. (Figure 2). Some parents (143/1137) reported that a broad variety of fruits and vegetables not listed on the questionnaire (non-core foods) caused putative FA symptoms. For more than half (59.7%) of the children with reported FA at 6 months no symptoms at the age of 1 year were reported. Of those with suspected FA at age 1 year, 58.1% did not report FA at age 2 years. Only 1.9%, 2.4% and 2.7% of parent-reported FA were diagnosed by a doctor at 6 months, age 1 and 2 years respectively. Diagnostic confirmation by oral food challenge was only obtained in 30.8% of the 6 months cases, in 21.1% of the 1-year cases and 10% of the 2-year cases.

In a univariate analysis we analyzed risk factors for FA suggested by previous studies for their association with FA at age 1 (supplementary Table 2a) and age 2 (supplementary Table 2b) in our dataset. The risk of parent-reported FA symptoms at 1 and 2 years of age was significantly higher if the child had other atopic diseases at the time. The presence of atopic diseases in first degree relatives was related to an increased risk for

food allergy at age 1 year, but not at age 2 years. Children with low birth weight had a decreased risk for FA at age 2 years while animal contact of the child or the mother during pregnancy (apart from farms) increased the risk for FA at this timepoint. Of note, passive smoke exposure of the mother was a strong predictor for FA at age 2 years. Only at age 1 year, a non-significant trend towards an increased risk to develop FA symptoms was observed in children of obese mothers compared to children of normal-weight mothers (measured before pregnancy). Of the 1006 infants enrolled into the SPATZ study, 4.1% of the infants had at least one food allergy reported either by the caring pediatrician or the parent (parental report of the doctor's diagnosis) within the first year of life. Common risk factors for FA were also investigated in the SPATZ study at age 1 and 2 years, whenever available (supplementary Tables 3a and 3b). Again, the presence of other allergies in the child was statistically significantly associated with FA (year 1 and 2) and atopic diseases in first degree relatives was a significant predictor for FA at age 1 year.

Next, we performed multiple regression modeling for the respective FA outcomes at age 1 and at age 2 in both cohorts based on risk factors which $p < 0.2$ in the univariate analyses. For KUNO Kids, atopic disease in the child was strongly associated with FA at age 1 and 2 years while a positive family history was associated only with FA at age 1 year (Table 2a and b). Gestational diabetes in the mother and exposure to traffic showed suggestive associations ($p < 0.1$) at age 1 and 2 years, respectively. Multivariate regression in the SPATZ study confirmed the association between another atopic disease in the child and FA at both ages (Table 3a and b). Furthermore, associations between male gender for FA at age 1 and exposure to cigarette smoking with FA at age 1 (suggestive) and age 2 (significant) were identified.

Finally, we compared the data from the KUNO Kids and SPATZ studies with results from EuroPrevall acquired approximately 10 years earlier. (Table 4). On average, mothers in the KUNO Kids study were approximately 4 years older when compared to mothers from the EuroPrevall and SPATZ studies. The frequency of caesarean section birth was lower in the KUNO Kids and SPATZ studies, while antibiotic use during delivery was considerably higher in KUNO Kids compared to EuroPrevall and SPATZ. Also, the prevalence of maternal smoking and passive smoking varied significantly between

Accepted Article

studies. Data on overall parent-reported and doctor diagnosed FA frequencies in EuroPrevall were extracted from the manuscript by Schoemaker et al.¹⁷. Frequencies were in a similar range in the two recent birth cohort studies KUNO Kids and SPATZ as well as the reported German and overall EuroPrevall data for parent-reported FA symptoms and doctor's diagnosis of FA at age 2 years.

Discussion

Data from KUNO Kids and SPATZ show that the prevalence of parent-reported FA and doctor's diagnosis of FA in infants at age 1 and 2 years did not differ substantially from results obtained by the EuroPrevall study 10 years ago. Non-core foods seem to play an important role in FA now. When classic risk factors for FA were investigated in the current studies, family history for atopy was associated with FA at age 1 but less so at age 2, while maternal passive smoke exposure was a risk factor for FA at age 2. Our study demonstrated that FA seems to be a highly dynamic entity with various changes in point prevalence, risk factors and predictors across over the first two years of life.

Some differences between study populations were observed, but most of the factors that have been associated with an increased risk of FA in previous studies were more pronounced in KUNO Kids and SPATZ. Due to the population structure of KUNO Kids with older, highly educated mothers an over-reporting of food allergy symptoms may be expected and consequently, FA would be overestimated. However, compared to data available in EuroPrevall at age 2 years for Germany and Europe, no difference was observed. Even though methodology differed, and the two study regions investigated here are Regensburg (KUNO Kids) and Ulm (SPATZ) and not Berlin (EuroPrevall), results seem comparable enough to assess trends over time. For the prevalence of doctor's diagnosis of food allergy, reports at age 1 and 2 years are very similar for KUNO Kids and SPATZ (table 3). For EuroPrevall, prevalence data for Europe and Germany were taken from the publication of Schoemaker¹⁷. Since the focus of that study is on cow's milk allergy, the calculated rate of doctor's diagnosed FA may be underestimated for EuroPrevall Germany. Taking all these factors into account, there is no indication of a significant rise in the prevalence of FA in Germany over the last 10 years. This is of interest, because at the turn of the century a worldwide trend towards an increasing FA prevalence was suggested⁶ when hospital admissions for food allergy quintupled³ and the number of children with FA increased by twelve times between 1995 and 2006². Especially in children below 2 years, an increase in food-induced anaphylaxis was reported⁷. Our data now suggests that the prevalence of food allergies reached a plateau in Germany, similar to what has been observed for many other atopic diseases, such as asthma, years earlier¹⁸⁻²⁰.

In KUNO Kids and SPATZ, assessments of FA were based on questionnaires only, while in EuroPrevall questionnaires and a double-blind placebo-controlled food challenge (DBPCFC) were applied. Several studies suggest that the prevalence of parent-reported FA overestimates the actual FA prevalence^{21,22,10}. In our study, parents suspected a multitude of different foods, especially fruit and vegetables, to be responsible for adverse reactions. While known allergens like cow's milk, wheat, hen's egg, soy and peanut are responsible for IgE mediated allergy, these putative causative agents seem to represent a fear of parents rather than actual food allergy. Citrus fruit and histamine-rich foods can cause perioral itching and redness (especially in children with eczema) which is not due to IgE-mediated allergy. Further investigations into that observation are needed.

The results of KUNO Kids showed that FA in infants is often not diagnosed with DBPCFC: Two thirds at the age of 6 months, 79% at the age of 1 year and 90% at the age of 2 years were diagnosed with other methods than DBPCFC which suggests that better education of doctors and parents along with increasing resources for DBPCFC may be necessary and needed to avoid unnecessary and even harmful dietary restrictions in children based on FA assumptions. However, results can be comparable between studies for questionnaire derived data, especially as KUNO Kids used the same questions as EuroPrevall (for symptoms of FA) and SPATZ (for doctor-diagnosed FA). Simple, questionnaire-based assessment tools are key for replication and comparisons between populations as previously demonstrated by the worldwide ISAAC project²³.

Some previously suggested risk factors²⁴⁻²⁶ were assessed here by univariate analysis in KUNO Kids and SPATZ and followed up in a multivariate analysis. Interestingly, a family history of atopic diseases in first degree relatives was associated with an increased risk for FA at age 1 but not age 2 in both studies. This may either indicate that the genetic impact on FA decreases with age or that pathogenesis of food allergy at age two years differs from FA at age one year. Accordingly, we demonstrated an extensive turnaround in presence and absence of FA symptoms over that time span. A (changeable) risk factor for FA at age 2, but less so at age 1, is passive smoke exposure either through maternal

smoking during pregnancy or via postnatal environmental exposure in both birth cohorts. This may present a valuable opportunity for future efforts of intervention and prevention.

In conclusion, our study suggests that the prevalence of FA in Germany has reached a plateau in infants and young children. Numerous allergens other than core foods are associated with FA in infants now. While better diagnosis of FA with DBPCFC in clinical practice is needed, questionnaire data can provide valuable comparisons between different populations and inform about time trend and regional differences. Risk factors for the development of FA seem to be age specific and that needs to be taken into account more carefully for further FA prediction analysis, when personalized medicine may be applied to FA.

Impact statement:

Based on data from two current birth cohorts we suggest that FA prevalence reached a plateau in Germany when compared to EuroPrevall data from 10 years ago. Numerous non-core foods contribute to FA nowadays and risk factors for FA may contribute differently to FA occurring at various timepoints over the course of infancy.

Acknowledgment:

We are grateful to all participants in the KUNO Kids and the Ulm SPATZ Health Studies and to their parents for filling out questionnaires over years. We thank all the students who contributed time and effort in the collection of data and all the partners in the study for their input and support.

Authors' contributions:

Study design: MK, JG

Data collection: MK, DVR, JCP, MM, CA, AP, CT, VP, JR, LN, JP, JG, DR

Statistical analysis and data interpretation: SB, VDG, DVR, LS, SH, CA, JG, MK

Manuscript writing: DVR, VDG, DR, SB, SH, MK

References

1. Pearce N, Ait-Khaled N, Beasley R, et al. Worldwide trends in the prevalence of asthma symptoms: phase III of the International Study of Asthma and Allergies in Childhood (ISAAC). *Thorax*. 2007;62(9):758-766. doi:10.1136/thx.2006.070169
2. Mullins RJ. Paediatric food allergy trends in a community-based specialist allergy practice, 1995-2006. *Med J Aust*. 2007;186(12):618-621.
3. Gupta R, Sheikh A, Strachan DP, Anderson HR. Time trends in allergic disorders in the UK. *Thorax*. 2007;62(1):91-96. doi:10.1136/thx.2004.038844
4. Schuh C, Fritscher LG, Chapman KR, Fritscher CC. The prevalence of asthma and atopy in schoolchildren from Porto Alegre, Brazil, has plateaued. *Respir Med*. 2015;109(3):308-311. doi:10.1016/j.rmed.2015.01.014
5. Braun-Fahrlander C, Gassner M, Grize L, et al. No further increase in asthma, hay fever and atopic sensitisation in adolescents living in Switzerland. *Eur Respir J*. 2004;23(3):407-413. doi:10.1183/09031936.04.00074004
6. Prescott SL, Pawankar R, Allen KJ, et al. A global survey of changing patterns of food allergy burden in children. *World Allergy Organ J*. 2013;6(1):21. doi:10.1186/1939-4551-6-21
7. Motosue MS, Bellolio MF, van Houten HK, Shah ND, Campbell RL. National trends in emergency department visits and hospitalizations for food-induced anaphylaxis in US children. *Pediatr Allergy Immunol*. 2018;29(5):538-544. doi:10.1111/pai.12908
8. McBride D, Keil T, Grabenhenrich L, et al. The EuroPrevall birth cohort study on food allergy: baseline characteristics of 12,000 newborns and their families from nine European countries. *Pediatr Allergy Immunol*. 2012;23(3):230-239. doi:10.1111/j.1399-3038.2011.01254.x

9. Burney P, Summers C, Chinn S, Hooper R, van Ree R, Lidholm J. Prevalence and distribution of sensitization to foods in the European Community Respiratory Health Survey: a EuroPrevall analysis. *Allergy*. 2010;65(9):1182-1188. doi:10.1111/j.1398-9995.2010.02346.x
10. Xepapadaki P, Fiocchi A, Grabenhenrich L, et al. Incidence and natural history of hen's egg allergy in the first 2 years of life-the EuroPrevall birth cohort study. *Allergy*. 2016;71(3):350-357. doi:10.1111/all.12801
11. Brandstetter S, Toncheva AA, Niggel J, et al. KUNO-Kids birth cohort study: rationale, design, and cohort description. *Mol Cell Pediatr*. 2019;6(1):1. doi:10.1186/s40348-018-0088-z
12. Logan C, Zittel T, Striebel S, et al. Changing Societal and Lifestyle Factors and Breastfeeding Patterns Over Time. *Pediatrics*. 2016;137(5). doi:10.1542/peds.2015-4473
13. Braig S, Weiss JM, Stalder T, Kirschbaum C, Rothenbacher D, Genuneit J. Maternal prenatal stress and child atopic dermatitis up to age 2 years: The Ulm SPATZ health study. *Pediatr Allergy Immunol*. 2017;28(2):144-151. doi:10.1111/pai.12680
14. Gupta M, Sicherer SH. Timing of food introduction and atopy prevention. *Clin Dermatol*. 2017;35(4):398-405. doi:10.1016/j.clindermatol.2017.03.013
15. van Ginkel CD, van der Meulen GN, Bak E, et al. Retrospective observational cohort study regarding the effect of breastfeeding on challenge-proven food allergy. *Eur J Clin Nutr*. 2018;72(4):557-563. doi:10.1038/s41430-018-0117-y
16. Ben-Shoshan M, Soller L, Harrington DW, et al. Eczema in early childhood, sociodemographic factors and lifestyle habits are associated with food allergy: a nested case-control study. *Int Arch Allergy Immunol*. 2015;166(3):199-207. doi:10.1159/000381829
17. Schoemaker AA, Sprickelman AB, Grimshaw KE, et al. Incidence and natural history of challenge-proven cow's milk allergy in European children--EuroPrevall birth cohort. *Allergy*. 2015;70(8):963-972. doi:10.1111/all.12630
18. Prescott S, Allen KJ. Food allergy: riding the second wave of the allergy epidemic. *Pediatr Allergy Immunol*. 2011;22(2):155-160. doi:10.1111/j.1399-3038.2011.01145.x
19. Asher MI, Montefort S, Björkstén B, et al. Worldwide time trends in the prevalence of symptoms of asthma, allergic rhinoconjunctivitis, and eczema in childhood: ISAAC Phases One and Three repeat

multicountry cross-sectional surveys. *Lancet*. 2006;368(9537):733-743. doi:10.1016/S0140-6736(06)69283-0

20. Björkstén B, Clayton T, Ellwood P, Stewart A, Strachan D. Worldwide time trends for symptoms of rhinitis and conjunctivitis: Phase III of the International Study of Asthma and Allergies in Childhood. *Pediatr Allergy Immunol*. 2008;19(2):110-124. doi:10.1111/j.1399-3038.2007.00601.x
21. Venter C, Pereira B, Grundy J, et al. Incidence of parentally reported and clinically diagnosed food hypersensitivity in the first year of life. *J Allergy Clin Immunol*. 2006;117(5):1118-1124. doi:10.1016/j.jaci.2005.12.1352
22. Nwaru BI, Hickstein L, Panesar SS, Roberts G, Muraro A, Sheikh A. Prevalence of common food allergies in Europe: a systematic review and meta-analysis. *Allergy*. 2014;69(8):992-1007. doi:10.1111/all.12423
23. Flohr C, Weinmayr G, Weiland SK, et al. How well do questionnaires perform compared with physical examination in detecting flexural eczema? Findings from the International Study of Asthma and Allergies in Childhood (ISAAC) Phase Two. *Br J Dermatol*. 2009;161(4):846-853. doi:10.1111/j.1365-2133.2009.09261.x
24. Koplin JJ, Dharmage SC, Ponsonby A-L, et al. Environmental and demographic risk factors for egg allergy in a population-based study of infants. *Allergy*. 2012;67(11):1415-1422. doi:10.1111/all.12015
25. Koplin JJ, Peters RL, Ponsonby A-L, et al. Increased risk of peanut allergy in infants of Asian-born parents compared to those of Australian-born parents. *Allergy*. 2014;69(12):1639-1647. doi:10.1111/all.12487
26. Koplin JJ, Allen KJ, Tang MLK. Important risk factors for the development of food allergy and potential options for prevention. *Expert Rev Clin Immunol*. 2019;15(2):147-152. doi:10.1080/1744666X.2019.1546577

Tables

Table 1: Characteristics of the study population (KUNO Kids study)

	Total study population		Study sample with data	
	% or		on FA	
	mean $\pm$ SD		% or	
		n	mean $\pm$ SD	n
Child				
Male sex	51.2	2653	49.9	1134
Duration of pregnancy (wk)	39.5 $\pm$ 1.6	2625	39.6 $\pm$ 1.6*	1128
Caesarean section	28.6	2679	27.8	1139
Birth weight (g)	3359 $\pm$ 509	2638	3371 $\pm$ 500	1130
Antibiotic use	2.7	2600	2.2	1113
Older siblings	43.7	2636	38.5*	1130
Migration background ^a	28.0	1824	14.2*	1065
Mother				
Age (yrs)	32.7 $\pm$ 4.5	2651	33.1 $\pm$ 4.1*	1132
Pre-pregnancy BMI (kg/m ²)	23.8 $\pm$ 5.0	2614	23.7 $\pm$ 4.7	1121
Regular animal contact ^b	54.9	2601	53.8	1116
Higher educational level (>10 years)	56.5	2597	62.8*	1116
Smokers in the household	22.4	2601	17.0*	1114

FA – food allergy; SD - standard deviation; BMI - body mass index;

^a mother or father with non-German nationality or born not in Germany

^b regular contact (more than once per month) to any pet during pregnancy

*comparison with total study population yielded statistically significant difference ($p < 0.001$)

Table 2: Multivariable logistic regression analysis in KUNO Kids of potential risk factors: outcome parent-reported symptoms of FA

a) at age 1

	Odds ratio (95% CI)	p value
Child		
Caesarean section	0.828 (0.562 – 1.219)	0.339
Birth weight < 2800 g	0.817 (0.466 – 1.432)	0.480
Duration of breastfeeding < 6 month	0.920 (0.640 – 1.322)	0.653
Daily animal contact ^a	1.091 (0.725 – 1.642)	0.676
High exposure to traffic, exhaust gases	1.038 (0.739 – 1.459)	0.828
Other atopic disease ^b	3.681 (2.504 – 5.412)	< 0.001
Mother		
Pre-pregnancy BMI		
- underweight	1.894 (0.637 – 5.633)	0.251
- normal weight	<i>reference</i>	
- overweight	1.041 (0.692 – 1.571)	0.841
- obese	1.377 (0.822 – 2.305)	0.224
Gestational diabetes	1.503 (0.960 – 2.354)	0.075
Passive exposure to cigarette smoking	1.049 (0.689 – 1.598)	0.823
Regular animal contact	0.969 (0.641 – 1.466)	0.883
School education		
- low (no degree)	1.463 (0.721 – 2.969)	0.292
- intermediate (< 10 years)	<i>reference</i>	
- high (> 10 years)	1.120 (0.755 – 1.662)	0.572
Atopic disease of 1st degree relatives ^d	1.531 (1.042 – 2.250)	0.030

b) at age 2

	Odds ratio (95 % CI)	p value
Child		
Caes		
Birth weight < 2800 g	0.425 (0.141 – 1.277)	0.127
Duration of breastfeeding < 6 month	0.832 (0.478 – 1.450)	0.517
Daily animal contacts ^a	1.685 (0.894 – 3.177)	0.107
High exposure to traffic, exhaust gases	1.608 (0.953 – 2.712)	0.075
Other atopic disease ^b	2.705 (1.472 – 4.972)	0.001
Mother		
Pre-pregnancy BMI		
- underweight	-/(no cases with FA)	-/

- normal weight	<i>reference</i>	
- overweight	1.305 (0.707 – 2.409)	0.395
- obese	1.107 (0.506 – 2.420)	0.799
Gestational Diabetes	0.569 (0.240 – 1.347)	0.199
Passive exposure to cigarette smoking	1.569 (0.878 – 2.804)	0.128
Regular animal contact ^c	1.318 (0.697 – 2.492)	0.395
School education		
- low (no degree)	1.999 (0.722 – 5.535)	0.182
- intermediate (< 10 years)	<i>reference</i>	
- high (> 10 years)	1.081 (0.576 – 2.028)	0.808
Atopic disease of 1st degree relatives ^d	1.470 (0.809 – 2.669)	0.206

FA – food allergy; BMI - body mass index; 95% CI - 95% confidence interval

^a daily contact to dog, cat, hamster, guinea pig, mouse, rat or rabbit

^b atopic eczema, allergic rhinitis, allergic conjunctivitis or asthma

^c regular contact (more than once per month) to any pet during pregnancy

^d atopic eczema, allergic rhinitis, allergic conjunctivitis, asthma or food allergy in parents or blood-related siblings

Risk factors were included if they were associated with the 1-year or the 2-year outcome ($p < .2$) in univariable analysis; N= 509; R^2 : 0.123

Table 3: Multivariable logistic regression analysis in Ulm SPATZ health study of potential risk factors: outcome parent-reported FA

a) at age 1

	Odds ratio (95% CI)	p value
Child		
Male sex	1.909 (1.003 – 13.223)	0.049
Other atopic disease ^a	2.99 (3.037 – 26.321)	<0.001
Mother		
Gestational diabetes	1.677 (0.699 – 11.303)	0.145
Cigarette smoking during pregnancy	1.572 (0.852 – 7.174)	0.096
Antibiotic use during pregnancy	1.513 (0.673 – 7.777)	0.185
Atopic disease of 1st degree relatives ^b	1.844 (0.932 – 12.413)	0.064

FA – food allergy; 95% CI - 95% confidence interval

^a atopic eczema, hay fever, asthma

^b atopic eczema, hay fever, asthma in parents and blood-related siblings

Risk factors were included if they were associated with the 1-year or the 2-year outcome ($p < .2$) in univariable analysis; N= 729; R : 0.044

b) at age 2

	Odds ratio (95% CI)	p value
Child		
Male sex	1.443 (0.841 – 5.16)	0.113
Other atopic disease ^a	2.012 (1.639 – 10.01)	0.002
Mother		
Gestational diabetes	1.279 (0.456 – 5.876)	0.450
Cigarette smoking during pregnancy	1.627 (1.099 – 6.371)	0.030
Antibiotic use during pregnancy	1.235 (0.493 – 4.716)	0.464
Atopic disease of 1st degree relatives ^b	1.417 (0.807 – 4.998)	0.134

FA – food allergy; 95% CI - 95% confidence interval

^a atopic eczema, hay fever, asthma

^b atopic eczema, hay fever, asthma in parents and blood-related siblings

Risk factors were included if they were associated with the 1-year or the 2-year outcome ($p < .2$) in univariable analysis
N=788; R²: 0.026

Table 4: Comparison of study results from KUNO Kids (recruitment 2015-2019), SPATZ (2012-2013) and EuroPrevall (2005-2010)

	KUNO Kids % or mean $\pm$ SD	SPATZ % or mean $\pm$ SD	EuroPrevall Germany % or mean $\pm$ SD	EuroPrevall total (Europe) % or mean $\pm$ SD
Study sample	1139	1006	1570	12049
Symptom-based FA				
6 months	14.2 (95% CI: 12.3-16.2)	-	-	-
12 months	13.2 (95% CI: 11.2-15.2)	-	-	-
24 months	13.9 (95% CI: 11.5-17.2)	-	14.5 ^a (95% CI: 13.1-16.3)	16.0 (95% CI: 15.4-16.6)
Doctor's diagnosed FA				
6 months	1.9 (95% CI: 1.2-2.8)		-	-
12 months	2.4 (95% CI: 1.6-3.4)	2.3 (95% CI: 1.3-3.6)	-	-
24 months	2.7 (95% CI: 1.2-4.3)	3.0 (95% CI: 2.0-4.5)	2.0 ^b (95% CI: 1.4-2.8)	3.8 ^b (95% CI: 3.5-4.2)
Male sex	49.9	52.1	51.7	51.4
Duration of pregnancy (wk)	39.6 $\pm$ 1.6	38.8 $\pm$ 1.7	-	39.6 $\pm$ 1.5
Caesarean section	27.8	27.9	31.1	24.0
Birth weight (g)	3371 $\pm$ 500	3157 $\pm$ 948.3	-	3404 $\pm$ 512
Pets				
Dog	9.9	-	9.0	16.0
Cat	19.4	-	15.0	15.1
Other	9.4	-	15.1	12.2
Maternal age (yr)	33.1 $\pm$ 4.1	32.7 $\pm$ 4.7	31.4 $\pm$ 5.4	30.7 $\pm$ 5.2
Pre-pregnancy BMI (kg/m ²)	23.7 $\pm$ 4.7	24.8 $\pm$ 5.1	-	23.3 $\pm$ 4.3
Maternal antibiotic intake				
During pregnancy	21.0 ^c	11.1 ^c	18.6	21.1

During delivery	54.3 ^c	-	32.4	15.4
Cigarette smoking during pregnancy				
Active	1.3	-	10.8	10.6
Passive	21.7	-	34.7	17.4
Maternal FA	27.1	-	30.2	14.3
Other maternal atopic disease ^d	31.5	34.1	35.2	26.2
Paternal FA	24.2	-	19.0	8.6
Other paternal atopic disease ^d	32.9	35.4	29.2	20.9
Other atopic disease of siblings ^d	17.1	15.9	19.9	26.0

FA – food allergy; SD – standard deviation; 95% CI - 95% confidence interval; BMI – body mass index

^a EuroPrevall documented this frequency for self-reported FA in siblings of their study children (Schoemaker et. al., 2015, *Allergy*)

^b calculated from data from Schoemaker et al., 2015, *Allergy*, as percentage of children eligible for DBPCFC

^c different methods: KUNO kids used a specific question („Did you take any antibiotics during pregnancy/delivery?“), SPATZ used an open question after delivery about any medication intake during pregnancy

^d Atopic eczema, allergic rhinitis, allergic conjunctivitis or asthma

Figure 1: Prevalence and characteristics of parent-reported food allergy related symptoms over the first 2 years in the KUNO Kids study





Figure 2: First occurrence of parent-reported adverse reactions to food allergens in the first 2 years of life (n = 189) and proportion of core foods causing the adverse reactions reported for specific months in the KUNO Kids study



Supporting information:

Table S1: Baseline characteristics of study population with entirely available FA questionnaires for all timepoints (KUNO Kids study)

	% or mean $\pm$ SD	n
Child		
Male sex	49.9	1134
Mean duration of pregnancy (wk)	39.6 $\pm$ 1.6	1128
Caesarean section	27.8	1139
Birth weight (g)	3371 $\pm$ 500	1130
Duration of breastfeeding > 6 months	66.0	960
Introduction of solid food at the age of 4 – 6 months	85.4	904
Antibiotic use	10.1	1137
Pet ownership		
Dogs	9.9	1134
Cats	19.4	1134
Other animals ^a	9.4	1134
Older siblings	38.5	1130
Other atopic disease ^b	17.5	1138
Migration background	14.2	1065
Mother		
Age (yr)	33.1 $\pm$ 4.1	1132
Pre-pregnancy BMI (kg/m ²)	23.7 $\pm$ 4.7	1121
Gestational diabetes	15.4	1106
Cigarette smoking		
during pregnancy	1.3	1107
passive exposure	21.7	1138
Antibiotic use		
during pregnancy	21.0	1123
during delivery	54.3	1118
while breastfeeding	10.3	948
Higher educational level (>10 years)	62.8	1116
FA	27.1	1088
Other atopic disease ^b	31.5	1139
Father		
FA	24.2	1032
Other atopic disease ^b	32.9	1129
Siblings		
Other atopic disease ^b	17.1	375

FA – food allergy; SD - standard deviation; BMI – body mass index

^a hamster, guinea pig, mouse, rat, rabbit

^b atopic eczema, allergic rhinitis, allergic conjunctivitis, asthma

^c mother or father with non-German nationality or born not in Germany

Table S2a: Univariable logistic regression analysis in KUNO Kids of potential risk factors: outcome parent-reported symptoms of FA at age 1

	Odds ratio (95% CI)	p value	n
Child			
Male sex	0.967 (0.721-1.296)	.822	1124
Preterm birth	0.755 (0.378-1.508)	.426	1118
Caesarean section	0.792 (0.565-1.111)	.177	1129
Birth weight < 2800 g	0.736 (0.450-1.203)	.222	1120
Duration of breastfeeding < 6 months	1.159 (0.843-1.610)	.379	952
Introduction of solid food at the age of			
- < 4 months	1.459 (0.383-5.565)	.580	
- 4 – 6 months	reference		898
- > 6 months	0.883 (0.538-1.448)	.621	
Antibiotic use	1.084 (0.673-1.745)	.740	1127
Daily animal contact ^a	1.153 (0.859-1.547)	.342	1128
Being on a farm regularly	1.125 (0.720-1.757)	.606	1129
Regular contact to hay	0.827 (0.436-1.567)	.560	1129
High exposure to traffic, exhaust gases	1.108 (0.824-1.489)	.496	1128
Other atopic disease ^b	3.594 (2.571-5.025)	<.001	1128
Older siblings	0.857 (0.631-1.163)	.322	1120
Migration background ^c	1.201 (0.792-1.820)	.389	1055
Mother			
Maternal age > 34 years	0.871 (0.641-1.185)	.379	1122
Pre-pregnancy BMI			
- underweight	1.953 (0.787-4.848)	.149	
- normal weight	reference		1111
- overweight	1.137 (0.798-1.619)	.478	
- obese	1.518 (0.983-2.344)	.060	
Gestational diabetes	1.326 (0.897-1.960)	.157	1096
Cigarette smoking			
- during pregnancy	1.114 (0.786-1.578)	.545	1128
- passive exposure	1.102 (0.305-3.986)	.882	1098
Antibiotic use			
- during pregnancy	1.068 (0.746-1.528)	.721	1113
- during delivery	0.836 (0.623-1.123)	.235	1108
- while breastfeeding	1.280 (0.784-2.090)	.323	939
Being on a farm regularly			
- during pregnancy	1.322 (0.738-2.369)	.348	1118
- while breastfeeding	1.417 (0.773-2.596)	.259	1116
Regular animal contact ^d	1.165 (0.865-1.568)	.316	1106
School education			
- low (no degree)	1.441 (0.801-2.592)	.222	
- intermediate (< 10 years)	reference		1106
- high (> 10 years)	1.131 (0.810-1.579)	.470	
Atopic disease of 1st degree relatives ^e	1.665 (1.186-2.338)	.003	1129

FA – food allergy; 95% CI - 95% confidence interval; BMI – body mass index

^a daily contact to dog, cat, hamster, guinea pig, mouse, rat or rabbit

^b atopic eczema, allergic rhinitis, allergic conjunctivitis and asthma

^c mother or father with non-German nationality or born not in Germany

^d regular contact (more than once per month) to any pet during pregnancy

^e atopic eczema, allergic rhinitis, allergic conjunctivitis, asthma or food allergy in parents or blood-related siblings

Table S2b: Univariable logistic regression analysis in KUNO Kids of potential risk factors: outcome parent-reported symptoms of FA at age 2

	Odds ratio (95% CI)	p value	n
Child			
Male sex	1.032 (0.661-1.611)	.891	644
Preterm birth	0.578 (0.173-1.933)	.374	641
Caesarean section	0.924 (0.559-1.527)	.758	646
Birth weight < 2800 g	0.337 (0.120-0.947)	.039	642
Duration of breastfeeding < 6 months	1.476 (0.905-2.409)	.119	533
Introduction of solid food at the age of			
- < 4 months	-/- (no cases with FA)	-/-	
- 4 – 6 months	reference		511
- > 6 months	0.715 (0.312-1.639)	.715	
Antibiotic use	0.834 (0.384-1.811)	.647	644
Daily animal contacts ^a	2.068 (1.298-3.294)	.002	645
Being on a farm regularly	0.702 (0.338-1.456)	.342	646
Regular contact to hay	0.634 (0.221-1.821)	.397	646
High exposure to traffic, exhaust gases	1.545 (0.988-2.415)	.056	645
Other atopic disease ^b	2.666 (1.599-4.444)	<.001	646
Older siblings	0.920 (0.578-1.464)	.724	642
Migration background ^c	1.410 (0.777-2.557)	.258	624
Mother			
Maternal age > 34 years	0.924 (0.576-1.482)	.924	641
Pre-pregnancy BMI			
- underweight	-/- (no cases with FA)	-/-	
- normal weight	reference		640
- overweight	1.272 (0.757-2.138)	.363	
- obese	1.333 (0.700-2.541)	.382	
Gestational Diabetes	0.639 (0.319-1.283)		626
Cigarette smoking			
- during pregnancy	2.522 (0.482-13.209)	.273	628
- passive exposure	1.979 (1.229-3.186)	.005	645
Antibiotic use			
- during pregnancy	1.152 (0.684-1.941)	.595	639
- during delivery	0.880 (0.562-1.379)	.577	634
- while breastfeeding	0.853 (0.371-1.964)	.709	525
Being on a farm regularly			
- during pregnancy	0.684 (0.237-1.974)	.483	640
- while breastfeeding	0.717 (0.248-2.072)	.539	641
Regular animal contact ^d	1.734 (1.085-2.770)	.021	643
School education			
- low (no degree)	2.088 (0.939-4.645)	.071	
- intermediate (< 10 years)	reference		634
- high (> 10 years)	1.047 (0.624-1.755)	.863	
Atopic disease of 1st degree relatives ^e	1.456 (0.872-2.430)	.151	646

FA - food allergy; 95% CI - 95% confidence interval; BMI – body mass index

^a daily contact to dog, cat, hamster, guinea pig, mouse, rat or rabbit

^b atopic eczema, allergic rhinitis, allergic conjunctivitis and asthma

^c mother or father with non-German nationality or born not in Germany

^d regular contact (more than once per month) to any pet during pregnancy

^e atopic eczema, allergic rhinitis, allergic conjunctivitis, asthma or food allergy in parents or blood-related siblings

Table S3a: Univariable logistic regression analysis in the Ulm SPATZ Health Study of potential risk factors: outcome parent-reported FA at age 1

	Odds ratio (95% CI)	p value	n
Child			
Male sex	2.172 (1.345 – 16.557)	.015	745
Preterm birth	1.225 (0.423 – 5.321)	.530	745
Caesarean section	0.715 (0.145 – 1.797)	.296	745
Birth weight < 2800 g	0.798 (0.144 – 2.821)	.553	745
Duration of breastfeeding < 6 months	0.838 (0.226 – 2.175)	.539	745
Introduction of solid food at the age of			
- < 4 months			
- 4 – 6 months	0.771 (0.169 – 2.089)	.417	745
- > 6 months	<i>reference</i>		
Other atopic disease ^a	3.260 (3.863 – 29.228)	<.0001	745
Older siblings	0.566 (0.041– 2.506)	.278	357
Migration background ^b	No cases with FA		722
Mother			
Maternal age > 34 years	1.077 (0.436 – 3.083)	.766	745
Pre-pregnancy BMI			
- underweight	1.01 (0.152 – 9.969)	.990	
- normal weight	<i>reference</i>		745
- overweight	0.734 (0.239 – 3.345)	.558	
- obese	1.642 (0.603 – 6.630)	.310	
Gestational diabetes	1.573 (0.691 – 8.863)	.164	745
Cigarette smoking			
- during pregnancy	1.595 (0.931 – 6.957)	.069	743
- passive exposure	0.743 (0.072 – 4.237)	.568	740
Antibiotic use			
- during pregnancy	1.568 (0.783 – 7.717)	.124	745
School education			
- low (no degree)	No cases with FA		
- intermediate (< 10 years)	<i>reference</i>		731
- high (> 10 years)	2.984 (0.674 – 5.216)	.967	
Atopic disease of 1st degree relatives‡	2.033 (1.178 – 14.510)	.0270	745

FA: food allergy; 95% CI - 95% confidence interval

^a atopic eczema, hay fever, asthma

^b non-German nationality

^c atopic exzema, hay fever, asthma in parents or siblings

Table S3b: Univariable logistic regression analysis in the Ulm SPATZ Health Study of potential risk factors: outcome parent-reported FA at age 2

	Odds ratio (95% CI)	p value	n
Child			
Male sex	1.55 (0.985 – 5.861)	.054	745
Preterm birth	0.999 (0.292 – 3.406)	.996	790
Caesarean section	0.806 (0.239 – 1.760)	.396	790
Birth weight < 2800 g	0.832 (0.204 – 2.356)	.556	790
Duration of breastfeeding < 6 month	0.744 (0.204 – 1.498)	.244	790
Introduction of solid food at the age of			
- < 4 months			
- 4 – 6 months	0.84 (0.260 – 1.916)	.495	790
- > 6 months	reference		
Other atopic disease ^a	2.623 (2.672 – 17.706)	<.001	746
Older siblings	0.942 (0.192 – 4.107)	.942	286
Migration background ^b	No cases with FA		765
Mother			
Maternal age > 34 years	0.984 (0.419 – 2.242)	.941	790
Pre-pregnancy BMI			
- underweight	0.760 (0.121 – 7.544)	.725	
- normal weight	reference		790
- overweight	1.099 (0.511 – 3.743)	.824	
- obese	1.505 (0.653 – 5.495)	.358	
Gestational Diabetes	1.285 (0.479 – 5.692)	.427	790
Cigarette smoking			
- during pregnancy	1.666 (1.173 – 6.563)	.020	788
- passive exposure	0.643 (0.055 – 3.123)	.392	744
Antibiotic use			
- during pregnancy	1.266 (0.535 – 4.799)	.340	790
School education			
- low (no degree)	No cases with FA		
- intermediate (< 10 years)	reference		776
- high (> 10 years)	1.462 (0.633 – 3.376)	.975	
Atopic disease of 1st degree relatives ^c	1.464 (0.878 – 5.225)	.094	790

FA – food allergy; 95% CI - 95% confidence interval

^a atopic eczema, hay fever, asthma

^b non-German nationality

^c atopic exzema, hay fever, asthma in parents or siblings