

Sequence Analysis and Biochemical Characterization of Two Non-specific Lipid Transfer Proteins from Tartary Buckwheat Seeds

Xiaodong Cui (✉ cuixiaodong@sxu.edu.cn)

Shanxi University

Wenhua Tian

Shanxi University

Wenhua Wang

Shanxi University

Jiao Li

Shanxi University

Zhuanhua Wang

Shanxi University

Chen Li

Shanxi University

Research Article

Keywords: Tartary buckwheat, Lipid transfer proteins, Antifungal activity, α -Amylase inhibitor, Food allergy

Posted Date: October 11th, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-2106951/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Plant non-specific lipid transfer proteins (nsLTPs) exert various biological activities that are valuable for agronomy, industrial manufacturing, and medicine. In this study, two nsLTPs derived from Tartary buckwheat, namely FtLTP1a and FtLTP1b, were produced by gene cloning, expression and purification. Sequence analysis and biochemical characteristics of the proteins, including lipid binding ability, α -amylase inhibition activity, antifungal activity, and allergenic activity, were investigated. Both FtLTP1a and FtLTP1b exhibited similar lipid binding and antifungal properties, but only FtLTP1b showed weak inhibitory activity against α -amylase from the porcine pancreas. Finally, the binding ability of the two protein to IgE antibodies in serum samples from nine patients allergic to buckwheat was assayed using ELISA to investigate their allergenic activity. The data suggested that FtLTP1b not only specifically bound to IgE in serum but also cross-reacted with artemisia pollen allergen (W6). These findings can give constructive suggestions for understanding of cross-reactivity and specificity of nsLTPs.

1. Introduction

Non-specific lipid transfer proteins (nsLTPs) belong to the prolamin superfamily with conserved cysteine residues, abundant α -helices, and low molecular mass [1]. Usually, nsLTPs contain four or five α -helices stabilized by four disulfide bonds formed by a motif with the conserved sequence C-Xn-C-Xn-CC-Xn-CXC-Xn-C-Xn-C. In addition, a tunnel-like hydrophobic cavity exists in the nsLTPs, which is responsible for various lipids transportation [2]. According to the protein sequence length, nsLTPs are classed into two families, namely nsLTP1s with about 90 amino acid residues and nsLTP2s with about 70 amino acids [3]. The two families also have different subfamilies based on the particular pairing of cysteines in disulfide bridges. For instance, nsLTP1s are characterized by four disulfide bridges formed by Cys residues 1-6, 2-3, 4-7, and 5-8 [4], while disulfide bridges in nsLTP2s are paired by Cys residues 1-5, 2-3, 4-7, and 6-8 [5]. nsLTPs also belong to PR-14, a pathogenesis-related protein [6]. They involve in defending against viral plant pathogens, bacterial, or fungal, by activating the plant immune system through a receptor-dependent mechanism [7].

Many nsLTPs have been isolated and characterized from various plant species. They are reported to play essential roles in cutin formation, embryogenesis and response to stresses [8]. nsLTPs can reversibly bind to lipid molecules, including sterols, acyl-coenzyme A (CoA) and fatty acids [3]. This effect endows nsLTPs roles in the transmembrane transportation of lipids and permeabilization of phytopathogen membranes [9]. Many nsLTPs also have been considered to be allergens from different sources such as pollen, food or latex, causing various severe symptoms, such as anaphylaxis [10]. For example, Pru p3, a nsLTP from peach, is the first plant food allergen found in the nsLTPs family [4].

Buckwheat, which belongs to the family *Polygonaceae*, can be found in many parts of the world, but is grown mainly in the northern hemisphere [11]. With the increase in buckwheat food consumption, the cases of instant hypersensitivity caused by food containing buckwheat additives have also increased. Some natural allergenic proteins from common buckwheat and Tartary buckwheat have been isolated

and identified. Their molecular weights range from 10 to 100 kDa. Among them, proteins with molecular weights of 24 kDa, 19 kDa, 16 kDa and 10 kDa are considered the main allergens in common buckwheat, and these allergens can cause serious allergic reactions, such as asthma and anaphylactic shock [12]. The 24-kDa protein has been identified as a small subunit of 11S globulins and named Fag e1 [13]. Moreover, the 16-kDa (Fag e2) and 19-kDa allergens have been identified as members of the 2S albumin family and 7S vicilin-like globulins, respectively. While, the 10-kDa allergen is the α -amylase inhibitor/trypsin inhibitor [14]. The 24-kDa allergen (Fag t1) isolated from Tartary buckwheat seeds is thermostable and homologous with the 24-kDa allergen from common buckwheat (Fag e1) [15]. A 56-kDa protein from buckwheat displayed allergic activity, and its nucleic acid sequence has been determined [16]. A 16-kDa allergenic protein (Fag t2) from Tartary buckwheat has been recombinantly expressed [17], and its core epitope has been identified [18]. A 9-kDa protein from buckwheat shows both allergic and trypsin inhibitory activity [12]. Research on a variety of allergens from buckwheat has been extensively carried out. However, nsLTPs from common buckwheat or Tartary buckwheat and their allergenic potentials have not been investigated.

In this article, the complete nucleic acid sequences of two nsLTPs from Tartary buckwheat seeds, namely FtLTP1a and FtLTP1b, were analyzed. The work aims to verify whether these nsLTPs are potential allergens. The two nsLTPs were recombinantly expressed in *Escherichia coli*. Their biochemical characteristics, including lipid binding ability, α -amylase inhibition and antimicrobial activities, were evaluated. The capacity of the two nsLTPs to bind IgE antibodies in serum of patients with allergy to the 24-kDa protein in buckwheat was also assessed.

2. Materials And Methods

2.1 Materials and reagents

The Tartary buckwheat strain Yunqiao No.1 was field-grown at the Science Academy of Agriculture in Yunnan Province, China. α -Amylase from porcine pancreas and Ni^{2+} -NTA agarose column were purchased from Sangon Biotech Co., Ltd. (Shanghai, China). The pET-32a vector was purchased from Promega Corporation (Madison, USA). T_4 DNA ligase and DNA restriction enzymes (*Xho* I and *Bam* HI) were obtained from Thermo Fisher Scientific (Waltham, USA). Sera from all recruited patients, aged 14 to 46 years, were tested for allergen-specific IgE using microarray immunoassay (Immuno CAP ISAC; Uppsala, Sweden). Information on the recruited patients, mugwort-specific IgE values, pollens causing respiratory symptoms, and other food allergen (based on CAP-system test), can be found in Table S1 and S2.

2.2 Sequence analysis

The open reading frame (ORF) of *FtLTP* genes was analyzed using the online software ORF finder (<http://www.ncbi.nlm.nih.gov/gorf/>). The identity of FtLTPs was determined by the BLAST tool available on the NCBI website. Amino acid sequence alignment of FtLTPs with other nsLTPs (or allergen) was

performed by ClustalW2 (<http://www.ebi.ac.uk/Tools/clustalw2/index.html>)[19]. Phylogenetic relationships were determined according to the Maximum Likelihood (ML) method with 1000 replicated bootstraps using MEGA7.0 Software [20]. The midpoint rooted tree was generated using Interactive Tree of Life (iTOL, version 3.2.317, <http://itol.emnl.de>).

2.3 Expression and purification of FtLTPs

The sequence of FtLTPs (removed putative signal peptides) was optimized based on the preferred codon of *E. coli*, synthesized and then cloned into a pUC-57 vector. In the construction of plasmid, the *FtLTP* genes were first amplified using the following primers: 5'-GGATCCTTTGAATGTGGTGCAGTTAGCAAA-3' and 5'-CTCGAGTTAATGAACTTTGCTACAATCGGT-3' for *FtLTP1a*, and 5'-GGATCCGCACTGACCTGTGGTACCGTTACC-3' and 5'-CTCGAGTTAATTAACTTTGCTACAATCGGTGCTC-3' for *FtLTP1b*. The primers contained *Bam* HI and *Xho* I restriction sites. The purified PCR product was digested by *Bam* HI and *Xho* I, and then cloned into a pET32a vector to construct pET-32a-*FtLTP1a* and pET-32a-*FtLTP1b* recombinant plasmids. After transformation into *E. coli* BL21 (DE3) competent cells, clones were selected by ampicillin screening, and the accuracy of the two recombinants was verified by DNA sequencing.

Transformed *E. coli* was inoculated in 5 mL of LB medium containing 100 µg/mL ampicillin, and cultured at 37°C and 180 rpm to attain log-phase growth. Then, the culture was transferred to 1000 mL of LB medium and incubated. When the OD₆₀₀ reached 0.6, protein expression was induced by IPTG (with a final concentration of 0.2 mM) at 16°C for 12 h. The cells were collected and subsequently sonicated. Finally, the cell lysate was analyzed by SDS-PAGE using a 12.5% separating gel.

The cell lysate supernatant was loaded onto a Ni²⁺-NTA agarose column pre-equilibrated with buffer A (10 mM Tris-HCl, 0.5 M NaCl, 10 mM imidazole). After being thoroughly washed with buffer B (10 mM Tris-HCl, 0.5 M NaCl, 50 mM imidazole), target proteins were washed with buffer C (10 mM Tris-HCl, 0.5 M NaCl, 200 mM imidazole). Collected samples were subsequently dialyzed against water and then stored at -80°C.

2.4 Lipid binding assays

Fluorescence of recombinant FtLTP1a and FtLTP1b were measured at 37°C (310K) at the excitation wavelength of 280 nm and emission wavelengths of 300–420 nm using an LS55 fluorescence spectrometer (Perkin Elmer, USA). Two milliliters of 1 µM FtLTP1a or FtLTP1b were added into the cuvette and then titrated with lipid by stepwise adding lipid into the cuvette. Titration curves, which showed the variation of fluorescence intensity ($F-F_0$) at 342 nm with the molar ratio of lipid/protein, R_i , were obtained.

2.5 α-Amylase inhibition assays

The enzymatic activity of α-amylase was assayed as per the reported method [21]. Briefly, the α-amylase activity was measured using starch as a substrate, and the product reducing sugar was quantified after reaction with 3, 5-dinitrosalicylic acid (DNS). The reaction mixtures containing 5 U of α-amylase and

diverse concentrations of FtLTP1a or FtLTP1b were preincubated for 15 min at 37°C. The reaction was initiated by adding a 400 µL substrate (1% starch; Sigma-Aldrich) and carried out for 15 min. After adding DNS, the reaction was terminated by boiling water treatment for 5 min. After cooling to room temperature (25°C), the absorbance at 540 nm of the mixture was measured. One unit of enzymic activity was defined as the amount of enzyme required for an increase in A_{540} nm by 0.1 in 30 min. The enzymatic activity of the control without FtLTP1a or FtLTP1b was considered to be 100%. The percent inhibition was calculated according to the following formula:

$[100 - (A_1 \times 100/A_2)]$, A_1 and A_2 are the absorbances of sample and control, respectively.

2.6 Antifungal assay

The antifungal activities of FtLTP1a and FtLTP1b were measured by spotting assay and a 2-fold microdilution assay [22, 23]. The hyphae of *Aspergillus niger*, *Aspergillus flavus*, or *Fusarium oxysporum* were added to sterilized ddH₂O to prepare mycelium suspension of 10^5 - 10^3 CFU/mL. A protein sample (100 µM) was added to the suspension and 10 µL of the resultant mixture was cultured on 1.2% potato dextrose agar (PDA) at 28°C. The colony growth was observed after 48 h. The minimum inhibitory concentrations (MIC) of FtLTP1a and FtLTP1b with respects to fungi were further determined by a 2-fold microdilution assay in RPMI 1640 medium. Briefly, mycelium suspensions with a turbidity of 0.5 McFarland standards were prepared. Cell suspension was diluted in RPMI 1640 medium, 50 µL was added to a 96-well plate, and protein samples with final concentrations of 0.25 to 256 µM were added. The plate was incubated at 28°C for 48 h and the fungal growth was observed. Medium alone and containing inoculum suspension were used as a negative control and positive control, respectively.

2.7 Enzyme-linked immunosorbent assay (ELISA)

An aliquot of purified FtLTPs (0.5 µg) was added to a 96-well plate containing 50 µL of TBS solution. After incubation for 1 h at 37°C, the plate was washed with TBST solution. Then, 1% BSA solution in TBS was added and incubated at 37°C for 2 h. After reaction with sera from allergic patients diluted in TBS at 37°C for another 2 h, the plate was washed with TBST. Anti-human IgE-horseradish peroxidase-conjugated antibody (Sino Biological Inc., China) was added to each well and carried out at 37°C for 1 h. The wells were rewashed with TBST, and bound antibodies were detected with TMB chromogen solution (Sigma). After termination by adding 4 N H₂SO₄, the absorbances at 450 nm were determined and analyzed.

2.8 Statistical analysis

In paper, all data were from at least three independent experiments and were presented as mean ± standard error of the mean (S.E.M). Analysis was evaluated using the SPSS version 20.0. One-way ANOVA test (Tukey's test) was used to determine the significant statistical differences. p value 0.05 or 0.01 were considered as significantly different.

3. Results And Discussion

3.1 Sequence and phylogenetic analyses

Full-length cDNA and amino acid sequences of FtLTP1a and FtLTP1b are shown in Fig. S1. Both FtLTP1a and FtLTP1a contained 117 amino acid residues. FtLTP1a contained a 91-residue mature peptide and a 26-residue putative signal peptide (GenBank accession number: OK340654), whereas FtLTP1b contained a 92-residue mature peptide and a 25-residue signal peptide (GenBank accession number: OK340655). The molecular masses of mature FtLTP1a and FtLTP1b were estimated using DNAMAN software to be 9228.76 Da and 9038.55 Da, respectively. It has been reported that nsLTPs consist of two main subfamilies, namely the 9-kDa nsLTP1 family and the 7-kDa nsLTP2 family [24]. Moreover, both the two purified proteins share high sequence homology with nsLTP1 family. Furthermore, they contain two conserved motifs (T/SXXDR/K and PYXIS), which is also a characteristic of the nsLTP1 family [3]. Therefore, based on the above results, we may conclude that FtLTP1a and FtLTP1b belong to the nsLTP1 family.

The phylogenetic tree is a framework indicating genetic diversities and distances between different biological species and other entities. To explore the phylogenetic relationships between FtLTP1a/b from Tartary buckwheat and those from other species, multiple sequence alignments were analyzed, and a phylogenetic tree of 114 protein sequences was generated using a neighbor-joining method available in MEGA7 software (Fig. 1A). We found that nsLTPs from Tartary buckwheat were grouped with those from *Chenopodium quinoa* (XP 021723225.1, XP 021766670.1), *Spinacia oleracea* (XP 021851885.1), and *Beta vulgaris* subsp. *Vulgaris* (XP 010690793.1). Additionally, FtLTP1a and FtLTP1b were closely related to many allergens, such as Amb a6 (O04004.1, *Ambrosia artemisiifolia*), Zea m14 (P19656.1, *Zea mays*), Tri a14 (P24296.2, *Triticum aestivum*), Bra o3 (Q9ZSL7, *Brassica oleracea* var. *oleracea*), Pla a 3.02 (ALF00099.1, *Platanus x hispanic*), and Pla o3 (ABY21307.1, *Platanus orientalis*).

The amino acid sequences of FtLTP1a and FtLTP1b were also compared with three unique proteins, Api g2, Pru p3 and Tri a14, available in the UniProt database (Fig. 1B). The IgE-binding epitopes in Pru p3, Tri a14, and Api g2, the homologous nsLTP from peach, wheat, and celery, respectively, have been identified. FtLTP1a shared 54.9%, 47.3% and 41.1% sequence identity with Api g2, Pru p3, and Tri a14, respectively, and FtLTP1b shared 64.8%, 52.7% and 48.9%, respectively. The comparison of amino acid sequences revealed that FtLTP1b is more similar to Api g2, Pru p3, and Tri a14 than FtLTP1a. Previous studies have identified the IgE binding epitopes in Pru p3 using different methodologies [25–27]. Comparing the amino acid sequences of Pru p3 and FtLTP1a, it was found that there was high degree of diversity in the regions potentially involved in IgE recognition (Fig. 1B). Moreover, FtLTP1b had a high sequence identity with Pru p3, including a highly conserved region (GKCGVSIPIYK) that can significantly contribute to IgE binding. This region in Api g2 and Tri a14 is also highly conserved. Therefore, the analysis above suggests that FtLTP1b is more likely to be a potential allergen than FtLTP1a.

3.2 Heterologous expression and purification of FtLTP1s

Recombinants of pET32a-FtLTP1a and pET32a-FtLTP1b were constructed, transformed into *E. coli* BL21(DE3) cells and induced expression. As shown in Fig. 2, the recombinant Trx-FtLTP1a and Trx-

FtLTP1a proteins were expressed in soluble form, and the target proteins were further purified by a Ni²⁺-NTA column. Using this single-step procedure, the purity of both Trx-FtLTP1a and Trx-FtLTP1b proteins was about 95% and migrated as a single band on SDS-PAGE (Fig. 2). The estimated yield of the FtLTP1a and FtLTP1b was 2.14 mg and 2.46 mg per liter of culture, respectively (Table S3). The purity of FtLTP1a and FtLTP1b were 94.25% and 96.33% by gel scan.

3.3 Both FtLTP1a and FtLTP1b could bind lipid

Two highly conserved regions (T/SXXDR/K and PYXIS) play an essential role in lipid binding, and the fluorescence of tyrosine (Y, in PYXIS) is sensitive to lipid binding. Fluorescence spectroscopy is a tool for investigating the quenching mechanism and the binding properties. Intrinsic fluorescence of FtLTP1a was compared with that of FtLTP1b. As shown in Fig. 3A, the maximal fluorescence emission wavelength of both FtLTP1a and FtLTP1b was 342 nm. However, the intensity of FtLTP1b was dramatically lower than that of FtLTP1a. The difference in intensity may be due to the fact that FtLTP1a contains one tyrosine residue and two phenylalanine residues, while FtLTP1b only contains two tyrosine residues. When the purified FtLTP1a and FtLTP1b were incubated with oleic acid, the fluorescence intensity was associated with the increase in oleic acid concentration (Figs. 3B and 3C). According to the apparent saturation levels, the molar ratio of oleic acid/FtLTP1a or oleic acid/FtLTP1b (*R_i*) was about 1.0 (Fig. 3D). The fluorescence of FtLTP1a and FtLTP1b with oleic acid was markedly increased by 160% and 144%, respectively, compared to that in the absence of oleic acid. In addition, the binding of FtLTTPs with other two types of lipids, stearic acid and linolenic acid, was also studied, from which similar results were observed under the same test conditions (data not shown). The tyr-79 residue of the present nsLTTPs and other corresponding residues of nsLTTPs from other plant species are involved in increasing of fluorescence intensity when binding to lipids. Tyr-79, located in the C-terminal fragment, a part of the extreme hydrophobic cavity, should be perturbed by lipid-binding [28]. Maize nsLTTP-fatty acid complex has shown that the carboxyl group in fatty acid is involved in forming hydrogen bonds with the hydroxyl group of Tyr81 [29]. Consequently, in a lipid-bound state, Tyr-79 in the protein may probably be less mobile, and thus could not be effectively quenched by a possible internal quencher.

3.4 FtLTP1b exhibits weak α -amylase inhibitory activity

According to the protein classification by NCBI, nsLTTPs belong to the AAL_LTSS superfamily (protease inhibitor/seed storage/LTP family). However, only a few nsLTTPs have been identified to exhibit α -amylase inhibitory activity till now. Thus, the α -amylase inhibitory activity of FtLTP1a and FtLTP1b was further tested. First, the secondary structures of FtLTP1a and FtLTP1b were verified by circular dichroism (CD) spectra. The CD spectra of two proteins showed typical peaks of α -helix, namely two minimum peaks at 208 and 222 nm (Fig.S2). This result is consistent with the CD spectrum of nsLTTPs from red wine and grapes [30], peach, cherry, hazelnut [31], and mugwort pollen [32], which are all rich in α -helices. Thus, we conclude that recombinant FtLTP1a and FtLTP1a folded correctly and formed a natural conformation. As shown in Fig. 4, both FtLTP1a and FtLTP1b were unable to inhibit α -amylase from *Bacillus licheniformis*, and FtLTP1b could weakly inhibit α -amylase from porcine pancreas. It has been reported that lipid transfer proteins from *Coffea canephora* and *Vigna unguiculata* L. Walp showed inhibitory activity

against α -amylase from human saliva and *C. maculatus*, respectively [33, 34]. One study has reported that rVu-LTP purified from *Vigna unguiculata* L. Walp can inhibit the activity of HAS and intestinal α -amylase from *C. maculatus*, but cannot inhibit intestinal α -amylase from *T. maculatus* [35].

Fragment Asp-Glu-Asp appears in the active site of almost all α -amylases [36]. Some studies have suggested that positively charged (arginine and lysine) amino acids in inhibitors can non-covalently interact with the negatively charged residues of Asp-Glu-Asp, resulting in an ultimate inhibition of the enzyme [37]. Two peptides from Vu-LTP, N₂₉-I₄₇ (NGVKNILNGARTTADRRGI) and α G₇₄-N₉₁ (GVNIPWKISSSTNaNTIN), have been identified and found to be responsible for HAS (human salivary α -amylase) inhibition [21]. The two regions of FtLTP1a did not contain positively charged amino acids, while there was a lysine (K) residue in one region (G₇₄-N₉₁) of FtLTP1b. This may explain why FtLTP1b exhibited weak α -amylase inhibitory activity, whereas FtLTP1a did not.

3.5 Both FtLTP1a and FtLTP1b exhibit antimicrobial activity

Plant nsLTPs belong to pathogenesis-related protein, which are reported to display broad-spectrum antimicrobial activity. The antifungal activities of FtLTP1a and FtLTP1b against *Aspergillus niger*, *Aspergillus flavus* and *Fusarium oxysporum* were evaluated. As shown in Fig. 5, the two proteins exhibited inhibitory activity against *A. niger*. The MIC (minimum Inhibitory Concentration) values of the proteins were determined and tabulated in Table S4. The MIC values of FtLTP1a to *Aspergillus niger*, *Aspergillus flavus* and *Fusarium oxysporum* were 8 μ M, 8 μ M, and 64 μ M, respectively, and those of FtLTP1b were 16 μ M, 16 μ M, and 64 μ M, respectively.

The antimicrobial activities of nsLTPs from different plant species have been studied. Although the nsLTPs can inhibit both bacterial and fungal growths, the inhibition against bacterial growth is more potent than that against fungi growth [38]. Nevertheless, the mechanism of this occurrence remains unclear. The findings to date have indicated the ability to interact with lipids contributes to the antimicrobial activity of plant nsLTPs. Although the peptide Ace-AMP1 (from onion), also characterized as an nsLTP, cannot transfer phospholipid from liposome to mitochondria, it possesses antimicrobial activity [8]. It has been speculated this activity was due to their interaction with cell membranes, causing the loss of membrane integrity and permeabilization [39].

3.6 Immunological characteristics of FtLTP1a and FtLTP1b

Sera from 9 Chinese patients who tested positive for buckwheat allergen were used to screen IgE reactivity of FtLTP1a and FtLTP1b using ELISA. The clinical data of the nine patients who participated in this study are summarized in Table S1. All patients tested positive for IgE antibodies-specific 24-kDa buckwheat allergen and experienced allergic reactions after consuming buckwheat food. The P/N values > 2.1 were used to evaluate the allergenic activity of nsLTPs from buckwheat. As shown in Fig. 6A, all the 2-fold diluted sera from nine individuals (100%) were sensitive to FtLTP1b (P/N values > 2.1) but not FtLTP1a (P/N values < 2.1). In addition, only three 10-fold diluted sera were sensitive to FtLTP1b (Fig. 6B).

The fact that patients who are allergic to a particular allergen tend to be allergic to other agents indicates that the same IgE antibodies may recognize other homologous allergens which possess similar epitopes. As shown in Table S1, eight out of these nine patients were also allergic to the artemisia pollen (W6). FtLTP1a and FtLTP1b were tested on three sera that were sensitive to W6 and polyvalent grass pollen (WX7), to determine whether they share the same IgE epitopes with other allergens and further evaluate their potential cross-reactivity. As shown in Fig. 6C, FtLTP1b was able to bind IgE in these sera ($P/N > 2.1$). The majority (89%) of *Artemisia vulgaris* pollen-sensitized patients is also allergic to Art v3, a mugwort allergen [40]. Clinical studies have shown that Art v3 could cross-react with another homologous protein from peach (Pur p3). Patients with peach allergy and are primarily sensitive to Pru p3 can experience allergic response when exposed to mugwort pollen because of cross-reactivity with Art v3 [41]. In this study, certain cross-reactivity occurred between FtLTP1b and pollen allergen (W6 or WX7). However, clinical symptoms caused by this cross-reactivity may not be observable.

4. Conclusion

In this work, two recombinants of nsLTP1s (FtLTP1a and FtLTP1b) were prepared and purified by affinity chromatography. Moreover, their chemical properties and biological activities were characterized, including lipid binding ability, α -amylase inhibition, antifungal, and IgE-binding activities. The results showed that FtLTP1b from Tartary buckwheat is a novel allergenic member of the lipid-transfer protein 1 family, which can cross-react with homologous protein from pollen. The data may further help clinicians manage of patients allergic to buckwheat by avoiding being in unnecessary contact with pollen.

Declarations

Ethical Approval Not applicable.

Consent to participate All authors consent to participate.

Consent for publication All authors consent to publication.

Competing interests All authors declare that there are no conflicts of interests.

Authors' contributions Xiaodong Cui and Chen Li wrote the manuscript text. Wenhua Tian and Wenhua Wang prepared figures and formal analysis. Zhuanhua Wang and Jiao Li were responsible for supervision and methodology. All authors reviewed the manuscript.

Funding The research was supported by the Applied Basic Research Programs of Shanxi Province (No. 20210302123465), the Shanxi Scholarship Council of China (No. 2020-015) and the Open Project Program of Xinghuacun College of Shanxi (Shanxi Institute of Brewing Technology and Industry (Preparation) (No. XCS-UK-202018).

Availability of data and materials All data generated or analyzed during this study are included in this article and its supplementary materials files are available.

References


1. Shewry PR, Beaudoin F, Jenkins J, Griffiths-Jones S, Mills ENC (2002) Plant protein families and their relationships to food allergy. *Biochem Soc T* 30:906–910
2. Nazeer M, Waheed H, Saeed M, Ali SY, Choudhary MI, Ul-Haq Z, Ahmed A (2019) Purification and Characterization of a Nonspecific Lipid Transfer Protein 1 (nsLTP1) from Ajwain (*Trachyspermum ammi*) Seeds. *Sci Rep* 9:4148
3. Salminen TA, Blomqvist K, Edqvist J (2016) Lipid transfer proteins: classification, nomenclature, structure, and function. *Planta* 244:971–997
4. Pasquato N, Berni R, Folli C, Folloni S, Cianci M, Pantano S, Helliwell JR, Zanotti G (2006) Crystal structure of peach Pru p 3, the prototypic member of the family of plant non-specific lipid transfer protein pan-allergens. *J Mol Biol* 356:684–694
5. Hoh F, Pons JL, Gautier MF, de Lamotte F, Dumas C (2005) Structure of a liganded type 2 non-specific lipid-transfer protein from wheat and the molecular basis of lipid binding. *Acta Crystallogr Sect D Biol Crystallogr* 61:397–406
6. Melnikova DN, Finkina EI, Bogdanov IV, Ovchinnikova TV (2019) Plant Pathogenesis-Related Proteins Binding Lipids and Other Hydrophobic Ligands. *Russ J Bioorg Chem* 44:586–594
7. Finkina EI, Melnikova DN, Bogdanov IV, Ovchinnikova TV (2017) Plant Pathogenesis-Related Proteins PR-10 and PR-14 as Components of Innate Immunity System and Ubiquitous Allergens. *Curr Med Chem* 24:1772–1787
8. Maximiano MR, Franco OL (2021) Biotechnological applications of versatile plant lipid transfer proteins (LTPs). *Peptides* 140:170531
9. Tam JP, Wang S, Wong KH, Tan WL (2015) Antimicrobial Peptides from Plants. *Pharmaceuticals* (Basel) 8:711–757
10. Florido Lopez JF, Quiralte Enriquez J, Arias de Saavedra Alias JM, Saenz de San Pedro B, Martin Casanez E (2002) An allergen from *Olea europaea* pollen (Ole e 7) is associated with plant-derived food anaphylaxis. *Allergy* 57:53–59
11. Fabjan N, Rode J, Kosir IJ, Wang Z, Zhang Z, Kreft I (2003) Tartary buckwheat (*Fagopyrum tataricum* Gaertn.) as a source of dietary rutin and quercitrin. *J Agric Food Chem* 51:6452–6455
12. Park JW, Kang DB, Kim CW, Koh SH, Yum HY, Kim KE, Hong CS, Lee KY (2000) Identification and characterization of the major allergens of buckwheat. *Allergy* 55:1035–1041
13. Yoshioka H, Ohmoto T, Urisu A, Mine Y, Adachi T (2004) Expression and epitope analysis of the major allergenic protein Fag e 1 from buckwheat. *J Plant Physiol* 161:761–767
14. Rolla G (2014) Buckwheat Allergy: An Emerging Clinical Problem in Europe. *Journal of Allergy & Therapy*; 05

15. Yang Z, Li Y, Li C, Wang Z (2012) Synthesis of hypoallergenic derivatives of the major allergen Fag t 1 from tartary buckwheat via sequence restructuring. *Food and chemical toxicology: an international journal published for the British Industrial Biological Research Association* 50:2675–2680
16. Zhang X, Yuan JM, Cui XD, Wang ZH (2008) Molecular cloning, recombinant expression, and immunological characterization of a novel allergen from tartary buckwheat. *J Agric Food Chem* 56:10947–10953
17. Chen P, Guo YF, Yan Q, Li YH (2011) Molecular cloning and characterization of Fag t 2: a 16-kDa major allergen from Tartary buckwheat seeds. *Allergy* 66:1393–1395
18. Zheng B, Zhang H, Shen W, Wang L, Chen P (2021) Core epitope analysis of 16 kDa allergen from tartary buckwheat. *Food Chem* 346:128953
19. Thompson JD, Higgins DG, Gibson TJ (1994) Clustal-W - Improving the Sensitivity of Progressive Multiple Sequence Alignment through Sequence Weighting, Position-Specific Gap Penalties and Weight Matrix Choice. *Nucleic Acids Res* 22:4673–4680
20. Kumar S, Stecher G, Tamura K, MEGA7 (2016) Molecular Evolutionary Genetics Analysis Version 7.0 for Bigger Datasets. *Mol Biol Evol* 33:1870–1874
21. da Silva FCV, do Nascimento VV, Machado OLT, Pereira LdS, Gomes VM, de Oliveira Carvalho A (2018) Insight into the α -Amylase Inhibitory Activity of Plant Lipid Transfer Proteins. *J Chem Inf Model* 58:2294–2304
22. Cui X, Du J, Li J, Wang Z (2018) Inhibitory site of alpha-hairpinin peptide from tartary buckwheat has no effect on its antimicrobial activities. *Acta Biochim Biophys Sin* 50:408–416
23. Lambert RJ, Pearson J (2000) Susceptibility testing: accurate and reproducible minimum inhibitory concentration (MIC) and non-inhibitory concentration (NIC) values. *J Appl Microbiol* 88:784–790
24. Douliez JP, Michon T, Elmorjani K, Marion D (2000) Structure, Biological and Technological Functions of Lipid Transfer Proteins and Indolines, the Major Lipid Binding Proteins from Cereal Kernels. *J Cereal Sci* 32:1–20
25. Garcia-Casado G, Pacios LF, Diaz-Perales A, Sanchez-Monge R, Lombardero M, Garcia-Selles FJ, Polo F, Barber D, Salcedo G (2003) Identification of IgE-binding epitopes of the major peach allergen Pru p 3. *J Allergy Clin Immunol* 112:599–605
26. Borges JP, Barre A, Culerrier R, Granier C, Didier A, Rouge P (2008) Lipid transfer proteins from Rosaceae fruits share consensus epitopes responsible for their IgE-binding cross-reactivity. *Biochem Biophys Res Commun* 365:685–690
27. Tordesillas L, Pacios LF, Palacin A, Quirce S, Armentia A, Barber D, Salcedo G, Diaz-Perales A (2009) Molecular basis of allergen cross-reactivity: non-specific lipid transfer proteins from wheat flour and peach fruit as models. *Mol Immunol* 47:534–540
28. Gincel E, Simorre J-P, Caille A, Marion D, Ptak M, Vovelle F (1994) Three-Dimensional Structure in Solution of a Wheat Lipid-Transfer Protein from Multidimensional ¹H-NMR Data. A New Folding for Lipid Carriers. *Eur J Biochem* 226:413–422

29. Shin DH, Lee JY, Hwang KY, Kim KK, Suh SW (1995) High-resolution crystal structure of the non-specific lipid-transfer protein from maize seedlings. *Structure* 3:189–199
30. Jaeckels N, Tenzer S, Rosfa S, Schild H, Decker H, Wigand P (2013) Purification and structural characterisation of lipid transfer protein from red wine and grapes. *Food Chem* 138:263–269
31. Hartz C, Lauer I, del Moncin MSanM, Cistero-Bahima M, Foetisch A, Lidholm K, Vieths J, Scheurer S (2010) Comparison of IgE-binding capacity, cross-reactivity and biological potency of allergenic non-specific lipid transfer proteins from peach, cherry and hazelnut. *Int Arch Allergy Immunol* 153:335–346
32. Gadermaier G, Harrer A, Girbl T, Palazzo P, Himly M, Vogel L, Briza P, Mari A, Ferreira F (2009) Isoform identification and characterization of Art v 3, the lipid-transfer protein of mugwort pollen. *Mol Immunol* 46:1919–1924
33. Diz MS, Carvalho AO, Ribeiro SF, Da Cunha M, Beltramini L, Rodrigues R, Nascimento VV, Machado OL, Gomes VM (2011) Characterisation, immunolocalisation and antifungal activity of a lipid transfer protein from chili pepper (*Capsicum annum*) seeds with novel alpha-amylase inhibitory properties. *Physiol Plant* 142:233–246
34. Zottich U, Da Cunha M, Carvalho AO, Dias GB, Silva NC, Santos IS, do Nascimento VV, Miguel EC, Machado OL, Gomes VM (2011) Purification, biochemical characterization and antifungal activity of a new lipid transfer protein (LTP) from *Coffea canephora* seeds with alpha-amylase inhibitor properties. *Biochim Biophys Acta* 1810:375–383
35. da Silva FCV, do Nascimento VV, Fernandes KV, Machado OLT, da Silva Pereira L, Gomes VM, Carvalho AdO (2018) Recombinant production and α -amylase inhibitory activity of the lipid transfer protein from *Vigna unguiculata* (L. Walp.) seeds. *Process Biochem* 65:205–212
36. Matsuura Y (2002) A possible mechanism of catalysis involving three essential residues in the enzymes of α -amylase family. *Biologia*; 11
37. Svensson B, Fukuda K, Nielsen PK, Bønsager BC (2004) Proteinaceous α -amylase inhibitors. *Biochimica et Biophysica Acta (BBA) - Proteins and Proteomics*. 1696:145–156
38. Kader JC (1996) Lipid-Transfer Proteins in Plants. *Annu Rev Plant Physiol Plant Mol Biol* 47:627–654
39. Regente MC, Giudici AM, Villalain J, de la Canal L (2005) The cytotoxic properties of a plant lipid transfer protein involve membrane permeabilization of target cells. *Lett Appl Microbiol* 40:183–189
40. Wildner S, Griessner I, Stemeseder T, Regl C, Soh WT, Stock LG, Volker T, Alessandri C, Mari A, Huber CG, Stutz H, Brandstetter H, Gadermaier G (2019) Boiling down the cysteine-stabilized LTP fold - loss of structural and immunological integrity of allergenic Art v 3 and Pru p 3 as a consequence of irreversible lanthionine formation. *Mol Immunol* 116:140–150
41. Sanchez-Lopez J, Tordesillas L, Pascal M, Munoz-Cano R, Garrido M, Rueda M, Vilella R, Valero A, Diaz-Perales A, Picado C, Bartra J (2014) Role of Art v 3 in pollinosis of patients allergic to Pru p 3. *J Allergy Clin Immunol* 133:1018–1025

Figures

A



B




Figure 1

Phylogenetic and sequence analyses.(A) Phylogenetic analysis of FtLTP1a and FtLTP1b. The analysis was carried out by MEGA7 using 114 amino acid sequences and the Neighbor-Joining method. The figure is available in color at JXB online. (B) Multiple amino acid sequence alignment. Two nsLTPs in this work

were compared with sequences of Api g2, Pru p3, and Tri a14. The alphabets A-C represent the epitopes identified in Pru p3 by (A) Garcia-Casado *et al.* (2003), (B) Borges *et al.* (2003), (C) Tordesillas *et al.* (2003).




Figure 2

SDS-PAGE analysis of FtLTP1a (A) and FtLTP1b (B). Mr: molecular markers; lane 1: soluble protein (pET32a); lane 2: insoluble protein (pET32a); lane 3: soluble protein without induction by IPTG (pET32a-FtLTP1a/b); lane 4: insoluble protein induced by IPTG (pET32a-FtLTP1a/b); lane 5: soluble protein induced by IPTG (pET32a-FtLTP1a/b); lane 6: insoluble protein induced by IPTG (pET32a-FtLTP1a/b); and lane 7: purified FtLTP1a or FtLTP1b containing Trx-His-tag.




Figure 3

Lipid binding activity of FtLTP1a and FtLTP1b. (A) Intrinsic fluorescence spectra. (B) Intrinsic fluorescence spectra of FtLTP1a in the presence of oleic acid. a: without oleic acid; b-k: different concentrations of oleic acid. (C) Intrinsic fluorescence spectra of FtLTP1b in the presence of oleic acid. a: without oleic acid; b-k: different concentrations of oleic acid. (D) Variation of fluorescence intensity of FtLTP1a and FtLTP1b as a function of oleic acid concentration.




Figure 4

α-Amylase activity of FtLTP1a and FtLTP1b.




Figure 5

Inhibitory effects of FtLTP1a and FtLTP1b against *Aspergillus niger*. FtLTP1a or FtLTP1b was incubated with *Aspergillus niger* at different concentrations (1: 10^5 CFU/mL; 2: 10^4 CFU/mL; and 3: 10^3 CFU/mL), and those incubated with sterilized distilled water were used as controls. Growth inhibition was measured after 24 h of incubation at 28°C. The concentration of FtLTP1a and FtLTP1b was 100 μ mol/L.




Figure 6

IgE-binding activity of purified FtLTP1a and FtLTP1b. (A) The binding of IgE in 2-fold diluted sera of patients allergic to buckwheat. (B) The binding of IgE in 10-fold diluted sera of patients allergic to buckwheat. (C) Cross-reactivity between FtLTP1a or FtLTP1b and pollen. The assay was performed using three replicate samples. P/N values > 2.1 were used as a benchmark for evaluating the presence of allergenic activity.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementaryInformationforPFHN.docx](#)
- [Supplementaryinformationforgelsandblotsimage.docx](#)

Supplementary Information

Figure S1

FtLTP1a

1	ATG	GCA	TTC	TCC	5	TAC	ACC	AAG	CTT	GCC	10	TGC	CTC	TTG	ATC	GTC	15	TGC	TTC	GCA	GTG	GCT	20	GTT	CCC	GCA	GCT	25	GAA
	M	A	F	S	Y	T	K	L	A	C	L	L	I	V	C	F	A	V	A	V	P	A	A	V	E				
26	GGG	TTC	GAA	TGC	30	GGT	GCC	GTC	TCG	AAG	35	AAC	CTC	GTC	CAG	TGC	40	ATC	ACT	TTC	CTT	AAA	45	GGC	GCT	GCT	GGC	50	AAC
	G	F	E	C	G	A	V	S	K	N	L	V	Q	C	I	T	F	L	K	G	A	A	G	P	N				
51	AAA	GCT	TGT	TGC	55	GGT	GGC	GTC	AGG	ACT	60	CTT	GCT	TCA	ATG	GCG	65	CAG	ACC	CCT	GCC	GAC	70	CGC	CAG	AAA	GCT	75	GGC
	K	A	C	C	G	G	V	R	T	L	A	S	M	A	Q	T	P	A	D	R	Q	K	A	C	G				
76	TGT	CTC	AAG	AAC	80	TCC	GCC	GCC	GCC	ATC	85	CCC	GGC	CTT	AAC	TTG	90	CAT	AAT	GCC	GCC	GCC	95	CTC	CCC	GGC	TTG	100	CAA
	C	L	K	N	S	A	A	A	I	P	G	L	N	L	H	N	A	A	A	L	P	G	L	C	Q				
101	ATC	AGT	GTT	CCC	105	TAC	CCC	ATT	AGC	CCC	110	AGC	ACC	GAC	TGC	TCC	115	AAG	GTT	CAC	TAA								
	I	S	V	P	Y	P	I	S	P	S	T	D	C	S	K	V	H	*											

FtLTP1b

1	ATG	GCT	TTC	TCC	5	TAC	ACC	AAG	CTC	GCC	10	TGC	CTC	TTG	ATG	GTG	15	TGC	ATG	GCA	GTG	GCA	20	GCT	CCG	GAA	GCC	GAG	25	GCC
	M	A	F	S	Y	T	K	L	A	C	L	L	M	V	C	M	A	V	A	A	P	E	A	E	A					
26	GCG	CTC	ACA	TGC	30	GGG	ACA	GTC	ACC	AAG	35	AAC	CTC	GTC	CCG	TGC	40	CTC	CCT	TAC	CTT	AAG	45	GGT	GGG	ACA	GGT	50	TCC	
	A	L	T	C	G	T	V	T	K	N	L	V	P	C	L	P	Y	L	K	G	G	T	G	P	S					
51	GTC	CCG	TGT	TGT	55	AAT	GGC	GTG	AAG	GCG	60	CTG	GCC	TCC	AGT	GCT	65	AAC	ACC	CCT	GCC	GAC	70	CGC	AAG	ATC	GCT	75	GGC	
	V	P	C	C	N	G	V	K	A	L	A	S	S	A	N	T	P	A	D	R	K	I	A	C	G					
76	TGC	CTC	AAG	CAG	80	TCC	GGT	GCT	ATT	TCC	85	GGC	CTT	AAC	TTA	GGC	90	GGC	AAT	GCC	GCC	GCC	95	CTC	CCC	GGC	AAG	100	GGC	
	C	L	K	Q	S	S	G	A	I	S	G	L	N	L	G	N	A	A	A	L	P	G	K	C	G					
101	GTC	AGT	GTT	CCC	105	TAC	AAA	ATC	AGC	CCC	110	AGC	ACC	GAC	TGC	TCC	115	AAG	GTT	AAC	TAA									
	V	S	V	P	Y	K	I	S	P	S	T	D	C	S	K	V	N	*												

Figure S1. cDNA sequences and amino acid sequences of FtLTP1a and FtLTP1b. Red font is predicted signal peptide sequence, and the green font is conserved regions for lipid binding.

Figure S2




Figure S2. Circular dichroism spectra of FtLTP1a and FtLTP1b.

Table

Table S1. Data on serum of specific buckwheat allergen

Serum No.	Sex	Ages	IgE specific for buckwheat (kU/L)	Other known allergens*
1	M	19	0.261	D1, H1, HX2
2	M	21	0.140	H1, HX2, MX2, TX4, W6, WX7
3	M	32	0.321	W6
4	M	27	0.790	W6
5	M	46	0.650	W6
6	F	15	0.173	T8, T12, TX4, W6, W22, WX7
7	F	23	0.151	TX4, W22, W6, WX7
8	M	14	0.216	W22, W6
9	M	27	0.133	TX4, W22, W6, WX7

*D1, house dust mite; D2, dermatophagoides farina; T8, elm pollen; T12, willow pollen; TX4, polyvalent tree pollen; H1, house dust; HX2, polyvalent mite; W6, artemisia pollen; W22, humulus; WX7, polyvalent grass pollen; MX2, polyvalent mold.

Table S2. Data on serum of pollen allergen

Serum No.	Sex	Ages	IgE specific for W6* (kU/L)	IgE specific for WX7* (kU/L)
1	F	23	3.169	2.053
2	M	37	3.341	2.477
3	F	14	8.919	6.523

*W6, artemisia pollen; WX7, polyvalent grass pollen;

Table S3. The purification process of FtLTP1a and FtLTP1b from per liter of culture

Steps	FtLTP1a	FtLTP1b
Wet cell mass(g)	17.12	19.29
Crude proteins(mg)	35.79	45.09
Target proteins(mg)	2.14	2.46

*Crude proteins were supernatant of *E.coli* after ultrasonic.

Table S4. MICs of the FtLTP1a and FtLTP1b

Microorganisms.	FtLTP1a (μmol/L)	FtLTP1b (μmol/L)
<i>Aspergillus niger</i>	8	16
<i>Aspergillus flavus</i>	8	16
<i>Fusarium oxysporum</i>	64	64

Uncropped gels image for figure 2

