https://doi.org/10.1093/nar/gkac446

AllerCatPro 2.0: a web server for predicting protein allergenicity potential

Minh N. Nguyen [1,2], Nora L. Krutz[3], Vachiranee Limviphuvadh[1,2], Andreas L. Lopata[1,4,5], G. Frank Gerberick[6] and Sebastian Maurer-Stroh [1,2,7,*]

[1]Bioinformatics Institute, 30 Biopolis Street, #07-01, Matrix 138671, Singapore, [2]IFCS Programme, Singapore Institute for Food and Biotechnology Innovation, Agency for Science, Technology and Research, Singapore, [3]NV Procter & Gamble Services Company SA, Strombeek-Bever, Belgium, [4]Molecular Allergy Research Laboratory, Australian Institute of Tropical Health and Medicine, James Cook University, Townsville, QLD, Australia, [5]Tropical Futures Institute, James Cook University, Singapore, [6]GF3 Consultancy LLC, West Chester, OH, USA and [7]Department of Biological Sciences, National University of Singapore, 14 Science Drive 4, 117543, Singapore

Received March 21, 2022; Revised May 07, 2022; Editorial Decision May 09, 2022; Accepted May 11, 2022

ABSTRACT

Proteins in food and personal care products can pose a risk for an immediate immunoglobulin E (IgE)-mediated allergic response. Bioinformatic tools can assist to predict and investigate the allergenic potential of proteins. Here we present AllerCatPro 2.0, a web server that can be used to predict protein allergenicity potential with better accuracy than other computational methods and new features that help assessors making informed decisions. AllerCatPro 2.0 predicts the similarity between input proteins using both their amino acid sequences and predicted 3D structures towards the most comprehensive datasets of reliable proteins associated with allergenicity. These datasets currently include 4979 protein allergens, 162 low allergenic proteins, and 165 autoimmune allergens with manual expert curation from the databases of WHO/International Union of Immunological Societies (IUIS), Comprehensive Protein Allergen Resource (COMPARE), Food Allergy Research and Resource Program (FARRP), UniProtKB and Allergome. Various examples of profilins, autoimmune allergens, low allergenic proteins, very large proteins, and nucleotide input sequences showcase the utility of AllerCatPro 2.0 for predicting protein allergenicity potential. The AllerCatPro 2.0 web server is freely accessible at https://allercatpro.bii.a-star.edu.sg.

GRAPHICAL ABSTRACT



INTRODUCTION

Evaluating the allergenic potential of proteins with the help of bioinformatic tools has gained increasing interest for foods, consumer and personal care products (1). However, the accurate prediction and evaluation of the allergenic potential of novel proteins is still a challenge. The computational approaches using the FAO/WHO guidelines of linear sequence window identity thresholds and short peptide hits for predicting protein allergenicity potential misclassify a large number of protein sequences as allergens (2). In addition, the FAO/WHO linear-window rule (3) and the current popular methods of PREAL (4), AllerHunter (5), AllergenFP (6), AllerTOPv2 (7) and AlgPred 2.0 (8) only achieved the overall accuracies from 51% (FAO/WHO rules (3)) to respectable 73% (7) on our benchmark datasets (2).

Many different factors can contribute to the overall allergenicity of proteins, such as: cleavage sites, protein stability,

*To whom correspondence should be addressed. Tel: +65 6478 8377; Fax: +65 6478 9542; Email: sebastianms@bii.a-star.edu.sg

© The Author(s) 2022. Published by Oxford University Press on behalf of Nucleic Acids Research.
This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial License (http://creativecommons.org/licenses/by-nc/4.0/), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

Downloaded from https://academic.oup.com/nar/advance-article/doi/10.1093/nar/gkac446/6595260 by guest on 02 June 2022

physico-chemical properties, and post-translational modifications (9). Nevertheless, a protein allergen still needs to be recognized by the immune system and thus, in order to sensitize an individual, induce a T cell and B cell response to produce allergen-specific Immunoglobulin (IgE) and/or, in order to elicit an IgE-mediated allergic response, interact with IgE on mast cells and basophils. These interactions and recognitions of allergenic proteins can be captured by their amino acid sequences and 3D structures. Here, we have developed AllerCatPro 2.0 web server using the computational method AllerCatPro established in our previous study (AllerCatPro 1.7) (2) to predict the allergenic potential of protein sequences based on similarity of both their amino acid sequences and 3D structures compared with the most comprehensive datasets of reliable proteins associated with allergenicity. Our AllerCatPro method was extensively benchmarked and compared with other popular methods for predicting protein allergenicity potential. Evaluation of the benchmark datasets used, AllerCatPro achieved an overall accuracy of 84%, significantly better than other methods (2).

Moreover, in this study we have improved AllerCatPro by enabling nucleotide alongside protein sequences as input and extending the dataset of 4180 (2) to 4979 proteins associated with allergenicity to improve the accuracy of AllerCatPro 2.0 for predicting protein allergenicity potential. We have also identified and added 165 human proteins associated with autoimmune diseases (in the following called 'autoimmune allergens') and 162 proteins of low allergenic potential ('low allergenic proteins'). The latter were identified from a dataset derived from six abundant and commonly consumed protein sources, resulting in a list of 178 characterized proteins with no evidence for allergenicity, despite opportunities for human exposure (10). The autoimmune allergens were extracted and curated from the databases of WHO/International Union of Immunological Societies (IUIS), Comprehensive Protein Allergen Resource (COMPARE), Food Allergy Research and Resource Program (FARRP), UniProtKB and Allergome. For our allergen database, we carefully selected annotated allergens from available sources that provide further evidence and/or have established procedures for manual expert curation, and hence the number of false annotations should be marginal and not having significant influence on performance estimates.

The prediction for the most similar protein allergen, similarity scores in protein sequence ('% identity, linear 80 aa window') and 3D structure ('% identity, 3D epitope'), Gluten-like repeats of Glutamine ('Gluten allergens (# of Q-repeats)'), and hexamer hits ('# of 3 × 6-mer overlaps') are based on the output of AllerCatPro 1.7 as presented earlier (2). For AllerCatPro 2.0, we now developed a more comprehensive output and extended the result to potential cross-reactivity, protein information (UniProt/NCBI) and functionality (Pfam, InterPro, SUPFAM), as well as the clinical relevance which refers to IgE prevalence data from Allergome and allergen information related to the most similar allergen (Figure 1). The allergen information refers to the allergenicity scoring used by Allergome and represents the number of positive evidence types annotated in the database of Allergome. Furthermore, we extended the applicability domain of AllerCatPro 2.0 to perform well with very long input sequences (>1000 amino acids), which was a limitation in the AllerCatPro 1.7. Various examples showcase the utility of the AllerCatPro 2.0 web server for predicting protein allergenicity potential from protein sequences as well as present the improvement of AllerCatPro 2.0 compared to AllerCatPro 1.7.

IMPLEMENTATION

Program overview

Our web server is developed using the AllerCatPro method established in our previous study (AllerCatPro 1.7) (2). The input of AllerCatPro 2.0 is the query protein/nucleotide sequences and the output of those queried sequences is the similarity to protein allergens and the resulting predicted strong, weak, or no evidence for allergenicity accompanied with a comment section to clarify the rule triggering the result. The allergenicity potential for the query protein is predicted using different measures of similarity of amino acid sequence and 3D structure compared with our dataset of known allergens. Since Gluten-like repeats of Glutamine (Gluten-like Q-repeats) are independent of any other similarity scores, the presence of Gluten-like Q-repeats is first evaluated in our workflow (Figure 2). The Gluten-like prediction of AllerCatPro 2.0 does not lead to 'strong evidence' unless there is similarity to a known protein allergens. Subsequently, AllerCatPro 2.0 checks the similarity of the queried sequence with representatives in our 3D model/structure database of known allergens. Currently, there are 714 representative protein allergens in our 3D model/structure database (2). If there is significant sequence similarity to the BLASTP search (11) for the input sequence against our 3D model/structure database of known allergens (E-value < 0.001), then the similarity of the 3D surface epitope is applied to predict as 'strong evidence' as the sequence identity of 3D surface epitope is >93% (this cutoff is 92% if Gluten-like Q-repeats are found in the query sequence) or as 'weak evidence' otherwise (Figure 2). If AllerCatPro 2.0 cannot find any structure hit, the linear-window approach is then applied to predict the queried sequence as protein allergen with strong evidence if the rule of 35% identity over 80 residues is found. After using the linear-window rule, and no hit is found, the hexamer hit approach with at least three short hexamers with known allergens is applied to evaluate the queried sequence (Figure 2). Finally, if AllerCatPro 2.0 still cannot find any hit, the prediction of 'no evidence' for allergenicity is assigned (Figure 2).

In addition to comparing the similarity of the query protein with the dataset of known allergens, our new web server AllerCatPro 2.0 now predicts the similarity of the query sequence to datasets of autoimmune allergens and low allergenic proteins separately. If a significant sequence similarity is found, then AllerCatPro 2.0 identifies hits of similar proteins associated with autoimmune diseases and/or similar proteins of low allergenic potential, and presents the sequence identity to the closest hit. As shown in Supplementary Figures S3A and S4, AllerCatPro 2.0 correctly predicts the query protein sequences as autoimmune allergens and low allergenic proteins while AllerCatPro misclassified

Downloaded from https://academic.oup.com/nar/advance-article/doi/10.1093/nar/gkac446/6595260 by guest on 02 June 2022



Figure 1. Submitting one or more protein/nucleotide sequences in FASTA format (**A**) leads to the AllerCatPro 2.0 output table with the result for strong, weak or no evidence for allergenicity per protein based on corresponding workflow decisions and, in case of a hit, the possibility to view the most similar allergens with the detailed results for cross-reactivity, protein information (UniProt/NCBI), functionality (Pfam, InterPro, SUPFAM), as well as clinical relevance of IgE prevalence (Allergome) and allergen information (**B**), the most similar 3D surface epitope via links with the structural view showing identical epitope residues as beads colored as blue for positive charges, red for negative charges and gray for all other amino acid types (**C**). AllerCatPro 2.0 also identifies all similar allergens that have significant sequence similarity to the query protein and refers to the number with the link in potential cross-reactivity of the output table (**D**), as well as all possible similar autoimmune allergens displayed in the link (**E**) and all possible similar low allergens in the link of output table.

these input sequences as allergenic proteins with 'strong' or 'no evidence' for allergenicity, respectively.

Recently, Sharma *et al.* have developed AlgPred 2.0 for predicting allergenic proteins based on a combination of different scores generated from machine learning (random forest) method, motif-emerging and with classes-identification, and BLAST search (8). Benchmark datasets are crucial to evaluate the performance of prediction models and they should include a set of allergens and a set of non-allergens with the same structure fold (2). Thus, AllerCatPro 2.0 was compared to AlgPred 2.0 on our benchmark datasets of 218 positive (known allergen and selected to be structurally non-redundant using our CLICK method (12–15)) and 212 negative (likely non-allergen) sequences. These benchmark datasets are available on the help page (https://allercatpro.bii.a-star.edu.sg/help.html) or via the help button at the AllerCatPro 2.0 website and are extracted from the benchmark datasets of 221 positive and 221 negative sequences (2) with the following adaptations: We removed one low allergenic protein ('Spi o RuBisCo' of *Spinacia oleracea*) and two proteins associated with autoimmune diseases in the positive set, and nine protein sequences related to the protein allergen 'Ory s 14' ('Ory s 14' of *Oryza sativa*) in the negative set due to its allergenic potential (16).

On our datasets, AllerCatPro 2.0 achieves an accuracy of 84.7% at 100% sensitivity and 68.9% specificity and Matthews correlation coefficient (MCC) of 0.727 (Supplementary Table S3) while AlgPred 2.0 only obtains an accuracy of 52.3% at 97.2% sensitivity and 6.1% specificity and MCC of 0.08. Also, we compare AllerCatPro 2.0 with other popular methods of PREAL (4), AllerHunter (5), AllergenFP (6), and AllerTOPv2 (7) on these benchmark datasets. We emphasize that these datasets are small but well representative of protein allergens with structures (2). The accuracy and MCC of AllerCatPro 2.0 are significantly better than those of the other methods with accuracies from 52.3% (AlgPred 2.0) to 75.0% (AllerTOPv2) and MCC from 0.08 (AlgPred 2.0) to 0.504 (AllerTOPv2) (Supplementary Figures S1A and B).

Downloaded from https://academic.oup.com/nar/advance-article/doi/10.1093/nar/gkac446/6595260 by guest on 02 June 2022



Figure 2. Decision workflow of AllerCatPro 2.0 from the query protein to the results of either strong, weak or no evidence for allergenic potential. AllerCatPro 2.0 checks the similarity of the query protein with 714 representatives in our 3D model/structure database of known allergens as well as the most comprehensive dataset of reliable proteins associated with allergenicity (4979 protein allergens). In addition to only comparing the similarity of the query protein with the dataset of known allergens in AllerCatPro 1.7, AllerCatPro 2.0 now predicts the similarity of the query sequence to datasets of 165 autoimmune allergens and 162 low allergenic proteins separately. If a significant sequence similarity is found, then AllerCatPro 2.0 identifies hits of similar proteins associated with autoimmune diseases and/or similar proteins of low allergenic potential and presents the sequence identity to the closest hit.

In addition, we perform AllerCatPro 2.0 on the larger validation datasets of 2003 positive (allergen) and 2015 negative (non-allergen) sequences from AlgPred 2.0 (8). These datasets are extracted from the validation datasets of 2015 positive and 2015 negative sequences of AlgPred 2.0 (https://webs.iiitd.edu.in/raghava/algpred2/stand.html) with the following adaptations: We remove one protein (P_10228) that is identical to low allergenic protein (*Phoenix dactylifera*) and eleven proteins (P_7089, P_7095, P_7102, P_7098, P_7101, P_7108, P_7125, P_7117, P_7114, P_7092, P_7111) associated with autoimmune diseases in the positive set. On these datasets, AllerCatPro 2.0 achieves high accuracy of 96.0% at 93.2% sensitivity and 98.8% specificity and MCC of 0.921 that are better than those of AllerCatPro 1.7 with accuracy of 93.0% at 91.1% sensitivity and 94.8% specificity and MCC of 0.860 (Supplementary Table S2). Since both AllerCatPro 2.0 and AllerCatPro 1.7 use the same algorithm and measures of similarity of amino acid sequence and 3D structure compared with the database of known allergens, the improvements of accuracy and MCC of AllerCatPro 2.0 compared to those of AllerCatPro 1.7 are due to the 1132 sequences of protein allergens, low allergenic proteins, and autoimmune allergens newly included into the database of AllerCatPro 2.0.

Downloaded from https://academic.oup.com/nar/advance-article/doi/10.1093/nar/gkac446/6595260 by guest on 02 June 2022

We also test AllerCatPro 2.0 on all recent IUIS allergens (twenty proteins) that has been created/modified from December 2021 to April 2022. For these new IUIS allergens, AllerCatPro 2.0 predicts their allergenicity potential correctly for fourteen proteins with 'strong evidence', five proteins with 'weak evidence' and incorrectly for only one protein with 'no evidence'.

Furthermore, we perform AllerCatPro 2.0 with and without using 3D structure similarity on the benchmark datasets of 218 allergens and 212 non-allergens with the same structure fold. On these benchmark datasets, AllerCatPro 2.0 achieves accuracy of 84.7% and MCC of 0.727 while AllerCatPro 2.0 without using 3D structure similarity only obtains accuracy of 52.8% and MCC of 0.148 (Supplementary Table S3). These results indicate that 3D structure similarity contributes significantly to the performance of AllerCatPro 2.0.

Our all datasets are provided on the help page (https://allercatpro.bii.a-star.edu.sg/help.html).

Server description

Input. Query sequences can be submitted by specifying their sequences in FASTA format, or by uploading a FASTA file of sequences (Figure 1A). The FASTA format has a description line starting with '>' for each entry followed by one or more lines of sequence. In our AllerCatPro 2.0 web server, we have improved the AllerCatPro method to allow users to input nucleotide sequences (Supplementary Figures S6A and B). In addition, AllerCatPro 2.0 has been developed to perform well with very long input sequences (>1000 amino acids) and allow users to submit more than 50 protein/nucleotide sequences at a time, which were limitations of AllerCatPro 1.7.

A detailed explanation of input sequences for predicting the allergenicity potential of AllerCatPro 2.0 is provided on the help page (https://allercatpro.bii.a-star.edu.sg/help.html).

Output. The output of AllerCatPro 2.0 for input protein/nucleotide sequences is a result table following the workflow in the 'Program Overview' section and allergenicity prediction for one protein per line (Figure 1). The output table presents the detailed results, i.e. Gluten-like Q-repeats, hexamer hits, the most similar allergen (in case there is significant sequence similarity), potential hits for cross-reactivity, protein information (UniProt/NCBI), functionality (Pfam, InterPro, SUPFAM), as well as the clinical relevance of IgE prevalence (derived from Allergome) and allergen information related to the most similar allergen (Figure 1B). The IgE prevalence result shows the total number of studied individuals used for IgE tests and provide also the link to the 'Epidemiology from Literature' page on the Allergome website. The allergen information for the most similar allergen refers to the allergenicity scoring used by Allergome and represents the number of positive evidence types annotated in the database of Allergome (e.g. Functional Test, Non-Functional Test, Skin Test, Conjunctival Provocation Test, Nasal Provocation Test, Bronchial Provocation Test, Oral Challenge, Epidemiology from Literature, ReTiME). In addition, this allergen information is accompanied with an annotation, if the most similar allergen is a known food or insect allergen and related to anaphylaxis.

AllerCatPro 2.0 also checks the similarity score of the query protein with the most similar allergen based on the linear-window approach and identifies the similarity on a 3D structure level, if there is significant sequence similarity based on the BLASTP search (11) for the query protein against the 3D model/structure database of known allergens. These results are shown in the output table in case of a 3D similarity hit (Figure 1B). If the 3D surface epitope similarity is found, AllerCatPro 2.0 presents a link to view the potential 3D epitope with the highest percent amino acid identity relative to the query and visualizes on the closest known 3D structure (Figure 1C). Since charge similarity can be important, positions of identical positively charged residues of 3D epitope are shown as blue, negatively charged as red and all other identical residues as grey beads. The 3D rendering of the epitope and the closest known structure is displayed using JSMol (http://www.jmol.org/). Finally, AllerCatPro 2.0 classifies the query protein based on the weight of evidence coming from the different levels of similarity to known allergens (Gluten-like Q-repeats, 3D epitopes, linear-window rule, and hexamer hits). A red label of 'strong evidence' presents that there is a clear similarity to known allergens. In contrast, a light green label of 'weak evidence' shows some similarity to known allergens but with clear divergence in relevant sequence and structure features (Figure 1B). The dark green label of 'no evidence' means that no significant similarity of sequence and structure features has been found against our current most comprehensive dataset of 4979 known allergens (Figure 1B).

Moreover, instead of providing only the most similar allergen for the query protein in AllerCatPro 1.7, AllerCatPro 2.0 now identifies all similar allergens within the AllerCatPro 2.0 dataset that have significant sequence similarity to the query protein (E-value of BLASTP search < 0.001) in the column 'Potential cross-reactivity of query protein (# and links)' as a number and a link to the according list of allergens (Figure 1B). The list, which opens in a separate browser tab, shows protein information (UniProt/NCBI), functionality (Pfam, InterPro, SUPFAM), sequence identity and E-value of all similar allergens (Figure 1D). In order to differentiate between typical environmental allergens versus autoimmune allergens as well as increasing the confidence in predicting the absence of allergenicity, AllerCatPro 2.0 now provides the similarity of the query proteins with the datasets of autoimmune allergens as well as low allergenic proteins. If there is significant sequence similarity, then AllerCatPro 2.0 finds hits of similar proteins associated with autoimmune diseases and/or similar proteins of low allergenic potential and presents the sequence identity to the closest hit with the link to a separate browser tab showing protein information, functionality, sequence identity, and E-value of all hits in the output table (Figure 1E).

The output table also includes a download link for the results of the query protein in comma-separated CSV format which can be opened by Excel or other popular spreadsheet programs. A detailed explanation of the output for predicting allergenicity potential of AllerCatPro 2.0 is provided

Downloaded from https://academic.oup.com/nar/advance-article/doi/10.1093/nar/gkac446/6595260 by guest on 02 June 2022

on the help page (https://allercatpro.bii.a-star.edu.sg/help.html).

RESULTS

Case study 1: differentiation of allergenic from low allergenic profilin proteins

The profilin protein family is one of the biggest families associated with food and pollen allergies, profilins are ubiquitous and highly conserved with sequence identities of 70% to 85% among allergenic plant profilins (17). Profilin is a clinically relevant aeroallergen and given its ubiquity, sensitized individuals might react to multiple profilin-containing sources with potentially even more severe symptoms, which makes this protein family highly relevant for both diagnosis and treatment of patients with plant food and pollen allergy (18).

In this example, we analyse and predict the allergenicity potential of different profilin proteins using AllerCatPro 2.0. Figure S2A in Supplementary shows all four pollen profilins from European white birch (*Betula pendula*), olive (*Olea europaea*), Timothy (*Phleum pratense*), and rice (*Oryza sativa*) are classified correctly as allergenic proteins with 'strong evidence' for allergenicity. Among these four proteins, three pollen profilins of *Betula pendula*, *Olea europaea*, *Phleum pratense* demonstrate high numbers of individuals tested for allergen-specific IgE (Supplementary Figure S2A), which indicates clinical relevance for these allergens. Figure S2B in Supplementary displays the 3D model of pollen profilin from Betula pendula based on the similarity of the 3D surface epitope. In this 3D model, blue, red and grey beads present positions of identical positively, negatively charged and the other identical residues of the 3D epitope, respectively (Supplementary Figure S2B).

As shown in Figure S2C in Supplementary, AllerCatPro 2.0 predicts correctly allergenic plant profilins from peach, apple, hazelnut, potato, spinach, and Para rubber tree with 'strong evidence' for allergenicity. All except the potato allergen Sola t 8, show high numbers of individuals tested for allergen-specific IgE.

For a set of five low allergenic profilins from yeast, human, cow, chicken, and fungus, AllerCatPro 2.0 predicts their allergenicity potential correctly with 'weak evidence' (Supplementary Figure S2D). On this set, AllerCatPro 1.7 overpredicted two profilins from human and cow with 'strong evidence' for allergenicity.

Case study 2: identification of autoimmune allergens

Many human proteins such as HLA (human leukocyte antigen)-DR and type I interferons importantly contribute to the pathogenesis of autoimmune diseases for type-1 diabetes, rheumatoid arthritis, and systemic lupus erythematosus (19,20). In this example, AllerCatPro 2.0 is used to identify the similarity to human proteins of HLA-DR alpha, alpha-galactosidase, nascent polypeptide-associated complex subunit alpha, granulocyte colony-stimulating factor, B-cell CLL/lymphoma 7 protein family member B, and type I interferon as autoimmune allergens. Since there are significant sequence similarity in BLASTP search for these input sequences against our dataset of autoimmune allergens, AllerCatPro 2.0 identified these proteins with 100% similarity to proteins associated with autoimmune diseases and shows in the output table that no significant hit ('no evidence') for environmental (non-human) allergens have been found (Supplementary Figure S3A). Compared to AllerCatPro 2.0, AllerCatPro 1.7 did not distinguish between environmental and autoimmune allergens, but assigned query proteins similar or identical to human proteins as allergens with 'strong evidence'. AllerCatPro 2.0 now can identify if a protein is similar to environmental allergens and/or similar to human proteins, which provides additional information for users.

In addition, AllerCatPro 2.0 identifies all proteins from the dataset of autoimmune allergens with significant sequence similarity to the query protein sequences. Figure S3B in Supplementary shows the list of related human protein sequences (all related to interferon alpha and beta) that are associated with autoimmune diseases and have significant sequence similarity to the input sequence of type I interferon.

Case study 3: identification of proteins with low allergenic potential

Since different levels of similarity are applied to identify evidence for allergenic potential (Gluten-like Q-repeats, 3D epitopes, linear-window rule, and hexamer hits) in AllerCatPro and the dataset of known allergenic proteins grows with each new version, the likelihood to predict similarity with weak evidence (especially due to the linear-window rule) to a query protein increases steadily. However, not all proteins predicted with weak evidence for allergenicity are necessarily of allergenic concern. In order to improve the output and prediction of low allergenic potential in AllerCatPro 2.0, we implemented a dataset of proteins derived from our previous publication (10) for which we have confidence that these proteins have only a low allergenic potential. In this example, we use AllerCatPro 2.0 to predict the similarity of 10 kDa prolamin from rice (*Oryza sativa*), a flower-specific gamma-thionin-like protein from tomato (*Solanum lycopersicum*), ribulose bisphosphate carboxylase from spinach (*Spinacia oleracea*), late embryogenesis abundant proteins 14, 17, 19 and group 3 from wheat (*Triticum aestivum*) to low allergenic proteins. The results show that AllerCatPro 2.0 provides the comment 'Identical to low allergenic protein' in addition to the 'weak evidence' for allergenic potential (Supplementary Figure S4), which is triggered if AllerCatPro 2.0 identifies that they have very high sequence similarity and identity to other proteins in our dataset of low allergenic proteins.

Nevertheless, this is the very first time that a prediction tool provides the feature of predicting low allergenic potential of proteins. Science is still at the early beginning to better understand what makes a protein not an allergen and thus, every user needs to be aware that such result needs to be interpreted cautiously. Here, there feature 'potential cross-reactivity of query protein', which provides additional results on sequence similarity to other known allergens, may be important to evaluate in addition to the result of similarity to low allergenic proteins.

Downloaded from https://academic.oup.com/nar/advance-article/doi/10.1093/nar/gkac446/6595260 by guest on 02 June 2022

Compared to AllerCatPro 2.0, the previous version AllerCatPro 1.7 inappropriately assigned e.g. the input protein of ribulose bisphosphate carboxylase, which can be considered as low allergenic protein (10), as protein with 'strong evidence' for allergenicity. Additionally, in case of 'no evidence' or 'weak evidence' for allergenicity, the version 1.7 did not provide any additional evidence to substantiate low allergenic potential. AllerCatPro 2.0 now provides the similarity to low allergenic proteins, which are found to be appropriate to be used as negative controls to aid the development and validation of alternative methods for more robust testing strategies (10).

Case study 4: Prediction of similarity to allergens for very large proteins

One of the limitations of AllerCatPro 1.7 was the low performance and high running time for very large proteins (>1000 amino acids). AllerCatPro 2.0 is now able to predict protein allergenicity potential for very long input sequences in shorter time. As shown in Figure S5A in Supplementary, AllerCatPro 2.0 predicts allergenic potential for a set of very large proteins with their lengths varying from 834 to 3967 residues. The average running time for each protein was approximately 8.5 seconds. Similarity towards allergenic proteins for these large proteins is predicted using the usual levels based on Gluten-like Q-repeats, 3D epitopes, linear-window rule, but excluding the hexamer rule as it is over-predicting similarity for large proteins and takes more time to finalize a run.

Among the input sequences shown in Figure S5A, eleven proteins (all except Query# 9, 11, 13) are predicted to be similar to an allergen. For ten proteins (Query# 1 to 8, 12, and 14) allergen information is shown for the most similar protein allergen with the number of positive evidence types indicating allergenicity annotated in the database of Allergome. Two proteins (Query# 2 and 6) are similar to allergens for which more than 100 individuals have been tested for allergen-specific IgE (Supplementary Figure S5A).

Figure S5B in Supplementary displays the 3D structure for the input protein with the UniProtID A0A5F4DEY2 (Query# 2) and its 3D similarity to the most similar protein allergen 'Cuc p AscO' with blue and grey beads showing positions of identical positively charged and the other identical residues, respectively.

Case study 5: extended input capability to nucleotide sequences

In the new version of AllerCatPro 2.0, we have improved AllerCatPro 1.7 by enabling the input of nucleotide sequences in addition to protein sequences. For this case study, we apply AllerCatPro 2.0 to predict the allergenic potential of three known allergens (ABD51778, CAB02154, AAA34278) from honey bee (*Apis mellifera*), European white birch (*Betula pendula*), and wheat (*Triticum aestivum*) using their nucleotide sequences as input (Supplementary Figure S6A). As shown in Figure S6B in Supplementary, AllerCatPro 2.0 classifies correctly as allergens with 'strong evidence' for all these nucleotide input sequences. The predicted most similar allergen for the nucleotide sequence CAB02154 from *Betula pendula* is annotated with a high number of individuals tested for IgE prevalence. AllerCatPro 2.0 predicts the nucleotide input sequence (AAA34278) from *Triticum aestivum* with strong evidence for allergenicity due to the presence of 122 Gluten-like Q-repeats and 100% sequence identity over the window of 80 amino acid residues are found towards the most similar protein allergen Prolamin (Supplementary Figure S6B).

CONCLUSION

Here, we present AllerCatPro 2.0, a web server that can be used to predict protein allergenicity potential for protein or nucleotide sequences with outperforming accuracies (84.7% at 100% sensitivity and 68.9% specificity on the difficult benchmark datasets) and a user-friendly interface with new features that provide detailed results for potential cross-reactivity, protein information (UniProt/NCBI), functionality (Pfam, InterPro, SUPFAM), as well as the clinical relevance with regards to IgE prevalence and allergen information related to the most similar allergen. To the best of our knowledge, this is the first web server for predicting protein allergenicity potential equipped with the capability for identifying the clinical relevance, allergen information, sequence similarities to autoimmune allergens and low allergenic proteins that help assessors making informed decisions.

AllerCatPro 2.0 is developed based on our previous established AllerCatPro method (AllerCatPro 1.7) (2). AllerCatPro 2.0 predicts protein allergenicity potential using similarity of both their amino acid sequences and 3D structures towards the most comprehensive datasets (4979 protein allergens, 162 low allergenic proteins, and 165 autoimmune allergens) of reliable proteins associated with allergenicity from the databases of WHO/IUIS, COMPARE, FARRP, UniProtKB and Allergome.

Various examples of profilins, autoimmune diseases, low allergenic proteins, very large proteins, and nucleotide input sequences showcase the utility of AllerCatPro 2.0 for predicting protein allergenicity potential from protein sequences as well as demonstrate the improvement of AllerCatPro 2.0 compared to our previous version AllerCatPro 1.7.

SUPPLEMENTARY DATA

Supplementary Data are available at NAR Online.

ACKNOWLEDGEMENTS

This study has been supported by the Agency of Science, Technology and Research (A*STAR) Singapore and the Procter & Gamble (P&G) company. We would like to acknowledge the Biomedical Research Council (BMRC)-Economic Development Board (EDB), Industry Alignment Fund, A*STAR, Singapore for support. Minh N. Nguyen would like to thank HBMS Domain Industry Alignment Fund Pre-Positioning (IAF-PP), A*STAR (H2001a0P14) for funding. Vachiranee Limviphuvadh would like to thank Industry Alignment Fund BMRC, A*STAR/P&G (APG2013/096) for funding.

Downloaded from https://academic.oup.com/nar/advance-article/doi/10.1093/nar/gkac446/6595260 by guest on 02 June 2022

FUNDING

Agency for Science, Technology and Research (A*STAR) HBMS Domain Industry Alignment Fund Pre-Positioning (IAF-PP) [H2001a0P14]; Industry Alignment Fund BMRC/P&G [APG2013/096]. Funding for open access charge: OC/EFSA/GMO/2021/04.
Conflict of interest statement. None declared.

REFERENCES

1. Krutz,N.L., Kimber,I., Maurer-Stroh,S. and Gerberick,G.F. (2020) Determination of the relative allergenic potency of proteins: hurdles and opportunities. *Crit. Rev. Toxicol.*, **50**, 521–530.
2. Maurer-Stroh,S., Krutz,N.L., Kern,P.S., Gunalan,V., Nguyen,M.N., Limviphuvadh,V., Eisenhaber,F. and Gerberick,G.F. (2019) AllerCatPro-prediction of protein allergenicity potential from the protein sequence. *Bioinformatics*, **35**, 3020–3027.
3. Organization,W.H., Nations,F. and A.O. of the U. and Joint FAO/WHO Expert Consultation on Allergenicity of Foods Derived from Biotechnology 2001: Rome, I (2001) In: *Evaluation of allergenicity of genetically modified foods: report of a Joint FAO/WHO Expert Consultation on Allergenicity of Foods Derived from Biotechnology*. 22–25 January 2001 Food and Agriculture Organization of the United Nations, Rome.
4. Wang,J., Zhang,D. and Li,J. (2013) PREAL: prediction of allergenic protein by maximum relevance minimum redundancy (mRMR) feature selection. *BMC Syst. Biol.*, **7**, S9.
5. Muh,H.C., Tong,J.C. and Tammi,M.T. (2009) AllerHunter: a SVM-Pairwise system for assessment of allergenicity and allergic cross-reactivity in proteins. *PLoS One*, **4**, e5861.
6. Dimitrov,I., Naneva,L., Doytchinova,I. and Bangov,I. (2014) AllergenFP: allergenicity prediction by descriptor fingerprints. *Bioinformatics*, **30**, 846–851.
7. Dimitrov,I., Bangov,I., Flower,D.R. and Doytchinova,I. (2014) AllerTOP v.2–a server for in silico prediction of allergens. *J. Mol. Model.*, **20**, 2278.
8. Sharma,N., Patiyal,S., Dhall,A., Pande,A., Arora,C. and Raghava,G.P.S. (2021) AlgPred 2.0: an. improved method for predicting allergenic proteins and mapping of IgE epitopes. *Brief. Bioinform.*, **22**, bbaa294.
9. Huby,R.D., Dearman,R.J. and Kimber,I. (2000) Why are some proteins allergens? *Toxicol. Sci.*, **55**, 235–246.
10. Krutz,N.L., Winget,J., Ryan,C.A., Wimalasena,R., Maurer-Stroh,S., Dearman,R.J., Kimber,I. and Gerberick,G.F. (2019) Proteomic and bioinformatic analyses for the identification of proteins with low allergenic potential for hazard assessment. *Toxicol. Sci.*, **170**, 210–222.
11. Altschul,S.F., Madden,T.L., Schäffer,A.A., Zhang,J., Zhang,Z., Miller,W. and Lipman,D.J. (1997) Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res.*, **25**, 3389–3402.
12. Nguyen,M.N. and Madhusudhan,M.S. (2011) Biological insights from topology independent comparison of protein 3D structures. *Nucleic Acids Res.*, **39**, e94.
13. Nguyen,M.N., Tan,K.P. and Madhusudhan,M.S. (2011) CLICK–topology-independent comparison of biomolecular 3D structures. *Nucleic Acids Res.*, **39**, W24–W8.
14. Nguyen,M.N., Sim,A.Y.L., Wan,Y., Madhusudhan,M.S. and Verma,C. (2017) Topology independent comparison of RNA 3D structures using the CLICK algorithm. *Nucleic Acids Res.*, **45**, e5.
15. Nguyen,M.N., Verma,C.S. and Zhong,P. (2019) AppA: a web server for analysis, comparison, and visualization of contact residues and interfacial waters of antibody-antigen structures and models. *Nucleic Acids Res.*, **47**, W482–W489.
16. Asero,R., Amato,S., Alfieri,B., Folloni,S. and Mistrello,G. (2007) Rice: another potential cause of food allergy in patients sensitized to lipid transfer protein. *Int. Arch. Allergy Immunol*, **143**, 69–74.
17. Radauer,C. and Breiteneder,H. (2006) Pollen allergens are restricted to few protein families and show distinct patterns of species distribution. *J. Allergy Clin. Immunol.*, **117**, 141–147.
18. Rodríguez Del Río,P., Díaz-Perales,A., Sánchez-García,S., Escudero,C., Ibáñez,M.D., Méndez-Brea,P. and Barber,D. (2018) Profilin, a change in the paradigm. *J. Investig. Allergol. Clin. Immunol.*, **28**, 1–12.
19. Gough,S.C.L. and Simmonds,M.J. (2007) The HLA region and autoimmune disease: associations and mechanisms of action. *Curr. Genomics*, **8**, 453–465.
20. Crow,M.K. (2016) Autoimmunity: interferon α or β: which is the culprit in autoimmune disease? *Nat. Rev. Rheumatol.*, **12**, 439–440.

Downloaded from https://academic.oup.com/nar/advance-article/doi/10.1093/nar/gkac446/6595260 by guest on 02 June 2022

Supplementary

AllerCatPro 2.0: A web server for predicting protein allergenicity potential

Minh N. Nguyen[1,2], Nora L. Krutz[3], Vachiranee Limviphuvadh[1,2], Andreas L. Lopata[1,4], G. Frank Gerberick[5], and Sebastian Maurer-Stroh[1,2,6]*

[1] Bioinformatics Institute, 30 Biopolis Street, #07-01, Matrix, Singapore 138671

[2] IFCS Programme, Singapore Institute for Food and Biotechnology Innovation, Agency for Science, Technology and Research, Singapore.

[3] NV Procter & Gamble Services Company SA, Strombeek-Bever, Belgium.

[4] Molecular Allergy Research Laboratory, Australian Institute of Tropical Health and Medicine, James Cook University, Townsville, QLD, Australia.

[5] GF3 Consultancy LLC, West Chester, OH, United States.

[6] Department of Biological Sciences, National University of Singapore, 14 Science Drive 4, Singapore 117543

Email: sebastianms@bii.a-star.edu.sg

Allergenicity potential

We define “allergenicity potential” or “allergenic potential” as the potential of a protein to cause/elicit immediate-type (IgE-mediated) allergic reactions in humans (1). We are not able (yet) to accurately predict sensitizing from non-sensitizing (but potentially cross-reactive) allergens (2). Therefore, we would consider “allergenicity potential” as the ability to sensitize an individual and/or potentially cross-react in pre-sensitized individuals.

Low allergenic protein

In the absence of methods to characterize the relative allergenic potential of proteins, the default assumption judges all proteins to be potentially allergenic unless there is evidence for the contrary. Nevertheless, it seems inappropriate to designate proteins as non-allergens in the absence of robust methods to identify the inherent lack of allergenic potential. However, individual proteins with a significant relative abundance in protein sources for which there are known to be significant opportunities for human exposure, but no evidence for allergenicity, can be considered with low allergenic potential. This paradigm to allow the identification of allergenic proteins that display low sensitizing potential has been proposed by Krutz *et al.* (3).

Gluten-like Q-repeats

Gluten-like Q-repeats are identified using the method in our previous study (4). From the FARRP AllergenOnline database, “Celiac disease peptides” are downloaded in our dataset. The amino acid frequencies are calculated for every 9-mer window within the peptides and a composition fingerprint score is computed by using a log odd ratio of the frequency in the “Celiac 9-mer” windows divided by a background database frequency (UniProtKB). This log odd score is used to score all 9-mers in a query sequence and it triggers a hit as Gluten-like Q-repeat if the score for a 9-mer is within one standard deviation of the average of the FARRP “Celiac disease peptides”.



Figure S1: The accuracy (**A**) and Matthews correlation coefficient (MCC) (**B**) of PREAL, AllerHunter, AllergenFP, AllerTOPv2, AlgPred 2.0 and AllerCatPro 2.0 on the benchmark datasets of 218 allergens and 212 non-allergens with the same structure fold.

Protein sequence of interest					Predicted most similar allergen (non-autoimmune allergens)		Potential cross-reactivity of query protein (# and links)	Protein info, functionality of most similar allergen				Clinical relevance of most similar allergen		Similarity to most similar allergen			Similarity to allergen and resulting predicted evidence for allergenicity		Similarity to autoimmune allergen (% identity to closest human hit)	Similarity to low allergenic protein (% identity to closest hit)
Query#	Name	Sequence Length	Gluten allergens (# of Q-repeats)	# of 3x6-mer overlaps	Protein allergen	Species		UniProt/NCBI	Functionality (SUPFAM)	Pfam	InterPro	IgE prevalence	Allergen info	% identity, linear 80 aa window	% identity, 3D epitope	Show 3D epitope	Result	Comment		
1	European white birch	133	0	-	Bet v 2	Betula pendula	250	P25816	Profilin (actin-binding protein) (SSF55770)	PF00235	IPR005455 IPR036140 IPR027310	247	AG(5)	100.0	100.0	show in structure	strong evidence	3Depi >93%	-	-
2	Olive	134	0	-	Ole e 2	Olea europaea	250	O24169	Profilin (actin-binding protein) (SSF55770)	PF00235	IPR005455 IPR036140 IPR027310	18546	AG(4)	100.0	100.0	show in structure	strong evidence	3Depi >93%	-	-
3	Timothy	131	0	-	Phl p 12	Phleum pratense	250	P35079	Profilin (actin-binding protein) (SSF55770)	PF00235	IPR005455 IPR036140 IPR027310	20847	AG(5)	100.0	100.0	show in structure	strong evidence	3Depi >93%	-	-
4	Rice	131	0	-	Ory s 12	Oryza sativa	250	Q9FUD1	Profilin (actin-binding protein) (SSF55770)	PF00235	-	-	-	100.0	100.0	show in structure	strong evidence	3Depi >93%	-	-

A

Query: European white birch
Reference model: 767_TRI_A_12_0102_C_P49233_AFTER_LOOP
Epitope: *D29D*, *S31S*, *P44P*, *Q45Q*, *E46E*, *I50I*, *M51M*, *K52K*, *D53D*, *F54F*, *E55E*, *E56E*, *H59H*, *L60L*, *T63T*, *G64G*, *G68G*, *Y72Y*, *V74V*, *G77G*, *E78E*, *G80G*, *I83I*, *K96K*
Hover with mouse over residue to see its position number. Right-click for more options.



B

Protein sequence of interest					Predicted most similar allergen (non-autoimmune allergens)		Potential cross-reactivity of query protein (# and links)	Protein info, functionality of most similar allergen				Clinical relevance of most similar allergen		Similarity to most similar allergen			Similarity to allergen and resulting predicted evidence for allergenicity		Similarity to autoimmune allergen (% identity to closest human hit)	Similarity to low allergenic protein (% identity to closest hit)
Query#	Name	Sequence Length	Gluten allergens (# of Q-repeats)	# of 3x6-mer overlaps	Protein allergen	Species		UniProt/NCBI	Functionality (SUPFAM)	Pfam	InterPro	IgE prevalence	Allergen info	% identity, linear 80 aa window	% identity, 3D epitope	Show 3D epitope	Result	Comment		
1	Peach	131	0	-	Pru p 4	Prunus persica	250	Q8GT40	Profilin (actin-binding protein) (SSF55770)	PF00235	IPR005455 IPR036140 IPR027310	455	AG(3), food/insect allergen, anaphylaxis (Low)	100.0	100.0	show in structure	strong evidence	3Depi >93%	-	-
2	Apple	131	0	-	Mal d 4	Malus domestica	250	Q9XF42	Profilin (actin-binding protein) (SSF55770)	PF00235	IPR005455 IPR036140 IPR027310	882	AG(4), food/insect allergen, anaphylaxis (Low)	100.0	100.0	show in structure	strong evidence	3Depi >93%	-	-
3	Hazelnut	131	0	-	Cor a 2	Corylus avellana	250	Q9AXH5	Profilin (actin-binding protein) (SSF55770)	PF00235	IPR005455 IPR036140 IPR027310	227	AG(2), food/insect allergen, anaphylaxis (Low)	100.0	100.0	show in structure	strong evidence	3Depi >93%	-	-
4	Potato	131	0	-	Sola t 8	Solanum tuberosum	250	Q38M47	-	-	-	-	-	100.0	100.0	show in structure	strong evidence	3Depi >93%	-	-
5	Spinach	133	0	-	Che a 2	Chenopodium album	250	C5J3U0	Profilin (actin-binding protein) (SSF55770)	PF00235	IPR005455 IPR036140 IPR027310	151	AG(3)	88.8	100.0	show in structure	strong evidence	3Depi >93%	-	-
6	Para rubber tree	131	0	-	Hev b 8	Hevea brasiliensis	250	O65812	Profilin (actin-binding protein) (SSF55770)	PF00235	IPR005455 IPR036140 IPR027310	18708	AG(4)	100.0	100.0	show in structure	strong evidence	3Depi >93%	-	-

C

Protein sequence of interest					Predicted most similar allergen (non-autoimmune allergens)		Potential cross-reactivity of query protein (# and links)	Protein info, functionality of most similar allergen				Clinical relevance of most similar allergen		Similarity to most similar allergen			Similarity to allergen and resulting predicted evidence for allergenicity		Similarity to autoimmune allergen (% identity to closest human hit)	Similarity to low allergenic protein (% identity to closest hit)
Query#	Name	Sequence Length	Gluten allergens (# of Q-repeats)	# of 3x6-mer overlaps	Protein allergen	Species		UniProt/NCBI	Functionality (SUPFAM)	Pfam	InterPro	IgE prevalence	Allergen info	% identity, linear 80 aa window	% identity, 3D epitope	Show 3D epitope	Result	Comment		
1	profilin_Baker's_yeast	126	0	-	Ama r 2	Amaranthus retroflexus	249	C3W2Q7	Profilin (actin-binding protein) (SSF55770)	PF00235	-	12	AG(2)	38.8	44.4	show in structure	weak evidence	3Depi <=93%	28.1	-
2	profilin_Human	140	0	-	Mus a 1	Musa acuminata	7	M0S9D0	Profilin (actin-binding protein) (SSF55770)	PF00235	IPR005455 IPR036140 IPR027310	16	AG(2), food/insect allergen, anaphylaxis (Low)	42.5	56.2	show in structure	weak evidence	3Depi <=93%	100.0	-
3	profilin_Cow	140	0	1	Der f 29	Dermatophagoides farinae	3	AIO08866	-	-	-	-	-	32.5	61.5	show in structure	weak evidence	3Depi <=93%	95.0	-
4	profilin_Chicken	140	0	-	Vit v 4	Vitis vinifera	45	A5ASF9	Profilin (actin-binding protein) (SSF55770)	PF00235	IPR005455 IPR036140 IPR027310	-	AG(1)	41.2	46.7	show in structure	weak evidence	3Depi <=93%	61.4	-
5	profilin_Fungus	130	0	-	Che a 2	Chenopodium album	250	C5J3U0	Profilin (actin-binding protein) (SSF55770)	PF00235	IPR005455 IPR036140 IPR027310	151	AG(3)	45.0	53.8	show in structure	weak evidence	3Depi <=93%	-	-

D

Figure S2: Pollen profilins from European white birch (*Betula pendula*), olive (*Olea europaea*), Timothy (*Phleum pratense*), and rice (*Oryza sativa*) are classified correctly as allergenic proteins with 'strong evidence' using AllerCatPro 2.0 (**A**), 3D model of pollen profilin from *Betula pendula* based on the similarity of 3D surface epitope (**B**). AllerCatPro 2.0 classifies correctly as allergenic proteins with 'strong evidence' for all the plant profilins of peach, apple, hazelnut, potato, spinach, and Para rubber tree (**C**) as well as the allergenicity potential with 'weak evidence' prediction for low allergenic profilins from yeast, human, cow, chicken, and fungus (**D**).

Protein sequence of interest					Predicted most similar allergen (non-autoimmune allergens)		Potential cross-reactivity of query protein (# and links)	Protein info, functionality of most similar allergen				Clinical relevance of most similar allergen		Similarity to most similar allergen			Similarity to allergen and resulting predicted evidence for allergenicity		Similarity to autoimmune allergen (% identity to closest human hit)	Similarity to low allergenic protein (% identity to closest hit)
Query#	Name	Sequence Length	Gluten allergens (# of Q-repeats)	# of 3x6-mer overlaps	Protein allergen	Species		UniProt/NCBI	Functionality (SUPFAM)	Pfam	InterPro	IgE prevalence	Allergen info	% identity, linear 80 aa window	% identity, 3D epitope	Show 3D epitope	Result	Comment		
1	HLA-DRA	254	0	-	No significant hit (similar to autoimmune allergen)	-	0	-	-	-	-	-	-	-	-	-	no evidence	no hits	100.0	-
2	Alpha-Galactosidase	429	0	-	No significant hit (similar to autoimmune allergen)	-	0	-	-	-	-	-	-	-	-	-	no evidence	no hits	100.0	-
3	Nascent polypeptide-associated complex subunit alpha	215	0	-	No significant hit (similar to autoimmune allergen)	-	0	-	-	-	-	-	-	-	-	-	no evidence	no hits	100.0	-
4	Granulocyte colony-stimulating factor	204	0	-	No significant hit (similar to autoimmune allergen)	-	0	-	-	-	-	-	-	-	-	-	no evidence	no hits	100.0	-
5	B-cell CLL/lymphoma 7 protein family member B	202	0	-	No significant hit (similar to autoimmune allergen)	-	0	-	-	-	-	-	-	-	-	-	no evidence	no hits	100.0	-
6	Type I interferon	187	0	-	No significant hit (similar to autoimmune allergen)	-	0	-	-	-	-	-	-	-	-	-	no evidence	no hits	100.0	-

A

Hit	Protein Name	Species	UniProt/NCBI	SUPFAM	Pfam	InterPro	% identity	E-value
1	Hom s IFN beta	Homo sapiens	Q5VWC9	4-helical cytokines (SSF47266)	PF00143	-	100.0	4e-138
2	Hom s IFN beta	Homo sapiens	Q15943	4-helical cytokines (SSF47266)	PF00143	-	96.3	4e-131
3	Hom s IFN alpha	Homo sapiens	Q86UP4	4-helical cytokines (SSF47266)	PF00143	IPR009079 IPR000471	39.2	2e-27
4	Hom s IFN alpha	Homo sapiens	Q6DJX8	4-helical cytokines (SSF47266)	PF00143	IPR009079 IPR000471	39.2	2e-27
5	Hom s IFN alpha	Homo sapiens	P01563	4-helical cytokines (SSF47266)	PF00143	IPR009079 IPR000471	39.2	2e-27

B

Figure S3: AllerCatPro 2.0 accurately finds the proteins of HLA-DR alpha, alpha-galactosidase, nascent polypeptide-associated complex subunit alpha, granulocyte colony-stimulating factor, B-cell CLL/lymphoma 7 protein family member B, and type I interferon to be associated with autoimmune diseases (**A**). AllerCatPro 2.0 identifies all similar proteins from the dataset of known human proteins for autoimmune diseases that have significant sequence similarity to the input sequence of type I interferon (**B**).

Protein sequence of interest					Predicted most similar allergen (non-autoimmune allergens)		Potential cross-reactivity of query protein (# and links)	Protein info, functionality of most similar allergen				Clinical relevance of most similar allergen		Similarity to most similar allergen			Similarity to allergen and resulting predicted evidence for allergenicity		Similarity to autoimmune allergen (% identity to closest human hit)	Similarity to low allergenic protein (% identity to closest hit)
Query#	Name	Sequence Length	Gluten allergens (# of Q-repeats)	# of 3x6-mer overlaps	Protein allergen	Species		UniProt/NCBI	Functionality (SUPFAM)	Pfam	InterPro	IgE prevalence	Allergen info	% identity, linear 80 aa window	% identity, 3D epitope	Show 3D epitope	Result	Comment		
1	10 kDa prolamin	134	0	-	Cuc ma 5	Cucurbita maxima	4	Q39649	Bifunctional inhibitor/lipid-transfer protein/seed storage 2S albumin (SSF47699)	PF00234	-	-	-	36.2	-	-	weak evidence	Identical to low allergenic protein	-	100.0
2	Flower-specific gamma-thionin-like protein	105	0	-	Api g 7	Apium graveolens	2	QUJ17885	-	-	-	-	-	36.2	-	-	weak evidence	Identical to low allergenic protein	-	100.0
3	Ribulose bisphosphate carboxylase	475	0	-	No significant hit (E-value threshold 0.001)	-	0	-	-	-	-	-	-	-	-	-	no evidence	no hits	-	100.0
4	Late embryogenesis abundant protein 14	215	0	-	Cic a 1	Cicer arietinum	1	QHW05434	-	-	-	-	-	43.0	-	-	weak evidence	Identical to low allergenic protein	-	100.0
5	Late embryogenesis abundant protein 17	344	0	-	Cic a 1	Cicer arietinum	1	QHW05434	-	-	-	-	-	50.0	-	-	weak evidence	Identical to low allergenic protein	-	100.0
6	Late embryogenesis abundant protein 19	200	0	-	Cic a 1	Cicer arietinum	1	QHW05434	-	-	-	-	-	41.2	-	-	weak evidence	Identical to low allergenic protein	-	100.0
7	Late embryogenesis abundant protein, group 3	224	0	-	Cic a 1	Cicer arietinum	1	QHW05434	-	-	-	-	-	46.9	-	-	weak evidence	Identical to low allergenic protein	-	100.0

Figure S4: AllerCatPro 2.0 is used to predict the similarity of 10 kDa prolamin from rice (*Oryza sativa*), a flower-specific gamma-thionin-like protein from tomato (*Solanum lycopersicum*), ribulose bisphosphate carboxylase from spinach (*Spinacia oleracea*), late embryogenesis abundant proteins 14, 17, 19, and group 3 from wheat (*Triticum aestivum*) to low allergenic proteins.

Protein sequence of interest					Predicted most similar allergen (non-autoimmune allergens)		Potential cross-reactivity of query protein (# and links)	Protein info, functionality of most similar allergen				Clinical relevance of most similar allergen		Similarity to most similar allergen			Similarity to allergen and resulting predicted evidence for allergenicity		Similarity to autoimmune allergen (% identity to closest human hit)	Similarity to low allergenic protein (% identity to closest hit)
Query#	Name	Sequence Length	Gluten allergens (# of Q-repeats)	# of 3x6-mer overlaps	Protein allergen	Species		UniProt/NCBI	Functionality (SUPFAM)	Pfam	InterPro	IgE prevalence	Allergen info	% identity, linear 80 aa window	% identity, 3D epitope	Show 3D epitope	Result	Comment		
1	A0A5F4CH33	2566	0	-	Gal d Ovomucin	Gallus gallus	13	Q98UI9	Serine protease inhibitors (SSF57567)	PF08742 PF01826 PF00094	IPR006207 IPR036084 IPR002919 IPR014853 IPR001007 IPR001846	-	AG(1)	45.0	-	-	strong evidence	win80lin >35%	-	-
2	A0A5F4DEY2	1073	0	-	Cuc p AscO	Cucurbita pepo	1	P37064	Cupredoxins (SSF49503)	PF00394 PF07731 PF07732	IPR008972 IPR001117 IPR011706 IPR011707 IPR033138 IPR002355 IPR034259 IPR034258 IPR034267 IPR017760	128	AG(2)	33.3	33.3	show in structure	weak evidence	3Depi <=93%	32.8	-
3	F1PHH5	3967	0	-	Gal d Ovomucin	Gallus gallus	5	Q98UI9	Serine protease inhibitors (SSF57567)	PF08742 PF01826 PF00094	IPR006207 IPR036084 IPR002919 IPR014853 IPR001007 IPR001846	-	AG(1)	66.2	-	-	strong evidence	win80lin >35%	-	-
4	A0A5F4C6T4	2726	0	-	Gal d Ovomucin	Gallus gallus	5	Q98UI9	Serine protease inhibitors (SSF57567)	PF08742 PF01826 PF00094	IPR006207 IPR036084 IPR002919 IPR014853 IPR001007 IPR001846	-	AG(1)	68.8	-	-	strong evidence	win80lin >35%	-	-
5	A0A1Y3BCS5	925	0	-	Ves v 6	Vespula vulgaris	9	G8IIT0	Lipovitellin-phosvitin complex, superhelical domain (SSF49431); Lipovitellin-phosvitin complex; beta-sheet shell regions (SSF56968)	PF09172 PF01347 PF00094	-	28	AG(2)	30.0	50.0	show in structure	weak evidence	3Depi <=93%	-	-
6	A0A1Y3BKH5	1658	0	-	Sch j Sj67	Schistosoma japonicum	2	Q5DAD7	Second domain of FERM (SSF47031); Moesin tail domain (SSF48678); Ubiquitin-like (SSF54236)	PF00769 PF09380 PF00373 PF09379	-	553	AG(2)	33.8	46.2	show in structure	weak evidence	3Depi <=93%	-	-
7	A0A2U1KS88	834	8	-	Str pn PspC	Streptococcus pneumoniae	9	B5AZD6	-	PF01473 PF19127 PF05062 PF04650	IPR018337 IPR007756 IPR038183 IPR005877	-	AG(1)	36.2	-	-	strong evidence	Gluten-like Q-repeats and win80lin >35%	-	-
8	A0A251VFN7	1220	0	-	Bla g RACK1	Blattella germanica	2	Q09KA1	WD40 repeat-like (SSF50978)	PF00400	IPR020472 IPR015943 IPR001680 IPR019775 IPR036322	-	AG(1)	42.5	41.2	show in structure	weak evidence	3Depi <=93%	-	-
9	A0A251V053	1530	0	-	No significant hit (E-value threshold 0.001)	-	0	-	-	-	-	-	-	-	-	-	no evidence	no hits	-	-
10	A0A2S3H4S4	975	0	-	Sch c 1 homolog	Arthroderma benhamiae	2	P0DN29	Six-hairpin glycosidases (SSF48208); Starch-binding domain-like (SSF49452)	PF00686 PF00723	-	-	-	32.5	61.5	show in structure	weak evidence	3Depi <=93%	-	-
11	A0A5N6RRW0	925	0	-	No significant hit	-	2	-	-	-	-	-	-	-	-	-	no evidence	no hits	-	-
12	A0A5N6RRQ1	971	0	-	Neu cr Glucoamylase	Neurospora crassa	3	P14804	Six-hairpin glycosidases (SSF48208); Starch-binding domain-like (SSF49452)	PF00686 PF00723	IPR008928 IPR012341 IPR013784 IPR034836 IPR002044 IPR011613 IPR000165 IPR008291 IPR013783	-	AG(1)	33.8	46.7	show in structure	weak evidence	3Depi <=93%	-	-
13	A0A5N6QXB7	1079	0	-	No significant hit	-	6	-	-	-	-	-	-	-	-	-	no evidence	no hits	-	-
14	B9RHB0	1217	0	-	Bla g RACK1	Blattella germanica	2	Q09KA1	WD40 repeat-like (SSF50978)	PF00400	IPR020472 IPR015943 IPR001680 IPR019775 IPR036322	-	AG(1)	40.0	46.2	show in structure	weak evidence	3Depi <=93%	-	-

A

Query: A0A5F4DEY2
Reference model: 9868_HOM_S_FACTR_8_C_1577_2351_AFTER_LOOP
Epitope: *F399F, P400P, H419H, H421H, H427H*
Hover with mouse over residue to see its position number. Right-click for more options.



B

Figure S5: AllerCatPro 2.0 predicts allergenicity potential for a set of very large proteins with their lengths varying from 834 to 3967 residues (**A**). AllerCatPro 2.0 displays the 3D model for the input protein of A0A5F4DEY2 based on the similarity of 3D surface epitope with blue and grey beads showing positions of identical positively charged and the other identical residues of 3D epitope, respectively (**B**).

A



Protein sequence of interest					Predicted most similar allergen (non-autoimmune allergens)		Potential cross-reactivity of query protein (# and links)	Protein info, functionality of most similar allergen				Clinical relevance of most similar allergen		Similarity to most similar allergen			Similarity to allergen and resulting predicted evidence for allergenicity		Similarity to autoimmune allergen (% identity to closest human hit)	Similarity to low allergenic protein (% identity to closest hit)
Query#	Name	Sequence Length	Gluten allergens (# of Q-repeats)	# of 3x6-mer overlaps	Protein allergen	Species		UniProt/NCBI	Functionality (SUPFAM)	Pfam	InterPro	IgE prevalence	Allergen info	% identity, linear 80 aa window	% identity, 3D epitope	Show 3D epitope	Result	Comment		
1	ABD51778	93	0	-	Api m 6	Apis mellifera	8	Q27SJ8	Serine protease inhibitors (SSF57567)	PF01826	IPR036084 IPR002919	43	AG(2)	100.0	-	-	strong evidence	win80lin >35%	-	-
2	CAB02154	161	0	-	Bet v 1	Betula pendula	250	P15494	-	PF00407	IPR000916 IPR024949 IPR023393	334	AG(9)	100.0	100.0	show in structure	strong evidence	3Depi >93%	-	-
3	AAA34278	320	122	-	Prolamin	Triticum aestivum	175	P04725	Bifunctional inhibitor/lipid-transfer protein/seed storage 2S albumin (SSF47699)	PF00234	-	-	-	100.0	-	-	strong evidence	Gluten-like Q-repeats and win80lin >35%	-	-

B

Figure S6: AllerCatPro 2.0 is applied to analyse and evaluate allergenicity potential of three known allergens (ABD51778, CAB02154, AAA34278) from *Apis mellifera*, *Betula pendula*, and *Triticum aestivum* using their nucleotide sequences as input (**A**). AllerCatPro 2.0 classifies correctly as allergens with 'strong evidence' for all these nucleotide input sequences (**B**).

References

1. Blackburn,K., N'jai,A.U., Dearman,R.J., Kimber,I. and Gerberick,G.F. (2015) Respiratory allergenic potential of plant-derived proteins: Understanding the relationship between exposure and potency for risk assessments. *Crit Rev Toxicol*, **45**, 799–811.
https://doi.org/10.3109/10408444.2015.1067876
http://www.ncbi.nlm.nih.gov/pubmed/26565768
2. Krutz,N.L., Kimber,I., Maurer-Stroh,S. and Gerberick,G.F. (2020) Determination of the relative allergenic potency of proteins: hurdles and opportunities. *Crit Rev Toxicol*, **50**, 521–530.
https://doi.org/10.1080/10408444.2020.1793895
http://www.ncbi.nlm.nih.gov/pubmed/32729356
3. Krutz,N.L., Winget,J., Ryan,C.A., Wimalasena,R., Maurer-Stroh,S., Dearman,R.J., Kimber,I. and Gerberick,G.F. (2019) Proteomic and Bioinformatic Analyses for the Identification of Proteins With Low Allergenic Potential for Hazard Assessment. *Toxicol Sci*, **170**, 210–222.
https://doi.org/10.1093/toxsci/kfz078
http://www.ncbi.nlm.nih.gov/pubmed/30903174
4. Maurer-Stroh,S., Krutz,N.L., Kern,P.S., Gunalan,V., Nguyen,M.N., Limviphuvadh,V., Eisenhaber,F. and Gerberick,G.F. (2019) AllerCatPro-prediction of protein allergenicity potential from the protein sequence. *Bioinformatics*, **35**, 3020–3027.
https://doi.org/10.1093/bioinformatics/btz029
http://www.ncbi.nlm.nih.gov/pubmed/30657872