

Primary immunodeficiencies: A rapidly evolving story

Nima Parvaneh, MD,^{a,b} Jean-Laurent Casanova, MD, PhD,^{c,d} Luigi Daniele Notarangelo, MD, PhD,^{e,f} and Mary Ellen Conley, MD^{g,h} *Tehran, Iran, New York, NY, Paris, France, Boston, Mass, and Memphis, Tenn*

The characterization of primary immunodeficiencies (PIDs) in human subjects is crucial for a better understanding of the biology of the immune response. New achievements in this field have been possible in light of collaborative studies; attention paid to new phenotypes, infectious and otherwise; improved immunologic techniques; and use of exome sequencing technology. The International Union of Immunological Societies Expert Committee on PIDs recently reported on the updated classification of PIDs. However, new PIDs are being discovered at an ever-increasing rate. A series of 19 novel primary defects of immunity that have been discovered after release of the International Union of Immunological Societies report are discussed here. These new findings highlight the molecular pathways that are associated with clinical phenotypes and suggest potential therapies for affected patients. (*J Allergy Clin Immunol* 2013;131:314-23.)

Key words: Primary immunodeficiencies, combined immunodeficiencies, well-defined syndromes with immunodeficiency, predominantly antibody defects, defects of immune dysregulation, congenital defects of phagocytes, defects in innate immunity, autoinflammatory disorders, mutation detection

Discuss this article on the JACI Journal Club blog: www.jaci-online.blogspot.com.

Known primary immunodeficiencies (PIDs) include individually rare but collectively diverse genetic defects that influence

Abbreviations used

ADAM17:	A disintegrin and metalloproteinase 17
BCR:	B-cell receptor
EV:	Epidermodysplasia verruciformis
HHV6:	Human herpesvirus 6
HPS:	Hermansky-Pudlak syndrome
HPV:	Human papilloma virus
HSE:	Herpes simplex virus encephalitis
HSV1:	Herpes simplex virus 1
ICA:	Isolated congenital asplenia
ICL:	Idiopathic CD4 ⁺ lymphopenia
IL-36Ra:	IL-36 receptor antagonist
ISG15:	Interferon-stimulated gene 15
IUIS:	International Union of Immunological Societies
LCK:	Lymphocyte-specific protein tyrosine kinase
LRBA:	LPS-responsive beige-like anchor
MCM4:	Minichromosome maintenance complex component 4
MSMD:	Mendelian susceptibility to mycobacterial disease
MST:	Mammalian Ste20-like kinase
NEMO:	NF- κ B essential modulator
NF- κ B:	Nuclear factor κ B
NK:	Natural killer
PI3K:	Phosphoinositide 3-kinase
PID:	Primary immunodeficiency
PLC γ 2:	Phospholipase C γ 2
RHOH:	Ras homolog gene family member H
TBK1:	TANK-binding kinase 1
TCR:	T-cell receptor
TLR3:	Toll-like receptor 3
TRIF:	Toll/IL-1 receptor domain-containing adaptor inducing IFN- β
UNC119:	Uncoordinated 119
WASP:	Wiskott-Aldrich syndrome protein
WIP:	WASP-interacting protein

From ^athe Pediatric Infectious Diseases Research Center and ^bthe Research Center for Immunodeficiencies, Children's Medical Center, Tehran University of Medical Sciences, Tehran; ^cSt Giles Laboratory of Human Genetics of Infectious Diseases, Rockefeller Branch, Rockefeller University, New York; ^dthe Laboratory of Human Genetics of Infectious Diseases, Necker Branch, Necker Medical School, University Paris Descartes and INSERM U980, Paris; ^ethe Division of Immunology and ^fthe Manton Center for Orphan Disease Research, Children's Hospital Boston, Harvard Medical School, Boston; ^gthe Department of Pediatrics, University of Tennessee College of Medicine, Memphis; and ^hLe Bonheur Children's Hospital, Memphis.

Disclosure of potential conflict of interest: J.-L. Casanova has received grants from the National Institutes of Health (NIH); has been a consultant for Pfizer, GlaxoSmith-Kline, NovImmune, BiogenIdec, Merck, and Sanofi-Aventis; and has grants/grants pending from Pfizer and Merck. L. D. Notarangelo is a board member for the Immune Disease Institute, is employed by Children's Hospital Boston, has grants/grants pending with the NIH, the March of Dimes, and the Jeffrey Modell Foundation; and has received payment for lectures, including service on speakers' bureaus for the WAS foundation. M. E. Conley is employed by the University of Tennessee, has received grants/grants pending from the NIH and the March of Dimes, has received payment for lectures from the Immune Deficiency Foundation, and is receiving royalties from the Southern Bio for monoclonal antibodies. N. Parvaneh declares no relevant conflicts of interest.

Received for publication October 2, 2012; revised November 6, 2012; accepted for publication November 29, 2012.

Corresponding author: Nima Parvaneh, MD, Pediatric Infectious Diseases Research Center, Children's Medical Center, 62 Gharib St, 14194 Tehran, Iran. E-mail: nparvaneh@tums.ac.ir.

0091-6749/\$36.00

© 2013 American Academy of Allergy, Asthma & Immunology
<http://dx.doi.org/10.1016/j.jaci.2012.11.051>

the development, function, or both of immunity. Taken in a broad sense, these disorders encompass both the hematopoietic and nonhematopoietic arms of host defense. They result in a wide range of clinical symptoms, including but not limited to susceptibility to infections, autoimmunity, inflammation, allergy, and malignancy.¹ Significant progress in the field has recently accelerated thanks to collaborative work around the world; improved techniques to analyze leukocyte subsets; attention paid to new phenotypes, such as infections selectively caused by certain pathogens in otherwise healthy children²; and use of high-throughput next-generation sequencing technology (eg, whole-exome sequencing).³ The International Union of Immunological Societies (IUIS) Expert Committee on PIDs published the biennial update of the classification of PIDs in a recent article dated November 2011.⁴ However, the continuously growing number and variety of PIDs make this type of classification difficult and perhaps calls for a revised approach.^{5,6} In this review we focus on the 19 new PIDs that appeared in the literature after the release of the last IUIS report. Some of these reports include

only a single case, making it difficult to confidently predict the clinical phenotype; however, each new disorder provides valuable insights into the pathways that normally ensure host defense from environmental challenges while not harming the host. We classify these new PIDs according to the latest IUIS classification (Table I).

COMBINED IMMUNODEFICIENCIES

T-cell receptor α gene mutation: T-cell receptor $\alpha\beta^+$ T-cell depletion

T cells comprise 2 distinct lineages that express either $\alpha\beta$ or $\gamma\delta$ T-cell receptor (TCR) complexes that perform different tasks in immune responses. During T-cell maturation, the precise order and efficacy of TCR gene rearrangements determine the fate of the cells.⁷ Productive β -chain gene rearrangement produces a pre-TCR on the cell surface in association with pre-T α invariant peptide (β -selection). Pre-TCR signals promote α -chain recombination and transition to a double-positive stage ($CD4^+CD8^+$).⁸ This is the prerequisite for central tolerance achieved through positive and negative selection of thymocytes. Additional insight into the development of TCR $\alpha\beta^+$ T cells was provided by the characterization of 2 patients from 2 unrelated Pakistani families who had the same homozygous mutation in the TCR α constant (*TRAC*) gene and shared increased susceptibility to infections, autoimmunity, and profound T-cell proliferative impairments but apparently had normal antibody responses.⁹ One of the cases showed predisposition to herpes virus infections (including protracted varicella zoster and chronic EBV/HHV6 viremia) and chronic lung disease. In addition to immunodeficiency, both children had evidence of immune dysregulation with a combination of eosinophilia, vitiligo, alopecia areata, autoimmune hemolytic anemia, eczema, and the presence of autoantibodies. T cells from affected subjects were devoid of surface expression of the TCR $\alpha\beta$ complex; all existing $CD3^+$ T cells expressed TCR $\gamma\delta$. Both patients were successfully treated with matched sibling bone marrow transplants. This study is of interest because it is the first pure TCR $\alpha\beta^+$ T-cell immunodeficiency, allowing a fine definition of the role of this major cell lineage. Previously described disorders affected all T-cell lineages alone or in combination with B-cell or natural killer (NK) cell deficits. This experiment confirms that $\gamma\delta$ T cells alone cannot ensure adequate host defense. It would be interesting to identify patients with a selective $\gamma\delta$ T-cell deficiency because the exact role of these T cells remains unknown.

Ras homolog gene family member H deficiency: Loss of naive T cells and persistent human papilloma virus infections

Human papilloma viruses (HPVs) are double-stranded DNA viruses with a tropism for keratinocytes, causing chronic epithelial lesions that can progress to cancer.¹⁰ Susceptibility to HPV infection can be genetically determined, as seen in epidermodysplasia verruciformis (EV). EV is characterized by early development of widespread, refractory flat warts and pityriasis-like lesions and occasional development of skin carcinomas caused by certain types of HPV, called EV-HPV (or β -HPV types), in otherwise healthy subjects.¹¹ Mutations in *EVER1* or *EVER2* have been identified in most but not all patients with EV.¹² To date, the exact mechanisms leading to persistent EV-HPV infections in EVER-deficient patients are unknown;

there is a clear keratinocyte phenotype¹³ but only mild T-cell anomalies in EVER-deficient patients.¹⁴ A recent study explored the genetic basis of an EV-like phenotype in 2 French siblings with persistent cutaneous EV-HPV infections and other clinical manifestations, including bronchopulmonary disease and Burkitt lymphoma in one of them, indicating that the phenotypic spectrum of the disease is not restricted to susceptibility to HPV.¹⁵ Therefore this condition is related to but distinct from EV. The patients were homozygous for a nonsense allele of the Ras homolog gene family member H (*RHOH*), which encodes an atypical Rho GTPase (RHOH) expressed predominantly in hematopoietic cells. RHOH is crucial for pre-TCR and TCR signaling and has a role during β -selection and positive selection in the thymus.^{16,17} The patients displayed a lack of circulating naive T cells, a lower than normal proportion of skin-homing $\beta7^+$ T cells, and impaired TCR signaling. The combination of these T-cell defects might explain the pathogenesis of susceptibility to cutaneous EV-HPVs. EV-like features were also recently reported in a child with mammalian Ste20-like kinase (MST) 1 deficiency (described below).¹⁸

MST1 deficiency: Loss of naive T cells

New insight into the role of MST1 as a critical regulator of T-cell homing and function was provided by the characterization of 8 patients from 4 unrelated families who had homozygous nonsense mutations in *STK4*, the gene encoding MST1.^{18,20} MST1 was originally identified as an ubiquitously expressed kinase with structural homology to yeast Ste20.²¹ MST1 is the mammalian homolog of the *Drosophila* Hippo protein, controlling cell growth, apoptosis, and tumorigenesis.²² It has both proapoptotic and antiapoptotic functions.¹⁹ Clinically, the patients had recurrent bacterial, viral, and candidal infections; lymphopenia; intermittent neutropenia; EBV-driven lymphoproliferation; lymphoma; autoimmune cytopenias; and subtle cardiac anomalies. Of note is the development of HPV (both EV-HPV and non-EV-HPV) infections in 3 of the patients.^{18,19} MST1-deficient patients demonstrated hypergammaglobulinemia and variable humoral responses. However, B-cell numbers (especially memory B-cell numbers) were significantly reduced in one report.¹⁹ Peripheral T cells displayed markedly impaired survival/proliferation to mitogens and antigens, a response that worsened with time.²⁰ Moreover, the T-cell compartment showed a restricted TCR repertoire and a severe reduction in circulating naive ($CD45RA^+$) cell numbers. Together with RHOH deficiency, MST1 deficiency seems to affect naive T-cell development and homing, predisposing to various infections in affected subjects.

Lymphocyte-specific protein tyrosine kinase deficiency: T-cell deficiency with $CD4^+$ lymphopenia

Defects in pre-TCR- and TCR-mediated signaling lead to aberrant T-cell development and function (Fig 1). One of the earliest biochemical events occurring after engagement of the (pre)-TCR is the activation of lymphocyte-specific protein tyrosine kinase (LCK), a member of the SRC family of protein tyrosine kinases.^{23,24} This kinase then phosphorylates immunoreceptor tyrosine-based activation motifs of intracellular domains of CD3 subunits. Phosphorylated immunoreceptor tyrosine-based activation motifs recruit ζ -chain associated protein kinase of 70 kDa, which, after being phosphorylated by LCK, is

TABLE I. Novel PID genes and their phenotypes

Gene	Protein	Inheritance	Phenotype
Combined immunodeficiencies			
<i>TRAC</i>	TCR α	AR	TCR $\alpha\beta^+$ T-cell deficiency, viral infections, autoimmunity
<i>RHOH</i>	RHOH	AR	Loss of naive T cells, HPV infection
<i>STK4</i>	MST1	AR	Loss of naive T cells, EBV infection, HPV infection, autoimmunity
<i>LCK</i>	LCK	AR	T-cell deficiency, CD4 $^+$ lymphopenia
<i>UNC119</i>	UNC119	AD, dominant negative	ICL
Well-defined syndromes with immunodeficiency			
<i>WIPF1</i>	WIP	AR	Wiskott-Aldrich syndrome-like
<i>PLCG2</i>	Phospholipase C γ 2	AD, dominant negative	Cold urticaria, humeral deficiency, autoimmunity, atopy (S707Y)
		AD, hypermorphic	Autoinflammatory syndrome (deletions)
Predominantly antibody defects			
<i>PIK3R1</i>	p85 α subunit of PI3K	AR	Agammaglobulinemia, absent B cells
<i>CD21</i>	CD21	AR	Hypogammaglobulinemia
<i>LRBA</i>	LRBA	AR	Hypogammaglobulinemia, autoimmunity, colitis
Defects of immune dysregulation			
<i>PLDN</i>	Pallidin	AR	HPS type 9, albinism, immunodeficiency
<i>CD27</i>	CD27	AR	EBV-associated lymphoproliferation, hypogammaglobulinemia
Congenital defects of phagocyte number, function, or both			
<i>ISG15</i>	ISG15	AR	MSMD
Defects in innate immunity			
<i>NKX2-5</i>	NKX2-5	AD, dominant negative	ICA
<i>TRIF</i>	TRIF	AR	Herpes simplex encephalitis
<i>TBK1</i>	TBK1	AD, dominant negative (G159A), haploinsufficiency (D50A)	Herpes simplex encephalitis
<i>MCM4</i>	MCM4	AR	NK cell deficiency, infection with herpesviruses, growth retardation, and adrenal insufficiency
Autoinflammatory disorders			
<i>ADAM17</i>	ADAM17	AR	Inflammatory skin and bowel disease, high IL-1 and IL-6 production
<i>IL36RN</i>	IL-36Ra	AR	Generalized pustular psoriasis

responsible for activation of critical downstream events. Major consequences include activation of the membrane-associated enzyme phospholipase C γ 1, activation of the mitogen-activated protein kinase, nuclear translocation of nuclear factor κ B (NF- κ B), and Ca $^{2+}$ /Mg $^{2+}$ mobilization. Through these pathways, LCK controls T-cell development and activation.²⁵ In mice lacking LCK, T-cell development in the thymus is profoundly blocked at an early double-negative stage.²⁶ Although 3 cases of combined immunodeficiencies with altered LCK protein expression had been reported in human subjects, the molecular defects at the genomic level were not documented.²⁷⁻²⁹ Recently, autosomal recessive LCK deficiency was described as the cause of profound T-cell immunodeficiency and immune dysregulation.³⁰ A French infant presented with early-onset protracted diarrhea, recurrent respiratory tract infections, failure to thrive, autoimmune thrombocytopenia, and skin/mucosal inflammatory disorders. Laboratory studies in the affected patient showed CD4 $^+$ T-cell lymphopenia and low Treg cell numbers. The residual T lymphocytes had an oligoclonal T-cell repertoire and exhibited a severe TCR signaling defect, with only weak tyrosine phosphorylation signals and no Ca $^{2+}$ mobilization after TCR stimulation. Moreover, the patient's T cells were resistant to activation-induced cell death. She was found to have a homozygous missense mutation of the *LCK* gene (c.1022T>C; L341P) resulting from maternal uniparental disomy. Anti-TNF therapy was partially effective in treating the serositis and skin inflammation. She underwent matched unrelated hematopoietic stem cell transplantation at 30 months, but unfortunately, she died of post-transplantation veno-occlusive disease. This study highlights the

importance of LCK for the development of a normal T-cell repertoire and also maintenance of central and peripheral T-cell tolerance.

Uncoordinated 119 deficiency: Idiopathic CD4 $^+$ lymphopenia

Idiopathic CD4 $^+$ lymphopenia (ICL) is a very heterogeneous clinical entity that is defined, by default, by persistent CD4 $^+$ T-cell lymphopenia (<300 cells/ μ L or <20% of total T cells) in the absence of HIV infection or any other known cause of immunodeficiency.³¹ The few studies examining the pathogenesis of ICL suggest that in some patients accelerated apoptosis and diminished proliferative capacity are partly due to disturbed TCR signaling.^{32,33} Uncoordinated 119 (UNC119), an activator of LCK, delivers LCK to the plasma membrane through an endosomal route and initiates the enzymatic activity of LCK on TCR stimulation.^{34,35} Through LCK, UNC119 regulates some T-cell functions, including immunologic synapse formation, proliferation, and differentiation into effector cells.³⁴⁻³⁷ Recently, a heterozygous dominant-negative missense mutation (V22G) of UNC119 was reported in a patient with ICL.³⁸ The patient was a 32-year-old woman with a history of recurrent respiratory tract infections, persistent fungal infections of the skin and nails, recurrent shingles, and oral herpes simplex lesions. The patient's cells showed reduced response to TCR stimulation, with impairment in LCK localization/activation resulting in decreased cell proliferation. Transduction of the mutant UNC119 but not wild-type UNC119 into normal T cells reproduced the signaling and




FIG 1. TCR signaling. Multiple signal transduction pathways are stimulated through the TCR. These pathways collectively activate transcription factors that organize T-cell survival, proliferation, differentiation, homeostasis, and migration. Mutant molecules in patients with TCR-related defects are indicated in red. AP1, Activator protein 1; BCL10, B-cell lymphoma/leukemia 10; CARMA, CARD-containing MAGUK protein; DAG, diacylglycerol; ER, endoplasmic reticulum; FOXO, forkhead box O; GAD, GRB2-related adaptor protein; IKK, IκB kinase; IP3, inositol trisphosphate; IP3R, inositol trisphosphate receptor; LAT, linker of activated T cells; MALT, mucosa-associated lymphoid tissue lymphoma translocation protein; MAPK, mitogen-activated protein kinase; mTOR, mammalian target of rapamycin; NCK, noncatalytic region of tyrosine kinase adaptor protein; NFAT, nuclear factor of activated T cells; PDK, phosphoinositide-dependent kinase; PIP2, phosphatidylinositol bisphosphate; PKB, protein kinase B; PKCθ, protein kinase C θ; RASGRP, RAS guanyl nucleotide-releasing protein; SLP-76, SH2 domain-containing leukocyte protein of 76 kDa; STIM1, stromal interaction molecule 1; ZAP70, ζ-chain associated protein kinase of 70 kDa.

proliferation defects, confirming the inhibitory function of G22V mutation. These findings shed light on the molecular mechanisms for a subset of patients with ICL.

WELL-DEFINED SYNDROMES WITH IMMUNODEFICIENCY

Wiskott-Aldrich syndrome protein-interacting protein deficiency: Wiskott-Aldrich syndrome-like phenotype

In hematopoietic cells Wiskott-Aldrich syndrome protein (WASP) is stabilized through forming a complex with WASP-interacting protein (WIP).³⁹ A female Moroccan infant presented in early infancy with recurrent infections, eczema, thrombocytopenia, T-cell lymphopenia, and decreased NK cell function reminiscent of WAS.⁴⁰ Despite normal WASP sequence and mRNA expression levels, WASP was not detected in the patient's cells, suggesting WASP protein degradation caused by the absence of WIP. Further studies showed that WIP could not be detected in the patient's T-cell blasts, and a homozygous nonsense mutation (S434X) was detected in *WIPF1*, which encodes WIP. Despite clinical similarities with patients with WAS, the WIP-deficient

patient displayed some immune abnormalities that have not been documented in patients with WAS. They include impaired T-cell response to IL-2, complete failure to proliferate in response to TCR ligation with anti-CD3, and complete abrogation of T-cell chemotaxis.⁴¹⁻⁴³ Moreover, the mean platelet volume was normal in the patient, as seen in WIP-deficient mice.⁴⁴ Unrelated cord blood transplantation was performed at 4.5 months, and the child is now doing well at age 21 months. This study is interesting because it reports the first autosomal recessive form of WAS, one of the most emblematic X-linked recessive PIDs and arguably one of the first PIDs described, first by Wiskott in 1937⁴⁵ and then again by Aldrich in 1954.⁴⁶

Phospholipase Cγ2 gain-of-function mutations: Cold urticaria, immunodeficiency, and autoimmunity/autoinflammatory

This is a unique phenotype, sharing features of antibody deficiency, autoinflammatory diseases, and immune dysregulatory disorders, making its classification difficult. Two recent studies validated the pleiotropy of genetic alterations in the same gene.^{47,48} In the first study genomic deletions in *PLCG2* were

detected in 3 distinct families with a dominantly inherited syndrome of cold urticaria and variable immune dysregulation.⁴⁷ *PLCG2* encodes phospholipase C γ 2 (PLC γ 2), a signaling molecule expressed in B cells, NK cells, mast cells, and platelets. PLC γ transmits information from activated receptors to downstream signal cascades by triggering receptor-mediated Ca²⁺ entry.^{49,50} Cold urticaria occurred in all 27 affected subjects. Interestingly, skin test results were positive for evaporative cooling but negative on ice-cube and cold-water immersion. Other variable manifestations included atopy, sinopulmonary infections, autoimmune diseases, and granulomatous rashes in order of frequency. Laboratory findings were also variable and included low serum IgM and IgA levels in some patients, diminished numbers of circulating CD19⁺ B cells and switched memory B cells, and low NK cell numbers, whereas IgE levels were increased in most of the subjects. The C-terminal SH2 deletions in PLC γ 2, which were seen in all 3 families, resulted in the failure of autoinhibition and caused constitutive phospholipase activity when transfected into COS7 cells.⁴⁷ Paradoxically, B cells and NK cells clearly had decreased PLC γ 2-dependent signaling and function; the exact mechanisms are not yet clear. Affected mast cells showed increased mast cell degranulation at cold temperatures without receptor stimulation, indicating that the altered function of mutant PLC γ 2 was the cause of cold urticaria. Another study by the same research group defined a hypermorphic dominant mutation within *PLCG2* (p.Ser707Tyr) as the cause of a novel autoinflammatory syndrome in 2 affected members of a family.⁴⁸ A father and his daughter both had epidermolysis bullosa-like eruptions in infancy, interstitial pneumonitis, arthralgia, eye inflammation, enterocolitis, cellulitis, and recurrent sinopulmonary infections. Immunologic workup showed low circulating IgM/IgA levels and absence of class-switched memory B cells. The inflammatory manifestations were refractory to treatment with non-steroidal anti-inflammatory drugs and TNF inhibitors but responsive to high-dose corticosteroids and, to a lesser extent, to IL-1 blockade. B cells showed increased PLC γ 2-dependent signaling after receptor cross-linking, as indicated by increased levels of inositol trisphosphate production, intracellular Ca²⁺ release, and extracellular signal-regulated kinase phosphorylation.

In conclusion, a central role for PLC γ in controlling Ca²⁺ signaling and thus a plethora of cellular responses could explain the various clinical features, with the involvement of the humoral immune system and various forms of inflammation as the common theme. Allelic heterogeneity at the *PLCG2* locus apparently further contributes to the diversity of cellular and clinical phenotypes.

PREDOMINANTLY ANTIBODY DEFECTS

Defect in the p85 α subunit of phosphoinositide

3-kinase: Agammaglobulinemia and absent B cells

Early B-cell development is under lineage-specific genetic control. Genetic defects of Bruton tyrosine kinase and components of the pre-B-cell receptor (BCR), such as the μ heavy chain, λ 5, Ig α , Ig β , and the downstream scaffold protein B-cell linker, have been described in more than 90% of patients with isolated defects in B-cell development and agammaglobulinemia.⁵¹

A report has recently described a homozygous premature stop codon in *PIK3R1*, which encodes 3 regulatory subunits of phosphoinositide 3-kinase (PI3K) by use of alternative splicing.⁵² The patient, a young woman with agammaglobulinemia and absent B cells, had early onset of infections and multiple

complications, including colitis, as a teenager.⁵² PI3Ks are a broadly expressed group of enzymes that respond to extracellular signals to activate a variety of cellular functions.⁵³ Although the mutation did not affect the expression of p50 α and p55 α , the other regulatory subunits encoded by *PIK3R1*, it did result in a marked decrease in levels of p110 δ , the catalytic subunit of PI3K. The developmental arrest in the patient was at the pro-B-cell stage, earlier than that seen in patients with defects in the BCR signaling pathway.⁵¹ The extracellular signal that requires activation of the PI3K pathway at this early stage of development is unknown; however, it was noted that the chemokine CXCR4 transduces through PI3K⁵⁴ and that defects in CXCR4 in mice cause a similar block in B-cell development.⁵⁵ The patient had neutropenia at the time of diagnosis that was not seen during follow-up evaluations. Patients with early defects of B-cell development might present with neutropenia before they are started on gammaglobulin replacement.⁵⁶⁻⁵⁸ Mice with p85 α deficiency show a wide range of defects, including a B-cell defect similar to that seen in mice with mutations in *Btk*, hypersensitivity to insulin, defective platelet function, and abnormal mast cell development.^{59,60} In contrast, in human subjects the absence of p85 α results in a severe isolated defect in development of early stages of B lineage cells without associated findings.

CD21 deficiency: Hypogammaglobulinemia

CD21, the complement receptor type 2, is expressed on mature B cells and follicular dendritic cells.⁶¹ In addition to binding the complement component C3d, CD21 also binds interferon α and CD23⁶² and serves as the main cellular entry receptor for EBV.^{63,64} CD21 forms a part of the CD19 complex, decreasing the threshold for antigen stimulation of B cells through the BCR.⁶⁵ A 28-year-old man presented with recurrent infections, splenomegaly, hypogammaglobulinemia, and reduced class-switched memory (IgD⁻CD27⁺) B-cell numbers but a normal antibody response to vaccine antigens.⁶⁶ Compound heterozygous mutations in the *CD21* gene prohibited CD21 receptor expression. Functional studies showed a complete loss of costimulatory activity of C3d in enhancing B-cell receptor stimulation. Vaccination response to pneumococcal polysaccharide vaccination was moderately impaired, but the responses to protein antigens were preserved. The patient's B cells were unable to bind to the EBVgp350 antigen; however, the patient underwent EBV seroconversion *in vivo*, and his B cells were easily immortalized by EBV *in vitro*. This indicates that CD21 is not required for EBV infection *in vitro* and *in vivo*. The patient's clinical condition improved on prophylactic antibiotics and intermittent gammaglobulin substitution. CD21 deficiency shares many features with CD19 and CD81 deficiency, although with less clinical severity.^{67,68} Preservation of specific antibody responses and normal responses to direct BCR stimulation in patients with CD21 deficiency probably contribute to this clinical variability.⁶⁶ This case suggests that patients with CD21 deficiency present with recurrent infections and low serum IgG levels but uneventful EBV infection.

LPS-responsive beige-like anchor deficiency: Hypogammaglobulinemia with autoimmunity and early colitis

Linkage analysis and whole-exome sequencing conducted by 2 groups led to the identification of homozygous, autosomal

recessive, LPS-responsive beige-like anchor (*LRBA*) gene mutations in 10 patients from 5 unrelated families.^{69,70} The clinical symptoms and laboratory findings in these patients were highly variable. Most had early onset of recurrent bacterial infections, autoimmune disease, or both. A subset had severe nonbloody diarrhea. One of the patients also had recurrent EBV-related lymphoproliferative disease.⁶⁹ Decreased serum IgG and IgA concentrations were seen in 8 of the 10 patients. CD19⁺ B-cell counts were variable, but switched memory B-cell counts were low in all patients who were tested. In addition, an increased susceptibility to apoptosis was documented in LRBA-deficient EBV immortalized B cells.⁷⁰ T cells were phenotypically normal, with variable proliferative responses to mitogens and anti-CD3.⁶⁹

The product of the *LRBA* gene (*LRBA*) is a cytosolic protein expressed in many tissues believed to be involved in endocytosis of ligand-activated receptors and also control of apoptosis.^{71,72} The authors suggest that the autoimmune features seen in LRBA-deficient patients were attributable to increased apoptosis and defective inhibitory receptor signaling.^{69,70} LRBA deficiency should be considered in the presence of early-onset hypogammaglobulinemia associated with colitis, autoimmune features, or both. The mechanism of disease remains to be deciphered.

DEFECTS OF IMMUNE DYSREGULATION

Pallidin deficiency: Hermansky-Pudlak syndrome type 9

Hermansky-Pudlak syndrome (HPS) subtypes present with some degree of albinism and variable bleeding diathesis.⁷³ However, some subtypes have additional features, such as neutropenia and defective lymphocyte-mediated cytotoxicity, as seen in HPS-2.⁷⁴ Two recent articles reported the same homozygous nonsense (c.232C>T) *PLDN* mutation in 2 unrelated patients with HPS-like phenotype.^{75,76} *PLDN* is mutated in the HPS mouse model *pallid* (or HPS-9) and encodes the protein pallidin that interacts with the early endosomal t-SNARE syntaxin-13.^{77,78}

A 17-year-old Italian girl with partial albinism, nystagmus, and normal neurologic development presented with recurrent cutaneous infections.⁷⁵ She had leukopenia and thrombocytopenia at presentation. NK cell degranulation and cytolysis were shown to be defective.⁷⁵ In addition, a 9-month-old Indian boy presented at birth with generalized hypopigmentation and respiratory distress but no history of infections or cytopenias.⁷⁶ Electron microscopy of platelets at 9 months showed absent platelet delta granules, which is consistent with HPS. Neither of the patients had a bleeding history, and results of platelet aggregation studies were normal.⁷⁶ The clinical picture of the Italian patient was reminiscent of the accelerated phase of hemophagocytic lymphohistiocytosis, a feature that was not substantiated in a *pallid* mouse model. The milder clinical phenotype of the second case might be attributed to his younger age. Identification of more patients with hemophagocytic lymphohistiocytosis presentation could list *PLDN* deficiency as a novel cause of albinism and immunodeficiency.

CD27 deficiency: Immune dysregulation and persistent EBV infection

In clinical practice CD27 is recognized as a marker for memory B cells and is used to subclassify patients with a variety of B-cell immunodeficiencies.⁷⁹ After binding its natural ligand, CD70, CD27 regulates differentiation and cellular activity in subsets of

T, B, and NK cells.^{80,81} Two independent reports have recently described a similar presentation of abnormal adaptive human immunity and persistent EBV viremia attributed to CD27 deficiency.^{82,83} Ten patients from 4 independent families (from Morocco, Turkey, and Lebanon) were confirmed to have homozygous mutations in the gene encoding CD27. The clinical picture varied from asymptomatic memory B-cell deficiency to persistent symptomatic EBV viremia and malignant lymphoma. After EBV infection, hypogammaglobulinemia developed in 3 of the affected subjects. T cell-dependent B-cell responses were abnormal,⁸² whereas antipolysaccharide antibodies were detectable.⁸³ Moreover, CD8⁺ T-cell function was disturbed, and invariant NK T-cell numbers were diminished. Three patients died, 2 others underwent successful allogeneic hematopoietic stem cell transplantations, and 2 received anti-CD20 therapy repeatedly. CD27 deficiency predisposes to symptomatic and potentially fatal EBV infection and hypogammaglobulinemia, a phenotype that is similar to that of X-linked lymphoproliferative disease.

CONGENITAL DEFECTS OF PHAGOCYTE NUMBER, FUNCTION, OR BOTH

Interferon-stimulated gene 15 deficiency:

Mendelian susceptibility to mycobacterial diseases

Mendelian susceptibility to mycobacterial diseases (MSMD) predisposes subjects to severe disease on infection by weakly virulent mycobacteria, such as nontuberculous environmental mycobacteria and BCG. Germline mutations in 6 autosomal and 2 X-linked genes involved in IFN- γ -mediated immunity have been reported to cause MSMD (Fig 2).⁸⁴⁻⁸⁶ Recently, another genetic cause of MSMD was described in 2 kindreds.⁸⁷ Three children from 2 distinct kindreds from Iran and Turkey presented with BCG-induced clinical disease starting in early infancy and were found to carry homozygous interferon-stimulated gene 15 (*ISG15*) mutations.⁸⁷ The clinical and immunologic phenotypes of these patients resembled those of IL-12 receptor β 1 deficiency,⁸⁸ with impaired IFN- γ immunity and relatively mild MSMD responsive to antimycobacterial therapy. In mice *ISG15* is induced by IFN- α/β , which is produced in response to viral infection and has been shown to play a role in antiviral defense.^{89,90} However, it seems that the antiviral function of *ISG15* is redundant in human subjects because the patients did not experience unusual viral infections. During phagocytosis, granulocytes are a major source of *ISG15* secretion.⁸⁷ Free *ISG15* then potentiates the secretion of IFN- γ , mostly by NK cells.⁹¹ *ISG15* is therefore not only an IFN- α/β -inducible, intracellular, ubiquitin-like molecule involved in ISGylation but also an IFN- γ -inducing secreted cytokine. This function complements the role of the IL-12-IFN- γ circuit in the control of mycobacterial infection. The IL-12-IFN- γ circuit involves mostly mononuclear phagocytes and T cells, whereas the *ISG15*-IFN- γ circuit involves mostly granulocytes and NK cells. In granulocytes *ISG15* was found to be expressed in secretory granules. The mechanisms by which *ISG15* stimulates NK and T cells are not yet known.

DEFECTS IN INNATE IMMUNITY

NKX2-5 deficiency: Isolated congenital asplenia

Congenital asplenia, which can be diagnosed based on the results of ultrasonography or the presence of Howell-Jolly bodies in blood smears, is often associated with complex visceral defects




FIG 2. Cross-talk between monocyte/macrophage cells and T/NK lymphocytes. Genes in the IL-12/IFN- γ pathway are particularly important for protection against mycobacterial disease. IRF8 is an IFN- γ -inducible transcription factor required for the induction of various target genes, including IL-12. The NF- κ B essential modulator (*NEMO*) mutations in the LZ domain impair CD40-NEMO-dependent pathways. Some gp91phox mutations specifically abolish the respiratory burst in monocyte-derived macrophages. ISG15 is secreted by neutrophils and potentiates IFN- γ production by NK/T cells. Genetic defects that preclude monocyte development (eg, *GATA2*) can also predispose to mycobacterial infections (not shown). Mutant molecules in patients with unusual susceptibility to infection are indicated in red. IFN γ R, IFN- γ receptor; IL-12R, IL-12 receptor; MB, mycobacterium; STAT, signal transducer and activator of transcription; TYK2, tyrosine kinase 2.

as part of heterotaxy syndromes.^{92,93} The causative mutations have been identified in various genes controlling left-right laterality.^{94,95} In contrast, isolated congenital asplenia (ICA; ie, in the absence of heterotaxy or cardiac anomalies) was first thought to be rare and sporadic.⁹⁶ Studies of case reports and rare national surveys suggested probable autosomal dominant, as well as spontaneous, occurrences.^{97,98} The most frequent pathogens in patients with ICA are encapsulated bacteria, with *Streptococcus pneumoniae* as the leading infectious agent.^{98,99} The infections associated with ICA can be fatal in childhood but tend to improve with age.⁹⁸ Using mouse models of spleen morphogenesis to help focus the exome sequencing data of a family with ICA led to the discovery of a heterozygous missense mutation in *NKX2-5*.¹⁰⁰ Pbx1, a prime regulator of the organogenesis of the spleen, governs spleen development through transactivation of the *Nkx2-5* gene product (*Nkx-2*).¹⁰¹ Moreover, both Pbx1 and *Nkx-2* control spleen growth by repression of CDK inhibitor p15Ink4b.¹⁰² Mice deficient for Pbx/*Nkx2-5*/p15 components exhibit disrupted spleen development.¹⁰⁰ Finding of a *NKX2-5* mutation as a possible cause of human ICA reinforces the central role of PBX target genes in the development of spleen. This study also paves the way for the investigation of ICA in other kindreds.

Toll/IL-1 receptor domain-containing adaptor inducing IFN- β and TANK-binding kinase 1 deficiencies: Herpes simplex encephalitis

Herpes simplex virus 1 (HSV1) is an ubiquitous pathogen that causes acute, self-limiting infection after primary exposure. However, it can trigger debilitating and fatal herpes simplex virus encephalitis (HSE) on rare occasions.¹⁰³ Thanks to recent studies, the Toll-like receptor 3 (TLR3)-interferon pathway

emerged as the major player in localized immunity against herpes simplex virus in the central nervous system.^{104,105} A recent study of patients with HSE explored the Toll/IL-1 receptor domain-containing adaptor inducing IFN- β (TRIF) protein as another component of the TLR3 antiviral pathway.¹⁰⁶ Two patients with HSE with TRIF deficiency were identified, one with a homozygous nonsense mutation, resulting in complete recessive TRIF deficiency, and another with a heterozygous missense mutation, leading to partial dominant (probably by negative dominance) TRIF deficiency. TRIF is an adaptor protein serving as the sole adaptor to TLR3 and as an alternative adaptor through TRIF-related adaptor molecule to TLR4.^{107,108} Once activated by agonist-induced TLR3 dimerization, TRIF activates downstream signaling events, culminating in production of IFN- α/β , IFN- λ , and proinflammatory cytokines.^{109,110} TRIF-deficient patients' fibroblasts did not produce interferons after stimulation with TLR3 agonists and showed increased susceptibility to HSV1 infections *in vitro*.¹⁰⁶

Another study also brings to light new facts about genetic susceptibility to HSE. Two different heterozygous missense mutations in TANK-binding kinase 1 (*TBK1*) were identified as causing HSE in 2 patients from Poland and France.¹¹¹ *TBK1* encodes TBK1, a noncanonical I κ B kinase of the NF- κ B signaling pathway, which controls the activity of transcription factors of the interferon regulatory factor family, mainly interferon regulatory factor 3.¹¹² Through this process, TBK1 exerts its antiviral role through regulation of the production of multiple interferons. Both mutant *TBK1* alleles were loss-of-function mutations underlying an autosomal dominant trait in heterozygous patients by haploinsufficiency (D50A) or negative dominance (G159A).¹¹¹ TLR3-related antiviral activities were blunted in both mutant fibroblasts; however, the defects were rescued by IFN- α 2b. The

elucidation of TRIF and TBK1 deficiencies substantiates the obligatory role of the TLR3 pathway in immunity against HSV1 in the central nervous system and its redundancy for protective immunity otherwise.

Minichromosome maintenance complex component 4 deficiency: NK cell deficiency associated with growth retardation and adrenal insufficiency

NK cells are cells of the innate immune system that exert antiviral and antitumor surveillance functions in mice.¹¹³ Some cases of selective quantitative circulating human NK cell deficiencies with specific susceptibility to viral infections have been reported.¹¹⁴⁻¹¹⁷ However, the mechanisms that control NK cell development in human subjects remain unclear. Two groups of researchers have shown that human NK cell deficiency is caused by a homozygous mutation in the minichromosome maintenance complex component 4 (*MCM4*) gene in patients from the Irish traveler community.^{118,119} In addition, patients with NK cell deficiency displayed growth retardation, increased chromosomal breakage, adrenal insufficiency, and, in 1 case, lymphoma. From an infectious perspective, the patients had unusual susceptibility to herpes viruses.¹¹⁸ The studied patients shared the same splice defect, probably because of a founder effect. *MCM4* is a highly conserved DNA helicase that is required for DNA replication and cell proliferation.¹²⁰ Patients' fibroblasts contained high numbers of DNA breaks and showed cell-cycle abnormalities.¹¹⁸ *MCM4* deficiency contributed to a developmental defect in transition of CD56^{bright} to CD56^{dim} NK cells, as evidenced by the lack of CD56^{dim} NK cells in the peripheral blood and the preservation of the small CD56^{bright} NK cell population.^{118,121}

AUTOINFLAMMATORY DISORDERS

A disintegrin and metalloproteinase 17 deficiency: Inflammatory skin and bowel disease

Two siblings born to consanguineous parents of Lebanese origin came to medical attention with early-onset pustular dermatitis, short and broken hair, paronychia, frequent cutaneous bacterial infections, and diarrhea.¹²² The younger sister died at 12 years of age after parvovirus B19-induced myocarditis, and the affected brother was found to have mild cardiomyopathy. A loss-of-function mutation in a disintegrin and metalloproteinase 17 (*ADAM17*) was identified as the cause of this syndrome. *ADAM17* is a major membrane-bound proteinase that cleaves cell-surface proteins, such as cytokines (eg, TNF- α), cytokine receptors (eg, IL-6 receptor and TNF receptor), growth factors (eg, TGF- α), and adhesion proteins (eg, L-selectin).¹²³ *In vivo*, *ADAM17* has a role in controlling inflammation and tissue regeneration. PBMCs obtained from the affected brother showed impaired release of TNF- α but high levels of LPS-induced production of IL-1 β and IL-6. Lack of TNF- α was considered partly responsible for their increased susceptibility to infection and development of cardiomyopathy.^{122,124}

IL-36 receptor antagonist deficiency: Generalized pustular psoriasis

Generalized pustular psoriasis is a life-threatening, multi-systemic inflammatory disease characterized by episodic,

widespread, diffuse erythematous pustular rash associated with high fever, malaise, and leukocytosis.^{125,126} By using homozygosity mapping and exome sequencing, a total of 19 patients were found to have homozygous or compound heterozygous mutations in the *IL36RN* gene, encoding IL-36 receptor antagonist (IL-36Ra).^{127,128} IL-36Ra (also known as IL-1F5) is an antagonist of 3 cytokines of the IL-1 family (IL-36A, IL-36B, and IL-36G) that have NF- κ B- and mitogen-activated protein kinase-activating properties.^{129,130} Defective expression and function of IL-36Ra thus could direct uncontrolled inflammatory processes on pathogen stimulation through mucosal surfaces. Generalized pustular psoriasis provides one more example that highlights the central role of IL-1 signaling in tissue inflammation,¹³¹ illuminating a rational target for its treatment.¹³²

CONCLUSION

The field of PIDs is advancing at full speed in 2 directions. New genetic causes of known PIDs are being discovered (eg, CD21 and TRIF). Moreover, new phenotypes qualify as PIDs with the identification of a first genetic cause (eg, generalized pustular psoriasis). Recent findings contribute fundamental knowledge about immune system biology and its perturbation in disease. They are also of considerable clinical benefit for the patients and their families. A priority is to further translate these new discoveries into improved diagnostic methods and more effective therapeutic strategies, promoting the well-being of patients with PIDs.

REFERENCES

1. Casanova JL, Abel L. Primary immunodeficiencies: a field in its infancy. *Science* 2007;317:617-9.
2. Casanova JL, Abel L. The human model: a genetic dissection of immunity to infection in natural conditions. *Nat Rev Immunol* 2004;4:55-66.
3. Bamshad MJ, Ng SB, Bigham AW, Tabor HK, Emond MJ, Nickerson DA, et al. Exome sequencing as a tool for Mendelian disease gene discovery. *Nat Rev Genet* 2011;12:745-55.
4. Al-Herz W, Bousfiha A, Casanova JL, et al. Primary immunodeficiency diseases: an update on the classification from the international union of immunological societies expert committee for primary immunodeficiency. *Front Immunol* 2011;2:54.
5. Al-Herz W, Notarangelo LD. Classification of primary immunodeficiency disorders: one-fits-all does not help anymore. *Clin Immunol* 2012;144:24-5.
6. Conley ME, Notarangelo LD, Casanova JL. Definition of primary immunodeficiency in 2011: a "trialogue" among friends. *Ann N Y Acad Sci* 2011;1238:1-6.
7. Davis MM, Bjorkman PJ. T-cell antigen receptor genes and T-cell recognition. *Nature* 1988;334:395-402.
8. Anderson G, Moore NC, Owen JJ, Jenkinson EJ. Cellular interactions in thymocyte development. *Annu Rev Immunol* 1996;14:73-99.
9. Morgan NV, Goddard S, Cardno TS, et al. Mutation in the TCR α subunit constant gene (TRAC) leads to a human immunodeficiency disorder characterized by a lack of TCR α beta+ T cells. *J Clin Invest* 2011;121:695-702.
10. Munger K, Howley PM. Human papillomavirus immortalization and transformation functions. *Virus Res* 2002;89:213-28.
11. Orth G. Host defenses against human papillomaviruses: lessons from epidermodysplasia verruciformis. *Curr Top Microbiol Immunol* 2008;321:59-83.
12. Ramoz N, Rueda LA, Bouadjar B, Montoya LS, Orth G, Favre M. Mutations in two adjacent novel genes are associated with epidermodysplasia verruciformis. *Nat Genet* 2002;32:579-81.
13. Lazarczyk M, Pons C, Mendoza JA, Cassonnet P, Jacob Y, Favre M. Regulation of cellular zinc balance as a potential mechanism of EVER-mediated protection against pathogenesis by cutaneous oncogenic human papillomaviruses. *J Exp Med* 2008;205:35-42.
14. Crequer A, Picard C, Pedergnana V, et al. EVER2 deficiency is associated with mild T-cell abnormalities. *J Clin Immunol* 2012 [Epub ahead of print].
15. Crequer A, Troeger A, Patin E, et al. Human RHOH deficiency causes T cell defects and susceptibility to EV-HPV infections. *J Clin Invest* 2012;122:3239-47.
16. Dorn T, Kuhn U, Bungartz G, et al. RhoH is important for positive thymocyte selection and T-cell receptor signaling. *Blood* 2007;109:2346-55.
17. Tybulewicz VL, Henderson RB. Rho family GTPases and their regulators in lymphocytes. *Nat Rev Immunol* 2009;9:630-44.

18. Crequer A, Picard C, Patin E, et al. Inherited MST1 deficiency underlies susceptibility to EV-HPV infections. *PLoS ONE* 2012;7:e44010.
19. Abdollahpour H, Appaswamy G, Kotlarz D, et al. The phenotype of human STK4 deficiency. *Blood* 2012;119:3450-7.
20. Nehme NT, Schmid JP, Debeurme F, et al. MST1 mutations in autosomal recessive primary immunodeficiency characterized by defective naive T-cell survival. *Blood* 2012;119:3458-68.
21. Creasy CL, Chernoff J. Cloning and characterization of a human protein kinase with homology to Ste20. *J Biol Chem* 1995;270:21695-700.
22. Zhao B, Li L, Lei Q, Guan KL. The Hippo-YAP pathway in organ size control and tumorigenesis: an updated version. *Genes Dev* 2010;24:862-74.
23. Straus DB, Weiss A. Genetic evidence for the involvement of the lck tyrosine kinase in signal transduction through the T cell antigen receptor. *Cell* 1992;70:585-93.
24. Veillette A, Horak ID, Horak EM, Bookman MA, Bolen JB. Alterations of the lymphocyte-specific protein tyrosine kinase (p56lck) during T-cell activation. *Mol Cell Biol* 1988;8:4353-61.
25. Salmond RJ, Filby A, Qureshi I, Caserta S, Zamoyska R. T-cell receptor proximal signaling via the Src-family kinases, Lck and Fyn, influences T-cell activation, differentiation, and tolerance. *Immunol Rev* 2009;228:9-22.
26. Molina TJ, Kishihara K, Siderovski DP, et al. Profound block in thymocyte development in mice lacking p56lck. *Nature* 1992;357:161-4.
27. Goldman FD, Ballas ZK, Schutte BC, et al. Defective expression of p56lck in an infant with severe combined immunodeficiency. *J Clin Invest* 1998;102:421-9.
28. Hubert P, Bergeron F, Ferreira V, et al. Defective p56Lck activity in T cells from an adult patient with idiopathic CD4+ lymphocytopenia. *Int Immunol* 2000;12:449-57.
29. Sawabe T, Horiuchi T, Nakamura M, et al. Defect of lck in a patient with common variable immunodeficiency. *Int J Mol Med* 2001;7:609-14.
30. Hauck F, Randriamampita C, Martin E, et al. Primary T-cell immunodeficiency with immunodysregulation caused by autosomal recessive LCK deficiency. *J Allergy Clin Immunol* 2012;130:1144-52.e11.
31. Unexplained CD4+ T-lymphocyte depletion in persons without evident HIV infection—United States. *MMWR Morb Mortal Wkly Rep* 1992;41:541-5.
32. Laurence J, Mitra D, Steiner M, Lynch DH, Siegal FP, Staiano-Coico L. Apoptotic depletion of CD4+ T cells in idiopathic CD4+ T lymphocytopenia. *J Clin Invest* 1996;97:672-80.
33. Roger PM, Bernard-Pomier G, Counillon E, Breittmayer JP, Bernard A, Dellamonica P. Overexpression of Fas/CD95 and Fas-induced apoptosis in a patient with idiopathic CD4+ T lymphocytopenia. *Clin Infect Dis* 1999;28:1012-6.
34. Gorska MM, Stafford SJ, Cen O, Sur S, Alam R. Unc119, a novel activator of Lck/Fyn, is essential for T cell activation. *J Exp Med* 2004;199:369-79.
35. Gorska MM, Liang Q, Karim Z, Alam R. Uncoordinated 119 protein controls trafficking of Lck via the Rab11 endosome and is critical for immunological synapse formation. *J Immunol* 2009;183:1675-84.
36. Gorska MM, Goplen N, Liang Q, Alam R. Uncoordinated 119 preferentially induces Th2 differentiation and promotes the development of asthma. *J Immunol* 2010;184:4488-96.
37. Gorska MM, Alam R. Consequences of a mutation in the UNC119 gene for T Cell function in idiopathic CD4 lymphopenia. *Curr Allergy Asthma Rep* 2012;12:396-401.
38. Gorska MM, Alam R. A mutation in the human Uncoordinated 119 gene impairs TCR signaling and is associated with CD4 lymphopenia. *Blood* 2012;119:1399-406.
39. de la Fuente MA, Sasahara Y, Calamito M, et al. WIP is a chaperone for Wiskott-Aldrich syndrome protein (WASP). *Proc Natl Acad Sci U S A* 2007;104:926-31.
40. Lanzi G, Moratto D, Vairo D, et al. A novel primary human immunodeficiency due to deficiency in the WASP-interacting protein WIP. *J Exp Med* 2012;209:29-34.
41. Gallego MD, de la Fuente MA, Anton IM, Snapper S, Fuhlbrigge R, Geha RS. WIP and WASP play complementary roles in T cell homing and chemotaxis to SDF-1alpha. *Int Immunol* 2006;18:221-32.
42. Haddad E, Zugaza JL, Louache F, et al. The interaction between Cdc42 and WASP is required for SDF-1-induced T-lymphocyte chemotaxis. *Blood* 2001;97:33-8.
43. Le BS, Massaad M, Koduru S, et al. WIP is critical for T cell responsiveness to IL-2. *Proc Natl Acad Sci U S A* 2009;106:7519-24.
44. Curcio C, Pannellini T, Lanzardo S, Forni G, Musiani P, Anton IM. WIP null mice display a progressive immunological disorder that resembles Wiskott-Aldrich syndrome. *J Pathol* 2007;211:67-75.
45. Wiskott A. Familiärer, angeborener Morbus Werlhofii? *Montasschr Kinderheilkd* 1937;68:212-6.
46. Aldrich RA, Steinberg AG, Campbell DC. Pedigree demonstrating a sex-linked recessive condition characterized by draining ears, eczematoid dermatitis and bloody diarrhea. *Pediatrics* 1954;13:133-9.
47. Ombrello MJ, Remmers EF, Sun G, et al. Cold urticaria, immunodeficiency, and autoimmunity related to PLCG2 deletions. *N Engl J Med* 2012;366:330-8.
48. Zhou Q, Lee GS, Brady J, et al. A Hypermorph missense mutation in PLCG2, encoding phospholipase Cgamma2, causes a dominantly inherited autoinflammatory disease with immunodeficiency. *Am J Hum Genet* 2012;91:713-20.
49. Patterson RL, van Rossum DB, Nikolaidis N, Gill DL, Snyder SH. Phospholipase C-gamma: diverse roles in receptor-mediated calcium signaling. *Trends Biochem Sci* 2005;30:688-97.
50. Wilde JI, Watson SP. Regulation of phospholipase C gamma isoforms in haematopoietic cells: why one, not the other? *Cell Signal* 2001;13:691-701.
51. Conley ME, Dobbs AK, Farmer DM, et al. Primary B cell immunodeficiencies: comparisons and contrasts. *Annu Rev Immunol* 2009;27:199-227.
52. Conley ME, Dobbs AK, Quintana AM, et al. Agammaglobulinemia and absent B lineage cells in a patient lacking the p85alpha subunit of PI3K. *J Exp Med* 2012;209:463-70.
53. Vanhaesebroeck B, Stephens L, Hawkins P. PI3K signalling: the path to discovery and understanding. *Nat Rev Mol Cell Biol* 2012;13:195-203.
54. Aiuti A, Tavian M, Cipponi A, et al. Expression of CXCR4, the receptor for stromal cell-derived factor-1 on fetal and adult human lympho-hematopoietic progenitors. *Eur J Immunol* 1999;29:1823-31.
55. Ma Q, Jones D, Borghesani PR, et al. Impaired B-lymphopoiesis, myelopoiesis, and derailed cerebellar neuron migration in CXCR4- and SDF-1-deficient mice. *Proc Natl Acad Sci U S A* 1998;95:9448-53.
56. Kanegane H, Taneichi H, Nomura K, Futatani T, Miyawaki T. Severe neutropenia in Japanese patients with x-linked agammaglobulinemia. *J Clin Immunol* 2005;25:491-5.
57. Farrar JE, Rohrer J, Conley ME. Neutropenia in X-linked agammaglobulinemia. *Clin Immunol Immunopathol* 1996;81:271-6.
58. Lopez GE, Porpiglia AS, Hogan MB, et al. Clinical and molecular analysis of patients with defects in mu heavy chain gene. *J Clin Invest* 2002;110:1029-35.
59. Fruman DA. Regulatory subunits of class IA PI3K. *Curr Top Microbiol Immunol* 2010;346:225-44.
60. Fukao T, Tanabe M, Terauchi Y, et al. PI3K-mediated negative feedback regulation of IL-12 production in DCs. *Nat Immunol* 2002;3:875-81.
61. Fearon DT, Carroll MC. Regulation of B lymphocyte responses to foreign and self-antigens by the CD19/CD21 complex. *Annu Rev Immunol* 2000;18:393-422.
62. Asokan R, Hua J, Young KA, et al. Characterization of human complement receptor type 2 (complement receptor type 2/CD21) as a receptor for IFN-alpha: a potential role in systemic lupus erythematosus. *J Immunol* 2006;177:383-94.
63. Szakonyi G, Guthridge JM, Li D, Young K, Holers VM, Chen XS. Structure of complement receptor 2 in complex with its C3d ligand. *Science* 2001;292:1725-8.
64. Lowell CA, Klickstein LB, Carter RH, Mitchell JA, Fearon DT, Ahearn JM. Mapping of the Epstein-Barr virus and C3dg binding sites to a common domain on complement receptor type 2. *J Exp Med* 1989;170:1931-46.
65. Carter RH, Fearon DT. CD19: lowering the threshold for antigen receptor stimulation of B lymphocytes. *Science* 1992;256:105-7.
66. Thiel J, Kimmig L, Salzer U, et al. Genetic CD21 deficiency is associated with hypogammaglobulinemia. *J Allergy Clin Immunol* 2012;129:801-10.
67. van Zelm MC, Reisli I, van der Burg M, et al. An antibody-deficiency syndrome due to mutations in the CD19 gene. *N Engl J Med* 2006;354:1901-12.
68. van Zelm MC, Smet J, Adams B, et al. CD81 gene defect in humans disrupts CD19 complex formation and leads to antibody deficiency. *J Clin Invest* 2010;120:1265-74.
69. Alangari A, Alsultan A, Adly N, et al. LPS-responsive beige-like anchor (LRBA) gene mutation in a family with inflammatory bowel disease and combined immunodeficiency. *J Allergy Clin Immunol* 2012;130:481-8.
70. Lopez-Herrera G, Tampella G, Pan-Hammarstrom Q, et al. Deleterious mutations in LRBA are associated with a syndrome of immune deficiency and autoimmunity. *Am J Hum Genet* 2012;90:986-1001.
71. de Souza N, Vallier LG, Fares H, Greenwald I. SEL-2, the *C. elegans* neurobeachin/LRBA homolog, is a negative regulator of lin-12/Notch activity and affects endosomal traffic in polarized epithelial cells. *Development* 2007;134:691-702.
72. Wang JW, Gamsby JJ, Highfill SL, et al. Deregulated expression of LRBA facilitates cancer cell growth. *Oncogene* 2004;23:4089-97.
73. Wei ML. Hermansky-Pudlak syndrome: a disease of protein trafficking and organelle function. *Pigment Cell Res* 2006;19:19-42.
74. Badolato R, Parolini S. Novel insights from adaptor protein 3 complex deficiency. *J Allergy Clin Immunol* 2007;120:735-41.
75. Badolato R, Prandini A, Caracciolo S, et al. Exome sequencing reveals a pallidin mutation in a Hermansky-Pudlak-like primary immunodeficiency syndrome. *Blood* 2012;119:3185-7.
76. Cullinane AR, Curry JA, Carmona-Rivera C, et al. A BLOC-1 mutation screen reveals that PLDN is mutated in Hermansky-Pudlak Syndrome type 9. *Am J Hum Genet* 2011;88:778-87.

77. Dell'Angelica EC. The building BLOC(k)s of lysosomes and related organelles. *Curr Opin Cell Biol* 2004;16:458-64.
78. Huang L, Kuo YM, Gitschier J. The pallid gene encodes a novel, syntaxin 13-interacting protein involved in platelet storage pool deficiency. *Nat Genet* 1999;23:329-32.
79. Wehr C, Kivioja T, Schmitt C, et al. The EUROclass trial: defining subgroups in common variable immunodeficiency. *Blood* 2008;111:77-85.
80. Borst J, Hendriks J, Xiao Y. CD27 and CD70 in T cell and B cell activation. *Curr Opin Immunol* 2005;17:275-81.
81. Nolte MA, van Oeffen RW, van Gisbergen KP, van Lier RA. Timing and tuning of CD27-CD70 interactions: the impact of signal strength in setting the balance between adaptive responses and immunopathology. *Immunol Rev* 2009;229:216-31.
82. van Montfrans JM, Hoepelman AI, Otto S, et al. CD27 deficiency is associated with combined immunodeficiency and persistent symptomatic EBV viremia. *J Allergy Clin Immunol* 2012;129:787-93.
83. Salzer E, Daschkey S, Choo S, et al. Combined immunodeficiency with life-threatening EBV-associated lymphoproliferative disorder in patients lacking functional CD27. *Haematologica* 2012 [Epub ahead of print].
84. Al-Muhsen S, Casanova JL. The genetic heterogeneity of Mendelian susceptibility to mycobacterial diseases. *J Allergy Clin Immunol* 2008;122:1043-51.
85. Bustamante J, Arias AA, Vogt G, et al. Germline CYBB mutations that selectively affect macrophages in kindreds with X-linked predisposition to tuberculous mycobacterial disease. *Nat Immunol* 2011;12:213-21.
86. Hambleton S, Salem S, Bustamante J, et al. IRF8 mutations and human dendritic-cell immunodeficiency. *N Engl J Med* 2011;365:127-38.
87. Bogunovic D, Byun M, Durfee LA, et al. Mycobacterial disease and impaired IFN-gamma immunity in humans with inherited ISG15 deficiency. *Science* 2012;337:1684-8.
88. de Beaucoudrey L, Samarina A, Bustamante J, et al. Revisiting human IL-12beta1 deficiency: a survey of 141 patients from 30 countries. *Medicine (Baltimore)* 2010;89:381-402.
89. Durfee LA, Lyon N, Seo K, Huijbregtse JM. The ISG15 conjugation system broadly targets newly synthesized proteins: implications for the antiviral function of ISG15. *Mol Cell* 2010;38:722-32.
90. Pitha-Rowe IF, Pitha PM. Viral defense, carcinogenesis and ISG15: novel roles for an old ISG. *Cytokine Growth Factor Rev* 2007;18:409-17.
91. D'Cunha J, Ramanujam S, Wagner RJ, Witt PL, Knight E Jr, Borden EC. In vitro and in vivo secretion of human ISG15, an IFN-induced immunomodulatory cytokine. *J Immunol* 1996;157:4100-8.
92. Ivemark BI. Implications of agenesis of the spleen on the pathogenesis of conotruncus anomalies in childhood; an analysis of the heart malformations in the splenic agenesis syndrome, with fourteen new cases. *Acta Paediatr Suppl* 1955;44(suppl 104):7-110.
93. Rose V, Izukawa T, Moes CA. Syndromes of asplenia and polysplenia. A review of cardiac and non-cardiac malformations in 60 cases with special reference to diagnosis and prognosis. *Br Heart J* 1975;37:840-52.
94. Belmont JW, Mohapatra B, Towbin JA, Ware SM. Molecular genetics of heterotaxy syndromes. *Curr Opin Cardiol* 2004;19:216-20.
95. Zhu L, Belmont JW, Ware SM. Genetics of human heterotaxias. *Eur J Hum Genet* 2006;14:17-25.
96. Myerson RM, Koelle WA. Congenital absence of the spleen in an adult; report of a case associated with recurrent Waterhouse-Friderichsen syndrome. *N Engl J Med* 1956;254:1131-2.
97. Gilbert B, Menetrey C, Belin V, Brosset P, de Lumley L, Fisher A. Familial isolated congenital asplenia: a rare, frequently hereditary dominant condition, often detected too late as a cause of overwhelming pneumococcal sepsis. Report of a new case and review of 31 others. *Eur J Pediatr* 2002;161:368-72.
98. Mahlaoui N, Minard-Colin V, Picard C, et al. Isolated congenital asplenia: a French nationwide retrospective survey of 20 cases. *J Pediatr* 2011;158:142-8.
99. Picard C, Puel A, Bustamante J, Ku CL, Casanova JL. Primary immunodeficiencies associated with pneumococcal disease. *Curr Opin Allergy Clin Immunol* 2003;3:451-9.
100. Koss M, Bolze A, Brendolan A, et al. Congenital asplenia in mice and humans with mutations in a Pbx/Nkx2-5/p15 module. *Dev Cell* 2012;22:913-26.
101. Berthelsen J, Zappavigna V, Ferretti E, Mavilio F, Blasi F. The novel homeoprotein Prep1 modulates Pbx-Hox protein cooperativity. *EMBO J* 1998;17:1434-45.
102. Latres E, Malumbres M, Sotillo R, et al. Limited overlapping roles of P15(INK4b) and P18(INK4c) cell cycle inhibitors in proliferation and tumorigenesis. *EMBO J* 2000;19:3496-506.
103. Whitley RJ. Herpes simplex virus in children. *Curr Treat Options Neurol* 2002;4:231-7.
104. Sancho-Shimizu V, Zhang SY, Abel L, et al. Genetic susceptibility to herpes simplex virus 1 encephalitis in mice and humans. *Curr Opin Allergy Clin Immunol* 2007;7:495-505.
105. Sancho-Shimizu V, Perez de Diego R, Jouanguy E, Zhang SY, Casanova JL. Inborn errors of anti-viral interferon immunity in humans. *Curr Opin Virol* 2011;1:487-96.
106. Sancho-Shimizu V, Perez de Diego R, Lorenzo L, et al. Herpes simplex encephalitis in children with autosomal recessive and dominant TRIF deficiency. *J Clin Invest* 2011;121:4889-902.
107. Yamamoto M, Sato S, Hemmi H, et al. Role of adaptor TRIF in the MyD88-independent toll-like receptor signaling pathway. *Science* 2003;301:640-3.
108. Hoebe K, Du X, Georgel P, et al. Identification of Lps2 as a key transducer of MyD88-independent TIR signalling. *Nature* 2003;424:743-8.
109. Funami K, Sasai M, Ohba Y, Oshiumi H, Seya T, Matsumoto M. Spatiotemporal mobilization of Toll/IL-1 receptor domain-containing adaptor molecule-1 in response to dsRNA. *J Immunol* 2007;179:6867-72.
110. Funami K, Sasai M, Oshiumi H, Seya T, Matsumoto M. Homo-oligomerization is essential for Toll/interleukin-1 receptor domain-containing adaptor molecule-1-mediated NF-kappaB and interferon regulatory factor-3 activation. *J Biol Chem* 2008;283:18283-91.
111. Herman M, Ciancanelli M, Ou YH, et al. Heterozygous TBK1 mutations impair TLR3 immunity and underlie herpes simplex encephalitis of childhood. *J Exp Med* 2012;209:1567-82.
112. Sharma S, Tenover BR, Grandvaux N, Zhou GP, Lin R, Hiscott J. Triggering the interferon antiviral response through an IKK-related pathway. *Science* 2003;300:1148-51.
113. Vivier E, Raulet DH, Moretta A, et al. Innate or adaptive immunity? The example of natural killer cells. *Science* 2011;331:44-9.
114. Etzioni A, Eidenschenk C, Katz R, Beck R, Casanova JL, Pollack S. Fatal varicella associated with selective natural killer cell deficiency. *J Pediatr* 2005;146:423-5.
115. Eidenschenk C, Dunne J, Jouanguy E, et al. A novel primary immunodeficiency with specific natural-killer cell deficiency maps to the centromeric region of chromosome 8. *Am J Hum Genet* 2006;78:721-7.
116. Biron CA, Byron KS, Sullivan JL. Severe herpesvirus infections in an adolescent without natural killer cells. *N Engl J Med* 1989;320:1731-5.
117. Eidenschenk C, Jouanguy E, Alcais A, et al. Familial NK cell deficiency associated with impaired IL-2- and IL-15-dependent survival of lymphocytes. *J Immunol* 2006;177:8835-43.
118. Gineau L, Cognet C, Kara N, et al. Partial MCM4 deficiency in patients with growth retardation, adrenal insufficiency, and natural killer cell deficiency. *J Clin Invest* 2012;122:821-32.
119. Hughes CR, Guasti L, Meimaridou E, et al. MCM4 mutation causes adrenal failure, short stature, and natural killer cell deficiency in humans. *J Clin Invest* 2012;122:814-20.
120. Bochman ML, Schwacha A. The Mcm complex: unwinding the mechanism of a replicative helicase. *Microbiol Mol Biol Rev* 2009;73:652-83.
121. Orange JS. Unraveling human natural killer cell deficiency. *J Clin Invest* 2012;122:798-801.
122. Blaydon DC, Biancheri P, Di WL, et al. Inflammatory skin and bowel disease linked to ADAM17 deletion. *N Engl J Med* 2011;365:1502-8.
123. Scheller J, Chalaris A, Garbers C, Rose-John S. ADAM17: a molecular switch to control inflammation and tissue regeneration. *Trends Immunol* 2011;32:380-7.
124. Wada H, Saito K, Kanda T, et al. Tumor necrosis factor-alpha (TNF-alpha) plays a protective role in acute viral myocarditis in mice: a study using mice lacking TNF-alpha. *Circulation* 2001;103:743-9.
125. Auguy F, Renaudier P, Nicolas JF. Generalized pustular psoriasis (Zumbusch): a French epidemiological survey. *Eur J Dermatol* 2006;16:669-73.
126. Zelikson BD, Muller SA. Generalized pustular psoriasis. A review of 63 cases. *Arch Dermatol* 1991;127:1339-45.
127. Marrakchi S, Guigue P, Renshaw BR, et al. Interleukin-36-receptor antagonist deficiency and generalized pustular psoriasis. *N Engl J Med* 2011;365:620-8.
128. Onoufriadi A, Simpson MA, Pink AE, et al. Mutations in IL36RN/IL1F5 are associated with the severe episodic inflammatory skin disease known as generalized pustular psoriasis. *Am J Hum Genet* 2011;89:432-7.
129. Sims JE, Smith DE. The IL-1 family: regulators of immunity. *Nat Rev Immunol* 2010;10:89-102.
130. Dinarello C, Arend W, Sims J, et al. IL-1 family nomenclature. *Nat Immunol* 2010;11:973.
131. Goldbach-Mansky R, Kastner DL. Autoinflammation: the prominent role of IL-1 in monogenic autoinflammatory diseases and implications for common illnesses. *J Allergy Clin Immunol* 2009;124:1141-9.
132. Viguier M, Guigue P, Pages C, Smahi A, Bachelez H. Successful treatment of generalized pustular psoriasis with the interleukin-1-receptor antagonist anakinra: lack of correlation with IL1RN mutations. *Ann Intern Med* 2010;153:66-7.