

Allergic contact dermatitis caused by benzisothiazolinone in a continuous positive airway pressure mask liquid soap

Claire Felmingham  | Rosemary Nixon | Amanda Palmer | Adriene Lee

Occupational Dermatology Research and Education Centre, The Skin and Cancer Foundation Inc., Carlton, Victoria, Australia

Correspondence

Dr Claire Felmingham, The Skin and Cancer Foundation Inc., Occupational Dermatology Research and Education Centre, Level 1/80 Drummond Street, Carlton, Victoria, Australia.

Email: clairefelmingham@gmail.com

KEYWORDS: allergic contact dermatitis, benzisothiazolinone, case report, isothiazolinone

As the important allergens methylisothiazolinone (MI) and methylchloroisothiazolinone (MCI) are removed from products, their replacements, including other isothiazolinones, may emerge as significant allergens. We report a case of allergic contact dermatitis (ACD) of the hands caused by benzisothiazolinone (BIT) after exposure to a liquid soap and a dishwashing concentrate.

1 | CASE REPORT

A 66-year-old retired cleaner presented with a longstanding history of pruritic hand dermatitis that had worsened in the last 2 years. She was currently being treated with methotrexate, and had previously been prescribed cyclosporin. She had a personal history of eczema and hay fever, and a family history of atopy. Her medical history included obstructive sleep apnoea, for which she used a continuous positive airway pressure (CPAP) mask. On examination, there was erythema with oedema, scaling and vesicles of both dorsal and ventral surfaces of her hands.

She had cleaned her CPAP mask daily with a liquid soap (CPAP wash) that she had purchased from a medical supplies store for the last 2 years. There were no ingredients listed on the liquid soap container, and we were unable to determine them on the first day of patch testing. She did not wear gloves when washing the mask. Furthermore, she made her own cleaning liquid, which she had used around the house for 5 years, comprising eucalyptus oil, dishwashing liquid, and water (own cleaner). The ingredients of the dishwashing

liquid were not available on the first day of patch testing. She would usually wear protective rubber gloves when cleaning with this mixture.

Patch testing with the Australian baseline series, a rubber series, eucalyptus oil, and her own products, including the CPAP wash (1% aq.) and her own cleaner (1% aq.), was undertaken. Testing was performed with allergEAZE skin patch test chambers, and allergens were obtained from allergEAZE (SmartPractice, Phoenix, Arizona) and Chemotechnique Diagnostics (Vellinge, Sweden), and applied for 2 days. Patch test readings were performed on day (D) 2 and D4 according to ICDRG guidelines. The results are shown in Table 1.

On D2, information on the ingredients of both the CPAP wash and the patient's own cleaner was not yet available. After extensive enquiries to the manufacturer, it was found that CPAP liquid soap

TABLE 1 Patch test readings and relevance on day (D) 2 and D4

Allergen	Patch test reading		Relevance
	D2	D4	
Methylisothiazolinone 0.2% aq.	+	+	Past
Methylisothiazolinone/ methylchloroisothiazolinone 0.02% aq.	+	+	Past
CPAP wash 1% aq.	+	+	Relevant
Hydroperoxides of linalool 1% pet.	?+	+	Unknown
Own cleaner 1% aq.	-	-	Relevant
Benzisothiazolinone 0.05% pet.	+		Relevant

CPAP, continuous positive airway pressure.

contained BIT. It was also found that the liquid detergent used as an ingredient in the patient's own cleaner contained BIT. Once this information was available, on D4, BIT 0.05% pet., supplied by Chemotechnique Diagnostics, was applied as an additional patch test.

As the patient lived some distance from the patch testing centre, she took photographs of her reactions to BIT on D2 and D4, and these showed a positive reaction on D2 (+). On the basis of this, and her history of a flare of her hand dermatitis in the 2 years since being exposed to the CPAP wash, she was diagnosed with ACD caused by BIT present in her CPAP wash and own cleaner, in addition to her pre-existing atopic hand eczema. MI, MCI and linalool were not found in the products that she was currently using, and may have been of past relevance. The patient was counselled to avoid all of these allergens.

2 | DISCUSSION

Preservatives are required in water-based products, including personal and household products, to prevent contamination with microorganisms and to protect the consumer. Some members of the family of isothiazolinones include MI, MCI, octylisothiazolinone (OIT), and BIT.

There has been an epidemic of ACD caused by isothiazolinones, particularly MI and MI/MCI, over the last decade.¹ International regulations from the EU indicate that MI/MCI and MI are currently only allowed in rinse-off personal care (cosmetic) products,² and that BIT and OIT are not allowed in personal care products.^{3,4} The allowed concentration of BIT in other products is, however, unregulated, and it has been found in relatively high concentrations.⁵ In this case, BIT was not labelled on the patient's CPAP wash, and the manufacturer needed to be contacted, as there was no material safety data sheet available for the product. BIT is found eg. in household cleaning products, in water-based paints, in polyvinyl chloride gloves, in glue, and as an industrial biocide, and has been reported to cause ACD as a result of all of these exposures.^{4,6,7}

In a recent study of 647 consecutively patch tested patients, 1.4% had positive reactions to BIT, and 56% of those were found to have BIT exposure.⁸ This is similar to the 1.6% of positive reactions to BIT found in the 7532 patch tested patients in another study.⁹ In older studies, the sensitization rates have been reported to be 0.8% to 1.2%.^{10,11} BIT is not generally included in baseline series, and is often found in products that may not be labelled appropriately, as in our case. It appears that there may be increasing use of BIT as an alternative to MI, and that it is an emerging allergen.

CONFLICTS OF INTEREST

The authors have no conflicts of interest to report.

ORCID

Claire Felmingham  <https://orcid.org/0000-0002-3443-8065>

REFERENCES

1. Lundov MD, Opstrup MS, Johansen JD. Methylisothiazolinone contact allergy—growing epidemic. *Contact Dermatitis*. 2013;69:271-275.
2. Regulation EC. No. 2016/1198 of 22 July 2016 amending Annex V to regulations (EC.) No. 1223/2009 of the European parliament and of the council on cosmetic products. *Official Journal of the European Union*, L198/10. 2016
3. Regulation EC. No. 1223/2009 of 30 November 2009 on cosmetic products. *Official Journal of the European Union*, L342/59. 2009
4. Garcia-Hidalgo E, Schneider D, von Goetz N, Delmaar C, Siegrist M, Hungerbühler K. Aggregate consumer exposure to isothiazolinones via household care and personal care products: probabilistic modelling and benzisothiazolinone risk assessment. *Environ Int*. 2018;118:245-256.
5. Garcia-Hidalgo E, Sottas V, von Goetz N, Hauri U, Bogdal C, Hungerbühler K. Occurrence and concentrations of isothiazolinones in detergents and cosmetics in Switzerland. *Contact Dermatitis*. 2017;76:96-106.
6. Ayadi M, Martin P. Pulpitis of the fingers from a shoe glue containing 1,2-benzisothiazolin-3-one (BIT). *Contact Dermatitis*. 1999;40:115-116.
7. Aalto-Korte K, Alanko K, Henriks-Eckerman ML, Jolanki R. Antimicrobial allergy from polyvinyl chloride gloves. *Arch Dermatol*. 2006;142:1326-1330.
8. Aalto-Korte K, Suuronen K. Patterns of concomitant allergic reactions in patients suggest cross-sensitization between octylisothiazolinone and methylisothiazolinone. *Contact Dermatitis*. 2017;77:385-389.
9. Geier J, Lessmann H, Schnuch A, Uter W. Concomitant reactivity to methylisothiazolinone, benzisothiazolinone, and octylisothiazolinone. International Network of Departments of Dermatology data, 2009-2013. *Contact Dermatitis*. 2015;72:337-339.
10. Andersen KE, Veien NK. Biocide patch tests. *Contact Dermatitis*. 1985;12:99-103.
11. Schnuch A, Geier J, Uter W, Frosch PJ. Patch testing with preservatives, antimicrobials and industrial biocides. Results from a multicentre study. *Br J Dermatol*. 1998;138:467-476.

How to cite this article: Felmingham C, Nixon R, Palmer A, Lee A. Allergic contact dermatitis caused by benzisothiazolinone in a continuous positive airway pressure mask liquid soap. *Contact Dermatitis*. 2019;81:152-153. <https://doi.org/10.1111/cod.13273>