


Journal Pre-proof

Clinical Communication: Regional and Temporal Awareness of Alpha-Gal Allergy – An Infodemiological Analysis Using Google Trends

Edward G.A. Iglesia, MD MPH, Cosby A. Stone, Jr., MD MPH, Mary Grace Flaherty, PhD MLS MS, Scott P. Commins, MD PhD



PII: S2213-2198(19)31023-2

DOI: <https://doi.org/10.1016/j.jaip.2019.12.003>

Reference: JAIP 2587

To appear in: *The Journal of Allergy and Clinical Immunology: In Practice*

Received Date: 13 August 2019

Revised Date: 24 October 2019

Accepted Date: 4 December 2019

Please cite this article as: Iglesia EGA, Stone CA Jr., Flaherty MG, Commins SP, Clinical Communication: Regional and Temporal Awareness of Alpha-Gal Allergy – An Infodemiological Analysis Using Google Trends, *The Journal of Allergy and Clinical Immunology: In Practice* (2020), doi: <https://doi.org/10.1016/j.jaip.2019.12.003>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2019 Published by Elsevier Inc. on behalf of the American Academy of Allergy, Asthma & Immunology

Clinical Communication: Regional and Temporal Awareness of Alpha-Gal Allergy – An Infodemiological Analysis Using Google Trends

Authors: Edward G.A. Iglesia, MD MPH¹, Cosby A. Stone, Jr. MD MPH², Mary Grace Flaherty, PhD MLS MS³, Scott P. Commins, MD PhD⁴

Author Affiliations: ¹Division of Pediatric Allergy, Immunology, and Rheumatology, University of North Carolina School of Medicine; ²Division of Allergy, Pulmonary, and Critical Care Medicine, Vanderbilt University Medical Center; ³University of North Carolina School of Information and Library Science; ⁴Division of Rheumatology, Allergy and Immunology, University of North Carolina School of Medicine

Short Title: Infodemiology of Alpha-Gal Allergy

Corresponding Author:

Edward G.A. Iglesia, MD MPH
Division of Pediatric Allergy, Immunology, and Rheumatology
University of North Carolina School of Medicine
3300 Thurston Bldg., CB# 7280
Chapel Hill, NC 27599-7280
Telephone: (919) 962-5136 Fax: (919) 962-4421
email: edward_iglesia@med.unc.edu

Grant Support: This research was funded by NIH Awards 5T32AI007062 (EAI), R01 AI135049 (SPC), 5T32 GM007569-41 (CAS), and 1K12HS026395-01 (CAS).

Potential competing interests: Dr. Commins reports honoraria from Genentech (speaker's bureau). Dr. Iglesia, Dr. Stone, and Dr. Flaherty report no conflicts of interest.

Word count: 1086

Specific author contributions (all authors approved the final version):

Iglesia: Project conception; data collection; data analysis/interpretation; manuscript drafting; critical revision

Stone: Project conception; data interpretation; critical revision

Flaherty: Data interpretation; critical revision

Commins: Supervision; data interpretation; critical revision

Keywords: alpha-gal; food allergy; galactose- α -1,3-galactose; red meat; epidemiology; infodemiology

Clinical Implications: Internet search query data via Google Trends demonstrate a growing regional and population-level interest in alpha-gal allergy. Clinicians should be prepared to direct patients to reputable online sources, as patients are seeking this information independently. Formal epidemiologic surveillance is needed.

To the Editor:

First officially described in 2009, alpha-gal allergy is a syndrome characterized by delayed allergic symptoms to the oligosaccharide epitope galactose-alpha-1,3-galactose present in mammalian meat.¹ Also known as mammalian meat allergy or red meat allergy, this syndrome is thought to be induced by bites of certain ticks, such as the lone star tick.¹ Alpha-gal allergy is increasingly reported in the southeastern United States and in other parts of the world, including Western and Northern Europe, South and Central America, Japan, South Africa, Western Africa, Australia,¹ though its true prevalence and epidemiology are unknown and difficult to ascertain. In the absence of population-level data or systematic surveillance of a rare disease such as alpha-gal allergy, “infodemiological” methods, such as online search traffic on Google, may provide a surrogate means to detect and monitor both disease interest and occurrence, including highlighting geographic areas where a disease may be emerging.²⁻⁷ We therefore aimed to characterize regional and temporal patterns of web-based search activity for alpha-gal allergy.

We conducted an infodemiological analysis using Google Trends, an open internet search engine-based surveillance tool that provides historical and present search volume information of Google users’ search queries.⁴ Google search volume

data are available since 2004 and can be analyzed by geographic regions. Google Trends reports its results as a “relative search volume” (RSV), which ranges from 1 to 100, and normalizes the search volume relative to a specific topic’s proportion of searches on all topics examined within a specified geography and specified time period. “Search term” queries are limited to defined keywords, while “Topic” queries encompass keywords in all search terms that share the same concept across languages. Up to 5 topics can be examined simultaneously. *A priori*, we selected keywords for analysis of online queries across all Google search categories: “Alpha-gal allergy” (topic), “alpha gal allergy” (search term), “alpha gal” (search term), “meat allergy” (search term). “Chronic urticaria” (search term) was included to serve as a control search term for normalization of data, as the prevalence of this allergic disease with similar clinical features to alpha-gal allergy is believed to be stable. We also analyzed “lone star tick” (search term). Suspecting *a priori* that news reports might contribute to search activity, we entered queries into Google News to identify reports directed to a national audience that were temporally associated with sudden increases in RSV. We retrieved query data from both the United States and worldwide from January 1, 2004 to July 28, 2019. Query data were analyzed as time series and correlations among time series were conducted using Stata 14.0 (College Station, TX).

There is interest in alpha-gal allergy (topic) in 47 out of 50 U.S. states (**Figure 1A**), with the highest RSV in Arkansas (100), Kentucky (79), Tennessee (65), Virginia (59), and North Carolina (54). When examined by specific search terms (alpha gal allergy, alpha gal, meat allergy), the highest RSV is similarly concentrated in the Southeast, the lower Great Plains/Midwest, Texas, Minnesota, and New York. These

regions also had the highest RSV for “lone star tick” (search term) (**Figure 1B**). There is interest in alpha-gal allergy on 6 continents, with Sweden (100), the U.S. (87), Australia (63), Canada (31), and South Africa (19) having the highest RSV among countries (**see Figure E1 in the Online Repository**). There is an upward, non-linear trend in RSVs for all selected alpha-gal allergy search queries in the U.S. since 2012, with peaks corresponding to news reports (**Figure 2**). RSV for the lone star tick (search term) was highly correlated with RSV for alpha-gal queries (alpha-gal allergy [topic], $R^2=0.83$ [$p<0.05$]; meat allergy [search term], $R^2=0.95$, [$p<0.05$]).

Google Trends has been previously been used to detect incident occurrences of influenza and thunderstorm-related asthma,^{5,6} as well as identifying seasonal patterns of allergic rhinitis.⁷ We used Google Trends to uncover regional and temporal patterns of internet searches related to alpha-gal allergy. The regional patterns of Google searches to date support existing observations that cases of alpha-gal allergy are concentrated in the southeastern U.S. and in countries where mammalian meat allergy has been reported.¹ RSV for alpha-gal terms correlates with RSV for the lone star tick and the regional search interest for these terms correspond to lone star tick endemic areas reflecting the suspected causal association between the lone star tick and alpha-gal allergy in the U.S. In addition to regional interest, there is a growing overall interest in alpha-gal allergy over time, as reflected by increasing RSV since 2009. This is particularly salient, as it is anecdotally reported that many patients identify their diagnosis with the aid of online resources prior to clinical presentation.⁸ Growing awareness appears to have been at least partially driven by media coverage; isolated spikes in alpha-gal specific RSV corresponded with discrete news reports (**Figure 2**). It

is unclear if increased public awareness is also driven by increased diagnosed cases or vice versa. In one clinical example, Pattanaik et al. have partly attributed the recognition of alpha-gal allergy as the leading identified cause of anaphylaxis in one southeastern U.S. emergency department to growing provider awareness of this condition.⁹

We acknowledge important limitations to this analysis. Alpha-gal-specific RSV has accelerated over time, but absolute Google search volume numbers are not publicly available. We also do not know what led internet users to inquire into alpha-gal allergy, and their demographic characteristics are unknown. There is potentially a bias toward seeking information on the lone star tick when also querying alpha-gal allergy, as the association between the lone star tick and alpha-gal allergy is well known. It is possible that increased overall internet usage could possibly account for the increase in alpha-gal allergy search volume, but we think it is unlikely given that searches for our control term of chronic urticaria were only slightly increased over the same time period. Nonetheless, there has been a substantial relative increase in alpha-gal allergy search interest, though such data only provide preliminary evidence of emerging health problems and more formal epidemiologic methods are needed to estimate disease incidence and prevalence.

In conclusion, Google search query data reflect a growing regional and population-level interest in alpha-gal allergy. By characterizing health-information seeking behaviors on the internet, Google Trends and other infodemiological methods may provide clues for diseases that are increasingly being recognized or whose epidemiology expresses a geographic pattern. Clinicians should be prepared to direct

123 patients to reputable web-based resources regarding alpha-gal allergy and
124 management, as they are already seeking this information independently. Systematic
125 case reporting and population-based surveillance methods are needed to understand
126 the true epidemiology of this emerging allergic disease.

References

1. Steinke JW, Platts-Mills TA, Commins SP. The alpha-gal story: lessons learned from connecting the dots. *J Allergy Clin Immunol.* 2015;135(3):589-96. doi: 10.1016/j.jaci.2014.12.1947.
2. Brownstein JS, Freifeld CC, Madoff LC. Digital disease detection--harnessing the Web for public health surveillance. *N Engl J Med.* 2009;360(21):2153-5. doi: 10.1056/NEJMp0900702.
3. Eysenbach G. Infodemiology and infoveillance: framework for an emerging set of public health informatics methods to analyze search, communication and publication behavior on the Internet. *J Med Internet Res.* 2009;11(1):e11. doi: 10.2196/jmir.1157.
4. Mavragani A, Ochoa G. Google Trends in Infodemiology and Infoveillance: Methodology Framework. *JMIR Public Health Surveill.* 2019;5(2):e13439. doi: 10.2196/13439.
5. Ginsberg J, Mohebbi MH, Patel RS, Brammer L, Smolinski MS, Brilliant L. Detecting influenza epidemics using search engine query data. *Nature.* 2009;457(7232):1012-4. doi: 10.1038/srep46469.
6. Bousquet J, O'Hehir RE, Anto JM, D'Amato G, Mosges R, Hellings PW, et al. Assessment of thunderstorm-induced asthma using Google Trends. *J Allergy Clin Immunol.* 2017;140(3):891-3 e7. doi: 10.1016/j.jaci.2017.04.042.

- 146 7. Kang MG, Song WJ, Choi S, Kim H, Ha H, Kim SH, et al. Google unveils a
147 glimpse of allergic rhinitis in the real world. *Allergy*. 2015;70(1):124-8. doi:
148 10.1111/all.12528.
- 149 8. Flaherty MG, Kaplan SJ, Jerath MR. Diagnosis of Life-Threatening Alpha-Gal
150 Food Allergy Appears to Be Patient Driven. *J Prim Care Community Health*.
151 2017;8(4):345-8. doi: 10.1177/2150131917705714.
- 152 9. Pattanaik D, Lieberman P, Lieberman J, Pongdee T, Keene AT. The changing
153 face of anaphylaxis in adults and adolescents. *Ann Allergy Asthma Immunol*.
154 2018;121(5):594-7. doi: 10.1016/j.anai.2018.07.017.

Figure Legends

Figure 1. Heat maps of relative search volume in the U.S., 2004 – 2019. (A) Alpha-gal allergy (topic). Highest RSV was in the following states: Arkansas, Kentucky, Tennessee, Virginia, North Carolina, and Missouri. (B) Lone star tick (search term). Highest RSV was in the following states: Arkansas, Maine, Tennessee, Kentucky, Delaware, Virginia, Missouri, and Kansas.

Figure 2. Relative search volume of alpha-gal allergy queries in the U.S., 2004 – 2019. Increasing relative search volume over time for alpha-gal allergy search queries. News media coverage coincides with search query peaks. A selected sampling of news reports are shown for clarity.

