Accepted Article

Title

Frequency of food allergy in school-aged children in eight European countries – the EuroPrevall-iFAAM birth cohort

Short title

Frequency of food allergy in European school children

Authors

Grabenhenrich L

Trendelenburg V

Bellach J

Yürek S

Reich A

Fiandor A

Rivero D

Sigurdardottir ST

Clausen M

Papadopoulos NG

Xepapadaki P

Sprikkelman AB

Dontje B

Roberts G

Grimshaw K

Kowalski ML

Kurowski M

Dubakiene R

Rudzeviciene O

Fernández-Rivas M

Couch P

Versteeg SA

van Ree R

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/ALL.14290
This article is protected by copyright. All rights reserved

Mills ENC
Keil T
Beyer K

Grabenhenrich L, Department for Methodology and Research Infrastructure, Robert Koch-Institut, Berlin, Germany and Department of Dermatology, Venerology and Allergology, Charité - Universitätsmedizin Berlin, Berlin, Germany
Trendelenburg V, Department of Paediatric Pneumology, Immunology and Intensive Care, Charité - Universitätsmedizin Berlin, Berlin, Germany
Bellach J, Department of Psychiatry and Psychotherapy, Ernst von Bergmann Academic Educational Hospital and Department of Paediatric Pneumology, Immunology and Intensive Care, Charité - Universitätsmedizin Berlin, Berlin, Germany
Yürek S, Department of Paediatric Pneumology, Immunology and Intensive Care, Charité - Universitätsmedizin Berlin, Berlin, Germany
Reich A, Epidemiology Unit, German Rheumatism Research Center, a Leibniz Institute, Berlin, Germany
Fiandor A, Department of Allergy, Hospital Universitario La Paz, Madrid, Spain
Rivero D, Department of Allergy, Hospital Universitario La Paz, Madrid, Spain
Sigurdardottir ST, Department of Immunology, Landspitali University Hospital, Reykjavik, Iceland and Faculty of Medicine, University of Iceland, Reykjavik, Iceland
Clausen M, Children's Hospital Reykjavik and Department of Allergy, Landspitali University Hospital and Faculty of Medicine, University of Iceland, Reykjavik, Iceland
Papadopoulos NG, Division of Infection, Immunity & Respiratory Medicine, University of Manchester, Manchester, United Kingdom and Allergy Department, 2nd Paediatric Clinic, National and Kapodistrian University of Athens, Athens, Greece
Xepapadaki P, Allergy Department, 2nd Paediatric Clinic, National and Kapodistrian University of Athens, Athens, Greece
Sprikkelman AB, Department of Pediatric Pulmonology & Pediatric Allergology, University Medical Center Groningen, Groningen, The Netherlands
Dontje B, Department of Pediatric Pulmonology & Pediatric Allergology, Emma Kinderziekenhuis AMC, Amsterdam, The Netherlands
Roberts G, Clinical and Experimental Sciences and Human Development and Health, Faculty of Medicine, University of Southampton, Southampton, United Kingdom and NIHR Southampton Biomedical

Accepted Article

This article is protected by copyright. All rights reserved

Research Centre, University Hospital Southampton NHS Foundation Trust, Southampton, United Kingdom

Grimshaw KEC, Clinical and Experimental Sciences Academic Units, Faculty of Medicine, University of Southampton, Southampton, United Kingdom and Department of Dietetics, Salford Care Organisation, Salford, United Kingdom

Kowalski ML, Department of Immunology, Allergy and Rheumatology, Medical University, Łódź, Poland

Kurowski M, Department of Immunology, Allergy and Rheumatology, Medical University of Łódź, Łódź, Poland

Dubakiene R, Faculty of Medicine, Vilnius University, Vilnius, Lithuania

Rudzeviciene O, Clinic of Children's Diseases, Faculty of Medicine, Vilnius University, Vilnius, Lithuania

Fernández-Rivas M, Department of Allergy, Hospital Clínico San carlos, Universidad Complutense de Madrid, IdISSC, Madrid, Spain

Couch P, Centre for Health Informatics, School of Health Sciences, University of Manchester, Manchester, United Kingdom

Versteeg SA, Departments of Experimental Immunology, Academic Medical Center, Amsterdam, The Netherlands

van Ree R, Departments of Experimental Immunology and of Otorhinolaryngology, Amsterdam University Medical Centers, Amsterdam, The Netherlands

Mills ENC, Division of Infection, Immunity and Respiratory Medicine, University of Manchester, Manchester, United Kingdom

Keil T, Institute for Clinical Epidemiology and Biometry, University of Würzburg, Würzburg, Germany and Institute for Health Resort Medicine and Health Promotion, Bavarian Health and Food Safety Authority, Bad Kissingen, Germany and Institute for Social Medicine, Epidemiology and Health Economics, Charité - Universitätsmedizin Berlin, Berlin, Germany

Beyer K, Department of Paediatric Pneumology, Immunology and Intensive Care, Charité - Universitätsmedizin Berlin, Berlin, Germany

Accepted Article

This article is protected by copyright. All rights reserved

Corresponding author

Pre-publication

Linus Grabenhenrich

Department for Methodology and Research Infrastructure

Robert Koch-Institut, Berlin, Germany

GrabenhenrichL@rki.de

+49-162-2164435

Post-publication

Kirsten Beyer

Department of Paediatric Pneumology, Immunology and Intensive Care

Charité - Universitätsmedizin Berlin, Berlin, Germany

Kirsten.Beyer@charite.de

Acknowledgement

We thank all families who participated in the birth cohort study and the medical and nursing staff of the participating hospitals, especially G. Christopoulou, I. Roumpedaki, and R. Stergiou (Greece); S. E. Sigurdardottir, K. L. Bjornsdottir, H. Ragnarsdottir, and prenatal care nurses at The Primary Health Care of the Reykjavik Capital Area (Iceland); S. Paschke-Goossens, G. Schulz, A. Rohrbach, S. Tschirner, T. Schrezenmaier, A. Scholz, D. McBride (Germany); A. Stanczyk-Przyłuska, K. Zeman, J. Wilczynski, and L. Podciechowski (Poland); S. Quirce, M. Reche, M. Martin-Esteban, R. Gabriel, J. I. Larco, I. Bobolea, and T. Cuevas (Spain); K. Foote, E. Oliver, A. Selby, L. Fairhead, Z. Dobson, S. Bhatt, S. Roberts, J. Martin, and NIHR Wellcome Trust Clinical Research Facility (United Kingdom); Midwives Zorggroep Almere, N. v. d. Berg, W. M. C. van Aalderen, L. Hulshof, N. C. M. Petrus, and H. C. Vriesendorp (The Netherlands); I. Butiene, D. Vaicekauskaite, A. Arlauskiene, J. Zakareviciene, and L. Stoskute-Malinauskiene (Lithuania).

Contributions

ENCM was overall coordinator of the collaborative research initiatives EuroPrevall and iFAAM and supervised food challenge meals; KB was initiator, principal investigator and iFAAM theme leader of the birth cohort study; TK was co-PI and iFAAM work package leader of the birth cohort; LG coordinated the iFAAM school-age follow-up of the birth cohort, carried out statistical analyses, and prepared the

This article is protected by copyright. All rights reserved

Accepted Article

manuscript; RvR was responsible for laboratory analyses for the whole project (data not used in this manuscript) and participated in the planning of the study design; SV conducted the laboratory analyses; MFR was responsible for laboratory analyses including skin prick tests and participated in the planning of the study design; PC was responsible for all central IT aspects, and participated in the planning of the school-age follow-up; AR was responsible for the central data management, participated in the cohort management and planning of the study; all authors participated in the planning and local implementation of the follow-up assessment including the clinical visits; all authors reviewed and commented the draft of the manuscript and approved the final version.

Accepted Article

This article is protected by copyright. All rights reserved

Abstract

Background. The prevalence of food allergy (FA) among European school children is poorly defined. Estimates have commonly been based on parent-reported symptoms. We aimed to estimate the frequency of FA and sensitization against food allergens in primary school children in eight European countries.

Methods. A follow-up assessment at age 6-10 years of a multi-centre European birth cohort, based was undertaken using an online parental questionnaire, clinical visits including structured interviews and skin prick tests (SPT). Children with suspected FA were scheduled for double-blind, placebo-controlled oral food challenges (DBPCFC).

Results. 6,105 children participated in in this school-age follow-up (57.8% of 10,563 recruited at birth). For 982 of 6,069 children (16.2%), parents reported adverse reactions after food consumption in the online questionnaire. Of 2,288 children with parental face-to-face interviews and/or skin prick testing, 238 (10.4%) were eligible for a DBPCFC. 63 foods were challenge-tested in 46 children. 20 food challenges were positive in 17 children, including seven to hazelnut and three to peanut. Another seventy-one children were estimated to suffer FA among those who were eligible but refused DBPCFC. This yielded prevalence estimates for FA in school-age between 1.4% (88 related to all 6,105 participants of this follow-up) and 3.8% (88 related to 2,289 with completed eligibility assessment).

Interpretation. In primary school children in eight European countries, the prevalence of FA was lower than expected even though parents of this cohort have become especially aware of allergic reactions to food. There was moderate variation between centres hampering valid regional comparisons.

Accepted Article

This article is protected by copyright. All rights reserved

Introduction

Validly measuring the frequency of food allergy (FA) in the general population has been challenging, mainly due to the disease being very heterogeneous in terms of eliciting food allergens and clinical signs and symptoms. Furthermore, it is difficult to differentiate FA from other food hypersensitivities, such as food intolerances, in the general population. The different study settings and designs, case definitions, individual interpretations of the medical history including observed appearances of FA and varying consumption habits have hampered sound comparisons of FA prevalence between research projects, geographic regions and time trends.[1-3] For example, studies aiming to estimate the prevalence of FA have applied different assessment techniques from questionnaires to double-blind placebo-controlled food challenges and have arrived at profoundly different estimates.[4-9] A cross-sectional study in Germany targeting children from birth to 17 years reported a prevalence of FA of 61.6% based on self- and parent-reported information. Subsequently, suspected cases of FA were examined clinically with food challenge tests, giving an estimated prevalence of confirmed FA of 2.2%.[7] In a Danish sample including children and adults aged 4 to 22 years, the prevalence of FA was estimated at 1.0% and 0.3% based on open (i.e. non-blinded) and double-blind, placebo-controlled food challenges (DBPCFC), respectively.[5] The prevalence of self-reported FA assessed in primary school-age children was estimated at almost 6% in Turkey, and 6% to 12% in the United Kingdom. However, using DBPCFC led to prevalence estimates of 0.7% to 1.4% in these studies.[4,8,10,11]

The methodological challenges of assessing reactions to foods have prompted the development of standards for the diagnosis of FA, mainly in clinical settings.[12] The birth cohort study from EuroPrevall (The prevalence, cost and basis of food allergy in eight European countries) agreed on a harmonized approach in all centres to confirm suspected FA using the diagnostic clinical gold-standard, i.e. DBPCFC, stringently in a large scale population-based study.[13,14] Based on PRACTALL recommendations, the documentation and interpretation of oral food challenges has been developed further for the school-age follow-up of the birth cohort.[12,15]

In this manuscript we report the range of frequencies of challenge-confirmed FA and sensitization against food allergens in primary school children from eight countries covering different European regions.

Accepted Article

This article is protected by copyright. All rights reserved

Methods

Study design and setting. Starting in 2005, the EuroPrevall birth cohort set out to recruit newborns from nine European countries, to prospectively trace the onset of food allergy (FA) from birth to 2.5 years.[14,16,17] Within the EU-funded iFAAM project (Integrated approaches to food allergen and allergy management), eight of the nine study centres (Iceland, United Kingdom, The Netherlands, Germany, Poland, Lithuania, Spain, and Greece) took part in a single follow-up assessment at early school age (6 to 10 years, between 2014 and 2017). This follow-up aimed to reach all children initially recruited at birth (10,563), and document any previous parent-reported reactions to food as well as the current FA status. Ethical approval was obtained separately for all participating country, as listed above.

Participants and sample definitions (denominators). Recruitment details have previously been described.[14] All recruited children of the EuroPrevall birth cohort (excluding the study centre in Italy) were re-invited for the current follow-up by invitation letters, electronic mail, and/or telephone calls, for up to seven approaches, as required.

Parents were asked to complete an online questionnaire at home, at the same time providing consent to participate in the follow-up. A very limited number of questionnaires were completed in the study centre or via telephone interviews, e.g. in cases where parents were unable or unwilling to access/complete the questionnaire by themselves. The questionnaire included items about previous reactions to food. Questionnaire data was reported for those with completed FA screening questions and the consumption history for a selection of commonly allergenic foods.

All children, irrespective of their FA history, were invited for a clinical visit to the local study centre, including a face-to-face interview on previous and current food reactions, consumption habits, and skin prick test (SPT) to a pre-defined panel of foods and aeroallergens, as well as foods reported to have previously caused reactions. The eligibility for one or more oral food challenges was defined using an algorithm based on interview data and SPT results (Figure 2). All forms and the diagnostic triage have been previously described.[15] Study outcomes based on interview data are reported for those who completed the clinical interview and have documented challenge eligibility.

Data sources and variables (case definitions, numerators). All outcome data reported in this manuscript were collected within the school-age follow-up of the birth cohort. Reports on previous or current reactions to food were derived from a single screening question (Q1, Figure 2), further differentiated by physician's diagnosed FA, challenge-proven FA, symptoms by organ system, exposure-symptom interval,

Accepted Article

This article is protected by copyright. All rights reserved

age at first occurrence (all Q3), tolerance development (Q4) and recent consumption (Q2, previous 3 months). Both the online-questionnaire and the clinical face-to-face interview covered all these aspects.[15]

The current FA status (period/current prevalence of the potential to react if exposed) was defined for a selected list of foods (termed core foods): cow's milk, hen's egg, wheat, soy, peanut, hazelnut, white and oily fish and crustacean shell-fish. The consumption history for non-core foods was not assessed for children who never reported problems to a specific food.

A multi-level outcome assessment for the likelihood of current FA was derived from the questionnaire/interview data using a decision algorithm.[15] This was further differentiated by details of recent symptoms, and avoidance behaviour. Food challenge eligibility was derived from the likelihood of current FA, complemented by SPT results i.e. individuals not currently consuming a suspected food with either previous symptoms following consumption of that food and/or a positive SPT (valid controls and largest wheal diameter ≥3mm, Figure 2) to that food.

Assessment by DBPCFC tests. These were conducted based on previously published methodology,[12] and documented and interpreted as previously described.[15] In brief, an escalating seven-dose protocol was followed to challenge children suspected to have FA. Children, families and medical staff were fully blinded to the order of the food/placebo given. Placebo days may have served as controls for more than one food, with each food/placebo tested on a separate day.

Assessment of differential non-response. With the expectedly high attrition, we used several sources of information to assess differential non-response at school age, aiming to cover important characteristics in question to impact and/or predict allergy development. From the baseline assessment of the birth cohort we included the following: the child's sex, delivery mode, season of birth, use of antibiotics in the first week of life, older siblings, mother's current and smoking during pregnancy, cat and dog in household, parental allergies, child's eczema (parental report of eczema symptoms) and child's FA (previously proven by DBPCFC within the study). From the school-age follow-up we included common atopy-associated diseases (asthma, allergic rhinitis and eczema), FA outcomes as described above, food-specific consumption habits and SPT results.

Statistics and Software. Study data was entered via a web interface, either by parents (questionnaire) or by study personnel (all other forms). The server architecture was specifically designed for data capturing in this project and to track completeness and congruency of the follow-up assessment. Cleaning and statistical analyses were performed using SAS 9.4 (SAS Institute Inc., Cary, NC, USA). The initial sample size was set to assess the incidence of FA up to age 2 years, as described before.[14] Outcome measures are

This article is protected by copyright. All rights reserved

reported by study centre, eliciting food and assessment method (questionnaire vs. interview). Raw frequencies were calculated as fractions of case definitions over corresponding samples, as described in the above subsections. Missing single items, besides those used to define sample membership, did not lead to the exclusion of the form or participant.

The potential for differential loss to follow-up was assessed comparing baseline characteristics and available outcome measures between groups of participants attending different numbers of follow-up assessments: (1) lost to follow-up, (2) questionnaire only, (3) eligible but not challenged, (4) eligible and challenged, and (5) not eligible, using group-to-group comparisons with a Chi-squared statistic. Comparing the characteristics in three different group-to-group assessments, comparisons with a p-value below 0.001 are highlighted. However, dichotomous test results were not used to guide the extrapolation strategy. As a manual weighting approach, the outcome frequencies in groups which were assessed were used as substitutes for groups where data for this outcome was not available. As an example, groups (3) and (4) were assumed to be quite similar, so the relative frequency of challenge-proven FA in those who completed the challenges (group (4)) was used to estimate the absolute number of potentially food allergic children in those who were not challenged. With the expectedly large differences between point estimates introduced by different but all justifiable sets of assumptions, we report ranges of frequency estimates. Confidence intervals for proportion estimates were calculated assuming that errors follow the beta distribution, and using sample sizes of actually assessed children for the respective outcome.

Role of the funding source

The birth cohort study was funded by the European Commission: (i) under the 6th Framework Programme (FOOD-CT-2005-514000) within the collaborative research initiative 'EuroPrevall', and (ii) under the 7th Framework Programme (FP7-KBBE-2012-6; grant agreement no. 312147) within the collaborative project 'iFAAM'. Additional funds were received by the Icelandic birth cohort centre from Landspitali University Hospital Iceland Science Fund, and from GlaxoSmithKline Iceland; by the United Kingdom birth cohort centre from the UK Food Standards Agency; by the Polish birth cohort center from the Ministry of Science and Higher education, by the Lithuanian birth cohort centre as unrestricted grants from Grida and MSD; and by the Dutch birth cohort centre as unrestricted grants from Nutricia Advanced Medical Nutrition Netherlands, AstraZeneca Netherlands, TEVA Netherlands, and GlaxoSmithKline Netherlands. None of the funding bodies had any influence on the study design, the collection and analysis of data, interpretation of results, manuscript preparation, or decision to submit the paper for publication.

Accepted Article

This article is protected by copyright. All rights reserved

Results

Participants. Parents completed the online-questionnaire for 6,069 out of the 10,563 children recruited at birth (57.5%) at age 6-10 years. Of these 38.3% (n=2,322) of those came to the study centres for the face-to-face interview and physical examination including skin prick testing (SPT), which was carried out in 2,188 participants (Figure 1). Questionnaire response differed by centre, ranging from 39.8% of those recruited in Southampton, UK to 70.5% in Reykjavik, Iceland (Table 1). The mean age at follow-up was 8.3 years (standard deviation 0.9). This report therefore covers a total observation period of 50,733 person-years. The online-questionnaire and the face-to-face interview allowed estimation of the frequency symptoms after food consumption until school age based on parental reports.

Lifetime prevalence of symptoms after food consumption and physician's diagnosis of food allergy reported by the parents. Of the 6,069 children who completed the online questionnaire, 982 (16.2% [15.3-17.1]) had previous adverse reactions after food consumption. Physician's diagnosis of FA was less frequent when compared to symptoms alone for all countries, both based on parent report. Cow's milk, hen's egg, peanut, hazelnut, and wheat were the five most commonly implicated food items as causing symptoms (Table 1. e-Table 2). Numbers were similar for the 2,322 children who came to the study centre for the face-to-face interview (e-Table 1). The majority (85.0 %) of the reported reactions occurred <2 hours after food consumption. Skin symptoms such as rash or pruritus (itching) and gastrointestinal reactions such as diarrhoea were the most frequently reported symptoms in all countries (e-Table 3, e-Table 4).

Food consumption and avoidance at school age. Based on 6,069 online-questionnaires almost all children consumed foods containing cow's milk, hen's egg and wheat. Among the "core foods", soy and crustacean shell-fish were consumed less often by school-aged children. The reason for not consuming a particular food was usually because that food was not part of the regular diet of the family, as assessed in the face-to-face interview. The foods most often avoided to prevent the development of FA were peanut and hazelnut (e-Table 5).

Food allergy at school age – patient history. 2,289 children completed the eligibility assessment, based on reaction history (face-to-face interview) and complemented with allergic sensitization status based on skin prick tests (Figure 1). The decision tree to define current FA status was applied to all core foods separately, e.g. for peanut, for 57 of 2,322 children's parents reported allergic symptoms after peanut consumption (Figure 2, Question 1 "yes"). If children either did not become tolerant as described in the patient history (Question 4 "no/don't know", n=40) or became tolerant but did not consume the food

This article is protected by copyright. All rights reserved

171 without symptoms recently (previous 3 months, Question 2 “no”, n=4), they were assessed further as
172 likely food allergic. The majority (2,265) of children had never had symptoms upon peanut consumption
173 (Figure 2, Question 1 “no”), and if peanuts were consumed in the previous 3 months without symptoms
174 (Question 2 “yes”, n=1,863), the child was not eligible for a challenge (Type A), i.e. not peanut allergic on
175 the basis of history alone. In children who did not consume peanuts in the last 3 months (n=402) FA had
176 to be considered possible as reactions upon exposure could not be ruled out (Figure 2, Table 2).

177 *Food allergy at school age - Skin Prick Test (SPT)*. A total of 223 of 2,188 children (10.2% [9.0-11.5]) had a
178 positive (≥3 mm) SPT to one or more “core foods”. Sensitization to peanut (5.6%) and hazelnut (5.2%) was
179 most frequent (Figure 3).

180 *Food allergy at school age - DBPCFC*. Of 2,289 children for which eligibility was documented, 238 (10.4%
181 [9.2-11.7]) were eligible for DBPCFC, either because they were likely to react based on the face-to-face
182 interview (e.g. 42 children for cow’s milk and 34 for hen’s egg) or because they had not consumed the
183 suspected food within the last 3 months and were sensitized to it (four children for cow’s milk and six for
184 hen’s egg). The latter was common for hazelnut and peanut with 42 and 36 children, respectively
185 (Table 2). In total, we performed 63 DBPCFC in 46 children, most often to assess hazelnut and peanut (16
186 and 15 challenges, respectively). The parents of 192 participants refused the DBPCFC (Figure 1).

187 Twenty DBPCFC days where the incriminated food was given to 17 children were rated positive (two with
188 less pronounced symptoms on the placebo test day, and three without a placebo test day as the parents
189 refused the placebo day after a clear positive reaction on the previous day where the incriminated food
190 was given). Seven placebo tests were rated as positive out of all placebo challenges conducted in this
191 study.

192 The 17 of 2,097 (0.8% [0.5-1.3]) completely assessed children with a positive DBPCFC included seven
193 children who reacted to hazelnut (0.3% [0.1-0.7]) and three to peanut (0.1% [0.0-0.4]). Only one child
194 reacted to hen’s egg (seven were challenged) and none to cow’s milk (six were challenged, Table 2).
195 Parents of all seven hazelnut-allergic children reported nasal symptoms in their children in the previous 12
196 months and two children had a physician’s diagnosed allergic rhinitis. Among non-core foods, cashew (2),
197 pine nut (2) and walnut (1) had positive DBPCFC tests (e-Table 6).

198 *Differential attrition.* Frequency of FA within the whole cohort sample was extrapolated as not all
199 participants completed all the necessary diagnostic steps to confirm or rule out current (prevalent) FA in
200 school-age. Children who did not take part in the school-age follow-up (group 1) came more often from
201 less educated families but were quite similar in terms of allergic family predisposition, eczema in infancy

Accepted Article

This article is protected by copyright. All rights reserved

and early childhood and other factors to those whose parents responded to the questionnaire but refused the clinical assessment of their child (group 2). Therefore, frequencies of FA in group 2 were used as substitutes for those not participating in the school-age follow-up (data and group labels in Table 3).

Children who came for a clinical visit, but were not eligible for a food challenge (group 5) had more parent-reported allergic rhinitis, eczema and more often an allergic family history than those only responding to the questionnaire (group 2). They were similar with regard to other potential indicators of FA such as sex and caesarean section (Table 3). Therefore, using FA frequencies from group 5 as substitutes for those not showing up for the clinical visit may yield only slightly upwards biased estimates.

Those children eligible, but whose parents refused the oral challenge testing (group 3), were quite similar to those eligible and successfully challenged (group 4). The detailed comparison of these two groups in terms of food-specific sensitization, current symptoms (12m), previous FA diagnosis, and specific food avoidance behaviour showed also very similar distributions. Specifically, the proportion of SPT positivity was similar in groups 3 and 4, with a considerable difference only in hen's egg sensitization (Table 4 and eFigure 1). Thus, FA frequency in group 4 was used to estimate the number of potential food allergic children in group 3 (Table 3).

Adjusted frequency of food allergy. A total of 238 children were eligible for an oral food challenge, of which 46 underwent a DBPCFC (group 4). In 192 children parents refused this diagnostic step (group 3). In addition to the 17 positively challenged children, 71 children were estimated to have FA in the group eligible but refusing the challenge procedure under the assumption of identical FA prevalence in those challenged and those eligible but not challenged (Tables 3 and 4), summing up to an estimated number of 88 cases (rounding for the all-country estimator). These 88 cases would lead to an extrapolated prevalence estimate of 3.8% in the group of all 2,289 children who completed the eligibility assessment. Under the assumption that all who participated in this follow-up but were not eligible for a food challenge had no FA, these estimated 88 cases would extrapolate to a school-age lower limit prevalence of current FA of 1.4% (of 6,105 children, Figure 4). Furthermore, considering the high similarity between those who participated only in the questionnaire assessment and those who were lost to follow-up by school-age, a lower-limit prevalence of current FA for the whole cohort would be 0.8% (88 of 10,563). Note that these estimates rely on a simple group-wise extrapolation approach only.

Accepted Article

This article is protected by copyright. All rights reserved

Discussion

Key results. In 17 of 2,097 completely assessed school age children from this European birth cohort study, food allergy (FA) to at least one allergen source was confirmed, yielding an average raw prevalence of 0.8% across all eight countries involved, as a lower limit estimate. Extrapolating to all children who completed the eligibility assessment, we estimated an adjusted FA prevalence between 1.4% (extrapolated to all children with questionnaire data) and 3.8% (extrapolated to those with completed eligibility assessment). Most of the positively challenged children reacted only mildly or moderately, except for five children with severe signs or symptoms during DBPCFC. However, more severe reactions might have been observed if those eligible but refusing to undergo the challenge were actually challenged. The most common allergens responsible for FA in school-age and for allergic sensitization assessed by skin tests were hazelnut and peanut.

The low absolute numbers per study centre hampered valid regional comparisons.

Comparison with other studies. Previous prevalence surveys were conducted only as single centre studies, and used different approaches hampering inter-country comparisons. As our measures were highly standardized, this is the first European multicentre study to report comparable estimates for the prevalence of FA in primary school-age. Results of our study were in line with the few previous studies of children with comparable age from Turkey with a prevalence of 0.7% at 6-9 years,[4] and 1.4% for 6,[8] and 1.3% for 10 year olds from the UK[10] though these studies did not report adjusted estimates. However, the MAAS study in the UK reported peanut allergy prevalence alone of about 2% at age 8 years of age.[11] The prevalence of FA in the HealthNuts study in Australia was estimated higher at 3.8% but in pre-school children at 4 years of age; however early life FAs against cow's milk and hen's egg are still more prevalent than among school-age children who have mostly become tolerant against these foods.[18] Compared to the previous studies from Europe we did not find an indication for an increase in the prevalence of FA in school-age.

Strengths and Limitations. Our birth cohort study that has been conducted in various European regions is the largest and first multi-national population-based investigation of DBPCFC-confirmed FA in childhood worldwide. However, the generalizability of the initial sample to the whole (regional or whole-country) populations has not been formally assessed. Due to a stringent, standardized, clinically-based methodology, it allows better comparisons of confirmed FA occurrence and influential factors in eight European countries than previous single-centre/-country studies. The diagnostic work-up included several possible approaches to identify suspected FA, also capturing measures of disease severity and impact. It

This article is protected by copyright. All rights reserved

261 also focussed on the need to thoroughly adjust for the different types of non-response, particularly the
262 refusal of a clinically indicated DBPCFC.[19]

263 The local consumption habits influence the likelihood that a dormant potential to react to a certain
264 allergen presents as an apparent reaction. This limits the comparability, e.g. of different regions and
265 parent-reported symptoms. We focussed the standardized assessment of consumption and sensitization
266 to a core list of major allergen sources and only for these foods the current prevalence estimates yield
267 valid results. For all other foods, for which the consumption history and sensitization assessments were
268 only available for children whose families reported problems ever, we cannot present valid denominators
269 for measures of disease frequency. We cannot exclude allergy to these foods in participants who may not
270 be consuming them. Furthermore, groups of foods containing similar (potentially allergenic) food
271 allergens, e.g., cow's milk in bakery products, would need to be differentiated in more detail. Including
272 more and more foods in the consumption and sensitization assessment would be likely to increase the
273 number of identified yet unnoticed food allergies and thus increase the estimated frequencies.

274 The overall participation in the questionnaire assessment was comparable to other birth cohorts on
275 allergy in Europe. There was considerable attrition in all stages of the assessment, form the questionnaire
276 and clinical assessment to the final diagnostic step. We thoroughly assessed the potential for differential
277 non-response and weighted after investigating a large set of background characteristics, potentially giving
278 a range of estimates that closer reflect the actual prevalence in the whole population, including those not
279 assessed. Compared to other one-time surveys, this prevalence assessment took place in a sample of
280 children recruited at birth and followed until 30 months and later until school-age for incident FA. This
281 sample's families are likely to be more aware of this specific disease and also far more examined and
282 assessed during infancy in search for food reactions.

283 A major weakness of the study was that a considerable number of families did not participate in the food
284 challenges, which would have been necessary (based on the predefined eligibility criteria) to confirm or
285 rule out FA. The willingness to undergo a time-consuming and stressful food challenge of at least two days
286 is much lower without the (felt) burden of the affected child and its family, usually the key trigger to
287 consent to this procedure in regular care.

288 In one approach, we extrapolated the proportion of food allergic children of all challenged to those
289 children who refused to undergo the procedure. There are many possible reasons for such refusal. Our
290 prevalence estimates would be too low if severe reactions and previously challenge-proven food allergy
291 were the main reason for the parents' hesitancy to allow food challenge. Thus, we reported a range of

Accepted Article

This article is protected by copyright. All rights reserved

frequency estimates reflecting several probably extreme but justifiable sets of assumptions. Providing a single and robust point estimate supported by a reliable measure of precision would require a close to complete assessment of a population-based sample. As the uncertainty due to other factors than random error are likely to outweigh the potential variation due to chance/sampling, we explicitly refrained from calculating confidence intervals (which mainly address random error) for extrapolated frequencies.

Conclusions. The occurrence of FA in European school children was estimated between 1-4%, depending on the strategy for weighting attrition, which was considerably lower than suspected from information based on parental reports. The prevalence of food allergy of the children who only completed the online questionnaire could only be estimated. Assuming they had the same prevalence of food allergy than those seen in the study centre likely overestimated the true frequency at 3.8%, whereas assuming they are all non-allergic certainly yields too low estimates (1.4%). The true prevalence is probably within this interval. The most common allergens responsible for FA and sensitization were peanut and hazelnut.

Accepted Article

This article is protected by copyright. All rights reserved

Declaration of interests

All authors report EU grant support as described under Role of the funding source; M Fernandez-Rivas reports grants from the Spanish government (MINECO, ISCIII), grants and personal fees from Aimmune Therapeutics, personal fees from Allergy Therapeutics, Fundación SEAIC, HAL, Thermo Fisher Scientific, Schreiber Foods, DBV, outside the submitted work; and patent PCT/ES2014/070634 issued; NG Papadopoulos reports grants from Gerolymatos, personal fees from Hal Allergy B.V., Novartis Pharma AG, Menarini, Hal Allergy B.V., Mylan, outside the submitted work; D Rivero reports employment as a MSL (medical science liaison) for genetic diseases at Shire pharmaceuticals; ST Sigurdardottir reports personal fees from Icepharma pharmaceutical company, outside the submitted work; V Xepapadaki reports personal fees from Uriach, Novartis, Nestlé, Nutricia, outside the submitted work; R van Ree reports personal fees from HAL Allergy BV, Citeq BV, ThermoFisher Scientific, outside the submitted work; ENC Mills reports grants from Reacta Biotech Ltd, outside the submitted work; and Chief Scientific Adviser and shareholder of Reacta Biotech Ltd, a start-up developed to commercialise foods for use in oral food challenges; K Beyer reports grants from European Commission, during the conduct of the study; grants and personal fees from Aimmune, Danone, Infectopharm, personal fees from Bencard, Hycor, Mabylon AG, Mylan, Nestle and grants from ALK, DBV, Hipp, outside the submitted work. All other authors have nothing to disclose.

Accepted Article

This article is protected by copyright. All rights reserved

Accepted Article

Tables

Table 1

Response at school age and lifetime prevalence (raw % and n) of parent-reported symptoms after food consumption and parent-reported physician diagnosed food allergy, based on online questionnaire (n=6,069).

Symptoms after consumption of "core foods" were asked specifically, all other foods are labelled "non-core foods".

*Only the most frequently reported non-core foods are listed here, rarely reported non-core foods are presented in e-Table 2.

	Reykjavik Iceland	Southampton UK	Amsterdam NL	Berlin Germany	Lodz Poland	Vilnius Lithuania	Madrid Spain	Athens Greece	Total
Baseline [n]	**1,341**	**1,140**	**976**	**1,570**	**1,513**	**1,556**	**1,387**	**1,080**	**10,563**
Response [%]	70.5	39.8	66.8	63.8	54.1	61.0	49.6	51.9	57.5
	945	**454**	**652**	**1,001**	**819**	**949**	**688**	**561**	**6,069**
n	**% (n)**	**% (n)**	**% (n)**	**% (n)**	**% (n)**	**% (n)**	**% (n)**	**% (n)**	**% (n)**
Parent-reported symptoms									
Any food	**20.1 (190)**	**20.7 (94)**	**19.8 (129)**	**17.2 (172)**	**22.0 (180)**	**12.3 (117)**	**11.2 (77)**	**4.1 (23)**	**16.2 (982)**
"Core foods"	**18.0 (170)**	**14.3 (65)**	**15.5 (101)**	**12.0 (120)**	**16.5 (135)**	**7.9 (75)**	**7.8 (54)**	**3.0 (17)**	**12.1 (737)**
Cow's milk	14.1 (133)	8.8 (40)	11.8 (77)	8.2 (82)	12.8 (105)	4.5 (43)	2.9 (20)	1.2 (7)	8.4 (507)
Hen's egg	3.8 (36)	4.2 (19)	4.1 (27)	1.9 (19)	5.5 (45)	3.1 (29)	3.5 (24)	0.7 (4)	3.3 (203)
Wheat	1.6 (15)	3.3 (15)	0.8 (5)	1.0 (10)	1.2 (10)	1.8 (17)	1.2 (8)	0.0 (0)	1.3 (80)
Soy	1.0 (9)	1.3 (6)	0.6 (4)	0.2 (2)	0.6 (5)	0.5 (5)	0.1 (1)	0.0 (0)	0.5 (32)
Peanut	1.6 (15)	2.2 (10)	2.0 (13)	1.4 (14)	2.4 (20)	1.1 (10)	1.6 (11)	0.5 (3)	1.6 (96)
Hazelnut	1.0 (9)	0.9 (4)	1.1 (7)	1.9 (19)	3.5 (29)	1.3 (12)	1.2 (8)	0.2 (1)	1.5 (89)
White fish	1.8 (17)	0.4 (2)	0.6 (4)	0.1 (1)	0.6 (5)	0.6 (6)	1.0 (7)	0.9 (5)	0.8 (47)
Oily fish	0.8 (8)	0.2 (1)	0.0 (0)	0.4 (4)	0.2 (2)	0.4 (4)	0.7 (5)	0.5 (3)	0.4 (27)
Crustaceans	1.0 (9)	0.7 (3)	0.6 (4)	0.1 (1)	0.0 (0)	0.5 (5)	0.4 (3)	0.4 (2)	0.4 (27)
"Non-core foods"*	**5.6 (53)**	**9.7 (44)**	**6.3 (41)**	**8.7 (87)**	**10.7 (88)**	**6.5 (62)**	**5.7 (39)**	**1.8 (10)**	**7.0 (424)**
Nuts (except hazelnut)	0.5 (5)	0.7 (3)	0.9 (6)	0.8 (8)	0.1 (1)	0.1 (1)	2.0 (14)	0.7 (4)	0.7 (42)
Tomato	0.3 (3)	1.8 (8)	1.2 (8)	0.5 (5)	0.7 (6)	0.3 (3)	0.0 (0)	0.0 (0)	0.5 (33)
Kiwi	0.3 (3)	0.4 (2)	0.9 (6)	0.2 (2)	0.9 (7)	0.1 (1)	1.0 (7)	0.0 (0)	0.5 (28)
Strawberry	0.0 (0)	0.7 (3)	0.6 (4)	0.9 (9)	1.1 (9)	0.2 (2)	0.0 (0)	0.0 (0)	0.4 (27)
Apple	0.1 (1)	0.2 (1)	0.3 (2)	0.7 (7)	1.0 (8)	0.3 (3)	0.0 (0)	0.0 (0)	0.4 (22)
Citrus fruits, not specified	0.2 (2)	0.2 (1)	0.2 (1)	0.5 (5)	1.2 (10)	0.6 (6)	0.0 (0)	0.0 (0)	0.4 (25)
Parent-reported doctors-diagnosed food allergy	**8.7 (82)**	**8.8 (40)**	**10.6 (69)**	**5.1 (51)**	**14.4 (118)**	**7.2 (68)**	**7.7 (53)**	**2.5 (14)**	**8.2 (495)**
"Core foods"	**8.0 (76)**	**8.1 (37)**	**10.0 (65)**	**4.6 (46)**	**12.1 (99)**	**5.7 (54)**	**5.8 (40)**	**2.1 (12)**	**7.1 (429)**
Cow's milk	4.9 (46)	5.3 (24)	7.4 (48)	2.1 (21)	9.8 (80)	3.4 (32)	1.9 (13)	1.1 (6)	4.4 (270)

This article is protected by copyright. All rights reserved

Hen's egg	3.7 (35)	3.3 (15)	3.7 (24)	1.3 (13)	4.6 (38)	2.7 (26)	3.3 (23)	0.5 (3)	2.9 (177)
Wheat	0.4 (4)	1.1 (5)	0.6 (4)	0.4 (4)	0.7 (6)	1.6 (15)	0.6 (4)	0.0 (0)	0.7 (42)
Soy	0.8 (8)	0.9 (4)	0.3 (2)	0.2 (2)	0.6 (5)	0.5 (5)	0.1 (1)	0.0 (0)	0.4 (27)
Peanut	1.4 (13)	1.8 (8)	0.8 (5)	1.2 (12)	1.2 (10)	0.6 (6)	1.6 (11)	0.4 (2)	1.1 (67)
Hazelnut	0.6 (6)	0.7 (3)	0.5 (3)	0.8 (8)	2.7 (22)	0.8 (8)	1.0 (7)	0.2 (1)	1.0 (58)
White fish	1.4 (13)	0.0 (0)	0.3 (2)	0.1 (1)	0.2 (2)	0.6 (6)	0.7 (5)	0.9 (5)	0.6 (34)
Oily fish	0.4 (4)	0.0 (0)	0.0 (0)	0.2 (2)	0.0 (0)	0.3 (3)	0.3 (2)	0.5 (3)	0.2 (14)
Crustaceans	0.6 (6)	0.4 (2)	0.2 (1)	0.0 (0)	0.0 (0)	0.3 (3)	0.4 (3)	0.0 (0)	0.2 (15)
"Non-core foods"*	**1.8 (17)**	**2.0 (9)**	**0.9 (6)**	**1.6 (16)**	**6.0 (49)**	**2.4 (23)**	**3.8 (26)**	**0.7 (4)**	**2.5 (150)**
Nuts (except hazelnut)	0.4 (4)	0.7 (3)	0.6 (4)	0.8 (8)	0.1 (1)	0.1 (1)	1.9 (13)	0.7 (4)	0.6 (38)
Tomato	0.0 (0)	0.2 (1)	0.2 (1)	0.0 (0)	0.5 (4)	0.2 (2)	0.0 (0)	0.0 (0)	0.1 (8)
Kiwi	0.2 (2)	0.0 (0)	0.2 (1)	0.1 (1)	0.4 (3)	0.0 (0)	0.6 (4)	0.0 (0)	0.2 (11)
Strawberry	0.0 (0)	0.2 (1)	0.0 (0)	0.2 (2)	0.5 (4)	0.0 (0)	0.0 (0)	0.0 (0)	0.1 (7)
Apple	0.0 (0)	0.0 (0)	0.0 (0)	0.0 (0)	0.7 (6)	0.3 (3)	0.0 (0)	0.0 (0)	0.1 (9)
Citrus fruits, not specified	0.0 (0)	0.0 (0)	0.0 (0)	0.0 (0)	0.7 (6)	0.2 (2)	0.0 (0)	0.0 (0)	0.1 (8)

Accepted Article

This article is protected by copyright. All rights reserved

Table 2

Parent-reported history of symptoms and consumption, eligibility assessment as judged by the supervising study physician, and results of food challenge tests. Percentages not adjusted for non-response.

Number of questions Q1-Q4 referring to Figure 2.

			Cow's milk	Hen's egg	Wheat	Soy	Peanut	Hazelnut	White fish	Oily fish	Crusta-ceans	Core foods	Any food
Parent-reported history of symptoms and consumption (based on interviews, n=2,322)													
Never symptoms (Q1) + Currently consumed (Q2)		n	2,008	2,183	2,274	857	1,863	2,019	2,131	1,846	1,043	-	-
Ever symptoms (Q1) + Tolerated again (Q4) + Currently consumed (Q2)		n	202	80	13	7	13	10	9	4	2	-	-
Never symptoms (Q1) + Currently not consumed (Q2)		n	10	16	17	1,448	402	245	162	463	1,265	-	-
Ever symptoms (Q1) + Tolerated again (Q4) + Currently not consumed (Q2)		n	2	2	0	1	4	0	0	0	1	-	-
Ever symptoms (Q1) + Never tolerated (Q4)		n	100	41	18	9	40	48	20	9	11	214	290
Eligibility assessment for food challenge as judged by physician (n=2,289)													
Currently consumed	**No** (Type A) No food allergy	n	2,199	2,166	2,255	918	1,886	1,831	2,130	1,866	1,119	-	-
Never symptoms, not consumed, SPT-	**No** (Type B) Food allergy possible	n	44	82	22	1,360	338	388	139	417	1,144	-	-
Never symptoms, not consumed, SPT+	**Yes** (Type C) Food allergy possible	n	4	6	3	8	36	42	8	2	17	186	238
Ever symptoms, not consumed	**Yes** (Type D) Food allergy likely	n	42	35	9	3	29	28	12	4	9		
Eligible (Type C + D)		**%**	**2.0**	**1.8**	**0.5**	**0.5**	**2.8**	**3.1**	**0.9**	**0.3**	**1.1**	**8.1**	**10.4**
Food challenge (fully completed assessments n=2,097)													
Conducted		n	6	7	0	0	15	16	2	1	7	39	46
Positive		n	0	1	0	0	3	7	1	1	2	13	17
Challenge-proven food allergy		**%**	**0**	**0.05**	**0**	**0**	**0.14**	**0.33**	**0.05**	**0.05**	**0.10**	**0.62**	**0.81**

Accepted Article

This article is protected by copyright. All rights reserved

Table 3

Basic characteristics of all recruited birth cohort participants, by follow-up status. Testing similarity between groups, Chi-square p-values.

*asthma, allergic rhinitis, or eczema.

#Cow's milk, Hen's egg, Wheat, Soy, Peanut, Hazelnut, White fish, Oily fish, Crustaceans

	Lost to follow-up	Questionnaire only	Eligible but not challenged	Eligible and challenged	Not eligible	Lost to follow-up vs. Questionnaire only	Questionnaire only vs. Not eligible	Elig. not challenged vs. Elig. Challenged
n	**4.458**	**3.816**	**192**	**46**	**2.051**			
Group number	**1**	**2**	**3**	**4**	**5**	**1 vs. 2**	**2 vs. 5**	**3 vs. 4**
	%	**%**	**%**	**%**	**%**	p-value	p-value	p-value
Birth & Family (assessed shortly after birth)								
Male sex	51,3	51,6	54,2	65,1	51,5	0,792	0,986	0,191
Cesarean section	22,7	23,1	24,6	14,6	23,8	0,713	0,537	0,168
Month of birth Jun-Nov	47,4	47,9	44,8	39,5	44,7	0,608	0,019	0,530
Antibiotics in first week	16,9	12,0	11,5	4,8	17,1	<0.001	<0.001	0,195
Older siblings	48,3	48,0	45,0	37,2	48,6	0,727	0,631	0,350
Mother smokes	15,2	10,2	7,8	4,7	10,3	<0.001	0,898	0,469
University/college degree father and/or mother	39,8	56,4	59,2	67,4	53,2	<0.001	0,018	0,315
Smoking in pregnancy	13,4	8,9	6,3	4,7	9,5	<0.001	0,415	0,689
Cat or dog in household	29,8	28,6	22,4	14,0	24,4	0,241	<0.001	0,218
Family's allergies (assessed shortly after birth)								
Self-reported physician's diagnosed allergy* (mother)	25,0	24,3	35,9	60,5	32,7	0,463	<0.001	0,003
Self-reported food allergy/hypersensitivity (mother)	13,3	14,8	18,8	16,7	17,4	0,042	0,011	0,752
Self-reported physician's diagnosed allergy* (father)	19,6	19,4	29,1	34,9	24,9	0,814	<0.001	0,456
Self-reported food allergy/hypersensitivity (father)	7,7	8,3	15,5	19,0	10,7	0,354	0,002	0,573
Child: 0-3 years (data from EuroPrevall)								
Eczema	34,5	32,2	61,4	66,7	44,8	0,041	<0.001	0,526
Challenge-proven FA (any)	1,3	1,6	14,9	31,7	1,6	0,355	0,998	0,012
...cow's milk allergy	0,4	0,5	4,0	15,0	0,7	0,765	0,464	0,009
...hen's egg allergy	0,8	0,9	8,6	26,8	1,0	0,695	0,571	0,001
...peanut allergy	0,1	0,1	1,2	7,5	0,0	0,786	0,204	0,017
Child: 6-10 years (assessed by online questionnaire)								
Parent-reported asthma (in previous 12 months)		6,4	27,7	11,6	9,2		<0.001	0,027
Parent-reported allergic rhinitis (in previous 12 months)		17,8	50,8	46,5	28,8		<0.001	0,613
Parent-reported eczema (in previous 12 months)		17,8	47,1	37,2	28,6		<0.001	0,238
Ever any reaction to food		12,7	74,9	62,8	16,2		<0.001	0,108
Consumed cow's milk in last 3 months		98,7	96,9	93,0	97,1		<0.001	0,234
Consumed hen's egg in last 3 months		98,6	91,1	95,3	96,9		<0.001	0,361

Accepted Article

This article is protected by copyright. All rights reserved

Consumed peanut in last 3 months	82,0	62,5	53,5	76,4	<0.001	0,274
Child: 6-10 years (assessed by face-to-face interview)						
Likely current FA (any food)		83,9	83,3	4,0		0,934
SPT positive (any core food#)		59,1	71,8	5,2		0,140
Eligible for cow's milk challenge		19,8	18,6			0,859
Eligible for hen's egg challenge		16,1	20,9			0,451
Eligible for peanut challenge		24,0	44,2			0,007

Accepted Article

This article is protected by copyright. All rights reserved

Table 4

Food-specific characteristics of challenge-eligible children, by challenge conduct. Only foods with more than 5 conducted challenges.

	challenge conducted	challenge not conducted
Eligible for cow's milk challenge (n = 47)	**6**	**41**
of these…		
…cow's milk SPT positive	17%	18%
…symptoms in previous 12m	83%	56%
…earlier physician's diagnosis	83%	39%
…current avoidance of cow's milk	50%	17%
Eligible for hen's egg challenge (n = 42)	**7**	**35**
of these…		
…hen's egg SPT positive	0%	57%
…symptoms in previous 12m	14%	20%
…earlier physician's diagnosis	43%	77%
…current avoidance of hen's egg	43%	40%
Eligible for peanut challenge (n = 66)	**15**	**51**
of these…		
…peanut SPT positive	85%	88%
…symptoms in previous 12m	0%	2%
…earlier physician's diagnosis	36%	39%
…current avoidance of peanut	64%	49%
Eligible for hazelnut challenge (n = 70)	**16**	**54**
of these…		
…hazelnut SPT positive	80%	73%
…symptoms in previous 12m	7%	7%
…earlier physician's diagnosis	33%	33%
…current avoidance of hazelnut	47%	46%
Eligible for crustacean challenge (n = 26)	**7**	**19**
of these…		
…crustacean SPT positive	71%	88%
…symptoms in previous 12m	0%	5%
…earlier physician's diagnosis	14%	21%
…current avoidance of crustaceans	43%	37%

Accepted Article

This article is protected by copyright. All rights reserved

Figure legends

Figure 1

Number of children participating assessments and diagnostic steps and number of children lost to follow-up.

Figure 2

Decision tree used to define current food allergy status, using peanut as an example (absolute numbers refer to peanut reactions).

Figure 3

Skin Prick Test (SPT) including allergens from the core food list in 2,188 European children aged 6-10 years. A total of 223 (10.2%) children had a positive SPT. Error bars indicate 95% confidence intervals for the proportion of positive tests (≥3 mm).

Figure 4

Adjusted prevalence of challenge-proven food allergy, by follow-up status.

Lost to follow-up at school age (Group 1) were 4,458 out of 10,563 subjects recruited at birth in 8 centres.

Adjustment as follows, based on similarity of groups regarding baseline and follow-up information as outlined in Table 3:

(i) disease frequency of those challenged (Group 4) as substitute for those eligible but refusing the challenge (Group 3), with an additional estimated number of 71 cases;

Accepted Article

This article is protected by copyright. All rights reserved

(ii) disease frequency of those not eligible (Group 5) as substitute for those who only answered the questionnaire (no estimated cases added);

Extrapolated numbers rounded to the next integer.

Accepted Article

This article is protected by copyright. All rights reserved

Accepted Article

References

1. Zuidmeer L, Goldhahn K, Rona RJ, et al. The prevalence of plant food allergies: a systematic review. *J Allergy Clin Immunol* 2008; **121**(5): 1210-8 e4.
2. Rona RJ, Keil T, Summers C, et al. The prevalence of food allergy: a meta-analysis. *J Allergy Clin Immunol* 2007; **120**(3): 638-46.
3. Nwaru BI, Hickstein L, Panesar SS, et al. The epidemiology of food allergy in Europe: a systematic review and meta-analysis. *Allergy* 2014; **69**(1): 62-75.
4. Orhan F, Karakas T, Cakir M, Aksoy A, Baki A, Gedik Y. Prevalence of immunoglobulin E-mediated food allergy in 6-9-year-old urban schoolchildren in the eastern Black Sea region of Turkey. *Clin Exp Allergy* 2009; **39**(7): 1027-35.
5. Osterballe M, Hansen TK, Mortz CG, Host A, Bindslev-Jensen C. The prevalence of food hypersensitivity in an unselected population of children and adults. *Pediatr Allergy Immunol* 2005; **16**(7): 567-73.
6. Pereira B, Venter C, Grundy J, Clayton CB, Arshad SH, Dean T. Prevalence of sensitization to food allergens, reported adverse reaction to foods, food avoidance, and food hypersensitivity among teenagers. *J Allergy Clin Immunol* 2005; **116**(4): 884-92.
7. Roehr CC, Edenharter G, Reimann S, et al. Food allergy and non-allergic food hypersensitivity in children and adolescents. *Clin Exp Allergy* 2004; **34**(10): 1534-41.
8. Venter C, Pereira B, Grundy J, Clayton CB, Arshad SH, Dean T. Prevalence of sensitization reported and objectively assessed food hypersensitivity amongst six-year-old children: a population-based study. *Pediatr Allergy Immunol* 2006; **17**(5): 356-63.
9. Venter C, Pereira B, Grundy J, et al. Incidence of parentally reported and clinically diagnosed food hypersensitivity in the first year of life. *J Allergy Clin Immunol* 2006; **117**(5): 1118-24.
10. Venter C, Patil V, Grundy J, et al. Prevalence and cumulative incidence of food hyper-sensitivity in the first 10 years of life. *Pediatr Allergy Immunol* 2016; **27**(5): 452-8.
11. Nicolaou N, Poorafshar M, Murray C, et al. Allergy or tolerance in children sensitized to peanut: prevalence and differentiation using component-resolved diagnostics. *J Allergy Clin Immunol* 2010; **125**(1): 191-7 e1-13.
12. Sampson HA, Gerth van Wijk R, Bindslev-Jensen C, et al. Standardizing double-blind, placebo-controlled oral food challenges: American Academy of Allergy, Asthma & Immunology-European Academy of Allergy and Clinical Immunology PRACTALL consensus report. *J Allergy Clin Immunol* 2012; **130**(6): 1260-74.

This article is protected by copyright. All rights reserved

13. McBride D, Keil T, Grabenhenrich L, et al. The EuroPrevall birth cohort study on food allergy: baseline characteristics of 12,000 newborns and their families from nine European countries. *Pediatr Allergy Immunol* 2012; **23**(3): 230-9.
14. Keil T, McBride D, Grimshaw K, et al. The multinational birth cohort of EuroPrevall: background, aims and methods. *Allergy* 2009.
15. Grabenhenrich LB, Reich A, Bellach J, et al. A new framework for the documentation and interpretation of oral food challenges in population-based and clinical research. *Allergy* 2016.
16. Schoemaker AA, Sprikkelman AB, Grimshaw KE, et al. Incidence and natural history of challenge-proven cow's milk allergy in European children - EuroPrevall birth cohort. *Allergy* 2015; **70**(8): 963-72.
17. Xepapadaki P, Fiocchi A, Grabenhenrich L, et al. Incidence and natural history of hen's egg allergy in the first 2 years of lifethe EuroPrevall birth cohort study. *Allergy* 2016; **71**(3): 350-7.
18. Peters RL, Koplin JJ, Gurrin LC, et al. The prevalence of food allergy and other allergic diseases in early childhood in a population-based study: HealthNuts age 4-year follow-up. *J Allergy Clin Immunol* 2017; **140**(1): 145-53 e8.
19. Grabenhenrich LB, Reich A, McBride D, et al. Physician's appraisal vs documented signs and symptoms in the interpretation of food challenge tests: The EuroPrevall birth cohort. *Pediatr Allergy Immunol* 2018; **29**(1): 58-65.

Accepted Article

This article is protected by copyright. All rights reserved

Accepted Article

Figure 1

Number of children participating assessments and diagnostic steps and number of children lost to follow-up.



This article is protected by copyright. All rights reserved

Accepted Article

Figure 2

Decision tree used to define current food allergy status, using peanut as an example (absolute numbers refer to peanut reactions).



This article is protected by copyright. All rights reserved

Figure 3

Skin Prick Test (SPT) including allergens from the core food list in 2,188 European children aged 6-10 years. A total of 223 (10.2%) children had a positive SPT. Error bars indicate 95% confidence intervals for the proportion of positive tests (≥3 mm).



Accepted Article

This article is protected by copyright. All rights reserved

Figure 4

Adjusted prevalence of challenge-proven food allergy, by follow-up status.

Lost to follow-up at school age (Group 1) were 4,458 out of 10,563 subjects recruited at birth in 8 centres.

Adjustment as follows, based on similarity of groups regarding baseline and follow-up information as outlined in Table 3:

(i) disease frequency of those challenged (Group 4) as substitute for those eligible but refusing the challenge (Group 3), with an additional estimated number of 71 cases;

(ii) disease frequency of those not eligible (Group 5) as substitute for those who only answered the questionnaire (no estimated cases added);

Extrapolated numbers rounded to the next integer.



Accepted Article

This article is protected by copyright. All rights reserved

Cow's Milk



Hen's Egg



Peanut



Hazelnut



Crustaceans



Online Appendix

Tables in this document.

e-Table 1	Page 2	Lifetime prevalence of parent-reported symptoms and doctor's diagnoses
e-Table 2	Page 3	Rare foods: numbers, questionnaire
e-Table 3	Page 4	Characteristics of food problems: n and %, questionnaire
e-Table 4	Page 5	Characteristics of food problems: n and %, interview
e-Table 5	Page 6	Consumption: n and %, interview and questionnaire
e-Table 6	Page 7	Positive DBPCFCs

The following abbreviations for country names will be used within this document.

IS	Iceland
GB	United Kingdom
NL	The Netherlands
DE	Germany
PL	Poland
LT	Lithuania
ES	Spain
GR	Greece

e-Table 1

Lifetime prevalence (raw % and n) of parent-reported symptoms after food consumption and parent-reported doctors diagnosed food allergy, based on face-to-face interview during study centre visit (n=2,322).

Symptoms after consumption of “core foods” were asked specifically, all other foods are labelled “non-core foods”.

*Only the most frequently reported non-core foods listed.

	Reykjavik Iceland	Southampton UK	Amsterdam NL	Berlin Germany	Lodz Poland	Vilnius Lithuania	Madrid Spain	Athens Greece	Total
	494	**221**	**201**	**332**	**407**	**142**	**323**	**202**	**2,322**
Study centre visit (n)	**% (n)**	**% (n)**	**% (n)**	**% (n)**	**% (n)**	**% (n)**	**% (n)**	**% (n)**	**% (n)**
Parent-reported symptoms									
Any food	**16.8 (83)**	**23.5 (52)**	**27.4 (55)**	**30.7 (102)**	**35.9 (146)**	**23.2 (33)**	**16.1 (52)**	**5.9 (12)**	**23.0 (535)**
"Core foods"	**16.0 (79)**	**16.3 (36)**	**25.9 (52)**	**19.6 (65)**	**30.2 (123)**	**21.8 (31)**	**13.0 (42)**	**4.5 (9)**	**18.8 (437)**
Cow's milk	12.3 (61)	9.5 (21)	20.4 (41)	13.6 (45)	24.3 (99)	9.2 (13)	5.9 (19)	2.5 (5)	13.1 (304)
Hen's egg	4.9 (24)	6.3 (14)	9.0 (18)	3.6 (12)	6.9 (28)	9.2 (13)	3.7 (12)	1.0 (2)	5.3 (123)
Wheat	0.8 (4)	3.2 (7)	1.0 (2)	2.4 (8)	1.5 (6)	1.4 (2)	0.6 (2)	0.0 (0)	1.3 (31)
Soy	1.6 (8)	0.5 (1)	0.5 (1)	0.3 (1)	0.5 (2)	2.1 (3)	0.3 (1)	0.0 (0)	0.7 (17)
Peanut	2.0 (10)	1.4 (3)	3.0 (6)	2.4 (8)	3.2 (13)	8.5 (12)	1.5 (5)	0.0 (0)	2.5 (57)
Hazelnut	0.6 (3)	0.5 (1)	1.0 (2)	3.3 (11)	4.4 (18)	14.8 (21)	0.3 (1)	0.5 (1)	2.5 (58)
White fish	2.0 (10)	0.9 (2)	0.5 (1)	0.3 (1)	0.7 (3)	3.5 (5)	1.9 (6)	0.5 (1)	1.2 (29)
Oily fish	0.8 (4)	0.5 (1)	0.0 (0)	0.3 (1)	0.2 (1)	0.7 (1)	1.5 (5)	0.0 (0)	0.6 (13)
Crustaceans	1.0 (5)	0.9 (2)	0.5 (1)	0.0 (0)	0.0 (0)	0.7 (1)	0.9 (3)	1.0 (2)	0.6 (14)
"Non-core foods"*	**1.4 (7)**	**9.5 (21)**	**5.0 (10)**	**14.5 (48)**	**11.3 (46)**	**4.9 (7)**	**4.3 (14)**	**2.0 (4)**	**6.8 (157)**
Nuts (except hazelnut)	0.6 (3)	1.4 (3)	2.0 (4)	1.8 (6)	0.5 (2)	0.0 (0)	1.5 (5)	0.5 (1)	1.0 (24)
Tomato	0.2 (1)	2.3 (5)	0.5 (1)	3.0 (10)	1.5 (6)	0.0 (0)	0.0 (0)	0.0 (0)	1.0 (23)
Kiwi	0.6 (3)	0.9 (2)	0.5 (1)	1.2 (4)	1.2 (5)	0.7 (1)	0.9 (3)	0.5 (1)	0.9 (20)
Strawberry	0.0 (0)	0.9 (2)	1.0 (2)	1.8 (6)	1.2 (5)	0.0 (0)	0.0 (0)	0.0 (0)	0.6 (15)
Apple	0.2 (1)	0.5 (1)	0.0 (0)	2.7 (9)	1.2 (5)	1.4 (2)	0.0 (0)	0.0 (0)	0.8 (18)
Citrus fruits	0.0 (0)	0.5 (1)	0.0 (0)	0.6 (2)	0.2 (1)	0.0 (0)	0.0 (0)	0.0 (0)	0.2 (4)
Parent-reported doctors-diagnosed food allergy	**8.3 (41)**	**10.0 (22)**	**20.9 (42)**	**6.3 (21)**	**21.1 (86)**	**14.8 (21)**	**7.7 (25)**	**3.0 (6)**	**11.4 (264)**
"Core foods"	**7.9 (39)**	**9.0 (20)**	**20.4 (41)**	**6.0 (20)**	**20.1 (82)**	**14.8 (21)**	**6.5 (21)**	**2.5 (5)**	**10.7 (249)**
Cow's milk	4.9 (24)	5.4 (12)	14.9 (30)	2.4 (8)	16.5 (67)	9.2 (13)	2.8 (9)	1.0 (2)	7.1 (165)
Hen's egg	4.5 (22)	5.9 (13)	7.0 (14)	2.1 (7)	4.9 (20)	8.5 (12)	2.5 (8)	1.0 (2)	4.2 (98)
Wheat	0.8 (4)	0.9 (2)	0.5 (1)	0.3 (1)	1.0 (4)	1.4 (2)	0.0 (0)	0.0 (0)	0.6 (14)
Soy	1.4 (7)	0.5 (1)	0.5 (1)	0.0 (0)	0.5 (2)	0.0 (0)	0.3 (1)	0.0 (0)	0.5 (12)
Peanut	2.0 (10)	0.9 (2)	2.0 (4)	1.8 (6)	1.7 (7)	3.5 (5)	1.2 (4)	0.0 (0)	1.6 (38)
Hazelnut	0.4 (2)	0.5 (1)	1.0 (2)	1.8 (6)	2.0 (8)	6.3 (9)	0.3 (1)	0.5 (1)	1.3 (30)
White fish	1.2 (6)	0.5 (1)	0.0 (0)	0.0 (0)	0.5 (2)	0.7 (1)	1.2 (4)	0.5 (1)	0.6 (15)
Oily fish	0.4 (2)	0.0 (0)	0.0 (0)	0.0 (0)	0.0 (0)	0.0 (0)	0.3 (1)	0.0 (0)	0.1 (3)
Crustaceans	0.8 (4)	0.0 (0)	0.5 (1)	0.0 (0)	0.0 (0)	0.0 (0)	0.3 (1)	0.0 (0)	0.3 (6)
"Non-core foods"*	**0.8 (4)**	**1.8 (4)**	**1.0 (2)**	**2.1 (7)**	**2.7 (11)**	**0.7 (1)**	**1.9 (6)**	**0.5 (1)**	**1.6 (36)**
Nuts (except hazelnut)	0.4 (2)	1.4 (3)	1.5 (3)	1.8 (6)	0.5 (2)	0.0 (0)	1.5 (5)	0.5 (1)	0.9 (22)
Tomato	0.0 (0)	0.5 (1)	0.0 (0)	0.0 (0)	0.2 (1)	0.0 (0)	0.0 (0)	0.0 (0)	0.1 (2)
Kiwi	0.2 (1)	0.0 (0)	0.0 (0)	0.0 (0)	0.5 (2)	0.0 (0)	0.3 (1)	0.0 (0)	0.2 (4)
Strawberry	0.0 (0)	0.5 (1)	0.0 (0)	0.3 (1)	0.0 (0)	0.0 (0)	0.0 (0)	0.0 (0)	0.1 (2)
Apple	0.2 (1)	0.0 (0)	0.0 (0)	0.0 (0)	0.2 (1)	0.7 (1)	0.0 (0)	0.0 (0)	0.1 (3)
Citrus fruits	0.0 (0)	0.0 (0)	0.0 (0)	0.0 (0)	0.0 (0)	0.0 (0)	0.0 (0)	0.0 (0)	0.0 (0)

e-Table 2

Number of children for parent-reported trouble/illness with specific food earlier in life, based on online questionnaire (n=6,069).

Listed are food items that were only rarely reported and not covered in Table 1.

Nuts	
Almond	4
Cashew	12
Macadamia	1
Pine nut	1
Pistachio	6
Walnut	8
Nuts, not specified	13
Animal products	
Caviar	1
Chicken	10
Beef	1
Goat's milk	2
Honey	10
Meat	4
Pork	4
Poultry	1
Squid	1
Plant products	
Apricot	3
Banana	13
Barley	3
Beans	6
Bell pepper	11
Blackcurrant	1
Blueberry	1
Broccoli	2
Buckwheat	1
Cabbage	1
Camomile	1
Carrot	17
Celery	4
Cherry	2
Chestnut	1
Chives	1
Citrus fruit, not specified	25
Cocoa	10
Coconut	1
Corn	2
Eggplant	3
Figs	1
Grapefruit	1
Grapes	3
Lemon	2
Lentil	2
Liquorice	2
Mango	1
Melon	4
Nectarine	1
Oats	1
Olives	1
Onions	3
Orange	10
Peach	7
Pear	2
Peas	2
Pineapple	9
Potato	6
Raspberry	2
Rice	5
Rye	2
Sesame	6
Spinach	1
Squash	1
Sweet potato	1
Tangerines	9
Watermelon	2

e-Table 3

Parent-reported symptoms after food consumption (ever), based on online questionnaire (n=6,069). A single child may report problems with several foods: relative frequencies refer to the number of children reporting problems.

* manually derived from free text answers

	Total	Cow's milk	Hen's egg	Peanut	Hazelnut	Reykjavik Iceland	Southampton UK	Amsterdam NL	Berlin Germany	Lodz Poland	Vilnius Lithuania	Madrid Spain	Athens Greece
n	**1,703**	**507**	**203**	**96**	**89**	**331**	**155**	**196**	**271**	**352**	**219**	**142**	**37**
	% (n)	**% (n)**	**% (n)**	**% (n)**	**% (n)**	**% (n)**	**% (n)**	**% (n)**	**% (n)**	**% (n)**	**% (n)**	**% (n)**	**% (n)**
Symptoms													
Itching, tingling or swelling in the mouth, lips or throat	26.5 (437)	10.4 (52)	28.9 (57)	52.3 (46)	48.2 (40)	22.2 (72)	31.6 (48)	23.6 (46)	35.8 (95)	16.1 (55)	12.7 (25)	58.7 (81)	40.5 (15)
Rash, nettle sting like rash or itchy skin	48.8 (805)	34.9 (174)	66.5 (131)	64.8 (57)	55.4 (46)	36.1 (117)	46.1 (70)	30.3 (59)	20.8 (55)	74.2 (253)	79.2 (156)	58.0 (80)	40.5 (15)
Eczema*	1.9 (31)	2.6 (13)	4.1 (8)	1.1 (1)	1.2 (1)	0.9 (3)	0.0 (0)	0.5 (1)	2.3 (6)	2.3 (8)	6.6 (13)	0.0 (0)	0.0 (0)
Diarrhoea or vomiting (other than food poisoning)	38.8 (640)	50.4 (251)	41.6 (82)	30.7 (27)	21.7 (18)	54.3 (176)	53.3 (81)	61.0 (119)	37.0 (98)	24.3 (83)	12.2 (24)	37.0 (51)	21.6 (8)
Gastrointestinal discomfort*	15.9 (262)	28.5 (142)	4.6 (9)	2.3 (2)	4.8 (4)	32.4 (105)	17.8 (27)	9.7 (19)	31.7 (84)	4.4 (15)	2.5 (5)	4.3 (6)	2.7 (1)
Runny or stuffy nose	13.9 (229)	14.1 (70)	17.3 (34)	19.3 (17)	19.3 (16)	7.4 (24)	8.6 (13)	27.2 (53)	0.8 (2)	25.2 (86)	14.2 (28)	16.7 (23)	0.0 (0)
Red, sore or runny eyes	8.1 (133)	4.2 (21)	10.2 (20)	17.0 (15)	12.0 (10)	8.6 (28)	2.0 (3)	8.7 (17)	4.2 (11)	10.0 (34)	6.6 (13)	18.8 (26)	2.7 (1)
Difficulty swallowing or breathlessness	7.0 (116)	2.4 (12)	8.1 (16)	26.1 (23)	12.0 (10)	7.1 (23)	3.9 (6)	8.2 (16)	6.4 (17)	4.1 (14)	4.6 (9)	20.3 (28)	8.1 (3)
Asthma*	8.9 (146)	4.4 (22)	9.1 (18)	28.4 (25)	14.5 (12)	9.9 (32)	4.6 (7)	8.2 (16)	7.2 (19)	6.7 (23)	8.1 (16)	21.0 (29)	10.8 (4)
Fainting or dizzyness	1.3 (21)	0.6 (3)	2.0 (4)	3.4 (3)	1.2 (1)	0.9 (3)	0.7 (1)	7.2 (14)	0.4 (1)	0.0 (0)	0.0 (0)	1.4 (2)	0.0 (0)
Headaches	5.2 (85)	6.0 (30)	2.5 (5)	4.5 (4)	2.4 (2)	9.3 (30)	3.9 (6)	1.0 (2)	4.2 (11)	3.5 (12)	2.5 (5)	13.8 (19)	0.0 (0)
generalized, more than 1 organ system	38.2 (630)	32.7 (163)	48.2 (95)	62.5 (55)	42.2 (35)	40.1 (130)	43.4 (66)	47.2 (92)	23.0 (61)	40.2 (137)	30.5 (60)	55.8 (77)	18.9 (7)
Immediate-type reaction (<2 hours)	85.0 (1,401)	83.3 (415)	83.2 (164)	93.2 (82)	92.8 (77)	84.0 (272)	88.2 (134)	87.7 (171)	90.2 (239)	77.1 (263)	83.2 (164)	94.9 (131)	73.0 (27)

e-Table 4

Parent-reported symptoms after food consumption (ever), based on face-to-face interviews during study centre visit (n=2,322). A single child may report problems with several foods.

*manually derived from free text answers

	Total	Cow's milk	Hen's egg	Peanut	Hazelnut	Reykjavik Iceland	Southampton UK	Amsterdam NL	Berlin Germany	Lodz Poland	Vilnius Lithuania	Madrid Spain	Athens Greece
n	**868**	**304**	**123**	**57**	**58**	**139**	**78**	**85**	**159**	**237**	**78**	**73**	**19**
	% (n)	**% (n)**	**% (n)**	**% (n)**	**% (n)**	**% (n)**	**% (n)**	**% (n)**	**% (n)**	**% (n)**	**% (n)**	**% (n)**	**% (n)**
Symptoms													
Pruritus (hard continuous scratching leading to excoriations)	23.6 (205)	18.8 (57)	32.5 (40)	28.1 (16)	20.7 (12)	10.9 (15)	3.8 (3)	24.7 (21)	13.8 (22)	39.7 (94)	21.8 (17)	32.9 (24)	47.4 (9)
Rash (more than few areas of faint erythema)	31.7 (275)	33.7 (102)	42.3 (52)	28.1 (16)	27.6 (16)	21.0 (29)	16.7 (13)	44.7 (38)	20.8 (33)	57.0 (135)	5.1 (4)	27.4 (20)	15.8 (3)
Urticaria (more than 2 hives)	18.1 (157)	12.9 (39)	30.9 (38)	24.6 (14)	15.5 (9)	21.7 (30)	30.8 (24)	25.9 (22)	14.5 (23)	11.4 (27)	3.8 (3)	35.6 (26)	10.5 (2)
Angioedema on lip and/or face	11.5 (100)	3.0 (9)	16.3 (20)	17.5 (10)	10.3 (6)	11.6 (16)	24.4 (19)	9.4 (8)	8.8 (14)	6.8 (16)	5.1 (4)	30.1 (22)	5.3 (1)
Oral allergy syndrome*	21.2 (184)	3.6 (11)	17.1 (21)	36.8 (21)	51.7 (30)	15.2 (21)	26.9 (21)	10.6 (9)	29.6 (47)	8.4 (20)	50.0 (39)	32.9 (24)	15.8 (3)
Worsening of eczema	20.2 (175)	26.7 (81)	35.8 (44)	12.3 (7)	10.3 (6)	15.9 (22)	12.8 (10)	47.1 (40)	6.3 (10)	26.6 (63)	33.3 (26)	4.1 (3)	5.3 (1)
Frequent sniffing and/or bursts of sneezing	5.7 (49)	3.0 (9)	13.0 (16)	7.0 (4)	8.6 (5)	10.1 (14)	0.0 (0)	8.2 (7)	0.6 (1)	9.7 (23)	0.0 (0)	5.5 (4)	0.0 (0)
Reddening or swelling of eyes	4.2 (36)	2.3 (7)	9.8 (12)	7.0 (4)	5.2 (3)	9.4 (13)	1.3 (1)	9.4 (8)	2.5 (4)	3.0 (7)	1.3 (1)	1.4 (1)	5.3 (1)
Wheezing	3.1 (27)	1.3 (4)	6.5 (8)	5.3 (3)	6.9 (4)	5.8 (8)	3.8 (3)	2.4 (2)	1.3 (2)	4.2 (10)	0.0 (0)	1.4 (1)	5.3 (1)
Stridor, cough (> 3 episodes) and/or persistent throat tightness	5.7 (49)	1.3 (4)	4.9 (6)	10.5 (6)	12.1 (7)	13.0 (18)	1.3 (1)	4.7 (4)	3.8 (6)	4.2 (10)	0.0 (0)	12.3 (9)	5.3 (1)
Emesis	17.0 (147)	20.5 (62)	19.5 (24)	10.5 (6)	8.6 (5)	23.2 (32)	34.6 (27)	11.8 (10)	14.5 (23)	12.7 (30)	0.0 (0)	30.1 (22)	15.8 (3)
Diarrhea	18.2 (158)	32.0 (97)	16.3 (20)	8.8 (5)	3.4 (2)	36.2 (50)	28.2 (22)	15.3 (13)	18.2 (29)	13.9 (33)	0.0 (0)	11.0 (8)	15.8 (3)
Gastrointestinal discomfort*	14.0 (121)	25.1 (76)	6.5 (8)	0.0 (0)	3.4 (2)	20.3 (28)	16.7 (13)	1.2 (1)	26.4 (42)	10.5 (25)	0.0 (0)	12.3 (9)	15.8 (3)
Change in mental status (unexplained drowsiness or agitation)	4.7 (41)	6.9 (21)	6.5 (8)	5.3 (3)	5.2 (3)	2.9 (4)	2.6 (2)	1.2 (1)	2.5 (4)	12.2 (29)	0.0 (0)	1.4 (1)	0.0 (0)
generalized, more than 1 organ system	26.8 (232)	25.1 (76)	38.2 (47)	26.3 (15)	20.7 (12)	36.2 (50)	37.2 (29)	28.2 (24)	14.5 (23)	31.6 (75)	1.3 (1)	38.4 (28)	10.5 (2)
Immediate-type reaction (<2 hours)	80.7 (700)	78.2 (237)	82.1 (101)	86.0 (49)	89.7 (52)	88.4 (122)	82.1 (64)	91.8 (78)	90.6 (144)	59.9 (142)	82.1 (64)	93.2 (68)	94.7 (18)

e-Table 5

Consumption of and reasons for not consuming "core foods" at school age. Based on face-to-face interview (n=2,322, top) and online questionnaires (n=6,069, bottom).

*Percentage of those currently and/or never consuming.

1 e.g. fresh cow's milk, yoghurt, cheese; 2 e.g. rolls, muffins, biscuit, cake; 3 e.g. soft-boiled, scrambled or fried egg; 4 e.g. cake, muffin, biscuit

		Cow's milk		**Hen's egg**					**Hazelnut**		**White**	**Oily**	
Based on interview (2,322)	n	not baked[1]	baked[2]	not baked[3]	baked[4]	**Wheat**	**Soy**	**Peanut**	raw	roasted	**fish**	**fish**	**Crustaceans**
Consumed in last 3 months													
Reykjavik, Iceland	494	98.4 (486)	99.0 (489)	90.1 (445)	98.6 (487)	99.6 (492)	24.7 (122)	82.6 (408)	77.3 (382)	77.1 (381)	98.6 (487)	83.0 (410)	54.3 (268)
Southampton, UK	221	99.1 (219)	99.1 (219)	84.6 (187)	99.5 (220)	98.6 (218)	76.5 (169)	58.4 (129)	47.1 (104)	72.9 (161)	95.0 (210)	80.5 (178)	46.6 (103)
Amsterdam, NL	201	94.5 (190)	96.0 (193)	86.1 (173)	96.5 (194)	99.0 (199)	32.8 (66)	87.6 (176)	68.7 (138)	89.6 (180)	94.5 (190)	64.2 (129)	47.8 (96)
Berlin, Germany	332	98.5 (327)	100.0 (332)	95.2 (316)	99.7 (331)	99.7 (331)	37.7 (125)	86.4 (287)	58.7 (195)	95.8 (318)	90.4 (300)	67.5 (224)	33.4 (111)
Lodz, Poland	407	97.8 (398)	98.5 (401)	96.3 (392)	99.3 (404)	100.0 (407)	51.4 (209)	92.4 (376)	90.9 (370)	93.6 (381)	88.5 (360)	85.5 (348)	17.2 (70)
Vilnius, Lithuania	142	88.7 (126)	86.6 (123)	84.5 (120)	86.6 (123)	93.7 (133)	16.9 (24)	43.0 (61)	60.6 (86)	64.8 (92)	59.9 (85)	45.1 (64)	14.1 (20)
Madrid, Spain	323	99.7 (322)	100.0 (323)	99.4 (321)	99.7 (322)	98.8 (319)	11.5 (37)	79.6 (257)	60.4 (195)	91.3 (295)	99.1 (320)	96.9 (313)	72.8 (235)
Athens, Greece	202	100.0 (202)	100.0 (202)	98.5 (199)	99.5 (201)	99.0 (200)	55.4 (112)	92.1 (186)	92.6 (187)	99.5 (201)	95.5 (193)	93.1 (188)	70.8 (143)
All countries	**2,322**	**97.8 (2,270)**	**98.3 (2,282)**	**92.7 (2,153)**	**98.3 (2,282)**	**99.0 (2,299)**	**37.2 (864)**	**81.0 (1,880)**	**71.4 (1,657)**	**86.5 (2,009)**	**92.4 (2,145)**	**79.8 (1,854)**	**45.0 (1,046)**
Never eaten	2,322	0.4 (10)	0.3 (8)	1.2 (29)	0.3 (6)	0.3 (7)	22.8 (530)	2.8 (64)	5.3 (123)	2.4 (55)	0.9 (22)	3.3 (76)	25.6 (594)
Reason for not consuming*													
Parent's suspicion of food allergy		25.0 (13)	17.5 (7)	5.3 (9)	10.0 (4)	4.3 (1)	0.5 (8)	6.6 (29)	6.0 (40)	10.2 (32)	4.0 (7)	1.5 (7)	0.7 (9)
Parent's report of positive food challenge		21.2 (11)	20.0 (8)	11.8 (20)	37.5 (15)	8.7 (2)	0.1 (2)	5.0 (22)	2.1 (14)	4.2 (13)	4.0 (7)	0.4 (2)	0.2 (3)
Prevent allergy (never symptoms)		7.7 (4)	5.0 (2)	0.6 (1)	2.5 (1)	8.7 (2)	0.6 (9)	5.9 (26)	2.6 (17)	2.9 (9)	1.7 (3)	0.6 (3)	0.5 (7)
Dislike		17.3 (9)	17.5 (7)	55.6 (94)	10.0 (4)	4.3 (1)	10.3 (150)	25.1 (111)	24.4 (162)	25.2 (79)	47.5 (84)	50.6 (237)	40.4 (516)
Not in regular diet		9.6 (5)	12.5 (5)	17.2 (29)	17.5 (7)	39.1 (9)	88.1 (1,284)	52.7 (233)	62.7 (417)	53.7 (168)	36.2 (64)	44.0 (206)	58.7 (749)
Vegetarian, vegan, religious		1.9 (1)	2.5 (1)	0.6 (1)	2.5 (1)	0.0 (0)	0.0 (0)	0.0 (0)	0.0 (0)	0.0 (0)	2.8 (5)	1.1 (5)	0.5 (6)
Based on questionnaire (6,069)													
Consumed in last 3 months													
Reykjavik, Iceland	945	97.5 (921)	98.4 (930)	83.2 (786)	98.7 (933)	99.5 (940)	14.5 (137)	66.0 (624)	39.7 (375)	59.7 (564)	97.5 (921)	77.2 (730)	44.2 (418)
Southampton, UK	454	98.5 (447)	98.5 (447)	82.6 (375)	97.6 (443)	97.6 (443)	25.6 (116)	67.2 (305)	41.9 (190)	79.5 (361)	91.6 (416)	77.5 (352)	41.4 (188)
Amsterdam, NL	652	96.2 (627)	98.0 (639)	92.0 (600)	98.6 (643)	99.1 (646)	48.3 (315)	93.6 (610)	68.4 (446)	92.9 (606)	93.3 (608)	78.8 (514)	58.7 (383)
Berlin, Germany	1,001	97.8 (979)	98.9 (990)	89.6 (897)	98.5 (986)	99.3 (994)	26.9 (269)	89.0 (891)	67.0 (671)	94.2 (943)	73.6 (737)	68.0 (681)	32.8 (328)
Lodz, Poland	819	92.8 (760)	91.3 (748)	87.1 (713)	91.8 (752)	94.5 (774)	9.3 (76)	71.2 (583)	58.6 (480)	74.2 (608)	65.9 (540)	59.3 (486)	12.2 (100)
Vilnius, Lithuania	949	94.3 (895)	97.2 (922)	94.7 (899)	97.6 (926)	98.3 (933)	49.7 (472)	77.8 (738)	83.5 (792)	90.1 (855)	85.7 (813)	83.2 (790)	45.6 (433)
Madrid, Spain	688	98.5 (678)	98.5 (678)	92.2 (634)	97.7 (672)	97.5 (671)	18.6 (128)	81.4 (560)	47.4 (326)	88.7 (610)	98.0 (674)	93.6 (644)	70.8 (487)
Athens, Greece	561	98.8 (554)	98.6 (553)	96.4 (541)	99.3 (557)	98.9 (555)	38.0 (213)	94.5 (530)	92.7 (520)	96.1 (539)	95.5 (536)	93.2 (523)	73.6 (413)
All countries	**6,069**	**96.6 (5,861)**	**97.3 (5,907)**	**89.7 (5,445)**	**97.4 (5,912)**	**98.1 (5,956)**	**28.4 (1,726)**	**79.8 (4,841)**	**62.6 (3,800)**	**83.8 (5,086)**	**86.4 (5,245)**	**77.8 (4,720)**	**45.3 (2,750)**

e-Table 6

Characteristics of 20 positive double-blind placebo-controlled food challenges in 17 children using a 7 escalating doses protocol as developed in this birth cohort. If not otherwise stated, a placebo challenge was conducted w/o signs or symptoms.

*three children had two positive challenges, labelled A-C; Severity grading based on supervising physician's judgment.

Food	Age [years]	Country	Verum day: Symptoms after	Severity (0-10)	Emerg. drugs	Placebo day: Symptoms after
Hen's egg	7	NL	**dose 1**: itchy mouth **dose 5 (stopped)**: anaphylaxis with generalized urticaria and rash, continuous scratching, mild lip edema, expiratory wheezing, emesis, persistent rhinorrhea, and periocular swelling	10	Adrenaline, Salbutamol, Antihistamine	no placebo day
Peanut	9 (B*)	UK	**dose 1 (stopped)**: persistent sniffing	2	Antihistamine	no symptoms
Peanut	9 (C*)	LT	**dose 2**: itchy mouth **dose 3 (stopped)**: gastrointestinal discomfort with emesis (once), frequent dry cough, occasional sniffing, mild lip edema, faint erythema on face	7	Antihistamine	no symptoms
Peanut	8	IS	**dose 5 (stopped)**: emesis (once), hoarseness and sniffing	5	Antihistamine	no symptoms
Hazelnut	8 (A*)	DE	**dose 1 (stopped)**: periocular swelling, reddening of eyes, and persistent throat tightness	3	Antihistamine	no symptoms
Hazelnut	9 (B*)	UK	**dose 1 (stopped)**: throat itch, frequent sniffing	2	Antihistamine	no symptoms
Hazelnut	10	PL	**dose 2**: pale trunk erythema with scratching **dose 3 (stopped)**: itching behind ears, on face and eyes, increased erythema	2	Steroids, Antihistamine	**dose 3:** abdominal pain
Hazelnut	9	LT	**dose 2**: four hives, faint erythema, frequent sniffing, continuous rubbing of eyes, periocular swelling, gastrointestinal discomfort with emesis (once) **dose 3:** no new symptoms **(stopped)**	8	Antihistamine	no symptoms
Hazelnut	9 (C*)	LT	**dose 3**: occasional sniffing and scratching, rubbing of eyes **dose 4 (stopped)**: continuous scratching, five hives, frequent sniffing, gastrointestinal discomfort with itchy mouth	7	Antihistamine	no symptoms
Hazelnut	9	LT	**dose 1**: itchy mouth and throat **dose 3**: some cough **dose 4 (stopped)**: frequent dry cough	4	Antihistamine	no symptoms
Hazelnut	10	IS	**dose 1 (stopped)**: throat itch, cough, stridor with hoarseness, crampy chest pain	7	Adrenaline, Antihistamine	no symptoms
White fish	7	IS	**dose 1 (stopped)**: gastrointestinal discomfort with emesis (once), itchy mouth, faint erythema with occasional scratching	5	none	no symptoms
Oily fish	9	IS	**dose 3**: occasional sniffing and throat tightness **after final dose**: gastrointestinal discomfort with diarrhea (3h after final dose)	2	none	no symptoms
Crustaceans	8	DE	**dose 6**: abdominal pain and nausea with emesis (once) **after final dose**: gastrointestinal distress, generalized urticaria with continuous scratching (3h after final dose)	3	none	no placebo day
Crustaceans	9	UK	**dose 4**: erythematous rash **dose 5 (stopped)**: mild lip edema	2	Antihistamine	no placebo day, family/child not blinded
Cashew	9	UK	**dose 1:** rash, angioedema, oral itching	[not doc.]	none	no symptoms
Cashew	8	IS	**dose 3**: stomach ache, occasional sniffing **after final dose**: gastrointestinal distress	3	none	**after final dose:** sniffing and cough (also before placebo day)
Pine nut	8	IS	**dose 1**: gastrointestinal discomfort **dose 3 (stopped)**: expiratory wheezing, conjunctivitis, rhinitis, nausea, rash with scratching	8	Adrenaline, Steroids, Antihistamine	no symptoms
Pine nut	9	DE	**after final dose**: swelling of hands, erythema on feet, occasional scratching (1/2h after final dose); generalized urticaria, lip edema, periocular swelling (1 1/2h after final dose)	6	Antihistamine	no symptoms
Walnut	8 (A*)	DE	**dose 1**: itchy mouth, two hives **dose 2**: sniffing, swelling of eyes **dose 4**: blisters of oral mucosa **dose 6**: gastrointestinal discomfort with nausea **not stopped**	1	none	no symptoms