



Contents lists available at SciVerse ScienceDirect



Review Article

Pediatric sublingual immunotherapy efficacy: evidence analysis, 2009–2012

Désirée Larenas-Linnemann, MD^{*}; Michael Blaiss, MD[†]; Hugo P. Van Bever, MD, PhD[‡]; Enrico Compalati, MD, PhD[§]; and Carlos E. Baena-Cagnani, MD^{¶,||}

^{*} Hospital Médica Sur, Mexico City, Mexico

[†] Departments of Pediatrics and Medicine, University of Tennessee Health Science, Memphis, Tennessee

[‡] National University Singapore, Singapore

[§] Allergy & Respiratory Diseases Clinic, Department of Internal Medicine, University of Genoa, Genoa, Italy

[¶] Research Centre in Respiratory Medicine, Faculty of Medicine, Catholic University, Cordoba, Argentina

^{||} School of Specialization, Respiratory Medicine, University of Genoa, Genoa, Italy

ARTICLE INFO

Article history:

Received for publication January 3, 2013.

Received in revised form February 19, 2013.

Accepted for publication February 20, 2013.

ABSTRACT

Objective: To perform a structured analysis of the latest scientific evidence obtained for the clinical efficacy of sublingual immunotherapy (SLIT) in children.

Data Sources: PubMed, Embase, reference lists from reviews, and personal databases were reviewed for original articles on clinical trials with SLIT in patients younger than 18 years published from January 1, 2009, through December 31, 2012, using broad search and medical subject heading terms.

Study Selections: Clinical trials, irrespective of their design, of SLIT in the treatment of respiratory and food allergy in patients 18 years or younger were selected. Clinical outcomes (symptom scores, medication use, provocation tests, pulmonary function tests, skin prick tests, and adverse events) and immunologic changes were tabulated. Quality of each trial and total quality of compounded evidence was analyzed with the Grading of Recommendations Assessment, Development and Evaluation system.

Results: Of 56 articles, 29 met the inclusion criteria. New evidence is robust for the pre-seasonal tablet and drop grass pollen SLIT efficacy in allergic rhinitis and scarce for seasonal asthma. Some evidence for *Alternaria* SLIT efficacy is appearing. For house dust mite (HDM) SLIT in asthma, there is high-quality evidence for medication reduction while maintaining symptom control; evidence for HDM SLIT efficacy in allergic rhinitis is of moderate-low quality. There is moderate evidence for efficacy of dual grass pollen–HDM SLIT after 12 months of treatment and 1 year after discontinuation. Specific provocation test results (nasal, skin) improve with grass pollen and HDM SLIT but nonspecific bronchial provocation testing does not. Food oral immunotherapy is more promising than food SLIT. Possible new surrogate markers have been reported. No anaphylaxis was found among 2469 treated children.

Conclusion: Evidence for efficacy of SLIT in children with respiratory or food allergy is growing.

© 2013 American College of Allergy, Asthma & Immunology. Published by Elsevier Inc. All rights reserved.

Introduction

Allergen immunotherapy (AIT) is still the only treatment directed at correcting the deviated immune response, which has been found to be the cause of allergy. Moreover, because of its mechanisms of action, AIT is the only therapy that modifies the natural history of the disease. Several studies have reported on the preventive effect of immunotherapy in children with allergic rhinitis (AR) because it appears to reduce the development of new allergic sensitizations and/or new-onset asthma.^{1,2} Today, clear humoral, cellular, and tissue level changes have been documented with AIT,^{3–5} and its clinical efficacy leads to economic savings after

6 months of treatment.⁶ In 2011, the centenary of subcutaneous immunotherapy (SCIT) was celebrated^{7,8}; concurrently, it was 25 years ago that the first double-blind, placebo-controlled (DBPC) trial with SLIT was published.⁹ This alternative, less traumatic, and safer route of administration seems especially suitable for children, and after the first big conclusive trials in adults,^{10,11} many pediatric SLIT trials were conducted, and pediatric SLIT was appraised in several meta-analyses and reviews.^{12–16} However, published meta-analyses generally only include a selection of trials based on their design.

The Grading of Recommendations Assessment, Development and Evaluation (GRADE) tool¹⁷ has been introduced as a method to support health policy decision making built on clinical recommendations as a result of analysis of different aspects, one of which is the quality of evidence coming from research. As such, the GRADE system developed tools to define the scientific quality of

Reprints: Désirée Larenas-Linnemann, MD, Hospital Médica Sur, Torre 2, cons. 602 Puente de Piedra 150, Colonia Toriello Guerra, Delagación Tlalpan, 14050 México D.F., México; E-mail: Marlar1@prodigy.net.mx.

Disclosures: Authors have nothing to disclose.

clinical trials, taking into account internal and external validation, including the risk of bias. In GRADE all clinical trials, irrespective of their design, are considered and their quality of evidence is established according to defined parameters (eTable 1).¹⁸ Since 2004, the GRADE system has been adapted by many specialties as a useful tool for the formulation of guidelines.

In this review, we analyze all clinical trials published on pediatric SLIT since the World Allergy Organization position paper was published in 2009,¹⁹ assess their scientific quality with GRADE, and integrate this evidence on clinical aspects of SLIT in children.

Methods

Search Strategy

Literature searches were conducted in PubMed and Embase (D.L.L., H.V.B., M.B.) to identify original articles on clinical trials with SLIT in children published between January 1, 2009, and December 31, 2012, and written in English or Spanish. Search terms and limits were all combinations of *desensitization*, *immunologic* (medical subject heading [MeSH] terms) OR *allergen immunotherapy* AND *administration*, *sublingual* (MeSH terms) AND 2009/01/01-2012/11/15 AND (Randomized) (Controlled) (Clinical Trial) AND (English OR Spanish) AND the MeSH terms *humans*, *infant* OR *child*, OR *adolescent*. We identified additional articles by manually searching references from the obtained articles, review articles, and the authors' own literature database. Study design was not a restriction; only full-text articles were included.

Study Selection

In the first phase of screening, 3 reviewers (D.L.L., M.B. and H.V.B.) independently examined the titles and abstracts of the search results. The second phase of screening was based on full-text articles, which were obtained and assessed for inclusion with the predetermined selection criteria: AIT administered sublingually to children (0–18 years old) with confirmed allergic disease and language of publication. Only those trials were incorporated that reported clinical data and/or safety data and/or immunologic findings as outcome measures. Trials recruiting both adults and children were only included if the pediatric data were presented separately or if more than 50% of the active group were younger than 18 years.

Data Collection and Analysis

The data from the full-text articles were extracted independently by at least 2 reviewers per article (D.L.L., C.B.C., H.V.B., M.B.). Disagreement, if any, was resolved by discussion. Data on design, outcomes, and immunologic changes were abstracted in extraction tables (Table 1 and Table 2) and into the GRADE quality assessment sheet (eTable 1), as described in previous publications.^{18,51,52} Two of the authors independently performed GRADE quality assessment of the studies (D.L.L., E.C.). Information on safety was captured in a descriptive way. We tried to follow the World Allergy Organization grading system of systemic adverse events⁵³ in classifying these, whenever possible.

Results

Retrieved Articles

Fifty-six articles were identified as possible candidates for review (eTable 2). Of these, 22 were excluded because of age limits; 4 studies that included adults were kept in the analysis, because most participants were children.^{38,39,47,49} Five further articles were excluded because of administration route,⁵⁴ publication type,⁵⁵ and outcome measures outside the scope of this review.^{56–58} In all, 29 articles on SLIT in children will be analyzed in this review article.

Two manuscripts were on the same trial^{24,29}; thus, 28 clinical trials are reviewed.

Twenty-five articles reported clinical data (Table 1), pulmonary function test (PFT) results, specific and nonspecific bronchial challenge test results, and/or skin prick test (SPT) results (Table 2). Three trials studied exclusively safety data,^{48–50} and 1 trial studied only immunologic outcomes²¹ because the clinical results of this latter trial had been published previously.²² Sixteen other articles reported some provocation testing or immunologic outcomes (Table 2).

Design and Quality of the Studies

Thirteen trials had a DBPC design. One trial was double-blind, double-dummy with 2 active and 1 placebo arms. Six were randomized controlled trials (RCTs) and 3 were randomized trials with both groups receiving active treatment and no control group. One was an open controlled trial. The rest had an observational design.

The possible quality of the trials was analyzed with the GRADE system (eTable 1). Nine articles (8 studies) were assigned the maximum GRADE score of 4 for the whole trial or part of it, and 9 had a score of moderate quality (grade score of G3), leaving the rest with low or very low quality.

Allergen Extracts

The allergen extracts were preparations from European allergen manufacturers in 25 of the 28 studies analyzed (10 from ALK-Abelló, 8 from Stallergènes, 2 from Lofarma, and 2 from Allergopharma). In 2 DBPC trials^{26,44} and in 1 RCT⁴⁵ an aqueous extract was used from a US manufacturer (Greer Laboratories, Lenoir, North Carolina); 2 of these studies were with food allergens.

Fourteen trials administered pollen SLIT (11 grass, 3 tree) and 1 house dust mite (HDM) SLIT, including 1 trial with dual grass pollen–HDM immunotherapy.²⁶ *Alternaria*, peanut, milk, and mixed allergens were administered in one trial each.

SLIT allergen extracts are preparations in liquid form in 24 trials; in all but one⁴³ a glycerinated natural allergen was given. Four studies used SLIT grass tablets, one of them being an allergoid.³² No adjuvant extracts were used in the reviewed trials.

Clinical Outcomes

In most trials AR or rhinoconjunctivitis was the leading allergic disease, with some of the included patients also having mild asthma. In 5 trials the principal disease was allergic asthma, caused by HDM (n = 3), grass (n = 1), or tree pollen (n = 1), with this latter being a safety study.⁵⁰ Table 1 depicts details of all clinical trials performed from January 1, 2009, through December 31, 2012. In the right column differences found between active and placebo (control) groups are stated, with the corresponding statistical significance as reported in the articles. In some studies only intra-group differences were reported, comparing data before and after SLIT. The study order is according to the allergen administered, the allergic disease primarily treated (rhinitis or asthma), and the study quality. Findings of the studies are then discussed, adding quality of evidence to them (eg, G2, meaning GRADE score 2). Publications with only safety data are presented at the bottom of Table 1, and outcomes of provocation testing, SPT results, and immunologic responses can be found in Table 2. From 2009–2012 there were no studies published on preventive or pharmacoeconomic effects of SLIT in children.

Symptom and Medication Scores: Seasonal Allergens

Four high-quality trials (G4) show a reduction in symptoms and medication score with grass pollen SLIT. Three of these 4 trials were

Table 1
Clinical efficacy of SLIT in children: update (2009–2012)

Source	Q score	Age range, y	Active/ placebo (control), No.	Dropout, No.	Allergen, drops or tablet	Duration	Dose, μ g, dosing frequency	Dose vs SCIT	Disease	Manufacturer	Statistically significant differences	No statistically significant differences
Pollen												
Blaiss et al. ²⁰ 2011	4	5–17	175/169	33/29	Grass, tablet	6 m	15 μ g of Phl p 5 daily	NS	RC (A)	ALK	SLIT vs placebo: Daily symptom (25%), daily medication (81%), and total scores (26%) and QoL improved 18% ($P < .04$). Mechanistic study: see mechanistic table	SLIT-placebo: Asthma symptom score
Nieminen et al. ²¹ 2010 (subgroup of study by Valovirta et al. ²² 2006)	4	5–15	10 Low, 10 high, 10 placebo		Birch-alder-hazel mix, drops	2 y	24,000 SQ U/wk (3.6 μ g group 1), 200,000 SQ U/wk (30 μ g of group 1)	0.5 and 4.5	RC (A)	ALK		
Steinmach et al. ²³ 2012	4	6–18	20 Continuous, 20 preseasonal, 20 placebo	1/3/2	Grass, drops	2 y	10 μ g of group 5 daily continuously for 2 y; preseasonal: 2 for 6 mo	NS	RC (A)	Stallergènes	Both active groups vs placebo: significant improvement in medication and symptom scores. Preseasonal group vs placebo: significant reduction in medication score	Medication score continuous group: Asthma symptoms
Wahn et al. ²⁴ 2009	4	4–17	139/139	8/4	6-grass, tablet	8 m	25 μ g of group 5 daily, tablets	30	R (A)	Stallergènes	SLIT vs placebo: improved rhinitis sympt ($P = .01$) and medication ($P = .0064$) scores. Fewer days with medication intake ($P = .015$)	
Bufe et al. ²⁵ 2009	4	5–16	126/127	12/7	Phleum pratense, tablet	6 m	15 μ g of Phl p 5 daily	30	R (A)	ALK	Active vs placebo: Significant reduction in RC symptoms score (-24%), asthma score (-64%), randomized controlled medications (-34%), and well days ($+28\%$). All $P < .03$.	
Swamy et al. ²⁶ 2012	3	5–58	20/10	0/0	Dual grass and HDM, drops	12 m	15 μ g of Phl p 1	1–2.8 for each dose	RC	Greer	Active-placebo: Rhinoconjunctivitis symptom score, medication score and combined score reduced at 12 and 24 m (12 m after treatment discontinuation) ($P < .001$)	SLIT vs placebo: Mean number of well days
Wahn et al. ²⁷ 2012	3	4–12	158/49	26/2	6-grass drops	8mo	40 μ g of group 5 daily	NS	RC (A)	All manufacturers	SLIT vs placebo: Change in pre-post treatment higher for symptom-medication, symptom, and medication scores. SLIT higher rate of positive response ($\geq 40\%$ decrease of the AUC of the SMS).	
Pajno et al. ²⁸ 2011	3	8–16	40 Continuous, 40 coseasonal	3/5	Grass drops	CONT3yrs COS: 3x4m	8 μ g of group 5, 5 times per week	NS	RA	Stallergènes	Continuous vs coseasonal: first year: Symptom plus medication, symptom, chest symptom and medication scores improved more in continuous SLIT groups.	Third year: no difference in clinical outcomes between continuous vs coseasonal SLIT

(continued on next page)

Table 1 (continued)

Source	Q score	Age range, y	Active/ placebo (control), No.	Dropout, No.	Allergen, drops or tablet	Duration	Dose, μ g, dosing frequency	Dose vs SLIT	Disease	Manufacturer	Statistically significant differences	No statistically significant differences
Haiken et al. ²⁹ 2010 (additional data from the study by Wahn et al. ²⁴ 2009)	3	5–17	278 Total: 131 SLIT, 135 placebo	?	Grass tablet	6 m	25 μ g of Phl p 5 daily (300 IR)		RC	Stallergènes	Active-placebo: Total symptom score reduced at whole and peak pollen season. Nasal and ocular symptoms reduced. Less rescue medication during whole and peak pollen season.	
Panzner et al. ³⁰ 2011 (open extension of previous DBPC 12-month trial by Panzner et al. ³¹ 2008) Agostinis et al. ³² 2009	2.5	Mean, 17/6	26 SLIT, 25 supralingual	8	6-Grass, drops	12 m (+24 m open)	11.2 μ g of group 5 3 times per week	Total: 20 times	R	Sevapharma	Pre-post treatment supralingual and SLIT: symptom, medication, and combined score reduced year by year.	SLIT vs supralingual: trend for more symptom and medication reduction in SLIT (NS)
Almadiafar et al. ³³ 2012	0	5–18	12 SLIT, 12 placebo	2/2	Lolium, drops	6 m	900 IR 3 times per week	100?	RC	Stallergènes	SLIT vs control: VAS improved after first and second year (both $P < .05$). SLIT pre-post treatment: reduction in sympt ($P < .05$)	Control pre-post treatment: no reduction in symptoms
Stelmach et al. ³⁴ 2009	2	6–17	20 SLIT, 15 placebo	5/10	Grass, drops	Preseasonal for 2 y	10 μ g of group 5 grass drops daily	NS	A	Stallergènes	SLIT vs placebo: asthma symptoms ($P < .002$), nasal sympt ($P < .04$), nasal and asthma symptoms, asthma medication and nasal, asthma, and medication scores (both $P < .001$)	SLIT vs placebo: ocular symptoms, total asthma, nose, and eye symptoms
HDM Yonekura et al. ³⁵ 2010	3	7–15	20/11	1/2	Mite, drops	1 y	0.5 μ g of Der f 1 once a week	20	RC	TOR	Active-placebo: week 30: reduced symptom score. Initial (weeks 0–3) to end (weeks 37–40) active group: Decrease in symptoms and symptom-medication score	Active-placebo: combined symptom-medication score
De Bot et al. ³⁶ 2012	2	6–18	126/125	15/17	Mite, drops	2 y	2.03 μ g of Der p 1 2 times per week; total cumulative dose (2 y): 435 μ g	NS	RC	ART	Total nasal symptom score, QoL, medication score, well days	
Han et al. ³⁷ 2012	1	6–18 and ≥ 18	54 Children/22 adults	0/0 ^b	HDM, drops	1 y	Pangramin SLIT: 6 μ g/mL of Der 1 and Der 2	NS	R	ALK	Pre-post treatment pediatric group: improvement in total symptom score and in each rhinitis symptom ($P < .05$). Medication use reduced, but no statistical calculations made.	Pediatric vs adult patients: no difference in symptom (NS), med score trend more improvement in pediatric group ($P = .06$)
Lee et al. ³⁸ 2011	1	Mean, 14.7 (range, 4–53)	70 Mono-sensitized, 64 poly-sensitized	NS	HDM, drops	12 m (first 30 d = up-dose)	5 drops of 1,000 STU/mL of Der p and Der f 3 times per week	NS	R	ALK	Mono- and polysensitized symptom and medication scores: all improved	No difference between mono- and polysensitized in any variable
Trebuchon et al. ³⁹ 2012	1	5–18 (n = 735) (1289 patients total)			HDM, drops	≥ 2 y	Variable, most 300 IR daily	NS	70% R, 50% A	Stallergènes, some ALK	More descriptive study of how SLIT is given, dosing schedules, duration, etc. Treatment (very) effective, according to physician: 82% Reduction in asthma medication: 26% stopped taking ICSS	

(continued on next page)

Table 1 (continued)

Source	Q score	Age range, y	Active/ placebo (control), No.	Dropout, No.	Allergen, drops or tablet	Duration	Dose, μ g, dosing frequency	Dose vs SCIT	Disease	Manufacturer	Statistically significant differences	No statistically significant differences
Other Allergens												
Kim et al, ⁴⁴ 2011	4	1–11	11/7	0/0	Peanut, drops	12 m	2000 μ g daily (8 pumps)		Peanut allergy	Greer	DBPC food challenge: ingestion of median cumulative dose of peanut protein SLIT 1,710 mg; placebo: 85 mg ($P < .011$).	
Keet et al, ⁴⁵ 2012	4	6–11	10 SLIT, SLIT start then: 10 high-dose OIT, 10 low-dose OIT	0/1/1	Milk protein drops	14 m	SLIT, 7 mg; high-dose OIT, 2,000 mg; low-dose OIT, 1,000 mg of milk protein daily	NS	Cow milk allergy	Greer	DBPC food challenge passed by more OIT patients vs SLIT alone (1 SLIT, 6 SLIT and low-dose OIT, 8 SLIT and high-dose OIT)	Regained hyperreactivity after 6-wk milk avoidance: 3 of 6 desensitized low-dose OIT patients, 3 of 8 high-dose OIT patients
Acquistapace et al, ⁴⁶ 2009	0–1	6–18	90 SLIT/81 controls	NA	Several, drops	2 y	Varied	NS	RC (A)	ALK (SLIT)	SLIT vs controls: reduced symptoms, medication score, and new sensitizations	SLIT vs control: asthma symptoms
Pozzan et al, ⁴⁷ 2010	2	10–65	34 SLIT/18 controls	1/0	<i>Alternaria</i> drops	36 m	1 vial of SLIT once daily	NS	R (A)	ALK	Results of pediatric group not separated: Primary outcome: active vs control: symptom score reduced by VAS ($P = .0002$); ICS dose reduced ($P < .01$). Active pre-post: medication score significantly reduced in SLIT but not control group.	Active vs control: No medication score reduced
Trials With Only Safety Data												
Seidenberg 2009 ⁴⁸ SAFETY	1	5–17	193 SLIT	10 (+50 <4-m treatment)	Grass and/or tree, drops	4 m	Started with ultrarush up-dosing: 30–90–150–300 IR each 30 min (μ g?)	Final dose approximately 30 times the SCIT dose	RC (A)	Stallergènes	During up-dosing: 60 patients (31%) reported 117 predominantly mild and local AEs, which resolved within 150 min. During maintenance: 562 AEs; most frequent local AEs were oral pruritus, burning sensation, lip or tongue swelling, and GI symptoms; the most frequent systemic AEs were RC and A. One clinically significant asthma event in an 11-year-old asthmatic boy: SLIT was resumed after 4 days	
Roger et al, ⁴⁹ 2011 (total population, 218, 4–64 years old; safety trial)	2	4–15	122	None	HDM, drops	Up-dose	Every 30 min: 30–60–120–240 IR		R and/or A	Stallergènes	8 systemic reactions (3 moderate), all continued SLIT. Higher frequency of AEs in asthmatic patients. No difference in severity of AEs in patients younger than 15 y.	Slight increased frequency in AEs in patients younger than under 15 y (59.3% of AE were in pediatric patients, whereas only 53.7% of all patients were pediatric; NS)
Mösges et al, ⁵⁰ 2010 (safety trial)	4 2	6–14	27/27	0/0	Tree pollen, drops	Up-dosing	30–90–150–300 IR each 30 min (μ g?)		A	Stallergènes		During up-dosing: active-placebo: serious AEs: no difference. During up-dosing: Active-placebo: PFT change: no difference

Abbreviations: A, asthma; AE, adverse event; AUC, area under the curve; B2, β_2 -agonist; CET, cetirizine; DBPC, double-blind, placebo-controlled; GI, gastrointestinal; HDM, house dust mite; ICS, inhaled corticosteroid; IR, index of reactivity; MEF₂₅, midexpiratory flow at 25% pulmonary capacity; NS, not stated or not applicable; OIT, oral immunotherapy; PFT, pulmonary function testing; Q, quality assessment according to Grading of Recommendations Assessment, Development and Evaluation; QoL, quality of life; R, rhinitis; RC, rhinoconjunctivitis; SAE, serious adverse event; SCIT, subcutaneous immunotherapy; SLIT, sublingual immunotherapy; SMS, symptom medication score; SQ, subcutaneous; VAS, visual analog scale.

^aParental passive smoke (at least 20 cigarettes per day).

^bNo dropouts mentioned and report of symptom scores on all included patients suggesting no one dropped out.

Table 2

Immunologic and provocation testing

Source	Lung function	Inflammatory markers	Immunologic markers
Pollen			
Blaiss et al, 2011 ²⁰			SLIT vs placebo: Phl p 5 specific IgG4 and IgE-blocking factor levels were higher at peak and end of the grass pollen season.
Bufe et al, 2009 ²⁵			SLIT vs placebo: increase in IgG4 ($P < .001$) and in IgE blocking factor ($P < .001$). Seasonal IgE peak blunted in active (NS).
Nieminen et al, 2010 ²¹			Patients with elevated symptom and medication score: increase in allergen-induced PBMC mRNA IL-17 expression; a positive and dose-dependent correlation SMS and IL-17 production. High-dose group vs placebo at 2 y: increase in FOXP3 mRNA expression. FOXP3 mRNA changes correlate with IL-10 and TGF- β mRNA.
Pajno et al, 2011 ²⁸			First year: continuous SLIT: increase in grass IgE, second to third years: no change grass IgE Continuous: First to third years: increase in grass IgG4, first to second years: larger increase in grass IgG4 vs coseason, third year: no difference
Panzner et al, 2011 ³⁰		SLIT and supralingual before vs after treatment: both reduction in SPT ($P < .0001$).	SLIT vs supralingual: larger increase in specific IgG4
Stelmach et al, 2009 ³⁴	SLIT vs placebo: FEV ₁ improved ($P = .005$), FEF _{25%-75%} only trend	SLIT vs placebo: methacholine PC ₂₀ trend for improvement ($P = .058$), nasal provocation test: no difference	SLIT vs placebo: no difference in specific IgE or total IgG4
Stelmach et al, 2012 ²³	No changes in morning PEF, FEV ₁ , and methacholine PD ₂₀ within or among any of the 3 groups	Both active groups vs placebo: significant decrease in FeNO level comparable in both active groups	Peripheral blood: induction CD4CD25Foxp3-positive cells no difference between groups
Swamy et al, 2012 ²⁶		SLIT vs placebo: nasal provocation test (nasal disk challenge): $P < .0001$ for GP at 18 mo (6 mo after treatment) (HDM not performed). SLIT vs placebo at 12 mo: reduced SPT GP and HDM ($P < .05$)	SLIT vs placebo: specific IgE reduction and IgG4 increase (both $P < .05$) at 12, 18, and 24 m, no change in control Oak immunoglobulins. SLIT GP and HDM, pre-post treatment at 24 mo: Basophil activation after GP and HDM stimulation reduced pre-post treatment ($P < .0001$). No difference with Oak or in placebo group. Epigenetic modification of induced Treg cells in dual SLIT patients: decreased DNA methylation ^a in CD45RO1 memory Treg cells after 12-month dual SLIT ($P < .05$). Increase in Foxp3 transcript levels of memory Treg cells (DNA methylation was augmented and Foxp3 transcript reduced in allergic patients without SLIT compared with healthy controls.) Tolerant vs nontolerant dual SLIT patients: Already at baseline tolerant patients' memory Treg cells had increased expression of Foxp3 ($P < .05$), programmed cell death protein 1, and IL-10 (NS). Six months after treatment increased number memory or induced Treg cells in tolerant patients ($P < .05$).
Wahn et al, 2009 ²⁴			SLIT vs placebo: greater increase in specific IgG4. No change in IgE.
Wahn et al, 2012 ²⁷			SLIT vs placebo, pre-post treatment: specific IgE: no difference; SLIT increase in IgG1 and IgG4
HDM			
Eifan et al, 2010 ⁴¹	SCIT and SLIT each vs pharmaceutical group: improved nasal provocation test ($P = .005$ and $.01$, respectively). No difference in lung function nor methacholine PD ₂₀	SCIT and SLIT each vs pharmaceutical group: reduced skin prick test reactivity at 12 mo. SLIT: $P = .006$ for Der p and $P = .01$ for Der f.	SCIT and SLIT vs pharmaceutical group: reduction serum specific IgE. SLIT vs pharmaceutical group: IL-10 increase. No difference in other T _H 1-T _H 2 cytokines in PBMC cultured with recombinant Der p 1 and Bet v 1.
Han et al, 2011 ³⁷			Pre-post treatment pediatric group: Total IgE, no change. Eosinophils decreased (NS) and serum eosinophilic cationic protein reduced ($P < .05$). Pediatric vs adult group: no differences in any of these 3 immunologic markers.

(continued on next page)

Table 2 (continued)

Source	Lung function	Inflammatory markers	Immunologic markers
Keles et al, 2011 ⁴⁰	FEV ₁ increased in SCIT → SLIT vs pharmaceutical group; nonspecific bronchial provocation test: result turned negative in 4/7 SCIT → SLIT patients (NS). Specific nasal provocation test: improved in all active groups vs pharmaceutical group	SPT: SCIT reduced at 12 mo	IgE total and specific: no change at 12 mo. IgG4 and IgG4/IgE ratio: increase in SCIT and SCIT → SLIT vs pharmaceutical group. No change in SLIT. Der p 1–stimulated PBMC supernatant: TGF- β and IL-10: increase from 4 mo on in all 3 groups. IFN- γ : increase at 4 mo, back to baseline at 12 mo in all 3 groups. IL-17: NS reduction in all 3 active groups.
Marogna et al, 2011 ⁴³	CET and passive smoke pre-post treatment: FEV ₁ , MEF ₂₅ worse. SLIT and passive smoke: MEF ₂₅ improved; SLIT and nonsmoke pre-post treatment: FEV ₁ and MEF ₂₅ improved	CET and passive smoke: methacholine PD ₂₀ worse, eosinophils increased. SLIT and passive smoke: methacholine PD ₂₀ improved. SLIT and no smoke: methacholine PD ₂₀ improved and eosinophils reduced	
Yukselen et al, 2012 ⁴²	FEV ₁ improved in SCIT and SLIT vs baseline. HDM nasal challenge improved in SCIT and SLIT vs baseline. Bronchial challenge improved vs baseline in SCIT.	Titration skin prick tests: reduced in SCIT and SLIT vs baseline. Nasal eosinophils increment after challenges: SCIT and SLIT significantly reduced vs placebo. SCIT vs baseline: reduction BAL eosinophils after bronchial HDM challenge	SCIT vs SLIT: greater increase in IgG4. SLIT and SCIT vs baseline: IgE HDM reduction, IL-10 increase. SCIT vs baseline: IgG4 increase. IFN- γ : no differences
Food			
Keet et al, 2012 ⁴⁵		All groups: reduced end point titration skin prick testing	All groups: increased IgG4 levels and decreased constitutive CD63 and CD203c expression. OIT groups only: decreased CM-specific IgE and reduced spontaneous basophil histamine release
Kim et al, 2011 ⁴⁴		SPT wheal reduced at 12 mo in active vs placebo group	Active vs placebo: Lower percentage of CD63 ⁺ basophils after low-dose peanut stimulation ($P = .009$). Peanut specific IgE: increase at 4 mo and reduction at 12 mo. Peanut specific IgG4: increased at 12 mo. Active vs placebo: IL-5 decreased ($P = .015$), IL-13 decreased (NS), IL-10 and IFN- γ no difference, % Treg cells increased (NS).

Abbreviations: BAL, bronchoalveolar lavage; CET, cetirizine; CM, cow's milk; FEF_{25%-75%}, forced expiratory flow between 25% and 75%; FeNO, fractional exhaled nitric oxide; FEV₁, forced expiratory volume in 1 second; GP, grass pollen; HDM, house dust mite; IFN- γ , interferon γ ; IL, interleukin; MEF₂₅, midexpiratory flow at 25% pulmonary capacity; mRNA, messenger RNA; NS, not stated or not applicable; OIT, oral immunotherapy; PBMC, peripheral blood mononuclear cell; PC₂₀, provocation concentration that caused a decrease in forced expiratory volume in 1 second of 20%; PD₂₀, provocation dose that caused a decrease in forced expiratory volume in 1 second of 20%; PEF, peak expiratory flow; SCIT, subcutaneous immunotherapy; SLIT, sublingual immunotherapy; SMS, symptom medication score; SPT, skin prick test; TGF- β , transforming growth factor β ; Treg, T-regulatory.

^aDecreased CpG methylation within the Foxp3 locus is related to more stable suppressive activity of Foxp 3 Treg cells.

with SLIT tablets given pre-seasonally for 6 to 8 months. Similar efficacy was shown by the 4 moderate-quality trials with grass pollen SLIT. Interestingly, the findings of Stelmach et al²³ (G4) suggest that SLIT drops during 2 years might have a better result when given pre-seasonally instead of continuously. Meanwhile, a slightly lower dose of the same grass pollen SLIT administered only co-seasonally for 4 months each year did not show clear efficacy during the first 2 seasons to reach levels of clinical improvement similar to the continuously administered product only until the third year of treatment.²⁸ (G3) Because this latter trial did not include a control group, conclusions should be drawn with caution. The same holds true for a trial in which clinical score improvement was documented when comparing pretreatment and posttreatment values for sublingual with supralingual immunotherapy because a control group was included in this study design only during the first year of the trial.³⁰ (G2–3) Blaiss et al²⁰ separately analyzed pediatric (5–11 years) and adolescent subgroups (12–17 years), showing differences in symptom plus medication score in favor of grass AIT in both (32% and 16%, respectively).

Most data from asthma outcomes with pollen SLIT came from studies where seasonal AR was the leading disease and thus are studies not adequately designed or powered to detect changes in asthma symptoms or medication. The only grass pollen SLIT study in pediatric asthma reports encouraging data: asthma clinical parameters improved after 2 years of pre-seasonal treatment comparing the active with the placebo group, reaching statistical significance even though the study was underpowered³⁴ (G2).

Mold allergy was addressed in one RCT of *Alternaria* SLIT in respiratory allergy⁴⁷ (G2). After 3 years symptom scores and inhaled corticosteroid use reduced, although total medication scores did not show any difference between the active and control groups.

Symptom and Medication Scores: Perennial Allergens

In the time span of our review there was one moderate-quality study investigating HDM SLIT in pediatric AR and 4 studies of (very) low quality (G1–G2). From these trials no clear conclusions can be drawn because in a placebo group was included in only 2 trials and only some trials showed improvement in nasal symptom and/or medication scores^{35,37,38} (G1–3), whereas others did not³⁶ (G2). As such, the best quality evidence of HDM SLIT efficacy for AR symptoms comes from 4 pediatric asthma trials. The results are also not uniform in these trials because AR symptoms improved compared with a randomized control group in 2 studies,^{41,43} (G2 and G4) but not in the other 2 studies^{40,42} (G2.5–3).

However, none of these is a simple SLIT trial; each has its peculiarities worth commenting. Keles et al⁴⁰ divided 60 children randomly to receive HDM SCIT, SLIT, SCIT build-up followed by SLIT maintenance (SCIT → SLIT), or pharmacotherapy alone. In comparison to the pharmaceutical group in the SCIT → SLIT group rhinitis, asthma symptoms, asthma attacks, and medication all improved at 12 months, reaching statistical significance even though the groups were small. In the SCIT group no rhinitis symptom improvement was seen, and in the SLIT group only asthma medication scores

improved. Two pediatric SLIT-in-asthma trials compared SLIT with SCIT and an open control⁴¹ (G3–G4) or placebo⁴² (G2.5–G3.5). The former found that total rhinitis symptoms, asthma symptoms, and medication improved in both active groups compared with the pharmaceutical group, but in the latter no statistically significant benefit of SLIT over placebo was found. SLIT and SCIT were found to be equivalent for all parameters, with the exception of asthma symptom scores in the trial of Yukselen et al. However, neither of these 2 trials was adequately powered to show differences between both active groups, so no conclusion can be drawn in this respect. In the last pediatric SLIT in asthma trial, Marogna et al⁴³ randomized 68 children with AR and intermittent asthma and positive methacholine provocation test results to receive SLIT or cetirizine. Half of each group consisted of children with exposure to high levels of environmental tobacco smoke in their homes. After 3 years, in passive smokers the methacholine challenge greatly improved in the SLIT group vs the cetirizine group (G3). The other clinical outcomes improve in the passive smoking SLIT group, whereas in the cetirizine group all parameters deteriorate. In the non-passive smoking groups, SLIT improved clinical scores and medication use, whereas there was no change with cetirizine. Unfortunately, no clear between-group comparisons are reported in the published document.

The overall balance of the efficacy of SLIT with HDM as part of the integral treatment in pediatric asthma as studied in these trials is positive, but because the trials are small scientific quality is not optimal.

Symptom and Medication Scores: Dual SLIT With Combined Grass Pollen–HDM Extract

One trial deserves special mentioning because this is the first trial on dual SLIT in children. Swamy et al²⁶ (G3) conducted a DBPC-RCT of dual SLIT administering a grass pollen–HDM glycerinated solution during 12 months. The investigators were able to show a statistically significant improvement in the rhinoconjunctivitis symptom score, medication score, and combined score at 12 and 24 months (12 months after treatment discontinuation). Immunologic markers were also tested (Table 2).

Clinical Outcomes of SLIT With Food Allergens

We found 2 trials on SLIT for food allergy in children, both (partly) conducted at Duke University Medical Center, Durham, North Carolina. Kim et al⁴⁴ reported beneficial effects after 12 months of daily SLIT with a glycerinated peanut extract (Greer Laboratories): DBPC food challenges showed an increase in the median cumulative dose of peanut in the active group vs the placebo group (1,710 vs 85 mg, $P < .01$). The second study design was more refined: 30 children with milk allergy were randomized to receive SLIT or SLIT build-up followed by oral immunotherapy (OIT) at low or higher dose. At the end of this 140month trial a DBPC food challenge proved OIT to be superior to SLIT alone. Even so, 3 of 8 patients in the high-dose OIT group who performed best regained hyperreactivity after only 6 weeks of milk avoidance, putting in doubt if true tolerance can be obtained with milk OIT.

PFT and Nasal and Bronchial Provocation Testing

No provocation testing was performed in any of the tablet studies. The effect of grass pollen SLIT drops on nasal provocation testing was documented by 2 investigators,^{26,34} with the higher-quality trial (G3) recording a reduction in specific nasal hyperreactivity. The effect of grass pollen SLIT on lung function parameters was investigated in 2 studies^{23,34}: exhaled nitric oxide was reduced after grass pollen SLIT; however, in PFTs no clear signal could be detected and methacholine bronchial challenges showed no improvement.

HDM SLIT improved specific nasal hyperreactivity in all 3 high-quality trials that investigated this parameter. However, nonspecific methacholine provocation dose that caused a decrease in forced expiratory volume in 1 second of 20% augmented only in 1 of the 4 studies⁴³ that included this measurement and PFTs improved compared with pretreatment values but showed only a trend for superiority compared with placebo.

Skin Prick Testing

SPT reactivity was investigated in pediatric SLIT trials with grass pollen³⁰ and HDM,^{40,41} in the dual grass pollen–HDM trial,⁴⁶ and in both food SLIT trials. It improved in all but one HDM study.⁴⁰

Efficacy Summary

Table 3 summarizes the evidence concerning clinical efficacy in children with respiratory allergies of SLIT with grass pollen, *Alternaria*, and HDM. This summary table is based on all reviewed studies and their scientific quality (GRADE score).

Safety Data

Although in almost all trials safety outcomes were mentioned, 3 trials investigated exclusively safety issues (2 large observational studies and 1 with a DBPC design)⁵⁰ (G4) (eTable 3). All 3 studies used an ultrarush, 90-min build-up phase of high-dose SLIT. The latter explored the safety of tree pollen SLIT in asthmatic children. During up-dosing no differences in serious adverse events (G4) or in PFT results (G2) were found between the active and placebo groups. Roger et al⁴⁹ (G2) investigated HDM SLIT drops in patients with rhinitis and/or asthma. Eight mild-moderate systemic adverse events were reported, with a higher frequency among asthmatic patients, but none discontinued SLIT. The treatment was equally well tolerated by children younger than 15 years in comparison with adult patients. Meanwhile, Seidenberg et al⁴⁸ showed coseasonal rush build-up is relatively well tolerated by rhinitis patients.

Safety issues reported in the rest of the trials were frequent and mild, mostly consisting of local reactions in the oral cavity: oral pruritus, throat irritation, and stomatitis (32%–85% in the active group vs 2%–20% in the placebo group). There were also mild systemic symptoms as eye, nose, or ear pruritus. No life-threatening systemic adverse events were reported in any of the trials, with a total 2469 children receiving active treatment. However, in the milk SLIT and OIT trial, 1 SLIT and 4 OIT children received adrenaline. Treatment-related discontinuation ranged from 0% to 7.4%.²⁰ In the real-life retrospective study by Trebuchon et al,³⁹ this number was 8%. In the 2 trials with SLIT and SCIT, treatment-related discontinuations were only reported in the SCIT groups. In some trials, patients with abdominal symptoms were referred with higher frequency in the active group. Epinephrine was used only in the trial conducted by Blaiss et al.²⁰ During this US DBPC trial with grass pollen SLIT tablets, epinephrine was administered to 3 children (2 in the active group and 1 in the placebo group), with only one administration due to a reaction to the tablet: this patient experienced lip angioedema, slight dysphagia, and intermittent cough with no other symptoms immediately after the first dose; epinephrine administration resolved this moderate local reaction (as judged by the investigator) and the patient discontinued participation in the trial.

Immunologic Findings

Several studies have documented an increase in specific IgG4 and IgE-blocking factor; some also documented an increase in serum interleukin (IL) 10 levels. Specific IgE levels were generally reduced, but in some trials they showed an initial surge. In peripheral blood mononuclear cells Nieminen et al documented

Table 3

Summary of the evidence

Statistically significant difference for active vs placebo (control)	Studies without effect	Evidence ^{a,b}
Grass and/or Birch		
AR symptoms	None	
Blaiss et al, 2012 (G4)		
Wahn et al, 2009 (G4)		
Bufe et al, 2009 (G4)		
Halken et al, 2010 (G3)		
Total Tablets:		Yes: ●●●●
Swamy et al, 2012 (G3) (dual grass and HDM at 12 mo and 12 mo after treatment discontinuation)		
Wahn et al, 2012 (G3)		
Stelmach et al, 2012 (G4) (preseasonal and continuous SLIT)		
Pajno et al, 2011 (G3) (first year)		
Panzner et al, 2011 (G2.5) (pre-post treatment)		
Stelmach et al, 2009 (G2)		
Total Drops:		Yes: ●●●● Yes: 12 mo after discontinuation: ●●●○
Medications	Stelmach et al, 2012 (G3) (continuous SLIT)	
Blaiss et al, 2012 (G4)		
Wahn et al, 2009 (G4)		
Bufe et al, 2009 (G4)		
Total Tablets:		Yes: ●●●●
Swamy et al, 2012 (G3) (dual grass and HDM at 12 mo and 12 mo after treatment discontinuation)		
Halken et al, 2010 (G3)		
Wahn et al, 2012 (G3)		
Stelmach et al, 2012 (G4) (preseasonal SLIT)		
Pajno et al, 2011 (G3) (first year)		
Panzner et al, 2011 (G2.5) (pre-post treatment)		
Stelmach et al, 2009 (G2)		
Total Drops:		Yes: ●●●○ Yes: 12 mo after discontinuation: ●●●○
Symptoms and medications	None	
Blaiss et al, 2012 (G4)		
Total Tablets:		Yes: ●●●●
Swamy et al, 2012 (G3) (dual grass and HDM at 12 mo and 12 mo after treatment discontinuation)		
Wahn et al, 2012 (G3)		
Panzner et al, 2011 (G2.5) (pre-post treatment)		
Stelmach et al, 2012 (G4) (preseasonal and continuous)		
Pajno et al, 2011 (G3) (first year)		
Total Drops:		Yes: ●●●● Yes: 12 mo after discontinuation: ●●●○
Nasal provocation		
Swamy et al, 2012 (G3) (dual grass and HDM)	Stelmach et al, 2009 (G2)	Yes: 6 mo after SLIT: ●○○○
Conjunctival provocation	None	No data
Asthma symptoms	Blaiss et al, 2012 (for asthma G3), and Stelmach et al, 2012 (G3)	
Bufe et al, 2009 (G4)		
Total Tablets:		Yes: ●○○○
Pajno et al, 2011 (G3) (first year)		
Stelmach et al, 2009 (G2)		
Total Drops:		Yes: ●○○○
Asthma medication	None	
Stelmach et al, 2009 (G2)		Yes: ●●○○
Lung function tests and bronchial provocation	PFT: Stelmach et al, 2012 (G3), and methacholine: Stelmach et al, 2012 (G3)	
PFT: Stelmach et al, 2009 (G2)		
Methacholine: Stelmach et al, 2009 (G2) (trend $P = .058$)		
FeNO: Stelmach et al, 2012 (G3)		
Total Drops:		PFT: No: ●○○○ Methacholine: No: ●●○○ FeNO reduction: Yes: ●●●○

(continued on next page)

Table 3 (continued)

Statistically significant difference for active vs placebo (control)	Studies without effect	Evidence ^{a,b}
SPT reactivity Swamy et al, 2012 (G3) (dual grass and HDM at 12 mo) Panzner et al, 2011 (G2.5) (pre-post treatment) Total Drops:	None	Yes: ●●○
New sensitizations	None	No data
Alternaria		
Improvement in AR symptoms Pozzan et al, 2010 (G2)	None	Yes: ●●○
Improvement in medication score Pozzan et al, 2010 (G2) (pre-post treatment)	Pozzan et al, 2010 (G2)	No: active-control: ●●○ Yes: pre-post treatment: ●●○
Symptoms and medications	No data	No evidence
Asthma medication (ICSs) Pozzan et al, 2010 (G2)	None	Yes: ●●○
HDM		
AR symptoms Eifan et al, 2012 (G4) Swamy et al, 2012 (G3) (dual grass and HDM at 12 mo and 12 mo after treatment discontinuation) Yonekura et al, 2010 (G3) Total	Keles et al, 2011 (G3) (SLIT group), Yukselen et al, 2012 (G2.5), de Bot et al, 2012 (G2)	Yes: ●●○ Yes: 12 mo after: ●●○
Medications Swamy et al, 2012 (G3) (dual grass and HDM at 12 mo and 12 mo after treatment discontinuation) Symptoms and medications Swamy et al, 2012 (G3) (dual grass and HDM at 12 mo and 12 mo after treatment discontinuation)	Keles et al, 2011 (G3) (SLIT group), Yukselen et al, 2012 (G2.5), de Bot et al, 2012 (G2) Yonekura et al, 2010 (G3)	No: ●●○ Yes: 12 mo after: ●●○ Contradictory
Nasal provocation Keles et al, 2011 (G4) Eifan et al, 2012 (G4) Yukselen et al, 2012 (G3.5) Total	None	Yes: ●●●●
Asthma symptoms Eifan et al, 2012 (G4)	Keles et al, 2011 (G3) (SLIT group), Yukselen et al, 2012 (G2.5)	Yes: ●○○○
Asthma medication Keles et al, 2011 (G4) (SLIT group) Eifan et al, 2012 (G4) Total	Yukselen et al, 2012 (G2.5)	Yes: ●●●●
Pulmonary function tests No positive studies	Eifan et al, 2012 (G4) (methacholine), Keles et al, 2011 (G3) (SLIT group), Yukselen et al, 2012 (G2.5)	No: ●●○
Specific/specific bronchial challenge Marogna et al, 2011 (G3) (passive smokers)	Eifan et al, 2012 (G4) (methacholines), Keles et al, 2011 (G3) (SLIT group), Yukselen et al, 2012 (G2.5)	Methacholine: No: ●●○
SPT reactivity Eifan et al, 2012 (G4) (12 mo) Swamy et al, 2012 (G3) (dual grass and HDM at 12 mo) Total	Keles et al, 2011 (G3) (SLIT group)	Yes: ●●●●
New sensitizations	Not studied	No evidence
Prevention asthma	Not studied	No evidence

Abbreviations: AR, allergic rhinitis; FeNO, fractional exhaled nitric oxide; HDM, house dust mite; PFT, pulmonary function test; SLIT, sublingual immunotherapy; SPT, skin prick test.

^a●●●●, high; ●●○, moderate; ●○○, low; ●○○○, very low.

^bYes/no: there is scientific evidence to support there was statistically significant improvement (yes) or not (no) of the item stated in the left column of each line, in patients undergoing SLIT compared with patients from the placebo (control) group. In some studies only intragroup statistically significant improvement (pre- vs posttreatment) was documented: this was not considered acceptable evidence and is not documented in this table.

a increase in Foxp3 messenger RNA (mRNA) expression, which correlated with an increase in IL-10 and tumor growth factor β mRNA, and allergen-induced peripheral blood mononuclear cell IL-17 mRNA expression correlated with symptom medication scores. However, Stelmach et al found no difference between the active and placebo groups in the induction of CD4CD25Foxp3-positive cells. Both milk⁴⁵ and peanut⁴⁴ trials showed a reduction in the percentage of CD63⁺ basophils after SLIT.

The dual SLIT trial by Swamy et al intensively studied immunologic changes. Basophil activation after specific stimulation was reduced after treatment (grass pollen and HDM) but remained unchanged after stimulation with a third allergen. Decreased CpG methylation within the Foxp3 locus is believed to be related to a more stable suppressive activity of Foxp3 T-regulatory (Treg) cells. Thus, these investigators detected epigenetic modification of induced Treg cells after 12 months of dual SLIT: DNA methylation in CD45RO1 memory Treg cells was decreased and Foxp3 transcript levels of memory Treg cells was augmented, with opposite findings in allergic patients not receiving SLIT compared with healthy controls. Finally, comparing those SLIT patients who 6 months after the trial had become tolerant vs nontolerant SLIT patients, the memory Treg cells of the tolerant patients had an increased expression of Foxp3 ($P < .05$) and programmed cell death protein 1 already at baseline. The investigators suggest that these Treg cell markers might be predictive of clinical tolerance.²⁶ In trials where both SLIT and SCIT were given, immunologic changes were usually more marked after SCIT.^{40,42}

Discussion

We analyze articles published on SLIT in the pediatric age group published between 2009 and 2012. Without restricting for study design we found a total of 29 articles that met the inclusion criteria, corresponding to 28 trials; a total of 2469 patients were treated in the SLIT group (2127 were analyzed on efficacy and 2225 on safety). After analyzing articles on their scientific quality with the GRADE system, we composed a summary table in which all evidence for the efficacy of SLIT in children is expressed per allergic disease and per allergen. Only a statistically significant difference between the active and placebo (control) groups was considered valuable evidence because such intragroup improvements are not included in this table.

Compared with a similar analysis of data on pediatric SLIT up until 2008,⁵⁹ the efficacy of grass pollen SLIT drops in reducing AR symptoms and medication is reconfirmed and new evidence for efficacy of grass pollen tablets is added. High-quality new evidence shows efficacy of dual grass-HDM SLIT. Also, a prolonged effect, reducing the combined rhinitis symptom medication score still 1 year after treatment discontinuation, was shown. Only one asthma trial was conducted with pollen SLIT in children in the time span of our review.³⁴ Thus, for seasonal asthma there is moderate evidence of a reduction in exhaled nitric oxide, but the quality of evidence for medication reduction stays low and the effect of grass pollen SLIT on asthma symptoms, PFT results, and nonspecific bronchial hyperreactivity is uncertain.

There is some new stimulating evidence for SLIT with *Alternaria* in children with respiratory allergy⁴⁷ (G2), but further trials are needed to improve the strength of the evidence and give recommendations.

For HDM SLIT in children there were 9 new trials. However, most of these trials were with small groups of patients, and half of them did not directly investigate the efficacy of SLIT against placebo (control). Even so, evidence of moderate-high quality could be added to its efficacy in the control of nasal symptom, the reduction in nasal specific hyperreactivity, and the reduction in asthma

medication. No effect was documented in reducing rhinitis medication, asthma symptoms, PFTs, or nonspecific bronchial hyperreactivity. High-quality evidence shows SPT reactivity reduces with both grass pollen and HDM SLIT. We found no new data on the preventive effect of SLIT in children.

For peanut and milk allergy no SLIT trials existed 4 years ago. New evidence is added in this field, although OIT showed better results than SLIT in the milk allergy trial.⁴⁵

By the end of 2011 a similar evidence analysis was published for SCIT in children.⁵¹ With respect to pollen SCIT, evidence of a benefit in rhinitis symptoms and medication was scarce; only one high-quality trial showed combined symptom and medication score improvement.⁶⁰ For seasonal asthma there was very low-quality evidence of symptom reduction. However, the specific provocation tests (nasal, ocular, and bronchial) reported clear improvement (G4) with pollen SCIT, as opposed to the nonspecific testing performed with methacholine in the pollen SLIT trials, where no favorable effect could be documented. This probably points to the fact that when provocation testing is performed in a trial, it should be specific. Several positive trials with *Alternaria* SCIT (G1-G4)⁶¹ result in better quality evidence for this treatment in rhinitis than for *Alternaria* SLIT, whereas both have low-quality evidence for efficacy in asthma. Compared with SLIT, the evidence for HDM SCIT efficacy in asthma is superior: high-quality evidence exists for a reduction in asthma symptoms, medication and combined scores, and improved specific bronchial challenge testing. Interestingly, almost all trials in pediatric HDM allergy were in asthma, so evidence of HDM immunotherapy efficacy in AR was better in the SLIT trial review presented in this article.

Three randomized trials compared SLIT with SCIT, all for HDM allergic asthma (and AR).^{40–42} One had a double-blind, double-dummy design. However, all of them were underpowered, with only 10 to 16 patients in each group. Even so the tendency was clear: both treatments showed improvement of asthma (and rhinitis) symptoms and medication scores compared with the control groups, but changes only reached statistical significance with SCIT. An indirect meta-analysis—based comparison of SCIT and SLIT for seasonal AR, although not restricted to the pediatric population and with several other limitations, came to the same conclusion.⁶² Keles et al⁴⁰ revealed that a combination of both routes could give specific benefits.

In our analysis we tried to differentiate between the effect of SLIT given in drops or as tablets. Tablet SLIT has only been studied thoroughly in seasonal AR: here the evidence is slightly better than for drops, as was also commented on in a recent review of US trials.¹⁶

In conclusion although publication bias can never be discarded completely, collectively the presented data show grass pollen SLIT is effective in seasonal allergic rhinitis in children from 5 years of age onward and might be effective in 4-year-old children. Grass or HDM SLIT can be used for allergic rhinitis in children with asthma, and HDM SLIT is probably effective in children with asthma and allergic rhinitis but should never be used as monotherapy in children with active asthmatic symptoms. Immune mechanisms are better understood. Currently, there is not enough evidence to recommend *Alternaria* SLIT in children. Initial results with milk and peanut SLIT show up-dosing should be slow (even so it is not without risks), but finally some patients develop tolerance. No new data on the preventive effect of SLIT in children have been published after initial positive trials of low-moderate quality.⁵⁹ Thus, at this moment the best evidence to recommend SLIT in children with allergic rhinitis for the prevention of asthma development is maintained at the low-moderate level. More large randomized trials are needed, especially with HDM SLIT and mold SLIT in children.

Supplementary Data

Supplementary data related to this article can be found at <http://dx.doi.org/10.1016/j.anai.2013.02.017>

References

- [1] Des Roches A, Paradis L, Menardo JL, Bouges S, Daures JP, Bousquet J. Immunotherapy with a standardized *Dermatophagoides pteronyssinus* extract, VI: specific immunotherapy prevents the onset of new sensitizations in children. *J Allergy Clin Immunol*. 1997;99:450–453.
- [2] Jacobsen L, Niggemann B, Dreborg S, et al. Specific immunotherapy has long-term preventive effect of seasonal and perennial asthma: 10-year follow-up on the PAT study. *Allergy*. 2007;62:943–948.
- [3] Scadding G, Durham SR. Mechanisms of sublingual immunotherapy. *Immunol Allergy Clin North Am*. 2011;31:191–209. viii.
- [4] Novak N, Mete N, Bussmann C, et al. Early suppression of basophil activation during allergen-specific immunotherapy by histamine receptor 2. *J Allergy Clin Immunol*. 2012;130:1153–1158.e2.
- [5] Novak N, Bieber T, Allam JP. Immunological mechanisms of sublingual allergen-specific immunotherapy. *Allergy*. 2011;66:733–739.
- [6] Hankin CS, Cox L, Lang D, et al. Allergen immunotherapy and health care cost benefits for children with allergic rhinitis: a large-scale, retrospective, matched cohort study. *Ann Allergy Asthma Immunol*. 2010;104:79–85.
- [7] Durham SR. Allergen immunotherapy: 100 years on. *Clin Exp Allergy*. 2011;41:1171.
- [8] Larenas Linnemann DE. One hundred years of immunotherapy: review of the first landmark studies. *Allergy Asthma Proc*. 2012;33:122–128.
- [9] Scadding GK, Brostoff J. Low dose sublingual therapy in patients with allergic rhinitis due to house dust mite. *Clin Allergy*. 1986;16:483–491.
- [10] Didier A, Malling HJ, Worm M, et al. Optimal dose, efficacy, and safety of once-daily sublingual immunotherapy with a 5-grass pollen tablet for seasonal allergic rhinitis. *J Allergy Clin Immunol*. 2007;120:1338–1345.
- [11] Durham SR, Yang WH, Pedersen MR, Johansen N, Rak S. Sublingual immunotherapy with once-daily grass allergen tablets: a randomized controlled trial in seasonal allergic rhinoconjunctivitis. *J Allergy Clin Immunol*. 2006;117:802–809.
- [12] Penagos M, Passalacqua G, Compalati E, et al. Metaanalysis of the efficacy of sublingual immunotherapy in the treatment of allergic asthma in pediatric patients, 3 to 18 years of age. *Chest*. 2008;133:599–609.
- [13] Radulovic S, Wilson D, Calderon M, Durham S. Systematic reviews of sublingual immunotherapy (SLIT). *Allergy*. 2011;66:740–752.
- [14] La Rosa M, Lionetti E, Leonardi S, et al. Specific immunotherapy in children: the evidence. *Int J Immunopathol Pharmacol*. 2011;24(4 suppl):69–78.
- [15] Scala G, Di Rienzo Businco A, Ciccirelli A, Tripodi S. An evidence based overview of sublingual immunotherapy in children. *Int J Immunopathol Pharmacol*. 2009;22(4 suppl):23–26.
- [16] Park D, Daher N, Blaiss MS. Adult and pediatric clinical trials of sublingual immunotherapy in the USA. *Expert Rev Clin Immunol*. 2012;8:557–564.
- [17] Guyatt GH, Oxman AD, Vist GE, et al. GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ*. 2008;336:924–926.
- [18] Brozek JL, Akl EA, Alonso-Coello P, et al. Grading quality of evidence and strength of recommendations in clinical practice guidelines, part 1 of 3: an overview of the GRADE approach and grading quality of evidence about interventions. *Allergy*. 2009;64(5):669–677.
- [19] Canonica GW. Sublingual immunotherapy: World Allergy Organization Position Paper 2009. http://www.allergychoices.com/school1000162/genie59/images/files/sub_lingual_immunotherapy_world_allergy.11.pdf. Accessed June 19, 2010.
- [20] Blaiss M, Maloney J, Nolte H, Gawchik S, Yao R, Skoner DP. Efficacy and safety of timothy grass allergy immunotherapy tablets in North American children and adolescents. *J Allergy Clin Immunol*. 2011;127:64–71. e1–e4.
- [21] Nieminen K, Valovirta E, Savolainen J. Clinical outcome and IL-17, IL-23, IL-27 and FOXP3 expression in peripheral blood mononuclear cells of pollen-allergic children during sublingual immunotherapy. *Pediatr Allergy Immunol*. 2010;21(1 pt 2):e174–e184.
- [22] Valovirta E, Jacobsen L, Ljorring C, Koivikko A, Savolainen J. Clinical efficacy and safety of sublingual immunotherapy with tree pollen extract in children. *Allergy*. 2006;61:1177–1183.
- [23] Stelmach I, Kaluzinska-Parzyszek I, Jerzynska J, Stelmach P, Stelmach W, Majak P. Comparative effect of pre-coseasonal and continuous grass sublingual immunotherapy in children. *Allergy*. 2012;67:312–320.
- [24] Wahn U, Tabar A, Kuna P, et al. Efficacy and safety of 5-grass-pollen sublingual immunotherapy tablets in pediatric allergic rhinoconjunctivitis. *J Allergy Clin Immunol*. 2009;123:160–166.e3.
- [25] Bufer A, Eberle P, Franke-Beckmann E, et al. Safety and efficacy in children of an SQ-standardized grass allergen tablet for sublingual immunotherapy. *J Allergy Clin Immunol*. 2009;123:167–173.e7.
- [26] Swamy RS, Reshamwala N, Hunter T, et al. Epigenetic modifications and improved regulatory T-cell function in subjects undergoing dual sublingual immunotherapy. *J Allergy Clin Immunol*. 2012;130:215–224.e7.
- [27] Wahn U, Klimek L, Ploszczek A, et al. High-dose sublingual immunotherapy with single-dose aqueous grass pollen extract in children is effective and safe: a double-blind, placebo-controlled study. *J Allergy Clin Immunol*. 2012;130:886–893.e5.
- [28] Pajno GB, Caminiti L, Crisafulli G, et al. Direct comparison between continuous and coseasonal regimen for sublingual immunotherapy in children with grass allergy: a randomized controlled study. *Pediatr Allergy Immunol*. 2011;22:803–807.
- [29] Halken S, Agertoft L, Seidenberg J, et al. Five-grass pollen 300IR SLIT tablets: efficacy and safety in children and adolescents. *Pediatr Allergy Immunol*. 2010;21:970–976.
- [30] Panzner P, Petras M, Sykora T, Lesna IK, Liska M. Both sublingual and supralingual routes of administration are effective in long-term allergen-specific immunotherapy. *Allergy Asthma Proc*. 2011;32:142–150.
- [31] Panzner P, Petras M, Sykora T, Lesna I. Double-blind, placebo-controlled evaluation of grass pollen specific immunotherapy with oral drops administered sublingually or supralingually. *Respir Med*. 2008;102:1296–1304.
- [32] Agostinis F, Foglia C, Bruno ME, Falagiani P. Efficacy, safety and tolerability of sublingual monomeric allergoid in tablets given without up-dosing to pediatric patients with allergic rhinitis and/or asthma due to grass pollen. *Eur Ann Allergy Clin Immunol*. 2009;41:177–180.
- [33] Ahmadiashar A, Maarefvand M, Taymourzade B, Mazloomzadeh S, Torabi Z. Efficacy of sublingual swallow immunotherapy in children with rye grass pollen allergic rhinitis: a double-blind placebo-controlled study. *Iran J Allergy Asthma Immunol*. 2012;11:175–181.
- [34] Stelmach I, Kaczmarek-Wozniak J, Majak P, Olszowiec-Chlebna M, Jerzynska J. Efficacy and safety of high-doses sublingual immunotherapy in ultra-rush scheme in children allergic to grass pollen. *Clin Exp Allergy*. 2009;39:401–408.
- [35] Yonekura S, Okamoto Y, Sakurai D, et al. Sublingual immunotherapy with house dust extract for house dust-mite allergic rhinitis in children. *Allergol Int*. 2010;59:381–388.
- [36] de Bot CM, Moed H, Berger MY, et al. Sublingual immunotherapy not effective in house dust mite-allergic children in primary care. *Pediatr Allergy Immunol*. 2012;23:150–158.
- [37] Han DH, Choi YS, Lee JE, et al. Clinical efficacy of sublingual immunotherapy in pediatric patients with allergic rhinitis sensitized to house dust mites: comparison to adult patients. *Acta Otolaryngol*. 2012;132(suppl 1):S88–S93.
- [38] Lee JE, Choi YS, Kim MS, et al. Efficacy of sublingual immunotherapy with house dust mite extract in polyallergen sensitized patients with allergic rhinitis. *Ann Allergy Asthma Immunol*. 2011;107:79–84.
- [39] Trebuchon F, David M, Demoly P. Medical management and sublingual immunotherapy practices in patients with house dust mite-induced respiratory allergy: a retrospective, observational study. *Int J Immunopathol Pharmacol*. 2012;25:193–206.
- [40] Keles S, Karakoc-Aydiner E, Ozen A, et al. A novel approach in allergen-specific immunotherapy: combination of sublingual and subcutaneous routes. *J Allergy Clin Immunol*. 2011;128:808–815.e7.
- [41] Eifan AO, Akkoc T, Yildiz A, et al. Clinical efficacy and immunological mechanisms of sublingual and subcutaneous immunotherapy in asthmatic/rhinitis children sensitized to house dust mite: an open randomized controlled trial. *Clin Exp Allergy*. 2010;40:922–932.
- [42] Yukselen A, Kendirli SG, Yilmaz M, Altintas DU, Karakoc GB. Effect of one-year subcutaneous and sublingual immunotherapy on clinical and laboratory parameters in children with rhinitis and asthma: a randomized, placebo-controlled, double-blind, double-dummy study. *Int Arch Allergy Immunol*. 2012;157:288–298.
- [43] Marogna M, Massolo A, Colombo F, Isella P, Bruno M, Falagiani P. Children passive smoking jeopardises the efficacy of standard anti-allergic pharmacological therapy, while sublingual immunotherapy withstands. *Allergol Immunopathol (Madr)*. 2011;39:60–67.
- [44] Kim EH, Bird JA, Kulis M, et al. Sublingual immunotherapy for peanut allergy: clinical and immunologic evidence of desensitization. *J Allergy Clin Immunol*. 2011;127:640–646.e1.
- [45] Keet CA, Frischmeyer-Guerrero PA, Thyagarajan A, et al. The safety and efficacy of sublingual and oral immunotherapy for milk allergy. *J Allergy Clin Immunol*. 2012;129:448–455.
- [46] Acquistapace F, Agostinis F, Castella V, et al. Efficacy of sublingual specific immunotherapy in intermittent and persistent allergic rhinitis in children: an observational case-control study on 171 patients: the EFESO-children Multicenter Trial. *Pediatr Allergy Immunol*. 2009;20:660–664.
- [47] Pozzan M, Milani M. Efficacy of sublingual specific immunotherapy in patients with respiratory allergy to *Alternaria alternata*: a randomised, assessor-blinded, patient-reported outcome, controlled 3-year trial. *Curr Med Res Opin*. 2010;26:2801–2806.
- [48] Seidenberg J, Pajno GB, Bauer CP, La Grutta S, Sieber J. Safety and tolerability of seasonal ultra-rush, high-dose sublingual-swallow immunotherapy in allergic rhinitis to grass and tree pollens: an observational study in 193 children and adolescents. *J Investig Allergol Clin Immunol*. 2009;19:125–131.
- [49] Roger A, Justicia JL, Navarro LA, et al. Observational study of the safety of an ultra-rush sublingual immunotherapy regimen to treat rhinitis due to house dust mites. *Int Arch Allergy Immunol*. 2011;154:69–75.
- [50] Mosges R, Graute V, Christ H, Sieber HJ, Wahn U, Niggemann B. Safety of ultra-rush titration of sublingual immunotherapy in asthmatic children with tree-pollen allergy. *Pediatr Allergy Immunol*. 2010;21:1135–1138.
- [51] Larenas-Linnemann DE, Pietropaolo-Cienfuegos DR, Calderon MA. Evidence of effect of subcutaneous immunotherapy in children: complete and updated review from 2006 onward. *Ann Allergy Asthma Immunol*. 2011;107:407–416.e11.
- [52] Thalheimer W, Cook S. How to calculate effect sizes from published research articles: a simplified methodology. August 2002 http://education.gsu.edu/coshima/EPRS8530/Effect_Sizes_pdf4.pdf. Accessed November 16, 2012.

- [53] Cox L, Larenas-Linnemann D, Lockett RF, Passalacqua G. Speaking the same language: the World Allergy Organization Subcutaneous Immunotherapy Systemic Reaction Grading System. *J Allergy Clin Immunol*. 2010;125: 569–574.
- [54] Zielen S, Kardos P, Madonini E. Steroid-sparing effects with allergen-specific immunotherapy in children with asthma: a randomized controlled trial. *J Allergy Clin Immunol*. 2010;126:942–949.
- [55] Leonardi S, Spicuzza L, La Rosa M. High-dose sublingual immunotherapy in children at 8-year follow-up. *Ann Allergy Asthma Immunol*. 2009;102: 259–260.
- [56] Theodoropoulos DS, Katzenberger DR, Jones WM, et al. Allergen-specific sublingual immunotherapy in the treatment of migraines: a prospective study. *Eur Rev Med Pharmacol Sci*. 2011;15:1117–1121.
- [57] Nguyen TH, Stokes JR, Casale TB. Future forms of immunotherapy and immunomodulators in allergic disease. *Immunol Allergy Clin North Am*. 2011; 31:343–365.
- [58] Grouin JM, Vicaut E, Jean-Alphonse S, et al. The average Adjusted Symptom Score, a new primary efficacy end-point for specific allergen immunotherapy trials. *Clin Exp Allergy*. 2011;41:1282–1288.
- [59] Larenas-Linnemann D. Sublingual immunotherapy in children: complete and updated review supporting evidence of effect. *Curr Opin Allergy Clin Immunol*. 2009;9:168–176.
- [60] Roberts G, Hurley C, Turcanu V, Lack G. Grass pollen immunotherapy as an effective therapy for childhood seasonal allergic asthma. *J Allergy Clin Immunol*. 2006;117:263–268.
- [61] Kuna P, Kaczmarek J, Kupczyk M. Efficacy and safety of immunotherapy for allergies to *Alternaria alternata* in children. *J Allergy Clin Immunol*. 2011;127: 502–508.
- [62] Di Bona D, Plaia A, Leto-Barone MS, La Piana S, Di Lorenzo G. Efficacy of subcutaneous and sublingual immunotherapy with grass allergens for seasonal allergic rhinitis: a meta-analysis-based comparison. *J Allergy Clin Immunol*. 2012;130:1097–1107.

eTable 1
Quality of evidence of pediatric SLIT studies published in 2009–2012, according to the GRADE Approach

Source	Design (starting score)	Large effect	Confound annulated ^a	Dose-response gradient	Total positive	Limitations in design and execution	Inconsistency of results	Indirectness of evidence	Imprecision of results	Publication bias	Total negative	Quality of evidence
Rhinitis Studies												
Blaiss et al, 2011 ¹ : AR (and mild persistent asthma), 5–17 y, pre-seasonal for 1 season, SLIT 175, placebo 169, 15 µg of Phl p 5 daily	DBPC (4)	X	X	X	0	Good: ITT analysis	X	X	X	X	0	4, high
Nieminen et al, 2010 ² : mechanistic study, respiratory allergy, 5–15 y, SLIT low dose 10, high 10, placebo 10, low: 24,000 SQ U/wk, high: 200,000 SQ U/wk, 2 y	DBPC (4)	X	X	+1	+1	X	X	X	Small groups	X	–1	4, high
Wahn et al, 2009 ³ : SAR (21% mild asthma), 131 SLIT, 135 placebo; 4–17 y, pre-season, 25 µg of group 5 grass tablet per day	DBPC (4)	X	X	X	0	X	X	X	X	X	0	4, High
Bufe et al, 2009 ⁴ : SAR (42% mild asthma), 114 SLIT, 120 placebo; 5–16 y, pre-season, 15 µg of Phl p 5 tablet per day	DBPC (4), rhinitis	X	X	X	0	X	X	X	X	X	0	4, High
	DBPC (4), asthma	X	X	X	0	X	X	Only symptom + medication	Very small numbers (9 vs 3 days)	X	–2	2, Low
Swamy et al, 2012 ⁵ : AR (mild/moderate persistent asthma), 6–57 y (55% of SLIT group are children), dual SLIT 20, placebo 10, 15 µg of Phl p 1 and 20 µg Der f 1+2, daily for 12 months. Posttreatment evaluations 12 and 6 and 12 and 12 mo.	DBPC (4)	X	Small groups and even so statistically significant difference	X	+1	Randomization method not described, –1	X	X	Small groups	X	–2	3, moderate
Halken et al, 2010 ^{6,d} : moderate-severe AR (intermittent asthma), SLIT 131, placebo 135, 5–17 y, 25 µg of group 5 daily, pre-season 6 mo	DBPC (4)	+1	X	X	+1	No description of dropouts	X	X	Large CI	X	–2	3, moderate

(continued on next page)

eTable 1 (continued)

Source	Design (starting score)	Large effect	Confound annulated ^d	Dose-response gradient	Total positive	Limitations in design/execution and inconsistency of results	Indirectness of evidence	Imprecision of results	Publication bias	Total negative	Quality of evidence
Wahn et al. 2012 ⁷ ; ARC (and GINA grade I-II asthma), 4–12 y, preseasonal SLIT, placebo, 40 µg of group 5 daily in drops, 8 mo	DBPC (4)	X	X	X	0	No ITT mentioned, X –1, 16.5% dropout in active group, <20%	X	Change in lung sympt not reported	X	–1 (for RC)	3 (for RC)
Yonekura et al. 2010 ⁸ ; AR, 7–15 y, SLIT 21, placebo 10, 0.5 µg of Der f 1 once per week, 40 wk	DBPC (4)	X	X	X	0	Use of nonvalidated adapted symptom score	X	X	X	–1	3, moderate
Stelmach et al. 2012 ⁹ ; rhinitis (20 also asthma), 6–18 y, preseasonal 20, continuous 20, placebo 20, for 2 y, 10 µg of group 5 grass drops daily	DBPC (4)	X	X	X	0	Preestablished adjusted to 1,000 pollen grains/m ³ of symptom and medication scores; no reduction	X	Small groups, underpowered	X	–1	3, moderate
Panzner et al. 2011 ¹⁰ (24-mo open continuation of 12-mo DBPC ¹¹); AR, mean (SD) age 17.6 (10) y. No. of children unknown, 3-y study continuation after 1-y DBPC, SLIT 26, supralingual 25, 11.2 µg of group 5 (6-grass pollen extract) 3 times per week for 3 y	Randomized, no-controlled (4)	X	X	Year by year improvement in symptom and medication, +1	+1	No control group, X small groups; only statistically significant difference pre-post treatment, not between both active groups; no description dropout per group	Post hoc fusion of groups to improve statistics but does not seem to affect outcomes: –0.5	X	Incorrect conclusion: SLIT treatment 2.5; medication score in supralingual not improved	–2	2.5; SLIT: 2.5 supralingual –3.5
De Bot et al. 2012 ¹² ; rhinitis (54% + mild asthma), 6–18 y; 125 SLIT, 126 placebo, 700BU HDM twice per week for 2 y	DBPC (4)	X	X	X	0	Patient selection incorrect: (1) no SPT, (2) no clinical history of symptom exacerbation on exposure, (3) recall bias ^c	X	SD high (1.8 for sympt)	X	–2	2, low
Pozzan et al. 2010 ¹³ ; AR (mild-moderate asthma), mean of 20 (9) y; <i>Alternaria alternaria</i> extract, for 3 y, 34 SLIT, 18 controls; ? µg of Alt a 1 dose daily	Randomized controlled (4)	+1, large effect	X	X	+1	Once yearly evaluation of patients' VAS and medication score: –2	No complete symptom evaluation	X	X	–3	2, low
Agostinis et al. 2009 ¹⁴ ; AR (60% mild asthma), 4–16 y, SLIT 20, control 20, preseasonal for 2 y; 1000 AU drops 5 times week	Randomized controlled (4)	X	Small groups, even so statistically significant	X	+1	No medication score X	VAS well-being only X once a year: –2		X	–3	2, low

Han et al, 2012 ¹⁵ ; AR, 6–18 y (54 patients) and ≥18 y (22 patients), all HDM SLIT; Pangramin, 12 mo	Observational (2)	X	X	X	0	No control group; underpowered to show difference between groups	X	Symptom and medication scores only evaluated 2 times: before and after treatment	X	–2	1, very low
Trebuchon et al, 2012 ¹⁶ ; 735 patients 5–18 y (1289 patients total), 97% AR (54% + asthma), HDM SLIT; several doses, most Der p and Der f 300 IR daily	Retrospective observational (2)	X	X	X	0	Physician selected 10X patients; selection bias	X	X	X	–1	1, very low
Lee et al, 2011 ¹⁷ ; AR (and asthma), monosensitized (70 patients), polysensitized (64 patients), mean age (range) 14.3 (4–53) y, Der p to Der f 5 drops of 1,000 STU/mL every 3 wk for 1 year	Observational (2)	X	X	X	0	X	X	Nonvalidated outcome measure: Q filled in twice on sympt (pre-post treatment) and each 2 mo taking medication	X	–1	1, very low
Almadiafshar et al, 2012 ¹⁸ ; ABC, 5–18 y, SLIT 12, placebo 12, 900 IR Lolium drops, 3 times per wk for 6 mo	Randomized controlled (4)	X	X	X	0	Very small groups. No comparison of symptom-medication reduction SLIT vs placebo	X	Incomplete nasal symptom score (no congestion measured)	No SD or CI given for X	–4	0, very low
Acquistapace et al, 2009 ¹⁹ ; SAR/PAIR, 90 SLIT, 81 control; 7–16 y, low-dose SLIT daily for 2 y various allergens	OCT (2), symptom reduction 50% New asthma	X	X	X	+1	No blinding investigator; no blinding patient	X	No diary data, point X evaluation Asthma = reply yes X to wheezing question SPT of study compared with historical data	X	–2	1, Very low
	New low-dose SLIT daily for 2 y various allergens	X	X	X	0	blinding patient	X	Asthma = reply yes X to wheezing question SPT of study compared with historical data	X	–2	0, very low
	New sensitizations	X	X	X	+1	X	X	X	X	–2	1, very low
Asthma											
Yukselen et al, 2012 ²⁰ (see also above); asthma and rhinitis, mean 10 y, SCIT 10, SLIT 10, placebo 10; SLIT: 1,000 TU of HDM 28 drops every 3 wk for 1 y; SCIT: 3,365 TU each 4 wk, no µg dose stated (Allergopharma)	DBPC, double-dummy (4), SCIT vs SLIT	X	X	Small groups, even so statistically significant	+1	No combined symptom-medication score, no µg dose stated; small groups; underpowered to compare SLIT vs SCIT	X	X	X	–1.5	SLIT vs SCIT: 2.5 = low-mod (Sympt 3.5)
Pajno et al, 2011 ²¹ ; seasonal asthma and AR to grass, 72 children, 8–16 y, 8 µg of group 5 grass 5 times week, for 3 y continuous vs coseasonal	Randomized, no controls (4)	X	X	Small groups, even so statistically significant improved	+1	No control group (IRB: not allowed)	X	X	X	–1.5	SCIT and SLIT vs placebo: 3.5 = Mod-high.
	Randomized, no controls (4)	X	X	X	0	No control group (IRB: not allowed)	X	X	X	–1	3, moderate

(continued on next page)

eTable 1 (continued)

Source	Design (starting score)	Large effect	Confound annulated ^a	Dose-response gradient	Total positive	Limitations in design ^b	Inconsistency of results	Indirectness of evidence	Imprecision of results	Publication bias	Total negative	Quality of evidence
Keles et al, 2011 ²² ; asthma and AR, 5-12 y, (1) SCT, (2) SLIT, (3) build-up SLIT, (4) SCIT then SLIT, (4) pharmacotherapy: 15-15-15-15 patients, respectively; SCT: 13 µg of Der p and Der f once per mo; SLIT: 0.75 µg Der p and Der f 13 times per w for 18 mo	Randomized controlled, asthma outcome (4)	X	X	X	0	Dropout just <15% (13.8)	X	X	X	X	0	4 (very low dose SLIT)
Marogna et al, 2011 ²³ ; AR and intermittent asthma (positive methacholine challenge at inclusion), 68 children, 5-17 y, HDM allergy 34; passive smokers, 34 not. Each group 17-17: SLIT HDM allergoid 1,000 AU once per week or cetirizine for 3 y	Randomized controlled (4)	Methacholine challenge SLIT-no vs SLIT-yes, passive smoke	X	X	0 (+1 methacholines challenge)	Patient selection based on RAST class II+-. Poor statistical evaluation	X	X	Statistical calculations only vs baseline, not between groups	X	-2	2 (3 for methacholine challenge, SLIT-no versus SLIT-yes passive smoke)
Eifan et al, 2010 ²⁴ ; asthma (+AR), 5-10 y, HDM drops (dose stated is confusing), for 12 mo	Randomized controlled (4)	X	SLIT and SCIT vs pharmaceutical group: small groups but even so statistically significant difference	X	+1 (compared against pharmaceutical group)	SCIT vs SLIT: underpowered to show statistically significant difference	X	X	X	X	-1	SLIT vs SCT: 3. SLIT and SCT vs pharmaceutical group: 4
Stelmach et al, 2009 ²⁵ ; asthma mild-moderate persist, 6-17 y, SLIT 20, placebo 15, pre-seasonal for 2 y, 10 µg of group 5 grass drops daily	DBPC (4)	X	X	X	0	40% dropout in placebo group. Symptoms and medications adjusted for pollen count	X	X	No pollen count reported	X	-3	2, low
Safety Roger et al, 2011 ²⁶ ; AR, 4-64 y (122: 4-15 y), HDM drops safety of an ultratrash 90-min build-up 30-60-120-240 IR; 8 mild-moderate systemic AEs	Observational (2)	X	X	X	0		X	X	X	X	0	2, low

Mosges et al, 2010 ²⁷ ; mild-moderate asthma, 6–14 y, 27 SLIT, 27 placebo; tree pollen SLIT ultrarush build-up (in 90 min to 30–90–150–300 IR); no serious adverse events, PFR increase more than in placebo	DBPC (4)	X	X	X	0	X	PFR is supposed to decrease when SLIT is started: it increased, probably learning effect	One of the primary outcome measures: PFR: reflected learning effect instead of lung function	X	X	–2	4 for SAE, 2 for PFR
Seidenberg et al, 2009 ²⁸ ; rhinitis (58% mild-moderate asthma), high-dose daily coseasonal SLIT, build-up in 90 min, 4 mo; 5–17 y, varying allergens	Observational (2)	X	X	X	0	28% did not finish study	X	X	X	X	–1	1, very low
Other Indications												
Keet et al, 2012 ²⁹ ; SLIT up-dosing, then 10 SLIT, 10 low-dose OIT, 6–17 y, 7 mg of milk protein (SLIT), 1,000 mg (OIT-A), 2,000 (OIT-B) daily for 60 wk	Randomized, no controls (4) but DBPC food challenges	X		Small groups, even so stat sign difference	+1	Small groups	X	X	X	X	–1	4, high
Kim et al, 2011 ³⁰ ; 18 children, 1–11 y, 6 mo up-dosing, 6 mo maintenance, 2,000 µg of peanut drops daily, for 12 mo	DBPC (4) and DBPC food challenges	+1		Small groups, even so stat sign difference	+2	No DBPC food challenge at study start: –0.5, interim analysis: 11 in active, 7 placebo	X	X	X	X	–1	4+, high

Abbreviations: AE, adverse event; AR, allergic rhinitis; ARC, allergic rhinoconjunctivitis; CI, confidence interval; DBPC, double-blind, placebo-controlled; GINA, Global Initiative for Asthma; IR, index of reactivity; ITT, intent to treat; HDM, house dust mite; OCT, open controlled trial; OIT, oral immunotherapy; PAR, perennial allergic rhinitis; PFR, pulmonary flow reserve; RCT, randomized controlled trial; RR, relative risk; SAR, seasonal allergic rhinitis; SCIT, subcutaneous immunotherapy; SLIT, sublingual immunotherapy; SPT, skin prick test.

^aAll plausible confounding may be working to reduce the demonstrated effect or increase the effect if no effect was observed. ^bLarge effect RR < 0.5, very large effect RR < 0.2. RR has been calculated from the data given in the articles.

^cInclusion criterium: 3-month retrospective nose symptom score: recall bias.

^dSame study as Wahn et al, 2009, already analyzed in the original World Allergy Organization SLIT paper.³¹

eTable 2

Included and excluded pediatric SLIT studies

Reference	Included/excluded	Reason
Wahn U, Klimek L, Ploszczuk A, et al. High-dose sublingual immunotherapy with single-dose aqueous grass pollen extract in children is effective and safe: a double-blind, placebo-controlled study. <i>J Allergy Clin Immunol</i> . 2012;130:886-893.	Included	
Han DH, Choi YS, Lee JE, et al. Clinical efficacy of sublingual immunotherapy in pediatric patients with allergic rhinitis sensitized to house dust mites: comparison to adult patients. <i>Acta Otolaryngol</i> . 2012;132(suppl 1):S88-S93.	Included	
Ahmadifashar A, Maarefvand M, Taymourzade B, Mazloomzadeh S, Torabi Z. Efficacy of sublingual swallow immunotherapy in children with rye grass pollen allergic rhinitis: a double-blind placebo-controlled study. <i>Iran J Allergy Asthma Immunol</i> . 2012;11:175-181.	Included	
Swamy RS, Reshamwala N, Hunter T, et al. Epigenetic modifications and improved regulatory T-cell function in subjects undergoing dual sublingual immunotherapy. <i>J Allergy Clin Immunol</i> . 2012;130:215-224.	Included	
Stelmach I, Kaluzińska-Parzyszek I, Jerzynska J, Stelmach P, Stelmach W, Majak P. Comparative effect of pre-coseasonal and continuous grass sublingual immunotherapy in children. <i>Allergy</i> . 2012;67:312-320.	Included	
de Bot CM, Moed H, Berger MY, et al. Sublingual immunotherapy not effective in house dust mite-allergic children in primary care. <i>Pediatr Allergy Immunol</i> . 2012;23:150-158.	Included	
Keet CA, Frischmeyer-Guerrero PA, Thyagarajan A, et al. The safety and efficacy of sublingual and oral immunotherapy for milk allergy. <i>J Allergy Clin Immunol</i> . 2012;129:448-455.	Included	
Yukselen A, Kendirli SG, Yilmaz M, Altintas DU, Karakoc GB. Effect of one-year subcutaneous and sublingual immunotherapy on clinical and laboratory parameters in children with rhinitis and asthma: a randomized, placebo-controlled, double-blind, double-dummy study. <i>Int Arch Allergy Immunol</i> . 2012;157:288-298.	Included	
Trebuchon F, David M, Demoly P. Medical management and sublingual immunotherapy practices in patients with house dust mite-induced respiratory allergy: a retrospective, observational study. <i>Int J Immunopathol Pharmacol</i> . 2012;25:193-206.	Included	
Pajno GB, Caminiti L, Crisafulli G, et al. Direct comparison between continuous and coseasonal regimen for sublingual immunotherapy in children with grass allergy: a randomized controlled study. <i>Pediatr Allergy Immunol</i> . 2011;22:803-807.	Included	
Keles S, Karakoc-Aydiner E, Ozen A, et al. A novel approach in allergen-specific immunotherapy: combination of sublingual and subcutaneous routes. <i>J Allergy Clin Immunol</i> . 2011;128:808-815.	Included	
Lee JE, Choi YS, Kim MS, et al. Efficacy of sublingual immunotherapy with house dust mite extract in polyallergen sensitized patients with allergic rhinitis. <i>Ann Allergy Asthma Immunol</i> . 2011;107:79-84.	Included	
Marogna M, Massolo A, Colombo F, Isella P, Bruno M, Falagiani P. Children passive smoking jeopardises the efficacy of standard anti-allergic pharmacological therapy, while sublingual immunotherapy withstands. <i>Allergol Immunopathol (Madr)</i> . 2011;39:60-67.	Included	
Panzner P, Petrás M, Sýkora T, Lesná IK, Liška M. Both sublingual and supralingual routes of administration are effective in long-term allergen-specific immunotherapy. <i>Allergy Asthma Proc</i> . 2011;32:142-150.	Included	
Kim EH, Bird JA, Kulis M, et al. Sublingual immunotherapy for peanut allergy: clinical and immunologic evidence of desensitization. <i>J Allergy Clin Immunol</i> . 2011;127:640-646.	Included	
Roger A, Justicia JL, Navarro LA, et al. Observational study of the safety of an ultra-rush sublingual immunotherapy regimen to treat rhinitis due to house dust mites. <i>Int Arch Allergy Immunol</i> . 2011;154:69-75.	Included	
Blaiss M, Maloney J, Nolte H, Gawchik S, Yao R, Skoner DP. Efficacy and safety of timothy grass allergy immunotherapy tablets in North American children and adolescents. <i>J Allergy Clin Immunol</i> . 2011;127:64-71.	Included	
Yonekura S, Okamoto Y, Sakurai D, et al. Sublingual immunotherapy with house dust extract for house dust-mite allergic rhinitis in children. <i>Allergol Int</i> . 2010;59:381-388.	Included	
Pozzan M, Milani M. Efficacy of sublingual specific immunotherapy in patients with respiratory allergy to <i>Alternaria alternata</i> : a randomised, assessor-blinded, patient-reported outcome, controlled 3-year trial. <i>Curr Med Res Opin</i> . 2010;26:2801-2806.	Included	
Mösger R, Graute V, Christ H, Sieber HJ, Wahn U, Niggemann B. Safety of ultra-rush titration of sublingual immunotherapy in asthmatic children with tree-pollen allergy. <i>Pediatr Allergy Immunol</i> . 2010;21:1135-1138.	Included	
Halken S, Agertoft L, Seidenberg J, et al. Five-grass pollen 300IR SLIT tablets: efficacy and safety in children and adolescents. <i>Pediatr Allergy Immunol</i> . 2010;21:970-976.	Included	
Eifan AO, Akkoc T, Yildiz A, et al. Clinical efficacy and immunological mechanisms of sublingual and subcutaneous immunotherapy in asthmatic/rhinitis children sensitized to house dust mite: an open randomized controlled trial. <i>Clin Exp Allergy</i> . 2010;40:922-932.	Included	
Nieminen K, Valovirta E, Savolainen J. Clinical outcome and IL-17, IL-23, IL-27 and FOXP3 expression in peripheral blood mononuclear cells of pollen-allergic children during sublingual immunotherapy. <i>Pediatr Allergy Immunol</i> . 2010;21(1 pt 2):e174-e184.	Included	
Agostinis F, Foglia C, Bruno ME, Falagiani P. Efficacy, safety and tolerability of sublingual monomeric allergoid in tablets given without up-dosing to pediatric patients with allergic rhinitis and/or asthma due to grass pollen. <i>Eur Ann Allergy Clin Immunol</i> . 2009;41:177-180.	Included	
Acquistapace F, Agostinis F, Castella V, et al. Efficacy of sublingual specific immunotherapy in intermittent and persistent allergic rhinitis in children: an observational case-control study on 171 patients: the EFESO-children multicenter trial. <i>Pediatr Allergy Immunol</i> . 2009;20:660-664.	Included	
Stelmach I, Kaczmarek-Woźniak J, Majak P, Olszowiec-Chlebna M, Jerzynska J. Efficacy and safety of high-doses sublingual immunotherapy in ultra-rush scheme in children allergic to grass pollen. <i>Clin Exp Allergy</i> . 2009;39:401-408.	Included	
Bufe A, Eberle P, Franke-Beckmann E, et al. Safety and efficacy in children of an SQ-standardized grass allergen tablet for sublingual immunotherapy. <i>J Allergy Clin Immunol</i> . 2009;123:167-173.	Included	
Wahn U, Tabar A, Kuna P, Halken S, Montagut A, de Beaumont O, Le Gall M; SLIT Study Group. Efficacy and safety of 5-grass-pollen sublingual immunotherapy tablets in pediatric allergic rhinoconjunctivitis. <i>J Allergy Clin Immunol</i> . 2009;123:160-166.	Included	
Seidenberg J, Pajno GB, Bauer CP, La Grutta S, Sieber J. Safety and tolerability of seasonal ultra-rush, high-dose sublingual-swallow immunotherapy in allergic rhinitis to grass and tree pollens: an observational study in 193 children and adolescents. <i>J Invest Allergol Clin Immunol</i> . 2009;19:125-131.	Included	
Wessel F, Chartier A, Meunier JP, Magnan A. Safety and tolerability of an SQ-standardized GRass Allergy immunotherapy tablet (GRAZAX®) in a real-life setting for three consecutive seasons - the GRAAL trial. <i>Clin Drug Invest</i> . 2012;32:451-463.	Excluded	Adult
Durham SR, Emminger W, Kapp A, et al. SQ-standardized sublingual grass immunotherapy: confirmation of disease modification 2 years after 3 years of treatment in a randomized trial. <i>J Allergy Clin Immunol</i> . 2012;129:717-725.	Excluded	Adult
Sieber J, Neis M, Brehler R, et al. Increasing long-term safety of seasonal grass pollen sublingual immunotherapy: the ECRI study. <i>Expert Opin Drug Saf</i> . 2012;11:7-13.	Excluded	Adult
Didier A, Worm M, Horak F, et al. Sustained 3-year efficacy of pre- and coseasonal 5-grass-pollen sublingual immunotherapy tablets in patients with grass pollen-induced rhinoconjunctivitis. <i>J Allergy Clin Immunol</i> . 2011;128:559-566.	Excluded	Adult

eTable 2 (continued)

Reference	Included/excluded	Reason
Milani M, Pecora S; Rainbow Study Investigator Group. Clinical relevance of non-grass pollens respiratory allergies in Italy and effects of specific sublingual immunotherapy: The Rainbow Trial, a multicentre 3-year prospective observational study. <i>Eur Ann Allergy Clin Immunol</i> . 2011;43:111–116.	Excluded	Adult
Migueres M, Fontaine JF, Haddad T, et al. Characteristics of patients with respiratory allergy in France and factors influencing immunotherapy prescription: a prospective observational study (REALIS). <i>Int J Immunopathol Pharmacol</i> . 2011;24:387–400.	Excluded	Adult
Mauro M, Russello M, Incorvaia C, et al. Birch-apple syndrome treated with birch pollen immunotherapy. <i>Int Arch Allergy Immunol</i> . 2011;156:416–422.	Excluded	Adult
Nelson HS, Nolte H, Creticos P, Maloney J, Wu J, Bernstein DI. Efficacy and safety of timothy grass allergy immunotherapy tablet treatment in North American adults. <i>J Allergy Clin Immunol</i> . 2011;127:72–80.	Excluded	Adult
Cortellini G, Spadolini I, Patella V, et al. Sublingual immunotherapy for Alternaria-induced allergic rhinitis: a randomized placebo-controlled trial. <i>Ann Allergy Asthma Immunol</i> . 2010;105:382–386.	Excluded	Adult
Marogna M, Spadolini I, Massolo A, Canonica GW, Passalacqua G. Long-lasting effects of sublingual immunotherapy according to its duration: a 15-year prospective study. <i>J Allergy Clin Immunol</i> . 2010;126:969–975.	Excluded	Adult
Leonardi S, Arena A, Bruno ME, et al. Olea sublingual allergoid immunotherapy administered with two different treatment regimens. <i>Allergy Asthma Proc</i> . 2010;31:e25–e29.	Excluded	Adult
García BE, González-Mancebo E, Barber D, et al. Sublingual immunotherapy in peach allergy: monitoring molecular sensitizations and reactivity to apple fruit and Platanus pollen. <i>J Invest Allergol Clin Immunol</i> . 2010;20:514–520.	Excluded	Adult
Marogna M, Colombo F, Spadolini I, et al. Randomized open comparison of montelukast and sublingual immunotherapy as add-on treatment in moderate persistent asthma due to birch pollen. <i>J Invest Allergol Clin Immunol</i> . 2010;20:146–152.	Excluded	Adult
O'Hehir RE, Gardner LM, de Leon MP, et al. House dust mite sublingual immunotherapy: the role for transforming growth factor-beta and functional regulatory T cells. <i>Am J Respir Crit Care Med</i> . 2009;180:936–947.	Excluded	Adult
Ott H, Sieber J, Brehler R, et al. Efficacy of grass pollen sublingual immunotherapy for three consecutive seasons and after cessation of treatment: the ECRIT study. <i>Allergy</i> . 2009;64:1394–1401.	Excluded	Adult
Horak F, Ziegelmayer P, Ziegelmayer R, et al. Early onset of action of a 5-grass-pollen 300-IR sublingual immunotherapy tablet evaluated in an allergen challenge chamber. <i>J Allergy Clin Immunol</i> . 2009;124:471–477.	Excluded	Adult
Amar SM, Harbeck RJ, Sills M, Silveira LJ, O'Brien H, Nelson HS. Response to sublingual immunotherapy with grass pollen extract: monotherapy versus combination in a multiallergen extract. <i>J Allergy Clin Immunol</i> . 2009;124:150–156.	Excluded	Adult
Ventura MT, Carretta A, Tummolo RA, Buquicchio R, Arsieni A, Murgia N. Clinical data and inflammation parameters in patients with cypress allergy treated with sublingual swallow therapy and subcutaneous immunotherapy. <i>Int J Immunopathol Pharmacol</i> . 2009;22:403–413.	Excluded	Adult
Malling HJ, Montagut A, Melac M, et al. Efficacy and safety of 5-grass pollen sublingual immunotherapy tablets in patients with different clinical profiles of allergic rhinoconjunctivitis. <i>Clin Exp Allergy</i> . 2009;39:387–393.	Excluded	Adult
Horak F, Jaeger S, Worm M, Melac M, Didier A. Implementation of pre-seasonal sublingual immunotherapy with a five-grass pollen tablet during optimal dosage assessment. <i>Clin Exp Allergy</i> . 2009;39:394–400.	Excluded	Adult
Marogna M, Spadolini I, Massolo A, et al. Long-term comparison of sublingual immunotherapy vs inhaled budesonide in patients with mild persistent asthma due to grass pollen. <i>Ann Allergy Asthma Immunol</i> . 2009;102:69–75.	Excluded	Adult
Zielen S, Kardos P, Madonini E. Steroid-sparing effects with allergen-specific immunotherapy in children with asthma: a randomized controlled trial. <i>J Allergy Clin Immunol</i> . 2010;126:942–929.	Excluded	SCIT
Leonardi S, Spicuzza L, La Rosa M. High-dose sublingual immunotherapy in children at 8-year follow-up. <i>Ann Allergy Asthma Immunol</i> . 2009;102:259–260.	Excluded	Letter to the editor
Theodoropoulos DS, Katzenberger DR, Jones WM, Morris DL, Her C, Cullen NA, Morris DL. Allergen-specific sublingual immunotherapy in the treatment of migraines: a prospective study. <i>Eur Rev Med Pharmacol Sci</i> . 2011;15:1117–1121.	Excluded	Migraine
Nguyen SA, Schlosser RJ. Assessment of palatability of two sublingual diluents in allergic patients: a prospective pilot study. <i>Am J Rhinol Allergy</i> . 2011;25:342–345.	Excluded	Palatability
Grouin JM, Vicaut E, Jean-Alphonse S, et al. The average Adjusted Symptom Score, a new primary efficacy end-point for specific allergen immunotherapy trials. <i>Clin Exp Allergy</i> . 2011;41:1282–1288.	Excluded	Recalculation

Abbreviations: SCIT, subcutaneous immunotherapy; SLIT, sublingual immunotherapy.

eTable 3

Adverse events for SLIT, 2009–2012

Source	No. receiving SLIT	Treatment discontinuation due to AE (active/placebo)	Life-threatening systemic AE	Treatment-related systemic AE	Most common local AE
Blaiss et al, 2011 ^a	175	13 (7.4%)/5 (3%)	None	No SAE, urticaria 3/175 (1.7%)/0%	Oral pruritus (39% vs 3.4%), throat irritation (37.1% vs 3%), stomatitis (15% vs 1.2%)
Stelmach et al, 2012 ^a	20	0/0/0	None	No SAE, headache, stomachache	Sublingual pruritus (45% vs 15.3%)
Wahn et al, 2009 ^a	139	7 (5%)/2 (1.4%)	None	No SAE. 12.2% SAE in active	Oral pruritus (32.4% vs 1.4%), mouth edema (13% vs 0%), throat irritation (8% vs 5%)
Bufe et al, 2009 ^a	126	4 (3%)/2 (2%)	None	SAE: 2 SLIT, 2 placebo, cough	Oral pruritus (32% vs 2%), throat irritation (10% vs 2%), swollen lip (7% vs 0%)
Swamy et al, 2012 ^a	20	0/0	None	GI concerns 15%, urticarial 10%	Itchy mouth/throat (85% vs 20%), rhinitis/ sneezing (30% vs 20%)
Wahn et al, 2012 ^a	158	10 (6.3%)/0	None	No SAE, abdominal concerns equal in active-placebo	Oral administration concerns (71% vs 12.2%)
Pajno et al, 2011 ^a	40/40	5 (6.3%) = coseasonal 4, continuous 1	None	GI symptoms	mouth burning
Panzner et al, 2011 ^a	26	?	None	No SAE, 35% systemic adverse events: rhinitis, painful breathing, conjunctivitis (treatment related?)	Undesirable taste, difficult swallowing, local swelling, or burning
Agostinis et al, 2009	20	0	None	No SAE	?
Ahmadi-far et al, 2012	12	0	None	No SAE	Higher AE score in placebo group (no statistical analysis)
Stelmach et al, 2009 ^a	20	0/0/0	None	No SAE, headache, stomachache	Sublingual pruritus (50% vs 14.3%), less second year (35% vs 20%)
Yonekura et al, 2010	20	0	None	No SAE	Bitter taste.
De Bot et al, 2012	126	0	None	No SAE, rhinitis, conjunctivitis, shortness of breath (similar in active-placebo)	Oral pharyngeal irritation/ swelling
Han et al, 2012	76	No safety data			
Lee et al, 2011	134	No safety data			
Trebuchon et al, 2012	735	8%	None	Systemic AE 4%.	Local, mild
Keles et al, 2011	15	0 ^b	None	No SAE	None reported
Eifan et al, 2010	16	0 ^c	None	No SAE	None reported
Yukselen et al, 2012	11	0	None	No SAE	Local, mild
Marogna et al, 2011	34	No safety data reported			
Kim et al, 2011 ^a	11		None	No SAE, SLIT: after 11 doses (0.26%) antihistamine was needed, after 1 dose (0.02%) β_2 -agonist was needed	Oropharyngeal reactions (9.3% vs 1.5%)
Keet et al, 2012	10	SLIT-SLIT: 0, SLIT-OIT low: 1, SLIT-OIT high: 1	Epinephrine given: with SLIT: 1, with OIT: 4	Systemic AE more frequent in OIT vs SLIT ($P = .01$ – $P < .001$) and more need for β -agonist and antihistamine treatment	Local AE with SLIT and oral immunotherapy similar (29% vs 23%)

Abbreviations: AE, adverse event; GI, gastrointestinal; OIT, oral immunotherapy; SAE, serious adverse event; SLIT, sublingual immunotherapy.

^aStudies reporting clearly treatment-related and non-treatment-related events.^bKeles et al, 2011: 2 of 13 children in SCIT group discontinued because of AEs.^cEifan et al: 2 SCIT patients discontinued because of SAEs.

eReferences

- [1] Blaiss M, Maloney J, Nolte H, Gawchik S, Yao R, Skoner DP. Efficacy and safety of timothy grass allergy immunotherapy tablets in North American children and adolescents. *J Allergy Clin Immunol*. 2011;127:64–71. e1–4.
- [2] Nieminen K, Valovirta E, Savolainen J. Clinical outcome and IL-17, IL-23, IL-27 and FOXP3 expression in peripheral blood mononuclear cells of pollen-allergic children during sublingual immunotherapy. *Pediatr Allergy Immunol*. 2010;21(1 pt 2):e174–e184.
- [3] Wahn U, Tabar A, Kuna P, et al. Efficacy and safety of 5-grass-pollen sublingual immunotherapy tablets in pediatric allergic rhinoconjunctivitis. *J Allergy Clin Immunol*. 2009;123:160–166.
- [4] Bufer A, Eberle P, Franke-Beckmann E, et al. Safety and efficacy in children of an SQ-standardized grass allergen tablet for sublingual immunotherapy. *J Allergy Clin Immunol*. 2009;123:167–173.
- [5] Swamy RS, Reshamwala N, Hunter T, et al. Epigenetic modifications and improved regulatory T-cell function in subjects undergoing dual sublingual immunotherapy. *J Allergy Clin Immunol*. 2012;130:215–224.
- [6] Halken S, Agertoft L, Seidenberg J, et al. Five-grass pollen 300IR SLIT tablets: efficacy and safety in children and adolescents. *Pediatr Allergy Immunol*. 2010;21:970–976.
- [7] Wahn U, Klimek L, Ploszczuk A, et al. High-dose sublingual immunotherapy with single-dose aqueous grass pollen extract in children is effective and safe: a double-blind, placebo-controlled study. *J Allergy Clin Immunol*. 2012;130:886–893.
- [8] Yonekura S, Okamoto Y, Sakurai D, et al. Sublingual immunotherapy with house dust extract for house dust-mite allergic rhinitis in children. *Allergol Int*. 2010;59:381–388.
- [9] Stelmach I, Kaluzinska-Parzyszek I, Jerzynska J, Stelmach P, Stelmach W, Majak P. Comparative effect of pre-coseasonal and continuous grass sublingual immunotherapy in children. *Allergy*. 2012;67:312–320.
- [10] Panzner P, Petras M, Sykora T, Lesna IK, Liska M. Both sublingual and supralingual routes of administration are effective in long-term allergen-specific immunotherapy. *Allergy Asthma Proc*. 2011;32:142–150.
- [11] Panzner P, Petras M, Sykora T, Lesna I. Double-blind, placebo-controlled evaluation of grass pollen specific immunotherapy with oral drops administered sublingually or supralingually. *Respir Med*. 2008;102:1296–1304.
- [12] de Bot CM, Moed H, Berger MY, et al. Sublingual immunotherapy not effective in house dust mite-allergic children in primary care. *Pediatr Allergy Immunol*. 2012;23:150–158.
- [13] Pozzan M, Milani M. Efficacy of sublingual specific immunotherapy in patients with respiratory allergy to *Alternaria alternata*: a randomised, assessor-blinded, patient-reported outcome, controlled 3-year trial. *Curr Med Res Opin*. 2010;26:2801–2806.
- [14] Agostinis F, Foglia C, Bruno ME, Falagiani P. Efficacy, safety and tolerability of sublingual monomeric allergoid in tablets given without up-dosing to pediatric patients with allergic rhinitis and/or asthma due to grass pollen. *Eur Ann Allergy Clin Immunol*. 2009;41:177–180.
- [15] Han DH, Choi YS, Lee JE, et al. Clinical efficacy of sublingual immunotherapy in pediatric patients with allergic rhinitis sensitized to house dust mites: comparison to adult patients. *Acta Otolaryngol*. 2012;132(suppl 1):S88–S93.
- [16] Trebuchon F, David M, Demoly P. Medical management and sublingual immunotherapy practices in patients with house dust mite-induced respiratory allergy: a retrospective, observational study. *Int J Immunopathol Pharmacol*. 2012;25:193–206.
- [17] Lee JE, Choi YS, Kim MS, et al. Efficacy of sublingual immunotherapy with house dust mite extract in polyallergen sensitized patients with allergic rhinitis. *Ann Allergy Asthma Immunol*. 2011;107:79–84.
- [18] Ahmadiashar A, Maarefvand M, Taymourzade B, Mazloomzadeh S, Torabi Z. Efficacy of sublingual swallow immunotherapy in children with rye grass pollen allergic rhinitis: a double-blind placebo-controlled study. *Iran J Allergy Asthma Immunol*. 2012;11:175–181.
- [19] Acquistapace F, Agostinis F, Castella V, et al. Efficacy of sublingual specific immunotherapy in intermittent and persistent allergic rhinitis in children: an observational case-control study on 171 patients: The EFESO-children Multicenter Trial. *Pediatr Allergy Immunol*. 2009;20:660–664.
- [20] Yukselen A, Kendirli SG, Yilmaz M, Altintas DU, Karakoc GB. Effect of one-year subcutaneous and sublingual immunotherapy on clinical and laboratory parameters in children with rhinitis and asthma: a randomized, placebo-controlled, double-blind, double-dummy study. *Int Arch Allergy Immunol*. 2012;157:288–298.
- [21] Pajno GB, Caminiti L, Crisafulli G, et al. Direct comparison between continuous and coseasonal regimen for sublingual immunotherapy in children with grass allergy: a randomized controlled study. *Pediatr Allergy Immunol*. 2011;22:803–807.
- [22] Keles S, Karakoc-Aydiner E, Ozen A, et al. A novel approach in allergen-specific immunotherapy: combination of sublingual and subcutaneous routes. *J Allergy Clin Immunol*. 2011;128:808–815.
- [23] Marogna M, Massolo A, Colombo F, Isella P, Bruno M, Falagiani P. Children passive smoking jeopardises the efficacy of standard anti-allergic pharmacological therapy, while sublingual immunotherapy withstands. *Allergol Immunopathol (Madr)*. 2011;39:60–67.
- [24] Eifan AO, Akkoc T, Yildiz A, et al. Clinical efficacy and immunological mechanisms of sublingual and subcutaneous immunotherapy in asthmatic/rhinitis children sensitized to house dust mite: an open randomized controlled trial. *Clin Exp Allergy*. 2010;40:922–932.
- [25] Stelmach I, Kaczmarek-Wozniak J, Majak P, Olszowiec-Chlebna M, Jerzynska J. Efficacy and safety of high-doses sublingual immunotherapy in ultra-rush scheme in children allergic to grass pollen. *Clin Exp Allergy*. 2009;39:401–408.
- [26] Roger A, Justicia JL, Navarro LA, et al. Observational study of the safety of an ultra-rush sublingual immunotherapy regimen to treat rhinitis due to house dust mites. *Int Arch Allergy Immunol*. 2011;154:69–75.
- [27] Mosges R, Graute V, Christ H, Sieber HJ, Wahn U, Niggemann B. Safety of ultra-rush titration of sublingual immunotherapy in asthmatic children with tree-pollen allergy. *Pediatr Allergy Immunol*. 2010;21:1135–1138.
- [28] Seidenberg J, Pajno GB, Bauer CP, La Grutta S, Sieber J. Safety and tolerability of seasonal ultra-rush, high-dose sublingual-swallow immunotherapy in allergic rhinitis to grass and tree pollens: an observational study in 193 children and adolescents. *J Investig Allergol Clin Immunol*. 2009;19:125–131.
- [29] Keet CA, Frischmeyer-Guerrero PA, Thyagarajan A, et al. The safety and efficacy of sublingual and oral immunotherapy for milk allergy. *J Allergy Clin Immunol*. 2012;129:448–455.
- [30] Kim EH, Bird JA, Kulis M, et al. Sublingual immunotherapy for peanut allergy: clinical and immunologic evidence of desensitization. *J Allergy Clin Immunol*. 2011;127:640–646.
- [31] Canonica GW. Sub-lingual immunotherapy: World Allergy Organization position paper 2009. http://www.allergychoices.com/school1000162/genie59/images/files/sub_lingual_immunotherapy_world_allergy.11.pdf. Accessed June 19, 2010.