


Journal Pre-proof

Misdiagnosis of systemic allergic reactions to mRNA COVID-19 vaccines

John M Kelso MD

PII: S1081-1206(21)00257-X
DOI: <https://doi.org/10.1016/j.anai.2021.03.024>
Reference: ANAI 3546



To appear in: *Annals of Allergy, Asthma Immunology*

Received date: 19 February 2021
Revised date: 18 March 2021
Accepted date: 23 March 2021

Please cite this article as: John M Kelso MD , Misdiagnosis of systemic allergic reactions to mRNA COVID-19 vaccines, *Annals of Allergy, Asthma Immunology* (2021), doi: <https://doi.org/10.1016/j.anai.2021.03.024>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2021 American College of Allergy, Asthma & Immunology. Published by Elsevier Inc. All rights reserved.

Misdiagnosis of systemic allergic reactions to mRNA COVID-19 vaccines

John M Kelso, MD
Division of Allergy, Asthma and Immunology
Scripps Clinic
San Diego California

Corresponding Author:

John M Kelso, MD
Division of Allergy, Asthma and Immunology
Scripps Clinic
3811 Valley Centre Drive
San Diego CA 92130

857-764-9010

kelso.john@scrippshealth.org

Funding Source: none.

Keywords: Vaccine allergy; mRNA COVID-19 vaccines

Abbreviations/Acronyms:

COVID-19: coronavirus disease 2019
mRNA: messenger ribonucleic acid
PEG: polyethylene glycol

Conflicts of interest: none.

Two mRNA COVID-19 vaccines were granted emergency use authorization by the United States Food and Drug Administration in December 2020.^{1,2} Although no anaphylactic reactions were reported in their research trials, during the first few weeks of clinical use, such reactions have been reported at 2.5 to 11.1 per million first doses administered.^{3,4} It was surprising that

anaphylaxis could occur because the patients had not previously been exposed and because the vaccines contain no protein. The lipid nanoparticles that surround the mRNA do contain polyethylene glycol (PEG),^{1,2} to which patients may have been previously exposed and which has rarely been implicated in IgE-mediated allergic reactions,^{5,6} although the amount of PEG in the vaccines is very small and there is no evidence to date that PEG is responsible for the vaccine reactions. An adenovirus vector COVID-19 vaccine recently granted emergency use authorization⁷ does not contain PEG, but does contain the structurally related polysorbate-80.

We have had the opportunity to evaluate four patients who suffered early onset reactions after their first doses of mRNA COVID vaccines, and who were diagnosed and treated for presumed systemic allergic reactions. However, further evaluation suggested that the reactions were not allergic (Table 1). Although notes were not generated regarding physical examination at the vaccine clinics, all patients had only subjective complaints with normal vital signs and physical exams when subsequently evaluated in the emergency department or urgent care center, including one patient subsequently admitted to the intensive care unit. Subsequent vaccine skin testing suggested that the reactions were not IgE-mediated, and three of the four have received their second doses either without symptoms or with only mild transient symptoms. Despite negative skin testing results, one patient declined her second dose. The skin test results and subsequent vaccination outcomes in these patients indicate that their initial reactions were not allergic. The patients had symptoms that were triggered by the vaccination, but not by an allergic mechanism.

Other early onset reactions to vaccination can mimic anaphylaxis. Vasovagal reactions can also cause lightheadedness or syncope. Vocal cord spasm can cause stridor and dyspnea. Panic attacks can cause a globus sensation, palpitations, dyspnea and other symptoms. It is likely that

our patients' initial reactions represent some variation of such reactions. Patients and providers are sometimes reluctant to believe that physical reactions and sensations can be triggered by something other than an allergic reaction. However, the somatic element of a psychosomatic response is in fact a real physical phenomenon, and conversely, symptoms can be present without physical findings. It has proved helpful to review with these patients other circumstances where this may be the case. For example, embarrassment can cause obvious flushing (vasodilatation) triggered only by a thought. The sensation of throat or tongue swelling when none is present (globus sensation) is also common. Most patients are familiar with the experience that after local anesthesia for a dental procedure, they often have the distinct sensation that their lip or tongue is swollen, but looking the mirror reveals that it is not. Patients being able to see negative skin test results in contrast to positive histamine control tests can also provide reassurance that they are not allergic to the vaccine.

Although it is important for providers overseeing vaccination clinics to recognize and treat anaphylaxis after vaccine administration, they should also be aware that there is a differential diagnosis. Epinephrine should certainly not be delayed or withheld if anaphylaxis is suspected, but in the setting of minor and more subjective feelings, observation may be appropriate. Patients with possible allergic reactions after immunization should be evaluated by an allergist rather than simply being labeled "allergic" or recommending that they not receive additional doses.⁸

In patients who may have had a reaction to the first dose of an mRNA COVID vaccine, an alternative to second dose administration could be to assess for the presence of anti-SARS-CoV-2 spike protein IgG antibody as evidence of immune response, but no data exists to demonstrate that a single dose provides the same level or duration of protection from disease as the two dose

series. Another approach might be to suggest that the patient forego the second dose of an mRNA vaccine and receive an alternative vaccine with different technology or ingredients, such as the adenovirus vector vaccine.⁷ However, there are no data on the immune response or protection that this might provide.

Thus, the more appropriate approach would seem to be for patients who have had possible immediate-type allergic reactions to an mRNA COVID vaccine to undergo an allergy evaluation to determine if the nature and timing of the patient's symptoms are consistent with an anaphylactic reaction, and to perform vaccine skin testing. The stability of any potential allergens in the COVID vaccines is not known and so skin testing should be performed within the same six-hour timeframe from reconstitution used for vaccine administration, using residual volume from the multi-dose vials so as not to waste doses. Negative vaccine skin test results argue strongly that a reaction was not IgE-mediated, and consideration can be given to administration of the second dose in the usual manner under observation as in the cases described.

Other reported anaphylactic reactions attributed to the mRNA COVID vaccines^{3,4} may not have been systemic allergic reactions, but rather vasovagal or panic reactions. Whatever risk may be posed by receiving a second dose of COVID vaccine when the first dose may have triggered a reaction must be weighed against the risk of remaining inadequately vaccinated against a potentially fatal disease. The former risk in most cases is small and manageable, while the latter is substantial. For most patients, weighing the relative risks of these options will favor careful evaluation and subsequent vaccination.

References

1. EMERGENCY USE AUTHORIZATION (EUA) OF THE PFIZER-BIONTECH COVID-19 VACCINE TO PREVENT CORONAVIRUS DISEASE 2019 (COVID-19) Pfizer Inc., New York, NY. 2020.
2. EMERGENCY USE AUTHORIZATION (EUA) OF THE MODERNA COVID-19 VACCINE TO PREVENT CORONAVIRUS DISEASE 2019 (COVID-19) Moderna. 2020.
3. Allergic Reactions Including Anaphylaxis After Receipt of the First Dose of Pfizer-BioNTech COVID-19 Vaccine - United States, December 14-23, 2020. MMWR - Morbidity & Mortality Weekly Report 2021; 70:46-51.
4. Allergic Reactions Including Anaphylaxis After Receipt of the First Dose of Moderna COVID-19 Vaccine - United States, December 21, 2020-January 10, 2021. MMWR - Morbidity & Mortality Weekly Report 2021; 70:125-9.
5. Wenande E, Garvey LH. Immediate-type hypersensitivity to polyethylene glycols: a review. *Clinical & Experimental Allergy* 2016; 46:907-22.
6. Stone CA, Jr., Liu Y, Relling MV, Krantz MS, Pratt AL, Abreo A, et al. Immediate Hypersensitivity to Polyethylene Glycols and Polysorbates: More Common Than We Have Recognized. *The Journal of Allergy and Clinical Immunology: In Practice* 2019; 7:1533-40.e8.
7. EMERGENCY USE AUTHORIZATION (EUA) OF THE JANSSEN COVID-19 VACCINE TO PREVENT CORONAVIRUS DISEASE 2019 (COVID-19).

8. Kelso JM, Greenhawt MJ, Li JT, Nicklas RA, Bernstein DI, Blessing-Moore J, et al. Adverse reactions to vaccines practice parameter 2012 update. *Journal of Allergy and Clinical Immunology* 2012; 130:25-43.

Table 1. Four patients misdiagnosed as systemic allergic reactions after mRNA COVID vaccines.

Patient age (years) sex	Vaccine brand	Time to onset of symptoms (minutes)	Symptoms	Level of care: Treatment after first dose	Vaccine skin test results *	Symptoms with second dose #	Treatment after second dose
47 Female	Pfizer-BioNTech	5	sensation of throat swelling, shortness of breath	Emergency Department, Intensive Care Unit: Prednisone, cetirizine, epinephrine, famotidine methylprednisolone	Negative	sensation of throat swelling, shortness of breath	None
56 Female	Pfizer-BioNTech	2	hot and sweaty, headache, disoriented, pruritus	Emergency Department: diphenhydramine, dexamethasone, hydroxyzine	Negative	Headache	Acetaminophen
43 Female	Pfizer-BioNTech	15	itching, flushing	Urgent Care: diphenhydramine, epinephrine (self-administered), dexamethasone, famotidine	Negative	None	None
42 Female	Moderna	1	sensation of throat swelling, puffy eyelids,	Ambulance: diphenhydramine, intravenous epinephrine (which caused nausea,	Negative	Refused due to fear of epinephrine to treat a	Not applicable

			hives	shaking and paranoia), Emergency Department: diazepam		reaction should one occur	
--	--	--	-------	---	--	---------------------------------	--

* Using residual vaccine from multidose vials within six hours of reconstitution. Prick full strength, if negative intradermal diluted 1:100 with normal saline with positive (histamine) and negative (saline) controls. Prick test considered positive if ≥ 3 mm wheal and ≥ 10 mm flare. Intradermal test considered positive if ≥ 3 mm larger than initial wheal and ≥ 10 mm flare.

Administered in the usual manner with 1 hour observation period