Le sfide più grandi. La scienza più avanzata.

Siamo impegnati nel rispondere alle sfide più grandi in tema di salute.

Mettiamo in campo innovazione e passione, dove il bisogno è maggiore.

Come azienda biofarmaceutica globale, il nostro obiettivo è avere un impatto significativo sulla vita delle persone.

È con il contributo di tutti che i progressi della scienza si traducono in farmaci per milioni di persone.
Per questo collaboriamo con università e centri di ricerca, organizzazioni governative, associazioni di pazienti e no profit.

Insieme, costruiamo la medicina del futuro.

abbvie.it

abbvie

People. Passion. Possibilities.®

IT/ABV/1018/1141

Check for updates

Real-world data on the effectiveness, safety and drug survival of dupilumab: an analysis from the TREATgermany registry

DOI: 10.1111/bjd.21794

Dear Editor, Multiple randomized clinical trials (RCTs) have demonstrated robust efficacy and favourable safety of the monoclonal anti-interleukin-4Rα antibody dupilumab in moderate-to-severe atopic dermatitis (AD).[1–4] However, in RCTs selected patient populations are studied in a mostly shorter timeframe, and the results do not always reflect outcomes under real-life conditions. Real-world studies offer insights into the ways in which a drug is actually used, and outcomes regarding toxicity and effectiveness in daily medical practice. Real-world evidence both supplements and complements information gleaned from the prevailing standard of RCTs. The TREATgermany registry is one of the largest AD registries worldwide with currently more than 1400 patients with moderate-to-severe AD followed up prospectively.[5,6]

In this interim analysis we aimed to analyse the real-world long-term effectiveness and safety and the drug survival of dupilumab compared with ciclosporin. All of the ethics committees of the participating recruitment centres have approved TREATgermany.

For the current analysis, data from 1211 patients with active moderate-to-severe AD (mean age 40·7 years, 42·2% female) were available (data release July 2021). The mean (SD) objective baseline scores were objective Scoring Atopic Dermatitis [oSCORAD, 40·5 (16·3)], Eczema Area and Severity Index [EASI, 16·1 (12·9)], Investigator' Global Assessment [3·2 (1·1)], Physician's Global Assessment [3·1 (1·2)], average itch numerical rating scale over the last 3 days [5·7 (2·8)], Patient-Oriented Eczema Measure [16·8 (7·6)] and Dermatology Life Quality Index [DLQI, 11·8 (7·8)]. Among the patients, 924 (76·3%) had received at least one systemic treatment for AD during observation in the registry. With a proportion of 81·9% (n = 757) dupilumab was the most widely used systemic drug. In contrast, before 1 December 2017, when dupilumab was launched in Germany, only 36·8% (n = 91) of the patients enrolled had received at least one systemic therapy during observation in the registry, with ciclosporin being the most commonly used drug (48%, n = 44; Figure 1a).

For safety and effectiveness, we analysed 369 and 41 patients who received dupilumab and ciclosporin, respectively, and had complete data from at least one scheduled follow-up visit. The baseline characteristics between the two groups did not differ significantly. The respective mean ages at initiation of dupilumab and ciclosporin were 41·3 and 39·2 years, and the proportions of male patients were comparable (58·1% and 56% for dupilumab and ciclosporin, respectively). The respective mean baseline EASI, oSCORAD and DLQI of patients treated with dupilumab were 21·5, 46·5 and 14·0; the respective mean scores for the ciclosporin group were 18·2, 45·0 and 14·7. Likewise, effectiveness outcomes were comparable for the two drugs (Figure 1b; and further data at https://doi.org/10.6084/m9.figshare.20342475.v2). However, discontinuation rates of ciclosporin were considerably higher. While dupilumab treatment was discontinued in only 5·0% and 11·4% of the patients until month 12 and 24, ciclosporin was discontinued in 78% and 100%, respectively (Figure 1d). The most frequent reasons for discontinuation of ciclosporin were side-effects (31%) and insufficient efficacy (27%); in 56% no reasons were reported. Nineteen (46%) of those who discontinued therapy with ciclosporin received dupilumab afterwards. The proportions of patients with ≥ 50%, ≥ 75% and ≥ 90% improvement in EASI were 80%, 50% and 20% respectively after 3 months of dupilumab treatment.

Adverse events (AEs) were reported in 42·5% of all dupilumab-treated patients. The most frequent dupilumab-associated AEs were ocular complaints, which were reported in 29·8% of the patients (mostly conjunctivitis, in 20·7% of dupilumab patients) (Figure 1c), 5·5% of whom discontinued therapy. In 48·2% of the cases the ocular AEs were reported in the first 3 months of treatment.

The results from this interim analysis demonstrate that in Germany dupilumab has rapidly emerged as the most commonly used systemic treatment and shows robust long-term effectiveness and favourable safety under real-life conditions. Ocular side-effects were more common than in trials, but rarely led to treatment discontinuation. Drug survival was high and compares favourably with a previous report from the BioDay registry.[7] Potential explanations are a continued treatment benefit, but potentially also a lack of discontinuation due to persistently controlled disease, and the lack of availability of alternative new systemic treatment options, such as tralokinumab and Janus kinase inhibitors, which have only recently become available. Thus, a re-evaluation needs to be done in due course. When interpreting ciclosporin survival, it has to be considered that most guidelines do not recommend continuous therapy for > 1–2 years.

Acknowledgments: We would like to thank all of the participating patients, study physicians and study staff at the participating clinics and practices for their support of the German Atopic Eczema Registry TREATgermany. Open Access funding enabled and organized by Projekt DEAL.

Dora Stölzl,[1] Nicole Sander,[1] Annice Heratizadeh,[2] Eva Haufe,[3] Inken Harder,[1] Susanne Abraham,[4] Luise Heinrich,[3] Andreas Kleinheinz,[5] Andreas Wollenberg,[6] Elke Weisshaar,[7] Knut Schäkel,[8] Konstantin Ertner,[9] Franca Wiemers,[10] Julia Wildberger,[11] Margitta Worm,[12] Ralph von Kiedrowski,[13] Isaak Effendy,[14] Andrea Asmussen,[15] Matthias Augustin,[16] Mario Pawlak,[17] Michael Sticherling,[18] Alexander Zink,[19] Melanie Hilgers,[20] Christiane Handrick,[21] Sven Quist,[22] Beate Schwarz,[23] Petra Staubach-Renz,[24] Magnus Bell,[25] Sun-Hei Hong-Weldemann,[26]

© 2022 The Authors. British Journal of Dermatology published by John Wiley & Sons Ltd on behalf of British Association of Dermatologists.
This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

13652133, 2022, 6, Downloaded from https://onlinelibrary.wiley.com/doi/10.1111/bjd.21794 by CochraneItalia, Wiley Online Library on [20/12/2022]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License



Figure 1 (a) Proportion of patients in the registry who had received systemic therapy for atopic dermatitis and the type of systemic therapy up to December 2017 and July 2021. (b) Effectiveness of dupilumab (n = 369 at baseline) and ciclosporin (n = 41): percentages with ≥ 50%, ≥ 75% and ≥ 90% improvement in Eczema Area and Severity Index (EASI). (c) Most frequently (> 1%) reported adverse events under therapy with dupilumab and ciclosporin. (d) Drug survival of dupilumab and ciclosporin over 24 months, Kaplan–Meier curve.

© 2022 The Authors. *British Journal of Dermatology* published by John Wiley & Sons Ltd on behalf of British Association of Dermatologists.

13652133, 2022, 6, Downloaded from https://onlinelibrary.wiley.com/doi/10.1111/bjd.21794 by CochraneItalia, Wiley Online Library on [20/12/2022]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

Bernhard Homey,[27] Jens-Joachim BrÜcher,[28] Jochen Schmitt,[3] Thomas Werfel,[2] and Stephan Weidinger[1] the TREATgermany study group

[1]Center for Inflammatory Skin Diseases, Department of Dermatology and Allergy, University Hospital Schleswig-Holstein, Campus Kiel, Kiel, Germany; [2]Division of Immunodermatology and Allergy Research, Department of Dermatology and Allergy, Hannover Medical School, Hannover, Germany; [3]Center of Evidence-based Healthcare, University Hospital and Medical Faculty Carl Gustav Carus, TU Dresden, Dresden, Germany
Email: dstoelzl@dermatology.uni-kiel.de

D.S. and N.S. contributed equally as first authors.

J.S., T.W. and S.W. contributed equally as senior authors.

Funding sources: TREATgermany is an academia-led, investigator-initiated clinical registry that is financially supported by AbbVie GmbH & Co KG, Galderma SA, LEO Pharma GmbH, Lilly Deutschland GmbH, Pfizer Inc. and Sanofi Deutschland GmbH.

Conflicts of interest: T.W. is co-principal investigator of the German Atopic Dermatitis Registry TREATgermany and has received honoraria for lectures or scientific advice on atopic dermatitis from AbbVie, Almirall, Galderma, Janssen/JNJ, LEO Pharma, Leti, Lilly, Novartis, Pfizer and Regeneron/Sanofi. S.W. is co-principal investigator of the German Atopic Dermatitis Registry TREATgermany; has received institutional research grants from LEO, Pfizer and Sanofi; and has performed consulting work and lectures for AbbVie, Almirall, Kymab, LEO Pharma, Lilly, Pfizer, Sanofi, Regeneron, LEO, Eli Lilly, AbbVie, Pfizer, GSK and Kymab. J.S. is leading principal investigator of the German Atopic Dermatitis Registry TREATgermany; has received institutional grants from Novartis and Pfizer for scientifically initiated research; and has received honoraria for consulting from Sanofi, Lilly, Novartis and ALK. S.A. has received lecture and/or consultancy fees from Novartis, LEO Pharma, Amgen, Lilly, Sanofi, Beiersdorf, Janssen, UCB and AbbVie. All of the other authors declare they have no conflicts of interest.

Data availability statement: The data that support the findings of this study are available from the corresponding author upon reasonable request.

Supporting Information

Additional Supporting Information may be found in the online version of this article at the publisher's website:

Appendix S1 Full list of affiliations.

References

1 Simpson EL, Bieber T, Guttman-Yassky E *et al.* Two phase 3 trials of dupilumab versus placebo in atopic dermatitis. *N Engl J Med* 2016; **375**:2335–48.
2 Ratchataswan T, Banzon TM, Thyssen JP *et al.* Biologics for treatment of atopic dermatitis: current status and future prospect. *J Allergy Clin Immunol Pract* 2021; **9**:1053–65.
3 Stölzl D, Weidinger S, Drerup K. A new era has begun: treatment of atopic dermatitis with biologics. *Allergol Select* 2021; **5**:265–73.
4 Drucker AM, Morra DE, Prieto-Merino D *et al.* Systemic immunomodulatory treatments for atopic dermatitis: update of a living systematic review and network meta-analysis. *JAMA Dermatol* 2022; **158**:523–32.
5 Schmitt J, Abraham S, Trautmann F *et al.* Usage and effectiveness of systemic treatments in adults with severe atopic eczema: first results of the German Atopic Eczema Registry TREATgermany. *J Dtsch Dermatol Ges* 2017; **15**:49–59.
6 Heratizadeh A, Haufe E, Stölzl D *et al.* Baseline characteristics, disease severity and treatment history of patients with atopic dermatitis included in the German AD Registry TREATgermany. *J Eur Acad Dermatol Venereol* 2020; **34**:1263–72.
7 Spekhorst LS, Ariëns LFM, van der Schaft J *et al.* Two-year drug survival of dupilumab in a large cohort of difficult-to-treat adult atopic dermatitis patients compared to cyclosporine A and methotrexate: results from the BioDay registry. *Allergy* 2020; **75**:2376–9.

Patient and disease characteristics associated with psychiatric symptoms and impaired quality of life in pityriasis rubra pilaris

DOI: 10.1111/bjd.21792

Dear Editor, Pityriasis rubra pilaris (PRP) is a rare papulosquamous condition that is comorbid with depression in 15–20% of cases.[1–3] A survey of 85 patients with PRP showed that PRP was associated with a worse quality of life (QoL) than other common dermatological conditions.[4] Despite this, there are scarce data on the burden of and characteristics associated with psychiatric symptoms and QoL in PRP. Herein, we examine psychiatric symptoms, QoL, and their associations in a multinational PRP cohort.

Patients were recruited from the PRP Alliance, an international PRP support group, via a mailing list. Inclusion criteria were age ≥ 18 at survey completion and biopsy-proven PRP diagnosed by a dermatologist. Patients were excluded if they had a concurrent diagnosis of a PRP mimic (e.g. psoriasis, eczema). Data were collected through a Qualtrics survey (Qualtrics; Provo, UT, USA). Mean Dermatology Life Quality Index (DLQI)[5] and QoL scores were compared using Kruskal–Wallis tests. Univariate and multivariate logistic and linear regressions were used to estimate odds ratios (ORs), beta coefficients (β) and 95% confidence intervals (CIs) for associations. Sensitivity analyses were performed for all adjustment variables. A P value of $< 0{\cdot}05$ (two-sided) was considered statistically significant. The study was exempted from approval by the UCLA Institutional Review Board. Data are available upon reasonable request from the corresponding author.

Response rate was 76·5% (525 of 686), and 452 patients were included from different continents (North

© 2022 British Association of Dermatologists.

13652133, 2022, 6, Downloaded from https://onlinelibrary.wiley.com/doi/10.1111/bjd.21794 by CochraneItalia, Wiley Online Library on [20/12/2022]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License