



Omalizumab chronic spontaneous urticaria

Efficacy, safety, predictors of treatment outcome, and time to response

Eustachio Nettis, MD^{*}; Luca Cegolon, MD, PhD, MSc[†]; Elisabetta Di Leo, MD, PhD[‡]; Fabio Lodi Rizzini, MD, PhD[§]; Aikaterini Detoraki, MD, PhD[¶]; Giorgio Walter Canonica, MD^{||}; on behalf of the Italian OCUREL study group

^{*} Department of Emergency and Organ Transplantation, School of Allergology and Clinical Immunology, University of Bari Aldo Moro, Bari, Italy

[†] Scientific Directorate, IRCCS Burlo Garofolo, Trieste, Italy

[‡] Section of Allergy and Clinical Immunology, Unit of Internal Medicine—"F. Miulli" Hospital, Acquaviva delle Fonti, Bari, Italy

[§] Allergy Unit, University of Brescia, Spedali Civili, Brescia, Italy

[¶] Department of Clinical Immunology, Clinical Pathology, Infectious Diseases and Dermatology, Azienda Ospedaliera Universitaria Federico II, Naples, Italy

^{||} Department of Internal Medicine, Respiratory Disease Clinic, IRCCS Humanitas Clinical and Research Center, Humanitas University, Milan, Italy

ARTICLE INFO

Article history:

Received for publication April 26, 2018.

Received in revised form May 31, 2018.

Accepted for publication June 15, 2018.

ABSTRACT

Background: Omalizumab is a recombinant anti-immunoglobulin E (IgE) antibody used in the treatment of patients with chronic spontaneous urticaria (CSU).

Objective: This multicentric study assessed the safety and efficacy of omalizumab in patients (n = 322) with CSU refractory to second-generation antihistamines, also investigating predictors of poor treatment outcome and time lag to response to anti-IgE therapy by serum auto-reactivity.

Methods: This retrospective observational study comprised a 4-week pretreatment period, a 24-week treatment period with omalizumab (300 mg/month), and a 16-week follow-up period. Primary efficacy endpoints were mean and median change in 7-day urticaria activity score (UAS7), weekly itch severity score (ISS), and hive score from baseline to 4-, 12-, and 24-week values. Secondary endpoints included the proportion of patients (defined "responders") with well-controlled urticaria (UAS7 ≤ 6) and complete treatment response (UAS7 = 0). Safety in terms of side effects was also assessed.

Results: Omalizumab significantly and consistently reduced the mean UAS7, ISS, and hive score from baseline to weeks 4, 12, and 24, with a clear decreasing trend over time. At the end of the treatment period (week 24), 84.2% of patients had a UAS7 score of 6 or less and 66.7% had a UAS7 of 0. Higher pretreatment IgE levels were less likely to be associated with poor treatment response (ie, UAS7 > 6). Patients with a positive autologous serum skin test (ASST) were significantly more likely to be "slow responders" to omalizumab treatment (ie, response beyond 8 days since omalizumab administration) than ASST-negative patients (P < .001). No treatment-related adverse events were recorded.

Conclusion: Monitoring baseline characteristics of patients before introduction of omalizumab therapy may help to predict treatment outcome in CSU patients.

© 2018 American College of Allergy, Asthma & Immunology. Published by Elsevier Inc. All rights reserved.

Introduction

Chronic spontaneous urticaria (CSU) is the most common subtype of urticaria, characterized by recurrent episodes of spontaneous wheals or angioedema with unknown triggers, lasting for more than 6 weeks.¹ Chronic spontaneous urticaria generally affects 0.5% to 1.0% of people 20 to 40 years of age, with a disease duration of 1 to 5 years.²

Reprints: Aikaterini Detoraki, MD, PhD, Department of Clinical Immunology, Clinical Pathology, Infectious Diseases and Dermatology, Azienda Ospedaliera Universitaria Federico II, Naples, Italy. E-mail: caterina.detoraki@gmail.com.

Disclosures: E. Nettis, E. Di Leo, L. Cegolon, F. Lodi Rizzini, A. Detoraki, and W. G. Canonica declare that they have no conflict of interest.

Disclaimer: Members of the Italian OCUREL study group are listed in the Acknowledgments section at the end of the article.

Funding sources: Medical writing assistance was funded by Novartis Farma, Italy.

Patients with CSU often experience sleep deprivation and psychiatric comorbidities such as anxiety and depression, which compromise the health-related and overall quality of life of the affected individuals.^{2,3}

Treatment of CSU can be challenging in most cases, because it is often difficult to identify the underlying cause of the disease, and symptomatic therapies remain the mainstay of treatment in these patients. First-line treatment of CSU includes second-generation nonsedating antihistamines.^{1,4,5} However, the licensed doses of these drugs relieve symptoms in only 50% of cases, and the current guidelines suggest up dosing of H1-antihistamines up to fourfold.¹ Both clinical experience and clinical studies support this approach, showing higher efficacy in many, but not all, patients.⁶ Patients refractory to antihistamine therapy are generally treated with up to 4 times the recommended dose of second-generation antihistamines plus omalizumab, or cyclosporine A in severe cases.^{1,5}

Omalizumab is a recombinant anti-immunoglobulin (Ig) E antibody approved for the treatment of CSU refractory to antihistamines.^{7,8} It is administered subcutaneously at the recommended doses of 150 or 300 mg every 4 weeks. “The mechanism of action is still unclear, although omalizumab seems to block the binding of IgE to the FcεRI receptor on the surface of target cells, including mast cells and basophils, thus reducing receptor expression and the release of inflammatory mediators.”⁹

Because predictors of response to omalizumab treatment in CSU patients are still unclear, this study aims to evaluate efficacy and safety of this drug in patients with H1 antihistamine-refractory CSU, investigating also baseline factors influencing treatment outcome as well as time lag to response to omalizumab by serum autoreactivity.

Methods

Study Design and Patients

This multicentric retrospective observational study involved 23 Italian allergy and clinical immunology secondary care centers. The study comprised a 4-week pretreatment period, followed by a 24-week treatment period, and a 16-week follow-up period. Patients aged 12 to 85 years included in the study had a history of spontaneous urticaria for more than 6 weeks, not responding to treatment with licensed doses of non-sedating second-generation H1-antihistamines for at least 4 weeks, with a daily urticaria activity score (UAS) of 4 or higher assessed in clinic on 1 of the pretreatment checkup days (days –14, –7, or 1), and a 7-day urticaria activity score (UAS7) greater than 16 assessed during 7 days before the first study treatment. Baseline autologous serum skin test (ASST) was performed as per standard guidelines. The ASST was considered to be positive if the wheal induced by serum was 1.5 mm larger in diameter than that induced by saline.¹⁰ Total IgE serum levels were also measured, using immunofluorometric assay (ImmunoCAP; ThermoFisher, Uppsala, Sweden) and were expressed in kUA/L, according to the manufacturer's instructions.

The key exclusion criteria were as follows: patients with serious psychological disturbances; history of malignancy or hypersensitivity to omalizumab; or treatment with omalizumab within the previous 6 months.

All procedures complied with the Helsinki Declaration of 1964, subsequently revised in 2013. The study protocol was approved by the ethical committee of Naples University Hospital, Italy. Informed consent was obtained from all patients who agreed to participate to this study.

According to the recommendations of the Italian Drug Agency for CSU, omalizumab was administered subcutaneously every 4 weeks at doses of 300 mg for 24 weeks in total (first treatment course) and was administered again (second treatment course) at least 8 weeks after the end of the first treatment course if the UAS7 score showed values similar to those at pretreatment.¹¹

Throughout the study period, patients were required to maintain stable doses of their pretreatment therapy with H1 antihistamines at approved doses. During this time, patients were therefore allowed to use up to 4 times the licensed dose of H1 antihistamines or systemic glucocorticoids, as needed, for symptomatic relief.

Study End Points and Statistical Analysis

The primary efficacy end points included mean and median change in UAS7 score, weekly itch severity score (ISS), and hive score from baseline to respective values at weeks 4, 12, and 24. The *t* test was employed for comparison of means.

Secondary efficacy outcomes included the proportion of patients with well-controlled urticaria (UAS7 ≤ 6) and complete response to

omalizumab treatment (UAS7 = 0) at weeks 4, 12, 24, and 40. Safety was evaluated by monitoring and recording the frequency and severity of adverse events (AEs).

Various factors were then investigated as predictors of response to omalizumab treatment, contrasting patients with UAS7 score of greater than 6 (nonresponders¹²) with those with UAS7 ≤ 6 (responders¹²) at week 24. Logistic regression analysis was employed to investigate treatment response as binary endpoint (UAS7 > 6 vs UAS7 ≤ 6) by different baseline characteristics. Total levels of pretreatment IgE were categorized into quartiles. A multivariate logistic regression model was fitted by selecting factors significant at the univariate analysis. The results of the multivariable logistic regression model, expressed as odds ratio (OR) with 95% confidence interval (95%CI), were also adjusted for the effect of participating centers.

Patients responding within 8 days since omalizumab administration were considered “fast responders,” and those responding between 8 days and 6 months were considered “slow responders.”¹³

The speed of response to omalizumab (*fast responders vs slow responders*) was tested against serum reactivity based on ASST results (positive vs negative) in a univariable exact logistic regression analysis. Results were expressed as OR with 95%CI.

Statistical analyses were performed using Stata 14.2 package (Stata Corporation, College Station, TX). Level of significance was set at <.05. Missing values were excluded, and complete case analysis was performed.

Results

The study was conducted at 23 allergy and clinical immunology secondary care centers in Italy, all of which were members of the Italian Society of Allergy, Asthma and Clinical Immunology task force. A total of 322 patients with refractory CSU who received at least 1 dose of omalizumab at the study centers from September 2015 to August 2017 were included in the study and were examined by 56 different physicians.

The mean age of patients was 46.5 ± 14.3 years, 222 patients (68.9%) were female, and the mean total IgE at baseline was 231.4 ± 506.6 kUA/L (Table 1). Mean duration of CSU was 44.1 ± 64.3 months, and 167 patients (53.2%) had a history of angioedema. The mean baseline UAS7 score was 27.9 ± 8.3, and in-clinic UAS was 4.8 ± 0.8; the baseline hive score was 13.3 ± 5.2, whereas baseline ISS was 14.4 ± 4.7. Angioedema was reported by 111 (36.3%) patients in the week before treatment.

Efficacy

As seen in Table 2, UAS7 scores diminished from baseline to week 4 (mean change = –19.5), up to week 12 (mean change = –22.7) and week 24 (mean change = –24.6), with a significant decreasing trend over time (*P* trend < .001). Similar mitigated trends from baseline to weeks 4, 12, and 24 were also observed for mean values of weekly hive scores (*P* trend < .001) as well as ISS (*P* trend < .001).

Nineteen patients (5.9%) discontinued the study during the 24-week treatment period; 12 patients had disease progression, 3 had personal problems, 2 had AEs, and 2 were lost to follow-up. Eleven patients (3.4%) discontinued treatment between week 24 and week 40 and were lost to follow-up.

During the follow-up period, 119 of 292 (40.8%) patients repeated treatment with omalizumab (second treatment course) because of an increase in the mean UAS7 score. At the end of the study, 173 of 292 (59.2%) patients did not repeat omalizumab treatment.

At the end of the 40-week study, 107 of 173 patients (61.8%) reported well-controlled urticaria (UAS7 score ≤ 6), and complete response to omalizumab treatment (UAS7 score = 0) was observed in 75 of 173 treated patients (43.4%) (Table 2).

Table 1
Baseline Characteristics of Patients Included in the Study

Baseline characteristics	N	
Age, years		
Mean ± SD	322	46.5 ± 14.3
Median (range)		46.5 (15–83)
Sex, n (%)		
Male	322	100 (31.1)
Female		222 (68.9)
Duration of CSU, months		
Mean ± SD	322	44.1 ± 64.3
Median (range)		24 (2–588)
History of CSU medications, n (%)		
H1 antihistamines at regular dose	322	322 (100.0)
H1 antihistamines at high dose		225 (69.9)
Leukotriene receptor antagonists		66 (20.5)
Systemic steroids		237 (73.6)
Cyclosporines		43 (13.3)
History of angioedema, n (%)	313	167 (53.2)
Presence of angioedema, ^a n (%)	306	111 (36.3)
Total IgE levels, KUA/L		
Mean ± SD	193	231.4 ± 506.6
Median (range)		120 (0–5237)
Positive thyroid autoantibody test, n (%)	236	58 (24.7)
Positive ASST, n (%)	160	76 (47.5)
In clinic UAS		
Mean ± SD	322	4.8 ± 0.8
Median (range)		5 (4–6)
UAS7 ^a	322	
Mean ± SD		27.9 ± 8.3
Median (range)		27 (17–42)
Weekly ISS ^a		
Mean ± SD	322	14.4 ± 4.7
Median (range)		14 (3–21)
Weekly No. hive score ^a		
Mean ± SD	322	13.3 ± 5.2
Median (range)		13 (0–21)

ASST, autologous serum skin test; CSU, chronic spontaneous urticaria; ISS, itch severity score; kUA, kilo Unit/liter of IgE; SD, standard deviation; UAS, urticaria activity score; UAS7, 7-day urticaria activity score.

^aBased on data collected in patient's daily diary in the week before first treatment.

Based on these data, factors significantly associated with a UAS7 score greater than 6 24 weeks after treatment start with omalizumab at univariable analysis were baseline angioedema, cyclosporine use, angioedema history, IgE levels, and (to some extent) duration of CSU (Table 3). At multivariate analysis, only higher pretreatment IgE values—categories (48–119) kUA/L and (120–236) kUA/L—were significantly less likely to be associated with a UAS7 score greater than 6 at the end of the 24-week treatment period (Table 3).

Autologous serum skin test was performed on only 160 patients. The results of ASST performed on the 135 patients who responded to omalizumab treatment (of 255 responders) were analyzed.

Sixty-six patients responding within 8 days of omalizumab administration were categorized as “fast responders,” and the

Table 2
Efficacy of Omalizumab Treatment During the Study (n = 322)

Endpoint	Week 4	Week 12	Week 24	Week 40
Change from baseline in UAS7 score				
Mean ± SD	−19.5 ± 10.9 ^a	−22.7 ± 10.7 ^a	−24.6 ± 9.9 ^a	—
Median (range)	−19 (−42, 9)	−21 (−42, 18)	−23 (−42, 4)	—
Change from baseline in ISS				
Mean ± SD	−9.7 ± 6.0 ^a	−11.6 ± 6.1 ^a	−12.3 ± 5.8 ^a	—
Median (range)	−10 (−21, 5)	−12 (−21, 13)	−12 (−21, 6)	—
Change from baseline in hive score				
Mean ± SD	−9.4 ± 6.2 ^a	−11.1 ± 5.9 ^a	−12.0 ± 5.7 ^a	—
Median (range)	−9 (−21, 14)	−10 (−21, 10)	−11 (−21, 3)	—
Proportion of patients with well-controlled urticaria (UAS7 ≤ 6)	58.4% (188/322)	73.4% (232/316)	84.2% (255/303)	61.8% (107/173)
Proportion of patients with complete response (UAS7 = 0)	38.2% (123/322)	54.1% (171/316)	66.7% (202/303)	43.4% (75/173)

ISS, itch severity score; SD, standard deviation; UAS7, 7-day urticaria activity score.

^at test: $P < .001$ vs baseline.

Table 3
Logistic Regression Analysis for UAS7 Scores as Binary Endpoint (> 6 vs ≤ 6)^a

Variable	Univariate analysis		Multivariate analysis
	OR (95% CI)	P value	aOR (95% CI)
Age, years			
<65	Reference	0.685	—
> 65	0.81 (0.27, 2.41)		
Sex			
Female	Reference	0.971	—
Male	1.01 (0.52, 1.97)		
Duration of CSU, months	1.01 (1.00, 1.01)	0.011	1.00 (1.00, 1.01)
History of angioedema			
No	Reference	0.010	Reference
Yes	2.36 (1.20, 4.62)		1.50 (0.38, 5.96)
Presence of angioedema			
No	Reference	0.002	Reference
Yes	2.77 (1.46, 5.27)		1.70 (0.48, 6.01)
Thyroid antibodies			
No	Reference	0.937	—
Yes	1.04 (0.42, 2.58)		
ASST		0.371	—
No	Reference		
Yes	1.52 (0.60, 3.85)		
Cyclosporine use		0.012	Reference
No	Reference		
Yes	2.81 (1.31, 6.05)		2.60 (0.70, 9.66)
Steroid use		0.058	
No	Reference		
Yes	2.08 (0.93, 4.66)		
High-dose antihistamine therapy		0.587	—
No	Reference		
Yes	1.21 (0.61, 2.41)		
Total IgE, kUA/L		0.011	Reference
<48	Reference		
48–119	0.12 (0.03, 0.50)		0.11 (0.02, 0.69)
120–236	0.28 (0.09, 0.89)		0.24 (0.06, 0.92)
≥237	0.58 (0.22, 1.55)		0.33 (0.09, 1.21)

OR, odds ratio; aOR, adjusted odds ratio; ASST, autologous serum skin test; CSU, chronic spontaneous urticaria.

^aOdds ratios crude (OR) and adjusted (aOR) with 95% confidence interval (95%CI) and likelihood ratio test P value.

remaining 69 patients were classed as “slow responders.” Nineteen of the 66 (28.8%) “fast responders” were ASST positive, compared with 44 of the 69 (63.8%) “slow responders” ($P < .001$) (Table 4). The ASST-positive patients were 4.30 times more likely to have a slow response to omalizumab treatment as compared with ASST-negative patients.

Safety

The overall incidence of AEs during the 24-week treatment phase was 11.2%, and 31 patients (9.6%) reported at least 1 AE. Migraine, asthenia, and administration site conditions were the most common AEs (Table 5). No treatment-emergent AEs were reported during the study.

Table 4
Patients Responding to Omalizumab by ASST Result

ASST result	Slow responders (Total = 69) N (%)	Fast responders (Total = 66) N (%)	OR (95% CI)	P value
Negative	25 (36.2)	47 (71.2)	Reference	<.001
Positive	44 (63.8)	19 (28.8)	4.30 (1.99–9.60)	

NOTE. “Slow responders” = patients who responded to omalizumab between 8 days and 6 months after omalizumab administration.

“Fast responders” = patients who responded to omalizumab within 8 days of omalizumab administration.

Exact logistic regression analysis.

ASST, autologous serum skin test; CI, confidence interval; N, Number; OR, odds ratio.

Discussion

The results of this study showed that omalizumab effectively reduced the symptoms of CSU in antihistamine-refractory patients. After 24 weeks of treatment, omalizumab significantly decreased the mean UAS7, ISS, and weekly hive scores. Furthermore, 84.2% of patients reported well-controlled urticaria at the end of the 24-week treatment period, and complete response to treatment was seen in 66.7% of patients. Overall, the treatment was well tolerated.

The UAS7 is a validated and widely used patient-reported measure of CSU that captures the intensity of pruritus and number of hives.¹⁴ Studies with omalizumab have reported a reduction in UAS7 scores with treatment in patients with CSU.^{15,16} A randomized, double-blind, placebo-controlled study of omalizumab in patients with CSU reported that treatment with omalizumab decreased UAS7 scores during the 24-week treatment period.¹⁵ Another prospective open-label study with omalizumab reported a reduction in the UAS7 scores from baseline (5.7 vs 32.2) with treatment.¹⁶ The results of the current study are in line with these reports and show that omalizumab treatment effectively reduced the UAS7 scores in patients with CSU over the 24-week treatment period, thus improving the overall symptoms in these patients.

Currently, the predictors of response to omalizumab treatment in CSU are not well known. A study by Ghazanfar and colleagues¹⁷ reported that favorable response to omalizumab treatment can be predicted by the presence of CSU, no angioedema, negative histamine release test, older age, shorter duration of symptoms, and no history of treatment with systemic immunosuppressant drugs.¹⁷ Another prospective study with omalizumab reported that patients with lower baseline IgE values showed poor response to treatment and that the ratio of IgE values after treatment to baseline IgE levels of patients can help in predicting the treatment response in patients with CSU.¹⁸ Similarly, the current study confirmed poor treatment outcome in relation to some baseline characteristics such as cyclosporine administration, presence of angioedema, longer duration of CSU symptoms, and low level of IgE. These results are in line with previous reports identifying markers of response to omalizumab treatment in patients with CSU, which may be of significant value in clinical settings. However, after controlling for the simultaneous effect of all factors, the only important predictor of efficacious treatment response in our study was higher levels of pretreatment IgE.

Table 5
Summary of Adverse Events (AEs) Reported During the 24-Week Treatment Period

AEs, n (%)	N = 322
Overall	36 (11.2)
Migraine	10 (3.1)
Asthenia	8 (2.5)
Administration site conditions	7 (2.2)
Skin rash	4 (1.2)
Arthralgia	4 (1.2)
Nasopharyngitis	1 (0.3)
Dyspepsia	1 (0.3)
Fever	1 (0.3)
Any AE leading to discontinuation of study	2 (0.6)

The ASST is a simple in vivo clinical test for the detection of basophil histamine-releasing activity. The ASST has a sensitivity of approximately 70% and a specificity of 80%, and it may be used as a reasonably predictive clinical test to indicate the presence of functional circulating anti-FcεRIα and anti-IgE autoantibodies,¹⁰ but it can cause false-positive results because of the formation of a large amount of bradykinin and secretion of tryptases from neutrophils during the coagulation process.¹⁹

Our study also showed that patients who were ASST positive were less likely to be “fast responders” (ie, response within 8 days of omalizumab administration) to omalizumab treatment than were those who were ASST negative. This result confirms the findings of Gericke et al,¹³ who also identified a significant association between ASST positivity and slower time lag to symptom relief with omalizumab in CSU patients,¹³ suggesting that omalizumab reduced the number of FcεRI receptors on mast cells and basophils in these patients.

Treatment with omalizumab was well tolerated in the current study. The most common AEs were migraine, asthenia, and administration site conditions, which were not related to omalizumab treatment. The main limitation of this study is its design, being observational and retrospective, without controls. In addition, although current guidelines for the diagnosis of CSU still recommend the in vivo ASST to define a CSU subgroup as having autoreactive urticaria,¹ concerns have been raised regarding its interpretation and specificity.¹⁰

In conclusion, omalizumab was well tolerated and effective in the treatment of patients with CSU refractory to second-generation non-sedating antihistamines. Monitoring the baseline characteristics of patients before introduction of omalizumab therapy may help to predict treatment outcome in CSU patients.

Acknowledgments

We thank Nishad Parkar, PhD, of Springer Healthcare Communications, for writing the first draft of the manuscript, and Melanie Gatt, PhD, an independent medical writer, who provided editorial assistance on behalf of Springer Healthcare Communications. This medical writing assistance was funded by Novartis Farma, Italy.

Italian OCUREL (Omalizumab Chronic Urticaria in Real Life) study group: I. Baiardini (Genova), M. Cancian (Padova), S. Capretti (Firenze), G. Colombo (Milano), M. Conte (Pordenone), M.T. Costantino (Mantova), A. D'Angelo (Borgomanero), G. De Feo (Salerno), T. De Pasquale (Civitanova Marche), A. de Paulis (Napoli), M. Di Gioacchino (Chieti), E. Favero (Conegliano Veneto), S. Fichera (Catania), F. Gaeta (Roma), S. Gangemi (Messina), A. Gatta (Chieti), E. Heffler (Catania), L. La Rosa (Borgomanero), D. Macchia (Firenze), L. Macchia (Bari), E. Maggi (Firenze), C. Marasco (Bari), I. Martignago (Parma), M. Mineni (Brescia), A. Pannofino (Policoro), R. Parente (Salerno), S. Peveri (Piacenza), S. Pucci (Civitanova Marche), A. Radice (Firenze), L. Ricciardi (Messina), E. Ridolo (Parma), A. Romano (Roma), O. Rossi (Firenze), E. Savi (Piacenza), G. E. Senna (Verona), R. Senter (Padova), G. Spadaro (Napoli), G. Stefanizzi (Verona), A. Vacca (Bari), A. Vignoli (Mantova), D. R. Villalta (Pordenone), M. Yacoub (Milano), I. Zaza (Bari).

References

- [1] Zuberbier T, Aberer W, Asero R, et al. The EAACI/GA2LEN/EDF/WAO Guideline for the definition, classification, diagnosis, and management of urticaria: The 2017 revision and update. *Allergy*. 2018; Jan 15. <https://doi.org/10.1111/all.13397>.
- [2] Maurer M, Weller K, Bindslev-Jensen C, et al. Unmet clinical needs in chronic spontaneous urticaria: A GA(2)LEN task force report. *Allergy*. 2011;66:317–330.
- [3] Maurer M, Abuzakouk M, Berard F, et al. The burden of chronic spontaneous urticaria is substantial: Real-world evidence from ASSURE-CSU. *Allergy*. 2017;72:2005–2016.
- [4] Bernstein JA, Lang DM, Khan DA, et al. The diagnosis and management of acute and chronic urticaria: 2014 update. *J Allergy Clin Immunol*. 2014;133:1270–1277.
- [5] Powell R, Leech S, Till S, et al. BSACI guideline for the management of chronic urticaria and angioedema. *Clin Exp Allergy*. 2015;45:547–565.
- [6] Guillen-Aguinaga S, Jauregui Presa I, Aguinaga-Ontoso E, et al. Updosing non-sedating antihistamines in patients with chronic spontaneous urticaria: A systematic review and meta-analysis. *Br J Dermatol*. 2016;175:1153–1165.
- [7] Genentech Inc. USA. Xolair® Prescribing Information. https://www.gene.com/download/pdf/xolair_prescribing.pdf. Accessed December 21, 2017.
- [8] Novartis Pharma GmbH. Xolair® Summary of product characteristics. http://www.ema.europa.eu/docs/en_GB/document_library/EPAR_-_Product_Information/human/000606/WC500057298.pdf. 2015. Accessed December 21, 2017.
- [9] Chang TW, Chen C, Lin C-J, et al. The potential pharmacologic mechanisms of omalizumab in patients with chronic spontaneous urticaria. *J Allergy Clin Immunol*. 2015;135:337–342.
- [10] Konstantinou G, Asero R, Maurer M, et al. EAACI/GA2LEN task force consensus report: The autologous serum skin test in urticaria. *Allergy*. 2009;64:1256–1268.
- [11] L'Agenzia Italiana del Farmaco. Piano terapeutico (PT) AIFA per la prescrizione di Xolair (omalizumab) (orticaria cronica spontanea). http://www.gazzettaufficiale.it/do/atto/serie_generale/caricaPdf?cdimg = 15A0625400100010110001&dgu = 2015-08-21&art.dataPubblicazioneGazzetta = 2015-08-21&art.codiceRedazionale = 15A06254&art.num = 1&art.tiposerie = SG. Accessed March 16, 2017.
- [12] Maurer M, Kaplan A, Rosen K, et al. The XTEND-CIU study: Long-term use of omalizumab in chronic idiopathic urticaria. *J Allergy Clin Immunol*. 2018;141:1138–1139.
- [13] Gericke J, Metz M, Ohanyan T, et al. Serum autoreactivity predicts time to response to omalizumab therapy in chronic spontaneous urticaria. *J Allergy Clin Immunol*. 2017;139:1059–1061.
- [14] Mathias SD, Crosby RD, Rosén KE, et al. The minimal important difference for measures of urticaria disease activity: Updated findings. *Allergy Asthma Proc*. 2015;36(5):394–398.
- [15] Saini SS, Bindslev-Jensen C, Maurer M, et al. Efficacy and safety of omalizumab in patients with chronic idiopathic/spontaneous urticaria who remain symptomatic on H1 antihistamines: A randomized, placebo-controlled study. *J Invest Dermatol*. 2015;135:67–75.
- [16] Sussman G, Hebert J, Barron C, et al. Real-life experiences with omalizumab for the treatment of chronic urticaria. *Ann Allergy Asthma Immunol*. 2014;112:170–174.
- [17] Ghazanfar MN, Sand C, Thomsen SF. Effectiveness and safety of omalizumab in chronic spontaneous or inducible urticaria: Evaluation of 154 patients. *Br J Dermatol*. 2016;175:404–406.
- [18] Ertas R, Ozyurt K, Atasoy M, et al. The clinical response to omalizumab in chronic spontaneous urticaria patients is linked to and predicted by IgE levels and their change. *Allergy*. 2018;73(3):705–712.
- [19] Asero R, Tedeschi A, Riboldi P, et al. Plasma of patients with chronic urticaria shows signs of thrombin generation, and its intradermal injection causes wheal-and-flare reactions much more frequently than autologous serum. *J Allergy Clin Immunol*. 2006;117:1113–1117.