

Department of Clinical Microbiology and Immunology, Faculty
of Medicine, Tel Aviv University, Tel Aviv, Israel

Correspondence

Ariel Munitz, Department of Clinical Microbiology and
Immunology, The Sackler School of Medicine, Tel Aviv
University, Ramat Aviv 69978, Israel.
Email: arielm@tauex.tau.ac.il

ORCID

Ariel Munitz  <https://orcid.org/0000-0003-1626-3019>

REFERENCES

1. Jacobsen EA, Jackson DJ, Heffler E, et al. Eosinophil knock-out humans: uncovering the role of eosinophils through eosinophil-directed biological therapies. *Annu Rev Immunol*. 2021;39:719-757.

2. Dolitzky A, Shapira G, Grisaru-Tal S, et al. Transcriptional profiling of mouse eosinophils identifies distinct gene signatures following cellular activation. *Front Immunol*. 2021;12:802839.
3. Rosenberg HF, Dyer KD, Foster PS. Eosinophils: changing perspectives in health and disease. *Nat Rev Immunol*. 2013;13(1):9-22.
4. Mesnil C, Raulier S, Paulissen G, et al. Lung-resident eosinophils represent a distinct regulatory eosinophil subset. *J Clin Invest*. 2016;126(9):3279-3295.
5. Hassani M, Tak T, van Aalst C, et al. Differential effects of short- and long-term treatment with mepolizumab on eosinophil kinetics in blood and sputum in eosinophilic asthma. *iScience*. 2021;24(8):102913.
6. Porter L, Toepfner N, Bashant KR, et al. Metabolic profiling of human eosinophils. *Front Immunol*. 2018;9:1404.

SUPPORTING INFORMATION

Additional supporting information may be found in the online version of the article at the publisher's website.

DOI: 10.1111/all.15363

Similar IgE binding patterns in Gulf of Mexico and Southeast Asian shrimp species in US shrimp allergic patients

To the Editor,

Shellfish allergy (SA) is a leading cause of food-induced anaphylaxis¹ and one of the most common causes of adult-onset food allergy worldwide, with 1%–3% of the United States (US) population affected.^{2–4} Nearly half (45%) of US adults with SA report utilizing emergency services for SA symptoms over their lifetime,² remaining at-risk for lethal allergic reactions. Several allergenic proteins have been identified across shellfish species, including tropomyosin (TM), arginine kinase (AK), myosin light chain (MLC), sarcoplasmic calcium-binding protein (SCP), hemocyanin, troponin C, and triosephosphate isomerase.⁵ (Table 1.) However, there are a large number of shrimp allergens that have been detected, but not yet characterized.⁶ The allergens of major importance in SA are the muscle proteins TM and AK. TM, the major allergen with specific-IgE antibodies in ≤90% of SA patients, is associated with severe clinical reactivity. AK is a pan-allergen with cross-reactivity with crustaceans and cephalopods.⁵

Cross-reactivity has been observed clinically when SA patients ingest various invertebrate species with subsequent allergic reactions, but further study of shrimp sIgE binding between different shrimp species is needed.^{7,8} This study examined the sIgE binding

patterns to 2 shrimp species from the Gulf of Mexico and Southeast Asia in US SA patients.

Shellfish allergy patients with a history of shrimp-induced allergic reactions, allergic reaction with clinical oral food challenge and positive immediate skin prick testing (IHST) and/or shrimp sIgE ImmunoCAP™ levels were recruited from the Baylor College of Medicine (BCM) Allergy and Immunology Clinics. The study was approved by the BCM IRB and all participants provided written, informed consent. The patients underwent IHST to shrimp extract (mixture of *Penaeus borealis*, *Penaeus monodon*, *Metapenaeus barbata*, and *Metapenaeopsis joyneri*), *Dermatophagoides pteronyssinus* (Der p1, Der p10), *Dermatophagoides farinae* (Der f1), cockroach, codfish, crab, lobster, and oyster using extracts from Greer™. The patients underwent prick and prick IHST to raw fresh shrimp (*Penaeus aztecus*), and cooked fresh shrimp (*Penaeus aztecus*). ImmunoCAP™ and ISAC™ customized testing by ThermoFisher™ assessed total IgE as well as sIgE levels for shrimp, recombinant *Penaeus aztecus* (TM), Der p10, Der p1, Der p2, recombinant *Penaeus monodon* AK, MLC, SCP, troponin C, crab, lobster, cockroach, clam, and oyster. Western blot (WB) analysis of sIgE binding profile to

Abbreviations: Der f1, *Dermatophagoides farinae*; Der p1, *Dermatophagoides pteronyssinus*; rPen a, recombinant *Penaeus aztecus*; rPen m, recombinant *Penaeus monodon*; SA, shrimp allergy; sIgE, specific IgE.

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2022 The Authors. *Allergy* published by European Academy of Allergy and Clinical Immunology and John Wiley & Sons Ltd.

TABLE 1 Allergenic molecules present in crustacean species

Species frequently implicated		Allergen name (IUIS)	Biochemical name	Molecular weight	Heat stability	Physiological function	Route of exposure
<i>Penaeus monodon</i> (Black tiger prawn) – Pen m	1	Cha f 1, Cra c 1, Hom a 1, Lit v 1, Mel l 1, Met e 1, Pan b 1, Pen m 1, Por p 1	Tropomyosin	34–38 kDa	Highly heat-stable	Muscle contraction	Ingestion Inhalation
<i>Litopenaeus vannamei</i> (White leg shrimp) Lit v	2	Cra c 2, Lit v 2, Pen m 2	Arginine kinase	40–45 kDa	Stable	Energy metabolism in muscles	Ingestion Inhalation
<i>Penaeus aztecus</i> (Brown shrimp) Pen a	3	Hom a 3, Lit v 3, Pen m 3	Myosin light chain	17–20 kDa	Stable	Muscle contraction	Ingestion
<i>Homarus americanus</i> (American lobster) Hom a	4	Cra c 4, Lit v 4, Pen m 4, Pon l 4	Sarcoplasmic calcium-binding protein	20–25 kDa	Stable	Muscle contraction regulation	Ingestion
	5	Cra c 6, Hom a 6, Pen m 6	Troponin C	20–21 kDa	Unknown	Calcium dependent activation of muscle contraction	Ingestion
<i>Crangon crangon</i> (Sand shrimp) Cra c	6	Pen m 7	Hemocyanin	75 kDa	Stable	Copper-containing Oxygen transport protein, anti-microbial property	Ingestion Inhalation
<i>Charybdis feriatus</i> (Crucifix crab) Cha f	7	Arc s 8, Cra c 8	Triose-Phosphate isomerase	28 kDa	Labile	Glycolysis (energy metabolism)	Ingestion Inhalation
<i>Portunus pelagicus</i> (Blueswimmer crab) Por p	8	Pen m 13	Fatty acid-binding protein	20 kDa	Unknown	Transport protein for lipophilic molecules (fatty acids)	Ingestion

Note: Registered allergen names in accordance with WHO/IUIS Allergen Nomenclature.

Matricardi PM, et al., EAACI Molecular Allergy User's Guide. *Pediatr Allergy Immunol*, 2016. 27(Suppl.23): p. 1–236.

the major allergenic proteins TM, AK, MLC, and SCP in *Penaeus aztecus* (Brown shrimp) and *Penaeus monodon* (Black tiger shrimp) in raw and boiled extracts was performed.

PATIENT POPULATION

Fifteen SA patients between the ages of 2–17 years were enrolled. (Table 2) On oral food challenge, the most frequent symptoms were urticaria and angioedema. Forty percent had antihistamine treatment, one required steroids, and none required epinephrine or albuterol.

ANALYSIS OF ALLERGEN-sIgE

When examining sIgE binding patterns to the major allergens, WB revealed nearly identical sIgE binding profiles to tropomyosin, MLC, and SCP in both species in the heated shrimp extracts. (Figure 1) Eight subjects (53%) demonstrated sIgE binding to TM and six to SCP (40%). There was variability in the binding to other shrimp proteins,

particularly in the unheated shrimp.⁵ The proposed sIgE binding allergens have been identified by molecular weight,^{5,6} and mass spectrometric in-gel analysis, (Table S1.) but need to be confirmed in future molecular studies.^{9,10}

The presence of Southeast Asian Black tiger shrimp sIgE has been established and quantified through ImmunoCAP™ ISAC™ testing, with over 50% of patients having binding to TM, 33% to SCP, 27% to AK, and 13% each to MLC and troponin C.⁶ Nevertheless, each individual patient's binding pattern in our study was similar between the two species, both unheated/raw (Figure 1A,B) and heated/boiled proteins (Figure 1C,D). Inhibitory immunoblot was performed which confirmed the identification of MLC and SCP. (Figure S1A,B) These similar sIgE binding patterns to the major allergens suggested conservation of the epitopes between different species of shrimp.

According to the Food and Agricultural Organization of the United Nations, *Penaeus aztecus* is native to the north-western Atlantic Ocean and Gulf of Mexico, while *Penaeus monodon* is native to the coasts of Southeast Asia, South Asia, East Africa, and Australia. Our study demonstrates shared epitopes among the major shrimp proteins for Gulf Brown shrimp and Southeast

TABLE 2 Demographics of US Shrimp Allergic (SA) population

	Total	
Gender		
Male	9 (60%)	
Female	6 (40%)	
Age (years)	9 (2–17)	
Race		
African American	5 (33.3%)	
Asian	1 (6.7%)	
Caucasian	3 (20%)	
Hispanic	6 (40%)	
Age of diagnosis (years)	4 (1–8)	
Skin test	Wheal (mm)	Flare (mm)
Shrimp extract	8 (0–42)	25 (0–56)
Raw shrimp	2 (0–8)	4 (0–25)
Cooked shrimp	1 (0–20)	1 (0–40)
Der p	5 (0–22)	15 (0–30)
Der f	9 (0–16)	20 (0–29)
Cockroach	4 (0–10)	10 (0–25)
Codfish	2 (0–30)	2.5 (0–48)
Crab	3 (0–20)	3 (0–25)
Lobster	2 (0–10)	4 (0–30)
Oyster	1 (0–7)	1 (0–15)
IgE (kU/L)		
Shrimp	3.35 (0–42.9)	
rPen a 1 (TM)	0.65 (0–41.1)	
Der p 10	0.68 (0–27.3)	
Der p 1	0 (0–100)	
Der p 2	0 (0–100)	
rPen m 2 (AK)	0 (0–0.81)	
rPen m 3 (MLC)	0 (0–4.01)	
rPen m 4 (SCP)	0 (0–15.1)	
rPen m 6 (TC)	0 (0–3.15)	
Total IgE	225 (2.71–5000)	
Crab	1.82 (0–33.5)	
Lobster	2.44 (0–50.2)	
Cockroach	0.59 (0–20.7)	
Clam	0.11 (0–8.95)	
Oyster	0 (0–8.25)	
Atopy history		
Asthma	9 (60%)	
AR	13 (86.7%)	
AD	8 (53.3%)	
FA other than shellfish	7 (46.7%)	
EGID	0 (0%)	
Onset of symptoms (minutes)	5 (5–30)	

TABLE 2 (Continues)

	Total
Symptoms	
Urticaria	4 (26.7%)
Angioedema	5 (33.3%)
Ocular	2 (12.5%)
Respiratory	0 (0%)
GI	0 (0%)
Treatment	
Antihistamine	6 (40%)
Epinephrine	0 (0%)
Steroids	1 (6.7%)
Albuterol	0 (0%)

Note: SA patients with a history of shrimp-induced allergic reactions, reaction on clinical oral food challenge and positive immediate skin prick testing (IHST) and/or shrimp sIgE ImmunoCAP™ and component ISAC™ levels were included. Data for age of onset, skin test, sIgE (kU/L) and onset of symptoms (minutes) presented as median (range). Abbreviations: AD, atopic dermatitis; AR, allergic rhinitis; EGID, eosinophilic gastrointestinal disease; FA, food allergy.

Asian Black tiger shrimp in US shrimp allergic subjects, implicating shared diagnostic and therapeutic potential for SA to varied shrimp species.

AUTHOR CONTRIBUTIONS

Sara Anvari, Contributed analysis tools, performed the analysis and wrote the paper. Shea Brunner, Contributed data, analysis tools and wrote the paper. Karen Tuano, Collected the data. Brenda Bin Su, Contributed analysis tools. Shaymaviswanathan Karnaneedi, Collected the data and performed analysis, and wrote the paper. Andreas L. Lopata, Conceived and designed analysis, performed the analysis and wrote the paper. Carla M. Davis, Conceived and designed analysis, collected data, and wrote the paper.

KEYWORDS

anaphylaxis, food allergy, shrimp allergy, shrimp oral immunotherapy, tropomyosin

FUNDING INFORMATION

The project was supported by the NIH-NIAID grant 1R34AI57948-01 and ThermoFisher Scientific.

ACKNOWLEDGMENT

We acknowledge funding from the National Institute of Allergy and Infectious Disease for this work. (1R34AI57948-01).

CONFLICT OF INTEREST

Drs. Anvari and Davis receive research contract funding from DBV Technologies, Regeneron and Aimmune Therapeutics. Dr. Davis is a




FIGURE 1 (A, B, C, D) Western blot analysis of the major and minor shrimp allergen proteins for different shrimp species in shrimp allergic and healthy controls (A) *Penaeus aztecus* (Brown shrimp), (B) *Penaeus monodon* (Black tiger shrimp), (C) heated extract of *Penaeus aztecus* (Brown shrimp), (D) heated extract of *Penaeus monodon* (Black tiger shrimp). The location of sIgE to proposed allergens are highlighted with red boxes. HC, healthy control

consultant for Moonlight Therapeutics and received grant funding from ThermoFisher Scientific. Dr Lopata received research contract funding from Aimmune Therapeutics. The other authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Sara Anvari^{1,2}

Shea Brunner^{1,2}

Karen S. Tuano^{1,2}

Brenda Bin Su^{1,2}

Shaymaviswanathan Karnaneedi³

Andreas L. Lopata^{3,4}

Carla M. Davis^{1,2}

Correspondence

Carla M. Davis, Department of Pediatrics, Division of Immunology, Allergy and Retrovirology, Baylor College of Medicine, Texas Children's Hospital, 1102 Bates Ave, Ste 330, 77030 Houston, TX, USA.

Email: carlad@bcm.edu

Sara Anvari and Shea Brunner co-first authors.

ORCID

Sara Anvari <https://orcid.org/0000-0002-9942-6188>

Brenda Bin Su <https://orcid.org/0000-0002-7236-9857>

Carla M. Davis <https://orcid.org/0000-0003-0866-7822>

REFERENCES

- Conrado AB, Patel N, Turner PJ. Global patterns in anaphylaxis due to specific foods: a systematic review. *J Allergy Clin Immunol.* 2021;148(6):1515-1525.e.
- Warren CM, Aktas OZ, Gupta RS, Davis CM. Prevalence and characteristics of adult shellfish allergy in the United States. *J Allergy Clin Immunol.* 2019;144(5):1435-1438.
- Wang HT, Warren CM, Gupta RS, Davis CM. Prevalence and characteristic of shellfish allergy in the pediatric population of the United States. *J Allergy Clin Immunol Pract.* 2020;8(4):1359-1370.
- Wood RA. Food allergen immunotherapy: current status and prospects for the future. *J Allergy Clin Immunol.* 2016;137(4):973-982.

¹Department of Pediatrics, Section of Immunology, Allergy and Retrovirology, Baylor College of Medicine, Texas Children's Hospital, Houston, Texas, USA

²Baylor College of Medicine, William T. Shearer Center for Human Immunobiology, Houston, Texas, USA

³James Cook University, Australian Institute of Tropical Health and Medicine, Centre for Molecular Therapeutics, Douglas, QLD, Australia

⁴Tropical Futures Institute, James Cook University, Singapore, Singapore

5. Ruethers T, Taki AC, Johnston EB, et al. Seafood allergy: a comprehensive review of fish and shellfish allergens. *Mol Immunol*. 2018;100:28-57. doi:10.1016/j.molimm.2018.04.008
6. Asero R, Pravettoni V, Scala E, Villalta D. House dust mite-shrimp allergen interrelationships. *Curr Allergy Asthma Rep*. 2020;20(4):9. doi:10.1007/s11882-020-0902-2
7. Nugraha R, Kamath S, Johnston E, Karnaneedi S, Ruethers T, Lopata A. Conservation analysis of B-cell allergen epitopes to predict clinical cross-reactivity between shellfish and inhalant invertebrate allergens. *Front Immunol*. 2019;10:2676. doi:10.3389/fimmu.2019.02676
8. MacGinnite A. Update on potential therapies for IgE-mediated food allergy. *Curr Allergy Asthma Rep*. 2017;17(1):4. doi:10.1007/s11882-017-0671-8
9. Gelis S, Rueda M, Valero A, Fernández EA, Moran M, Fernández-Caldas E. Shellfish allergy: unmet needs in diagnosis and treatment. *J Investig Allergol Clin Immunol*. 2020;30(6):409-420. doi:10.18176/jiaci.0565
10. Wai CYY, Leung NYH, Chu KH, et al. Overcoming shellfish allergy: how far have we come? *Int J Mol Sci*. 2020;21(6):2234. doi:10.3390/ijms21062234

SUPPORTING INFORMATION

Additional supporting information may be found in the online version of the article at the publisher's website.

DOI: 10.1111/all.15366

Anti-inflammatory effect of siRNAs targeted *IL-4* and *IL-13* in a mouse model of allergic rhinitis

To the Editor,

Allergic rhinitis (AR) is a chronic inflammatory disease of upper airways characterized by sneezing, itching, nasal congestion, and rhinorrhea. AR pathogenesis includes the elevation of allergen specific IgE in serum, inflammatory cell infiltration to nasal mucosa, and overexpression of Th2-cytokines (IL-4, IL-5, and IL-13). Current pharmacotherapy using intranasal corticosteroids, antihistamines, and anti-leukotriene medications results in symptom relief, but fail to regulate the underlying pathogenesis. Anticytokine therapy mainly with monoclonal antibodies (Mab) is a promising approach for the treatment of asthma and other allergic conditions. Dupilumab targeting receptor subunit IL-4R α (common for both IL-4 and IL-13) is now approved for severe asthma and atopic dermatitis treatment and was found efficient in uncontrolled persistent asthma and comorbid AR.¹ However, the high cost of this drug limits its broad implementation into clinical practice, and there is a need to develop effective and available therapies to suppress Th2 cytokines. RNA interference provides promising approach for anti-cytokine therapy of inflammation though sequence-specific silencing of disease-causing genes using small interfering RNAs (siRNAs).² In the current study, we aimed to evaluate the anti-inflammatory effect of nasal administration of siRNAs targeted *il-4* and *il-13* in mouse model of AR.

The mouse anti-IL-4 and IL-13 siRNAs^{3,4} as well as branched (dendrimeric) cationic peptide LTP for nucleic acids delivery into mammalian cells, including lymphocytes (important source of IL-4 and IL-13)⁵ were designed and tested previously. In the current study, we confirmed the ability of siRNA/LTP complexes to transfect lymphocytes. The anti-IL-4 and IL-13 siRNAs/LTP complexes (at mass ratio 1/12.5) substantially reduced the production of these cytokines in concanavalin A (ConA)-stimulated thymocytes compared with non-specific siRNA (Figure 1A, B).

To reveal the therapeutic potential of the siRNA-based complexes mice with OVA-induced AR were intranasally treated with the siRNAs/LTP complexes at doses of 5 and 15 μ g/mouse. Corticosteroid budesonide which is widely used in the indications of AR and allergic asthma treatment served as a positive control. The group of intact mice treated with PBS only served as negative control. Intranasal instillation of the complexes containing siRNAs against IL-4 and IL-13 effectively suppressed the production of these Th2-cytokines by lymphocytes in the local lymph nodes (Figure 1C,D), while did not influence IFN γ level (Figure 1E). Moreover, we found decreased serum levels of OVA-specific IgE and IgG1, but not IgG2a (Figure 1F-H), that indicates the shift towards Th1 response after siRNA-mediated suppression of IL-4 and IL-13. It is important to note that non-specific siGFP did not exhibit effects on IL-4 and IL-13 expression neither in