

Review article

Allergen-specific IgG4 in atopic disease

Aalberse RC, Van Milligen F, Tan KY, Stapel SO. Allergen-specific IgG4 in atopic disease.

Allergy 1993; 48: 559-569. © Munksgaard 1993.

R. C. Aalberse¹, F. Van Milligen¹,
K. Y. Tan², S. O. Stapel¹

¹ Publication Secretariat, CLB, Amsterdam; and

² Department of Allergy, Holy Hospital, Rotterdam, The Netherlands

Prof. Dr R. C. Aalberse
P/O Publication Secretariat
CLB
PO Box 9406
1006 AK Amsterdam
The Netherlands

Accepted for publication 28 May 1993

Allergen-specific IgG4 has often been regarded as a two-headed creature: potentially harmful as well as potentially protective. As more is found out about these antibodies, the harmful effects are found to be hard to substantiate, at least in the allergy domain. The protective effects (with respect to allergy) are still debated, but particularly from the field of parasitology the evidence is accumulating that IgG4 does, under certain conditions, effectively interfere with allergen-induced, IgE-mediated, effector-cell triggering; i.e., it acts as a blocking antibody. It should be noted that, with respect to parasite immunity, this may be an undesirable effect (32, 36, 44, 45, 69, 70, 105).

Particularly, recent research in the regulation of B cells by T cells and T-cell-dependent cytokines helps to explain the similarities and differences between IgE and IgG4. It will be explained that a striking similarity has been found between IgE and IgG4 with respect to the type of antigen that triggers these immune responses. As will be explained in some detail later, it is now clear that this effect is a reflection of the similar T-cell help requirements: TH2-type cells need to be activated for both immunoglobulin isotypes.

In view of this similarity between IgE and IgG4 with respect to *antigen* specificity, it was unexpected to find a marked difference in *epitope* specificity be-

tween IgE and IgG4. The consequence of the differences in epitope specificity is that binding of IgG4 antibody does not necessarily interfere very effectively with the binding of IgE antibodies. In this review, we will argue that the level of allergen-specific IgG4 does not accurately reflect the level of protection obtained, because only a fraction of the allergen-specific IgG4 can interfere effectively with IgE binding.

The position with respect to the use of IgG4 antibodies to monitor immunotherapy that we will put forward in this review is that this use of IgG4 antibody assays is justifiable, but its value should not be overrated. The reasoning behind this position statement is that conventional immunotherapy depends on the immunologically specific activation or inactivation of allergen-specific T cells. Currently, this means stimulation of TH2 cells, and this process can be monitored conveniently by measuring allergen-specific IgG4. If no IgG4 antibody is induced by conventional immunotherapy, the therapy is likely to have been ineffective in reaching the allergen-specific immune system.

A future challenge for immunotherapy might well be to exploit the increasing knowledge about B-cell regulation and devise strategies to stimulate the production of IgG4 antibodies with the correct epitope specificities.

History

The interest in IgG4 in relation to atopic disease started approximately 20 years ago because of two unrelated observations. Firstly, particularly in the guinea pig, but also in other experimental animals, IgG4 antibodies proved to be important for some anaphylactic reactions. Because these sensitizing antibodies were less species-specific than what are now known as the IgE-type antibodies, the term "heterocytotropic", as opposed to "homocytotropic", was proposed for these antibodies. These non-complement-fixing, short-time-sensitizing antibodies had a relatively high electrophoretic mobility, just like IgG4 (13). In his investigations of sensitizing IgG antibodies in man, Parish (61) found suggestive evidence for involvement of IgG4 in this anaphylactic activity.

Secondly, three groups of investigators (Devey et al. (24), van der Giessen et al. (91), and Stanworth et al.; unpublished), all presenting their data at a workshop at the Second International Congress of Immunology (Brighton, UK, 1974), found a relative overrepresentation of IgG4 in patients receiving allergen immunotherapy.

Three questions have been debated by allergists ever since:

- 1) is IgG4 a mast cell/basophil-triggering antibody?
- 2) are IgE and IgG4 immune responses linked to the same type of antigen (allergen)?
- 3) are IgG4 antibodies particularly effective as allergen-blocking antibodies?

Is IgG4 a mast cell/basophil-triggering antibody?

The observations by Parish (61) generated considerable scientific activity (14, 57, 67, 86, 101–104), summarized in several reviews (2, 7, 20, 26, 30, 48, 55, 64, 74, 85, 87, 92, 93). Most investigators now accept that basophils cannot be triggered for histamine release by allergens via IgG4, but that basophils of some donors can be triggered by some anti-IgG4 antibodies (67, 101, 102). The dust has not completely settled on this topic, as evidenced by several recent papers on the subject of IgG4-induced histamine release. Beauvais et al. (12) present data suggesting that the basophils are triggered indirectly, via eosinophils. Jimeno et al. (39) found marked differences between two anti-IgG4 monoclonal antibodies. These authors suggest involvement of the only $\text{Fc}\gamma$ -receptor reported to be present on basophils (9, 90): the low-affinity type II γ -receptor CD32 (which is not to be confused with the low-affinity type II ϵ -receptor CD23). Interestingly, this low-affinity IgG receptor was recently also found to have a low affinity for IgE (8, 41, 88). The analysis of

basophil triggering in the investigations by Jimeno et al. (39) is complicated by the presence of soluble IgG4, because these experiments were performed in whole blood rather than with plasma-free cells. Lichtenstein et al. (47) have presented data suggesting that IgG-IgE complexes are responsible for anti-IgG-induced histamine release. Interestingly, these investigators found most histamine-releasing activity with anti-IgG3 and least with anti-IgG4 (two monoclonal antibodies tested for each human isotype).

Lichtenstein et al.'s paper (47) touches upon the interesting subject of IgG anti-IgE autoantibodies, which are, to a substantial extent, of the IgG4 isotype (e.g., references 17, 78). However, this topic is outside the scope of this review.

One of the recurrent arguments in favor of a pathogenic role for IgG4 antibodies is that they are occasionally found in patients with a history suggestive of immediate-type allergy without demonstrable specific IgE (e.g., reference 43). The value of these observations is, however, very limited, because IgG4 antibodies in nonallergic control subjects are usually also found, particularly if an exposure-matched control is tested. Statistically, however, patients with a food allergy have been found to have higher mean IgG4-antibody levels than nonallergic controls (10, 11, 34, 37, 40, 52, 56, 73, 75), possibly reflecting increased exposure to the allergen as a consequence of a permeability-increasing local allergic reaction in the gut. In very young food-allergic children (less than 18 months old) with a physiologically unresponsive IgG4 system, the IgG1- rather than the IgG4-antibody level to the food is often in the upper percentiles for that age group. A similar explanation presumably holds for the increased IgG-antibody levels found in atopic eczema patients (2, 19, 29, 31, 34, 51, 68, 72, 73, 76, 77, 106) or patients with celiac disease (37).

Biologic properties of IgG4 antibodies

In large antibody excess, IgG4 antibodies form multi-antigenic complexes just as IgG1 antibodies do. This can be shown, for example, by the binding of these complexes to the rheumatoid factor. In contrast to the IgG1 complexes, however, these IgG4 complexes do not bind to the first complement component C1q. IgG4 antibodies, therefore, do not activate the classical complement route. In fact, IgG4 antibodies were found to interfere with the complement activation by IgG1 antibodies (95).

Upon lowering the antibody/antigen ratio, researchers noted another difference between IgG1 antibodies and IgG4 antibodies. With IgG1 antibodies, the size of the immune complexes increases dramatically, whereas the size of the IgG4 complexes

decreases. Further experiments indicated that IgG4 antibodies are functionally monovalent (6, 80, 94). Yet, physicochemical evidence indicates the presence of two antigen-binding sites in IgG4. These results can be reconciled by assuming that binding of one antigen sterically prevents the binding of a second antigen. A practical consequence of this functional monovalency is that IgG4 antibodies are not detected in antibody assays that depend on bivalency, such as the reverse sandwich assay (100). A related issue is the apparently low avidity of IgG4 antibodies in some assays (23, 65). IgG4 complexes are too small to precipitate in polyethylene glycol, and this may result in spuriously low avidity values. Using anti-immunoglobulin reagents (protein A or anti-IgG4) and avoiding the pitfall of competing "invisible" antibodies in the other isotypes, a pitfall which also leads to an underestimation of the avidity, researchers have found the avidity of IgG4 antibodies to be of the same order of magnitude as that of IgG1 or IgE antibodies.

It is occasionally assumed that the functional monovalency of IgG4 makes it unsuitable to act as a cytophilic antibody, like IgE, to become cross-linked by antigen.

As indicated above, cross-linking of two IgG4 antibodies by a single antigen is not necessarily a problem, although the restricted hinge flexibility might impose some barriers. However, the engagement of Fc-receptors would be limited to two, whereas IgE-mediated cross-linking seems to be considerably more effective at slightly higher aggregation ratios. This is presumably an academic discussion, because the only Fc-receptor known to be present on basophils that bind IgG4, the type II Fc- γ -receptor, binds only polymeric, i.e., antigen-complexed, IgG4.

IgG4 as an indicator of chronic antigenic stimulation

The marked contribution of IgG4 to the immune response induced by repeated allergen injections has been known for a long time. This suggested some link with allergy. However, we soon found that also in conditions unrelated to allergy repeated antigen stimulation resulted in an IgG4-dominated antibody response. This was particularly clear in a study of the immune response in novice beekeepers (6). During the first few months of beekeeping, we found predominantly IgG1 antibodies. In more than 50% of the beekeepers, IgE antibodies also appeared, usually with large local reactions, but without severe generalized allergic reactions. IgG4 dominance became apparent particularly during the second year of beekeeping. Professional beekeepers had extremely high IgG4 antibody levels, sometimes even resulting in increased total IgG4 levels. Since these

professional beekeepers had virtually no reaction to bee stings, it is likely that the immune response was protective in the conventional way: antitoxic immunity comparable to immunity to diphtheria toxin.

Although these findings might be described as a switch from IgG1 (and possibly IgE) to IgG4, this would be confusing, since it suggests a class-switching event on a cellular level. It is improbable that the cells that originally produce IgG1 antibodies themselves switch to IgG4 production. It is more likely that repeated antigenic stimulation results in the maturation of previously resting B cells (presumably memory cells) that subsequently mature and undergo a class switch to become IgG4-secreting plasma cells.

Are IgE and IgG4 immune responses linked?

All IgE-inducing antigens investigated so far have proved to be excellent IgG4-inducers, including not only many different atopic allergens, but also parasite allergens and castor bean allergens. The reverse does not seem to be necessarily the case, as evidenced by a food protein that induces strikingly high IgG4 responses without a marked contribution to food allergy: the banana protein BanLec I (42). Analysis by Western blotting of complicated allergen extracts (22) or parasite extracts (35) reveals the IgG4 response to be more similar to the IgE response than the other immunoglobulin isotypes. On the level of the total allergen molecule, IgE and IgG4 responses are thus found to be associated (48). Presumably, the explanation is that both isotypes are IL-4 dependent and thus require the stimulation of TH2-type T-helper cells (27, 28, 38, 49, 59, 81–84).

Antigens that, for whatever reason, are effective TH2-inducers will thus promote both IgE and IgG4 responses, whereas antigens that induce TH1 cells are more likely to stimulate IgG1 and IgG3 responses. (Note the awkward nomenclature problem: the codes for the human isotypes do not correspond to the murine isotypes; human IgG4 is, to some extent, comparable to mouse IgG1, whereas human IgG1 is more like mouse IgG2a; human IgG2 behaves like mouse IgG3.)

Because of this similarity between IgE and IgG4 on immunoblots, it was tempting to suggest that IgE and IgG4 are descended from the same parental B cell, diverging upon class switching in the daughter cells. One possible scheme would be to postulate that nonatopic B cells switch to IgG4 because of a relatively low IL-4 production by TH2 cells, whereas a significant fraction of atopic B cells switch to IgE because of higher IL-4 production in these patients. The concentration of IL-4 required for switching to

human IgG4 (or mouse IgG1) is likely to be lower than that required for IgE switching.

It has even been suggested that IgG4 antibodies that are produced after immunotherapy result from switching IgE-producing cells to IgG4-producing cells. This is unlikely for several reasons. One reason is that the induction of IgG4 antibodies occurs even in patients in whom the IgE antibody level does not decrease. Another reason is that a switch from IgG4 to IgE rather than the other way around would be expected from what is currently known about isotype switching. Indeed, in mice there is good evidence for isotype switching from (murine) IgG1 to IgE (50, 79).

According to this common parental cell scheme, the epitope specificity of IgE and IgG4 antibodies would be expected to be very similar if not identical. Recent data from our laboratory, however, indicate a marked difference between IgE and IgG4 in epitope specificity. These results were obtained in three different test systems, all three using variants of natural allergen molecules.

In one approach, we analyzed *phylogenetic variants* of allergens. We compared the IgE and IgG4 responses to the major birch allergen *Bet v 1* and the homologous protein in apple (21). The cross-reactivity of these two proteins for IgE is well known (16). We found that in one patient clear IgG4 reactivity with apple was induced upon immunotherapy with birch pollen extract, but that, in contrast to the situation with IgE antibodies against apple, most IgG4 reactivity with apple extract was, in general, not related to birch pollen. These data suggested that IgE and IgG4 antibodies against the birch allergen did not fully overlap.

Similar, but more quantitative, data were obtained with another well-characterized major allergen, the cat allergen *Fel d 1* (99). In this case, we compared the response to the cat molecule with that to the ocelot homologue. The ocelot was more suited than some of the other wild members of the cat family we tested, e.g., lion and tiger, which proved to be too cross-reactive for the present purpose.

In order to exclude any contribution of feline allergens distinct from *Fel d 1* and its ocelot equivalent, we used affinity-purified human antibodies to study the cross-reactivity in a quantitative way. Allergen-specific antibody from sera of cat-allergic patients was purified on a *Fel d 1* affinity column. The purified antibodies were tested in IgE and IgG4 RAST for antibody activity to cat and ocelot. As with the birch-apple study, we found a higher cross-reactivity for IgE than IgG4.

A second approach was the use of *recombinant chains* of the same allergen, *Fel d 1*. The primary structure of this allergen has been established, and *E. coli*-derived recombinant protein has been pro-

duced (71). The allergen is a dimer of a two-chain molecule; it is called chain 1 and chain 2. We found more immune reactivity with chain 1 than with chain 2, but for both chains we found higher reactivity for IgE than for IgG4 (97).

The third approach was to compare IgE and IgG4 binding to small *chemically synthesized peptides* (98). By testing a panel of overlapping peptides, we first identified several IgE-binding peptides. These peptides were subsequently used to isolate by affinity chromatography peptide-reactive antibodies from human sera. Then, we measured IgE- and IgG4-antibody activity among these peptide-reactive antibodies. For this assay, we used purified radio-labeled allergen (i.e., ^{125}I -labeled *Fel d 1*). The ratio (IgE anti-allergen/IgG4 anti-allergen) was higher in the affinity-purified material than in the starting serum. We did not find indications for a lower recovery of IgG4 antibodies in such affinity purification procedures. These results indicate that IgE is more reactive with partially processed allergen than IgG4.

IgE responses to atopic allergens in atopic and nonatopic subjects

The nature of the immunologic difference between atopic and nonatopic subjects is one of the basic problems in allergy. Platts-Mills has shown that, for the grass pollen allergen *Lol p 1* (66) and for the house-dust mite allergen *Der p 1* (18), not only are the IgE antibody levels higher in atopics, but often also the IgG antibody levels. Nonallergic persons usually have little or no detectable antibody against these two allergens. This effect seems to be a reflection of the exposure level, because this difference at the allergen-specific IgG level between allergic and non-allergic subjects disappears at higher exposure levels. Natural exposure to *Lol p 1* or *Der p 1* is very low, at least 10-fold lower than exposure to the cat allergen *Fel d 1* when a cat is in the house. It is not uncommon to find substantial levels of IgG antibodies against the cat allergen without concomitant IgE antibodies (33, 92, 96) (Fig. 1). Even higher levels of exposure are found for experimental animals (e.g., rodents) in the case of occupational exposure. In this situation, virtually everyone develops IgG antibodies (46). The IgG isotype follows the usual pattern: it starts with an IgG1-dominated response and, upon continued exposure, gradually shifts, in most cases, towards an IgG4-dominated response.

These findings suggest that the difference between allergic and nonallergic responses might be antigen-dose-related: allergic responses might be induced at immunizing doses too low to stimulate effective IgE-control mechanisms, and atopic subjects might be more inclined towards an immune response (both




Fig. 1. IgG antibodies against two purified allergens in allergic and nonallergic donors. *Lol p 1* from grass pollen and *Fel d 1* from cat were iodinated and used in antigen-binding test with protein A Sepharose as IgG-binding reagent. Per test, 2.5 μ l serum was used. Allergic patients had not received immunotherapy. Under these conditions, no binding of allergen via IgE was found. These results illustrate that 1) IgG antibodies against purified allergens are more common in allergic than in nonallergic donors; 2) IgG antibodies against high-exposure allergen (*Fel d 1*) are more common in nonallergic donors than IgG antibodies against low-exposure allergens (*Lol p 1*). Contribution of IgG4 to IgG response in these sera is usually less than 50%.

IgG and IgE) at a low antigen dose. In other words, the antigen threshold may be lower in atopy.

Despite the fact that both IgE and IgG responses are often induced by allergens that enter the body via the mucosal membranes, these responses are not of mucosal type: the usual IgA domination of the classical mucosal immune response is absent and the antibodies are not preferentially excreted. The antibody levels in the secretions are usually a fairly close reflection of the plasma levels.

To accommodate the data currently available on the antiallergen immune response, we have proposed a two-compartment hypothesis (van Milligen (96)). According to this hypothesis, the normal immune response to allergens occurs when the antigen is taken up by a dendritic cell and transported to the lymphoid organs and there induces a well-orchestrated immune response. This response may include some IgE, but this IgE is relatively harmless because of the excess of IgG antibodies with the same specificity. An example is the human response to tetanus toxoid immunization. Under some conditions (particularly infestation by certain parasites), an unusually large amount of IL-4 is produced, resulting in a larger production of IgE. This IgE is still not as harmful as might have been expected, because of the presence of even larger quantities of IgG antibodies (particularly IgG4) with exactly the same epitope specificity.

An allergic immune response occurs (still according to this two-compartment hypothesis) when iso-

lated, solitary immune cells are confronted by an antigen outside the lymphoid organs. Usually such an encounter would not trigger an immune reaction, because the microenvironment is unsuitable. Occasionally, the local situation is permissive, probably in relation to some transient, special kind of inflammatory reaction that attracts and activates the types of cells required for the triggering of a few B and T cells. In many cases, these signals also activate the regulatory circuits, and then no harm is done.

Sometimes, however, the signals are not sufficiently strong to activate suppressive mechanisms, and IgE is produced in quantities quite similar to those of IgG antibodies that are locally induced. Compared to "ordinary" IgG responses, this is a negligible response. Most of the IgG produced by allergic patients is from the lymphoid compartment, and thus of different epitope specificity. This lymphoid IgG synthesis might be triggered via the IgE-mediated, CD23-dependent, antigen-presentation pathway (53, 54). The basic immunologic problem in atopy might then be related to an increased risk of mounting these "ectopic" immune responses; e.g., because of enhanced antigen trapping in extralymphoid sites. The preferential recognition by IgE antibodies of partially unfolded epitopes suggests that allergens are perhaps sticking to a tissue matrix component at the time of the first B-cell encounter. Another factor contributing to enhanced antigen trapping might be a preexisting IgG antibody response to cross-reacting structures on the allergen. This would fit with two other observations (15):

- 1) for some foods it has been shown that cross-reactivity exists with inhalant allergens
- 2) a higher than normal immune response to foods is a risk factor for the development of inhalant allergy.

Apparently, IgG4 production results not only from the activation of B-cell precursors that give rise to the IgE-producing B cells, but also from precursors that yield few or no IgE-producing B cells.

Can IgG4 antibody assays be used for diagnosing allergies?

As discussed above, IgG4 antibodies are found independently of atopic background in virtually everyone with sufficient exposure to TH2-stimulating antigens. Nevertheless, a statistical association is found between allergic symptoms and high IgG4 antibody levels to the offending allergen. For mite and pollen allergens, and, to a lesser extent, for cat allergens, more allergen-specific IgG4 antibodies are found in donors who also make IgE antibodies. These results are probably best explained by assuming that in atopy an increased reactivity of the immune system

to TH2-type responses exists. A contributing factor might be the enhanced, IgE-mediated antigen presentation (54). This difference is much more marked for the most potent allergenic components in an allergen extract than for nonallergenic components, and it is often (but not always, e.g., not in young children (40)) more marked for IgG4 than for other isotypes.

An additional problem is that in some cases the elevated IgG antibody titers are a consequence rather than a cause of the disease. It is well known that conditions such as wheat intolerance may result in elevated IgG antibody levels not only against gliadin but also chicken ovalbumin and other food antigens, presumably secondarily to malfunctioning of the intestinal mucosa.

Despite the statistical difference between allergic patients and nonallergic controls that can be found in some conditions, it is hard to define a condition where the assay of allergen-specific IgG4 would be advisable (52). In most of these conditions, a better discrimination can be found with specific IgE assays.

Investigations in which allergic patients without detectable allergen-specific IgE were identified with allergen-specific IgG4 did not include the two types of reference values that are essential for the correct interpretation of an IgG4 antibody assay: 1) a sufficient number of age- and exposure-matched controls; 2) responses of the same patient to other antigens with similar exposure characteristics.

Can IgG4 antibody assays be used to monitor immunotherapy?

The question of whether IgG4 antibody assays can be used to monitor immunotherapy brings us back to the last question mentioned in the beginning: is IgG4 antibody particularly effective as an allergen-blocking antibody? The current answer to the latter question is that, presumably, it is not, but it is the best we have. The next question then might be this: is an IgG4 antibody assay relevant to monitor the effects of conventional immunotherapy? (1, 3, 58, 60, 63, 89). There is a tendency to push aside completely the concept of blocking antibodies as a factor in immunotherapy. Undoubtedly, there is more to immunotherapy than blocking antibodies alone. It is now abundantly clear that the late inflammatory sequelae of the acute allergic reaction are a major factor in the pathology, and it is highly probable that T-cell activation plays an important role in this inflammatory process. Downregulation of these allergen-specific, proinflammatory T cells is an important issue in immunotherapy. It should be kept in mind that these T cells are antigen-specific; i.e., they have to be activated by the allergen. One of the possible effects of blocking antibody might be on

T-cell activation, or rather on prevention of antigen presentation to T cells. Antigen presentation to allergen-specific T cells has been shown to be greatly enhanced by allergen-specific IgE, probably via the low-affinity IgE receptor CD23 on the antigen-presenting cell (54). Competition between IgE and blocking antibody at this level (IgE on the low-affinity receptor on an antigen-presenting cell rather than IgE on the high-affinity receptor on the mast cell) could explain the effects of immunotherapy in the late reaction.

However, conventional blocking activity between mast-cell-bound IgE and soluble blocking antibody is also a mechanism to keep in mind. Immunotherapy usually does have an effect on the immediate skin reactivity that cannot be attributed to a decrease in allergen-specific IgE.

Furthermore, allergen-neutralizing activity can be shown to be present in post-treatment serum and in parasitic infections (36, 69), by *in vivo* (skin test; e.g., reference 62), *in vitro* (basophil histamine release), and RAST inhibition/RAST interference (1, 3–5). Binding of IgG4 antibody to an allergen does not necessarily interfere with IgE binding. This can easily be shown *in vitro* by adding a small amount of allergen to a postimmunotherapy serum that contains both IgE antibodies and IgG4 antibodies (IgG4 being in excess), and analyzing by RIA the immune complexes formed. A large fraction of the IgG4-containing immune complexes has been found to contain IgE. This IgG4 is unlikely to be bound directly to IgE (i.e., as IgG4 autoanti-IgE or as IgE anti-IgG4), because this binding is allergen-dependent (Fig. 2). The implication is that a large fraction of allergen-specific IgG4 induced by immunotherapy has an epitope specificity different from the epitope specificity of IgE. These results thus confirm the conclusion of the direct epitope-mapping data mentioned earlier.

What do we know about the association between efficacy of immunotherapy and IgG4 antibody levels? Few investigators would deny that the overall association is strong: conventional immunotherapy is usually effective and usually results in a marked increase in allergen-specific IgG4. On the other hand, the association is certainly not perfect, and, on an individual patient basis, many exceptions can be found. Some patients do very well without much IgG4 response, whereas other patients show an impressive allergen-specific IgG4 response without much effect on their symptoms.

The first category of patient should not come as a surprise, considering the appreciable placebo score found for any therapeutic modality in allergy. The issue of whether this category is larger than can be explained by the placebo effect has, to our knowledge, not been addressed adequately.




Fig. 2. IgG4 binding to allergen does not necessarily interfere with binding of IgE antibodies. Serum of grass-allergic patient with high titer of IgG4 antibodies against *Lol p 1* and other grass pollen allergens was incubated with graded doses of grass pollen extract. Allergen-serum mixture was incubated with monoclonal anti-IgG4 coupled to Sepharose. After washing, labeled anti-IgE was added. Results illustrate formation of immune complexes containing, apart from allergen, both IgG4 and IgE. Even at very low allergen doses, no indication was found for lag phase that would suggest formation of IgE-free immune complexes. This is indicated by results of parallel experiments with allergen bound either directly to CNBr-activated Sepharose ("direct RAST") or to rabbit anti-allergen antibodies coupled to Sepharose ("antigrass catching"). These results support view that, because of epitope specificity different from IgE antibodies, no large fraction of allergen-specific IgG4 antibodies interferes effectively with IgE binding. However, RAST-interference studies show that small fraction of IgG4 antibodies does interfere (1, 3, 5).

The second category, patients with high IgG4 titers without clinical benefit, needs more discussion. One subgroup of such patients was identified by Djurup (25): patients with a high IgG4 titer before the start of the immunotherapy. This investigator found that these patients did not improve on subsequent treatment. One possible explanation for this phenomenon might be related to the answer to the question of why these patients had high levels of preexisting IgG4. In view of what is now known about IgG4, it is likely that these patients were highly exposed to the allergen in their environment. It would not be unexpected if patients with high allergen exposure would have less benefit from immunotherapy.

The lack of a tight correlation between IgG antibody titers and a decrease in symptom scores is undoubtedly also caused by the imprecision of determination of the two variables. Symptom scoring is not only imprecise, but it is also easily influenced by other factors, such as infections and additional allergies. IgG antibodies can be measured with a high precision, but these results are often confounded by the inclusion of irrelevant specificities, in measuring the pan-IgG response to a total allergen extract. A far more informative assay is obtained by

Allergen-specific IgG4 in atopic disease




Fig. 3. Follow-up study of patients receiving immunotherapy with dog (patient no. 1), grass pollen (patient no. 2), or house-dust mites (patients nos. 3 and 4). For each patient, the IgG antibody assays are shown before and during immunotherapy. Open bars represent allergen-specific pan-IgG (i.e., strictly speaking, protein A-binding isotypes: IgG1, 2, 4) antibodies; filled bars represent allergen-specific IgG4. Assay was performed with radiolabeled purified allergens (*Can f 1* for dog, *Lol p 1* for grass pollen, *Der p 1* for house-dust mites) in combination with anti-immunoglobulin (protein A or monoclonal anti-IgG4) coupled to CNBr-activated Sepharose. Note that, provided one uses purified allergens, increase in antibody titer, particularly for IgG4, is well over 10-fold.

testing the IgG4 response to purified allergen. This is best illustrated by analyzing the "stimulation index" that can be found upon immunotherapy. With the first type of assay, a high background antibody response is found, which is present in allergic as well as in nonallergic patients. After immunotherapy, some increase can usually be found, but usually not more than three- to fivefold over pretherapy levels. This contrasts to the results obtained with the other procedure: IgG4 against a purified allergen (Fig. 3). Here the background is usually low, and upon successful treatment the antibody level usually increases more than 10-fold and occasionally more than 100-fold.

Using an assay with a high "stimulation index" enables a clear distinction between effective and ineffective treatment in the immunologic sense. Moreover, as indicated before, the response measured is an unmistakable sign that the TH2-arm of the immune system has been reached.

Conclusions

- 1) Allergen-specific IgG4 is unlikely to be a direct cause of allergic reactions.
- 2) IgG4 antibody responses are often associated with IgE antibody responses. This is presumably

a reflection of the requirements of IgE-inducing antigens to stimulate TH2-type immune responses.

- 3) Without immunotherapy, IgG antibodies against atopic allergens are not restricted to the IgG4 isotype: in many cases, IgG1 antibodies are predominant.
- 4) IgG4 antibodies against allergens are particularly prominent after prolonged immunotherapy, in line with the place of the IgG4 antibody response against other TH2-stimulating antigens.
- 5) IgG4 antibodies are often found also in non-allergic subjects, particularly after chronic exposure to protein antigens. Early in these responses, IgG1 antibodies usually predominate.
- 6) Despite the similarity between IgE and IgG4 with respect to the type of antigens recognized, the epitope specificity of IgG4 antibodies is often found to differ from that of IgE antibodies. These two apparently conflicting observations can be explained by assuming that the IgE and IgG4 responses are regulated by the same type of T cells but that distinct B cells may be involved.
- 7) From both the therapeutic and diagnostic points of view, it is important to be able to focus on IgG4 antibodies with the correct epitope specificity.
- 8) Classical immunotherapy can be considered to be immunologically effective if a substantial (10–100-fold) increase in allergen-specific IgG4 is induced.

Acknowledgments

The experimental results from this institute (CLB) presented in this review were obtained as part of projects supported by the Netherlands Asthma Fund, project 87.32.

References

1. AALBERSE RC. The use of radioimmunoassay procedures for the detection of IgG antibodies to allergens. In: RICCI M, FAUCI AS, ARCANGELI P, TORZUOLI P, eds. Developments in clinical immunology. London: Academic Press, 1978: 149–60.
2. AALBERSE RC. IgG subclasses in atopic eczema. In: RUZICKA T, RING J, PRZYBILLA B, eds. Handbook of atopic eczema. Berlin: Springer-Verlag, 1991: 166–72.
3. AALBERSE RC, DIEGES PH, KNUL-BRETLOVA V, VOOREN P, AALBERS M, VAN LEEUWEN J. IgG4 as a blocking antibody. In: HALPERN GM, ed. Clinical reviews in allergy. Vol. 1. New York: Elsevier, 1983: 289–302.
4. AALBERSE RC, REERINK-BRONGERS EE, VERMEULEN E. RAST-inhibiting factors in human serum. *Int Arch Allergy Appl Immunol* 1973; 45: 46–9.
5. AALBERSE RC, REERINK-BRONGERS EE, HOORWEG E. Methods for the assay of allergen-neutralizing antibodies based on competition with IgE antibodies for allergen. In: PEETERS H, ed. Protides of the biological fluids. 24th Colloquium. Oxford: Pergamon Press, 1976: 725–9.
6. AALBERSE RC, VAN DER GAAG R, VAN LEEUWEN J. Serologic aspects of IgG4 antibodies. I. Prolonged immunization results in an IgG4-restricted response. *J Immunol* 1983; 130: 722–6.
7. AALBERSE RC, VAN DER ZEE JS, VLUG A. IgG4 antibodies in atopic allergy. *Lab Manage* 1985; 23: 19–32.
8. ALBER G, KENT UM, METZGER H. Functional comparison of FCγRI, FcγRII, and FcγRIII in mast cells. *J Immunol* 1992; 149: 2428–36.
9. ANSELMINO LM, PERUSSIA B, THOMAS LL. Human basophils selectively express the FcγRII (CDw32) sub-type of IgG receptor. *J Allergy Clin Immunol* 1989; 84: 907–14.
10. BARNES RMR, JOHNSON PM, BLEARS J, HARVEY MM, FINN R. Human serum antibodies reactive with dietary proteins. Antigenic specificity. *Int Arch Allergy Appl Immunol* 188: 87: 189–93.
11. BARNES RMR, JOHNSON PM, HARVEY MM, BLEARS J, FINN R. Human serum antibodies reactive with dietary proteins. IgG subclass distribution. *Int Arch Allergy Appl Immunol* 1988; 87: 184–8.
12. BEAUVAIS F, HIEBLT C, BURTIN C, BENVENISTE J. Bimodal IgG4-mediated human basophil activation: role of eosinophils. *J Immunol* 1990; 14: 3881–90.
13. BINAGHI RA. Nonreaginic anaphylactic antibodies in animals. In: HALPERN GM, ed. Nonreaginic anaphylactic and/or blocking antibodies. Clinical reviews in allergy. Vol. 1, No. 2. New York: Elsevier, 1983: 237–47.
14. BRIGHTON WD. Frequency of occurrence of IgG(S-TS). *Clin Allergy* 1980; 10: 97–100.
15. CALKHOVEN PG. The role of food antigens in the development of inhalant allergy. Thesis, University of Amsterdam, Meppel: Krips Repro, 1989: 1–114.
16. CALKHOVEN PG, AALBERS M, KOSHTA VL, POS O, OEI HD, AALBERSE RC. Cross-reactivity among birch pollen, vegetables and fruits as detected by IgE antibodies is due to at least three distinct cross-reactive structures. *Allergy* 1987; 42: 382–90.
17. CARINI C, FRATAZZI C. Detection of IgG subclasses with anti-IgE activity in patients with atopic diseases. *Int Arch Allergy Appl Immunol* 1992; 98: 227–32.
18. CHAPMAN MD, PLATTS-MILLS TAE. Measurement of IgG, IgA and IgE antibodies to *Dermatophagoides pteromyssinus* by antigen-binding assay, using a partially purified fraction of mite extract (F4P1). *Clin Exp Immunol* 1978; 34: 126–36.
19. CHAPMAN MD, ROWNTREE S, MITCHELL EB, DI PRISCO DE FUENMAJOR MC, PLATTS-MILLS TAE. Quantitative assessments of IgG and IgE antibodies to inhalant allergens in patients with atopic dermatitis. *J Allergy Clin Immunol* 1983; 72: 27–33.
20. CONROY MC, ZHOU T, SPENGLER H, DE WECK AL. IgG4 as blocking or anaphylactic antibody. In: KERR JW, GANDERTON MA, eds. Proceedings XI International Congress of Allergy and Clinical Immunology, London, 1982. London: Macmillan, 1983: 341–5.
21. DE GROOT H. Allergens from cats and dogs. PhD thesis. Amsterdam: University of Amsterdam, 1990.
22. DESVAUX FX, PELTRE G, DAVID B. Characterization of grass pollen-specific IgE, IgA, IgM classes and IgG subclasses in allergic patients. *Int Arch Allergy Appl Immunol* 1989; 89: 281–7.
23. DEVEY ME, LEE SR, RICHARDS D, KEMENY DM. Serial studies on the functional affinity and heterogeneity of antibodies of different IgG subclasses to phospholipase A₂ produced in response to bee-venom immunotherapy. *J Allergy Clin Immunol* 1989; 84: 326–30.
24. DEVEY ME, WILSON DV, WHEELER AW. The IgG subclasses of antibodies to grass pollen allergens produced in hay fever patients during hyposensitization. *Clin Allergy* 1976; 6: 227–36.

25. DJURUP R. The subclass nature and clinical significance of the IgG antibody response in patients undergoing allergen-specific immunotherapy. *Allergy* 1985; 40: 469-86.
26. FAGAN DL, SLAUGHTER CA, CAPRA JD, SULLIVAN TJ. Monoclonal antibodies to immunoglobulin G4 induce histamine release from human basophils *in vitro*. *J Allergy Clin Immunol* 1982; 70: 399-404.
27. GASCAN H, GAUCHAT J, RONCAROLO M, YSSEL H, SPITS H, DE VRIES JE. Human B cell clones can be induced to proliferate and to switch to IgE and IgG4 synthesis by interleukin 4 and a signal provided by activated CD4+ T cell clones. *J Exp Med* 1991; 173: 747-50.
28. GASCAN H, GAUCHAT J-F, AVERSA G, VAN VLSSELAER P, DE VRIES JE. Anti-CD40 monoclonal antibodies or CD4+ T cell clones and IL-4 induce IgG4 and IgE switching in purified human B cells via different signaling pathways. *J Immunol* 1991; 147: 8-13.
29. GONDO A, SAEKI N, TOKUDA Y. IgG4 antibodies in patients with atopic dermatitis. *Br J Dermatol* 1987; 117: 301-10.
30. GOODWIN BFJ. Nonreaginic anaphylactic antibodies in man. In: HALPERN GM, ed. *Clinical reviews in allergy*. Vol. 1. New York: Elsevier, 1983: 249-58.
31. GWYNN CM, SHAKIB F. IgG subclasses in skin disorders. *Monogr Allergy* 1986; 19: 198-201.
32. HAGAN P, BLUMENTHAL UJ, DUNN D, SIMPSON AJG, WILKINS HA. Human IgE, IgG4 and resistance to reinfection with *Schistosoma haematobium*. *Nature* 1991; 349: 243-5.
33. HOMBURGER HA, MAUER K, SACHS MI, O'CONNEL EJ, JACOB GL, CARON JC. Serum IgG4 concentrations and allergen-specific IgG4 antibodies compared in adults and children with asthma and nonallergic subjects. *J Allergy Clin Immunol* 1986; 77: 427-34.
34. HUSBY S, LARSEN FS, AHLSTEDT S, SYEHAG SE. Humoral immunity to dietary antigens in atopic dermatitis. II. Analysis of IgE and IgG subclass antibodies. *Allergy* 1986; 41: 386-91.
35. HUSSAIN R, OTTESSEN EA. IgE responses in human filariasis. IV. Parallel antigen recognition of IgE and IgG4 subclass antibodies. *J Immunol* 1986; 136: 1859.
36. HUSSAIN R, POINDEXTER RW, OTTESSEN EA. Control of allergic reactivity in human filariasis: predominant localization of blocking antibody to the IgG4 subclass. *J Immunol* 1992; 148: 2731-7.
37. HYATUM M, SCOTT H, BRANDTZAEG P. Serum IgG subclass antibodies to a variety of food antigens in patients with coeliac disease. *Gut* 1992; 33: 632-8.
38. ISHIZAKA A, SAKIYAMA Y, NAKANISHI M, et al. The inductive effect of interleukin-4 on IgG4 and IgE synthesis in human peripheral blood lymphocytes. *Clin Exp Immunol* 1990; 79: 392-6.
39. JIMENO L, LOMBARDERO M, CARREIRA J, MOSCOSO DEL PRADO J. Presence of IgG4 on the membrane of human basophils. Histamine release is induced by monoclonal antibodies directed against Fab but not the Fc region of the IgG4 molecule. *Clin Exp Allergy* 1992; 22: 1007-14.
40. KEMENY DM, PRICE JF, RICHARDSON V, RICHARDS D, LESSOR MH. The IgE and IgG subclass antibody response to foods in babies during the first year of life and their relationship to feeding regimen and the development of food allergy. *J Allergy Clin Immunol* 1991; 87: 920-9.
41. KINET JP, ADAMCZEWSKI M, BRINI A, et al. Structure and function of FcγRI and of "new" receptors for immunoglobulin E. In: WÜTHRICH B, ed. *Highlights in allergy and clinical immunology*. Göttingen: Hogrefe & Huber, 1992: 28-9.
42. KOSHTA VL, AALBERS M, CALKHOVEN PG, AALBERSE RC. The potent IgG4-inducing antigen in banana is a mannose-binding lectin. *BanLec-I*. *Int Arch Allergy Appl Immunol* 1992; 97: 17-24.
43. KRAUT A, PENG Z, BECKER AB, WARREN CPW. Christmas candy maker's asthma: IgG4-mediated pectin allergy. *Chest* 1992; 102: 1605-7.
44. KURNIAWAN A, YAZDANBAKHSH M, VAN REE R, et al. Differential expression of IgE and IgG4 specific antibody responses in asymptomatic and chronic human filariasis. *J Immunol* 1993; 150: 1-10.
45. KWAN-LIM G-E, FORSYTH KP, MAIZELS RM. Filariasis-specific IgG4 response correlates with active *Wuchereria bancrofti* infection. *J Immunol* 1990; 145: 4298-305.
46. LEE D, EDWARDS RG, DEWDNEY JM. Measurement of IgG subclasses in the sera of laboratory workers exposed to rats. *Int Arch Allergy Appl Immunol* 1988; 87: 222-4.
47. LICHTENSTEIN LM, KAGEY-SOBOTKA A, WHITE JM, HAMILTON RG. Anti-human IgG causes basophil histamine release by acting on IgG-E complexes bound to IgE receptors. *J Immunol* 1992; 148: 3929-36.
48. LUCAS AH. IgG subclass-restricted immune responses to allergens. *Springer Semin Immunopathol* 1992; 12: 385-400.
49. LUNDGREN M, PERSSON U, LARSSON P, et al. Interleukin 4 induces synthesis of IgE and IgG4 in human B cells. *Eur J Immunol* 1989; 19: 1311-15.
50. MANDLER R, FINKELMAN FD, LEVINE AD, SNAPPER CM. IL-4 induction of IgE class switching by lipopolysaccharide-activated murine B cells occurs predominantly through sequential switching. *J Immunol* 1993; 150: 407-18.
51. MERRETT J, BARNETSON RS, BURR ML, MERRETT TG. Total and specific IgG4 antibody levels in atopic eczema. *Clin Exp Immunol* 1984; 56: 645-52.
52. MORGAN JE, DAUL CB, LEHRER SB. The relationships among shrimp-specific IgG subclass antibodies and immediate adverse reactions to shrimp challenge. *J Allergy Clin Immunol* 1990; 86: 387-92.
53. MUDDER GC, HANSEL TT, VAN REUSEN FC, OSTERHOFF BF, BRUIJNZEEL-KOOME CAFM. IgE: an immunoglobulin specialized in antigen capture. *Immunol Today* 1990; 11: 440-3.
54. MUDDER GC, VAN REUSEN FC, BOLAND GJ, DE GAST GC, BRUIJNZEEL PLB, BRUIJNZEEL-KOOME CAFM. Allergen presentation by epidermal Langerhans' cells from patients with atopic dermatitis is mediated by IgE. *Immunology* 1990; 69: 335-41.
55. NAKAGAWA T, DE WECK AL. Membrane receptors for the IgG4 subclass on human basophils. In: HALPERN GM, ed. *Clinical reviews in allergy*. Vol. 1. New York: Elsevier, 1983: 197-206.
56. NAKAGAWA T, MUKOYAMA T, BABA M, SAKAI Y, YAMASHITA N, MIYAMOTO T. Egg white-specific IgE and IgG4 antibodies in atopic children. *Ann Allergy* 1986; 57: 359-62.
57. NELSON HS, BRANCH LB. Incidence of IgG short-term sensitizing antibodies in an allergic population. *J Allergy Clin Immunol* 1977; 60: 266-70.
58. NORDVALL SL, RENCK B, EINARSSON R. Specific IgE and IgG antibody responses in children to timothy pollen components during immunotherapy. *Allergy* 1989; 44: 380-4.
59. NÜSSELIN HC, SPIEGELBERG HL. The role of lymphokines in IgE and IgG subclass formation. In: PICHLER WJ, STADLER BM, DAHINDEN CA, et al., eds. *Progress in allergy and clinical immunology*. Vol. 1. Toronto: Hans Huber, 1989: 149-51.
60. OEHLING A, SANZ ML, GARCÍA BE. Immunological parameters in the immunotherapy follow-up. *Int Arch Allergy Immunol* 1992; 99: 474-7.

61. PARISH WE. Skin sensitizing non-IgE antibodies. Association between human IgG S-TS and IgG4. In: BRENT L, HOLBOROW J, eds. Progress in immunology. II. Part 4. North Holland, 1974: 19-28.
62. PECQUET C, MURRIETA M, MICHELEN V, et al. Blocking activity of mite-specific IgG antibodies studied by skin-tests. *Allergy* 1989; 44: 427-31.
63. PENG Z, NACLERIO RM, NORMAN PS, ADKINSON NF Jr. Quantitative IgE- and IgG-subclass responses during and after long-term ragweed immunotherapy. *J Allergy Clin Immunol* 1992; 89: 519-29.
64. PERELMUTTER L. IgG4 and the immune system. In: HALPERN GM, ed. Clinical reviews in allergy. Vol. 1. New York: Elsevier, 1983: 267-87.
65. PERSSON MAA, BROWN SE, STEWARD MW, et al. IgG subclass-associated affinity differences of specific antibodies in humans. *J Immunol* 1988; 140: 3875-9.
66. PLATTS-MILLS TAE. Local production of IgG, IgA and IgE antibodies in grass pollen hay fever. *J Immunol* 1979; 122: 2218-25.
67. POULSEN LK, STAHL SKOV P, MOSBECH H, WEEKE B. Role of IgG4 in histamine release from human basophil leucocytes. I. Sensitization of cells from normal donors. *Int Arch Allergy Appl Immunol* 1988; 86: 383-90.
68. REINHOLD U, PAWELEC G, WEHRMANN W, HEROLD M, WERNER P, KREYSEL HW. Immunoglobulin E and immunoglobulin G subclass distribution *in vivo* and relationship to *in vitro* generation of interferon-gamma and neopterin in patients with severe atopic dermatitis. *Int Arch Allergy Appl Immunol* 1988; 87: 120-6.
69. RIHET P, DEMEURE CE, DESSEIN AJ, BOURGEOIS A. Strong serum inhibition of specific IgE correlated to competing IgG4, revealed by a new methodology in subjects from an *S. mansoni* endemic area. *Eur J Immunol* 1992; 22: 2063-70.
70. RIHET P, DEMEURE CE, OUATTARA M, et al. Immunological mechanisms of human resistance to schistosome infection: the protective effect of immunoglobulin E is balanced by a blocking action of immunoglobulin G4. In: GODARD P, BOUSQUET J, MICHEL FB, eds. Advances in allergology and clinical immunology (EAACI XV, Paris 1992). Camforth (UK): Parthenon, 1992: 83-91.
71. ROGERS BL, MORGENTHAU JP, GARMAN RD, BOND JF, KUO M-C. Expression of recombinant *Fc ϵ 1*: expression, purification, IgE binding and reaction with cat allergic human T cells. *Mol Immunol* 1993 (in press).
72. ROWNTREE S, MITCHELL EB, PLATTS-MILLS TAE. Assessment of skin reactivity and specific antibody to egg and milk proteins in patients with atopic dermatitis: comparison with responses to the house dust mite antigen. *Int Arch Allergy Appl Immunol* 1986; 79: 132-9.
73. ROWNTREE S, PLATTS-MILLS TAE, COGSWELL JJ, MITCHELL EB. A subclass IgG4-specific antigen-binding radioimmunoassay (RIA): comparison between IgG and IgG4 antibodies to food and inhaled antigens in adult atopic dermatitis after desensitization treatment and during development of antibody responses in children. *J Allergy Clin Immunol* 1987; 80: 622-30.
74. SHAKIB F. Basic and clinical aspects of IgG subclasses. Basel: Karger, 1987.
75. SHAKIB F, BROWN HM, REDHEAD R, PHELPS A. IgE and IgG4 antibodies to bovine milk fat globule membrane in atopic eczema patients: a study of their occurrence, relevance and antigenic specificity. *Clin Allergy* 1985; 265-71.
76. SHAKIB F, BROWN HM, STANWORTH DR. Relevance of milk- and egg-specific IgG4 in atopic eczema. *Int Arch Allergy Appl Immunol* 1984; 75: 107-12.
77. SHAKIB F, McLAUGHLIN P, STANWORTH DR, SMITH E, FAIRBAIRN E. Elevated serum IgE and IgG4 in patients with atopic dermatitis. *Br J Dermatol* 1977; 97: 59.
78. SHAKIB F, MILLS CS, POWELL-RICHARDS A. The recognition of a recombinant human Fc ϵ 1 fragment by the subclasses of IgG autoanti-IgE: disproportional subclass distribution of complexed autoantibody. *Int Arch Allergy Appl Immunol* 1992; 97: 243-8.
79. SNAPPER CM, MOND JJ. Towards a comprehensive view of immunoglobulin class switching. *Immunol Today* 1993; 14: 15-17.
80. SNOW ME, AMZEL LM. A molecular-mechanics study of the conformation of the interchain disulfide of human immunoglobulin G4 (IgG4). *Mol Immunol* 1988; 25: 1019-24.
81. SPIEGELBERG HL. Fc receptors for IgE and interleukin-4 induced IgE and IgG4 secretion. *J Invest Dermatol* 1990; 94 (Suppl.): 49S-52S.
82. SPIEGELBERG HL. The role of interleukin-4 in IgE and IgG subclass formation. *Springer Semin Immunopathol* 1990; 12: 365-83.
83. SPIEGELBERG HL, FALKOFF RJM, O'CONNOR RD, BECK L. Interleukin-2 inhibits the interleukin-4-induced human IgE and IgG4 secretion *in vitro*. *Clin Exp Immunol* 1991; 84: 400-5.
84. SPIEGELBERG HL, O'CONNOR RD, FALKOFF RJM, BECK L. Interleukin-4 induced IgE and IgG4 secretion by B cells from atopic dermatitis patients. *Int Arch Allergy Appl Immunol* 1991; 94: 181-3.
85. STANWORTH DR. IgG4 antibodies in atopics and their families. In: KERR JW, GANDERTON MA, eds. Proceedings XI International Congress of Allergy and Clinical Immunology, London, 1982. London: Macmillan, 1983: 337-40.
86. STANWORTH DR, SMITH K. Inhibition of reagin-mediated PCA reactions in baboons by the human IgG4 subclass. *Clin Allergy* 1973; 3: 37-41.
87. STAPEL SO, VAN DER ZEE JS, AALBERSE RC. Role of IgG4. EAACI proceedings. 1986; 13: 377-80.
88. TAKIZAWA F, ADAMCZEWSKI M, KINET J-P. Identification of the low affinity receptor for immunoglobulin E on mouse mast cells and macrophages as Fc ϵ gammaRII and Fc ϵ gammaRIII. *J Exp Med* 1992; 176: 469-76.
89. URBANEK R, COLLET M. Changes in skin and mucosa sensitivity correlated with specific IgE- and IgG-subclass antibodies in hyposensitized patients. In: PICHLER WJ, STADLER BM, DAHINDEN CA, et al, eds. Progress in allergy and clinical immunology. Vol. 1. Toronto: Hans Huber, 1989: 391-4.
90. VALENT P, BETELHEIM P. Cell surface structures on human basophils and mast cells: biochemical and functional characterization. In: DIXON FJ, ed. Advances in immunology. San Diego, CA: 1992: 333-423.
91. VAN DER GIESSEN M, HOMAN WL, VAN KERNEBEEK G, AALBERSE RC, DIEGES PH. Subclass typing of IgG antibodies formed by grass pollen-allergic patients during immunotherapy. *Int Arch Allergy Appl Immunol* 1976; 50: 625-40.
92. VAN DER ZEE JS, AALBERSE RC. The role of IgG in allergy. In: LESSOF MH, LEE TH, KEMENY DM, eds. Allergy: an international textbook. London: Wiley, 1987: 49-67.
93. VAN DER ZEE JS, AALBERSE RC. Immunochemical characteristics of IgG4 antibodies. *NER Allergy Proc* 1988; 9: 31-3.
94. VAN DER ZEE JS, VAN SWIETEN P, AALBERSE RC. Serologic aspects of IgG4 antibodies. II. IgG4 antibodies form small, nonprecipitating immune complexes due to functional monovalency. *J Immunol* 1986; 137: 3566-71.
95. VAN DER ZEE JS, VAN SWIETEN P, AALBERSE RC. Inhi-

96. VAN MILLIGEN FJ. The human antibody response to the cat allergen *Felis domesticus*. Enschede, The Netherlands: FEBO, 1993.
97. VAN MILLIGEN FJ, CRAIG S, ROGERS BL, VAN SWIETEN P, AALBERSE RC. Differences between specificities of IgE and IgG4 antibodies: studies using recombinant chain 1 and chain 2 of the major cat allergen *Fel d 1*. *J Allergy Clin Immunol* 1993 (in press).
98. VAN MILLIGEN FJ, VAN 'T HOF W, AALBERSE RC. IgE and IgG4 binding to synthetic peptides of the cat (*Felis domesticus*) major allergen *Fel d 1*. *Int Arch Allergy Immunol* 1993 (in press).
99. VAN MILLIGEN FJ, VAN SWIETEN P, AALBERSE RC. IgE and IgG4 to phylogenetic variants of the cat (*Felis domesticus*) major allergen *Fel d 1*. *Allergy* 1993 (in press).
100. VAN REE R, STAPELSO, AALBERSE RC. Reverse-sandwich ELISA may underestimate antibody titers. *J Allergy Clin Immunol* 1989; 84: 562-3.
101. VAN TOORENENBERGEN AW, AALBERSE RC. IgG4 and passive sensitization of basophil leukocytes. *Int Arch Allergy Appl Immunol* 1981; 65: 432-40.
102. VAN TOORENENBERGEN AW, AALBERSE RC. IgG4 and release of histamine from human peripheral blood leukocytes. *Int Arch Allergy Appl Immunol* 1982; 67: 117-22.
103. VAN TOORENENBERGEN AW, AALBERSE RC. Allergen-specific IgE and IgG4 antibody levels in sera and the capacity of these sera to sensitize basophil leukocytes *in vitro*. *Clin Allergy* 1982; 12: 451-8.
104. VIJAY HM, PERELMUTTER L. Inhibition of reagin-mediated PCA-reactions in monkeys and histamine release from human leukocytes by human IgG4 subclass. *Int Arch Allergy Appl Immunol* 1977; 53: 78-87.
105. WAMAE CN, ROBERTS JM, EBERHARD ML, LAMMIE PJ. Kinetics of circulating human IgG4 after diethylcarbamazine and ivermectin treatment of bancroftian filariasis. *J Infect Dis* 1992; 165: 1158-60.
106. WÜTHRICH B, BENZ A, SKVARIL F. IgE and IgG4 levels in children with atopic dermatitis. *Dermatologica* 1983; 166: 229-35.

This document is a scanned copy of a printed document. No warranty is given about the accuracy of the copy. Users should refer to the original published version of the material.