



Morphology changes in human lung epithelial cells after exposure to diesel exhaust micron sub particles (PM_{1.0}) and pollen allergens

V. Esposito^a, A. Lucariello^a, L. Savarese^a, M.P. Cinelli^b, F. Ferraraccio^c, A. Bianco^d, A. De Luca^a, G. Mazzearella^{e,*}

^a Department of Medicine and Public Health, Section of Human Anatomy, Second University of Naples, Italy

^b Department of Biomorphological and Functional Sciences, University of Naples Federico II, Naples, Italy

^c Department of Medicine and Public Health, Section of Pathologic Anatomy, Second University of Naples, Italy

^d Department of Health Sciences, University of Molise, Faculty of Medicine and Surgery, Italy

^e Department of CardioThoracic and Respiratory Sciences, Second University of Naples, 80138 Naples, Italy

ARTICLE INFO

Article history:

Received 22 March 2012

Received in revised form

27 June 2012

Accepted 1 July 2012

Keywords:

Asthma

Sub-micron PM particles

TEM

ABSTRACT

In the recent literature there has been an increased interest in the effects of particulate matter on the respiratory tract. The objective of this study was to use an in vitro model of type II lung epithelium (A549) to evaluate the cell ability to take up sub-micron PM_{1.0} particles (PM_{1.0}), *Parietaria officinalis* (ALL), and PM_{1.0} + ALL together.

Morphological analysis performed by Transmission Electron Microscope (TEM) showed that PM and ALL interacted with the cell surface, then penetrating into the cytoplasm. Each single treatment was able to point out a specific change in the morphology. The cells treated appear healthy and not apoptotic. The main effect was the increase of: multilamellar bodies, lysosomal enzymes, microvilli, and presence of vesicle/vacuoles containing particles.

These observations demonstrate morphological and functional alterations related to the PM_{1.0} and *P. officinalis* and confirm the induction of the inflammatory response in lung cells exposed to the inhalable particles.

© 2012 Published by Elsevier Ltd.

1. Introduction

In the recent literature there has been an increased interest in the effects of particulate matter on the respiratory tract (Anderson et al., 2011; Donaldson et al., 2005; Gehr et al., 2006; Kreyling et al., 2006; Mühlfeld et al., 2008). In particular epidemiological data outline a strict correlation between the exposure to particulate matter and several forms of pulmonary diseases (BruneKreef and Holgate, 2002; Knaapen et al., 2004; Turner et al., 2011).

Particulate matter is a mixture of organic and inorganic solid and liquid particles of different origin, size, and composition. Penetration of the tracheobronchial tract is related to particle size and the efficiency of airway defense mechanisms (Mazzearella et al., 2012).

Particulates can be classified according to their composition and size. Epidemiological study has demonstrated that only smaller size of particulate, commonly indicated as “fine particulate”, can reach

the lower airways and can play a role as major contributor to adverse health effects.

It has been demonstrated that urban fine particulate can penetrate deep into the airways and induce alveolar inflammation, which is responsible for variation in blood coagulability and release of mediators favoring acute episodes of respiratory and cardiovascular diseases (Mazzearella et al., 2012, 1998; Catena et al., 1993).

Particulate matter, according to the criteria of particle size, is traditionally classified as coarse, fine and ultrafine (Gualtieri et al., 2009). Particles with a diameter ranging from 2.5 µm to 10 µm belong to coarse PM, commonly known as PM_{10–2.5}; particles with a diameter less than 2.5 µm belong to the fine fraction, named PM_{2.5}; instead particles with a diameter less than 100 nm belong to the ultrafine fraction indicated as PM_{0.1} (Annesi-Maesano et al., 2007; Gualtieri et al., 2009; Mazzearella et al., 2007).

Diesel exhaust particles (DEPs) represent the greatest part of particulate (up 90%) given the increase in the number of new cars with diesel engines in industrialized countries (Diaz Sanchez et al., 1997; Mazzearella et al., 2007).

People that live in big cities are usually exposed to higher levels of vehicle emissions and several other forms of pollutants (due to

* Corresponding author.

E-mail address: gennaro.mazzearella@unina2.it (G. Mazzearella).

the economic and industrial growth) that mixed with aero-allergens, such as those derived from pollen grains, help to facilitate allergic sensitization of the airways and lead to bronchial obstruction in predisposed individuals (D'Amato et al., 2010; Gilmour et al., 2006; Cazzola et al., 1999).

Pollen particles, in high environmental humidity conditions, can be subjected to osmotic shock, resulting in the release of micro-particles or paucimicronic particles that may contain allergenic proteins (Suphioglu et al., 1992; Taketomi et al., 2006).

In the same way, through physical contact with the pollen particles, diesel exhaust particles can disrupt the former, leading to the release of paucimicronic particles and transporting them by air thus facilitating their penetration of the human airways (Churg and Brauer, 1997).

The interaction among cells, pollen particles and particulate has been deeply studied and several biochemical effects have been described upon size and quantity used. However few informations are present about morphological events following these treatments.

The objective of this study was to use an in vitro model of type II lung epithelium (A549) to evaluate the cell ability to take up sub-micron PM_{1.0} particles (PM_{1.0}), *Parietaria officinalis* (ALL), and PM_{1.0} + ALL together.

In the present study we used particles with a size of the 1.0 µm or less at a concentration of 0.1 mg/mL derived from filters of diesel engine exhausts. The concentration used is 1/10 compared to that which causes chronic inflammation of the airways.

The principal interest was to investigate if type II lung epithelium cells undergo morphological changes after treatment; in particular we analyzed the morphological distribution and the quantitative incorporation using a transmission electron microscope (TEM).

2. Materials and methods

2.1. Particulate sampling

Particulate matter was collected at the exhaust of a primary brand automobile equipped with a 2000 cm³ turbo-diesel engine, compliant to Euro IV emission regulations. Particulate matter was obtained from Diesel exhaust particles as described previously (Mazzarella et al., 2007). Briefly, Diesel PM is sampled by filtering air-diluted diesel car exhaust fumes at temperatures not higher than 52 °C. A dilution tunnel connected with a NOVA-MMB Messtechnik PDP-CVS unit has been used to produce the air/diesel combustion gases mixture; this configuration cleverly resembles conditions at which diesel exhaust PM is released from vehicles into the atmosphere.

Furthermore, in order to obtain PM_{1.0} the particulate sampling filters were conditioned, then weighted; after sampling, filters were weighed again, and 5.5 mg of DEP-PM were placed in a beaker containing HPLC-grade Et–OH (Aldrich CHROMASOLV®). The suspension so obtained was sonication-treated for 30 min on a Misonix XL-2020 device, at a frequency of 20 kHz. Most Et–OH was subsequently removed from sample using a vacuum rotary evaporator, in order to get stock samples with PM concentrated to 900 mg/mL. Aliquots so prepared were stored at 2 °C upon usage.

2.2. Epithelial cell culture

We used the A549 cell line and it was cultured as described previously (Mazzarella et al., 2007). Briefly, A549 cells were cultured in RPMI-1640 (Sigma, Poole, UK) medium supplemented with 10% fetal bovine serum (Sigma) and 10,000 units/mL penicillin, 10 mg/mL streptomycin and 25 mg/mL amphotericin B (Sigma), in humidified atmosphere of 5% CO₂, 95% air at 37.5 °C to confluence. At this stage, A549 monolayers were gently detached using ready prepared Trypsin solution (Sigma), cells were rinsed and pelleted; pellets were resuspended in serum-free RPMI-1640 medium only supplemented with antimicrobial agents and plated onto "Multi-G-Lass" chambers (Iwaki Asahi Techno Glass); incubation took place at 36.5 °C in humidified atmosphere containing 5% CO₂ for 48 h, where cells normally reached semiconfluence.

At this time (our "starting time") PM_{1.0} (0.1 mg/mL) and pollen extract *P. officinalis* (0.5 DBU/mL) (Diagnostic Biological Units by Lofarma). One DBU is one hundredth of the potency of one extract inducing a wheal equal to that induced by histamine chloride 10 mg/mL, were added, separately and in combination, to cell cultures.

2.3. Transmission Electron Microscope – TEM

For the fine ultrastructural investigation, confluent cell monolayers were directly fixed on dishes with 2.5% glutaraldehyde in 0.2 M in sodium phosphate buffer (PH 7.2), and post- fixed with 1% OsO₄ in the same buffer for 2 h. After repeated washings, they were dehydrated in a graded ethanol series and embedded in EPON[®] 812. Frontal 60 nm ultrathin sections of the cell layers were cut on a LAIKA Ultracut UCT and they were routinely stained with 4% uranyl acetate for 15 min and 3% lead citrate for 20 s. The grids were observed under a Transmission Electron Microscope (TEM) LEO 912 AB.

3. Results

A549 cell line was treated with sub-micron PM_{1.0} particles (PM_{1.0}), *P. officinalis* (ALL), and PM_{1.0} + ALL together. Cells following this treatment were examined at 24, 48, and 72 h and compared with control cells.

Cells, after treatment, in culture flask were observed using an inverted microscope. A morphological comparison among the different points shows that untreated cells appeared moderately large and exhibited different kinds of shapes. Most of the cells appear separated from one another nicely and neatly with a shape often hexagonal or spindle-like. Instead treated cells appear with clearly atypical nuclei and moderately big. The cytoplasmic compartment showed granular inclusions. Furthermore we investigated the different cell treatment with Transmission Electron Microscope (TEM).

A549 cells untreated show a rounded shape, with several distributed microvilli on the cell surface. Within the cytoplasm, the nucleus has an irregular shape. Several mitochondria are visible: they have a roundish/oval shape and are uniformly distributed in the cytoplasm. Granules with a low electrodensity, probably of a lipidic type, are present. There are also numerous multilamellar bodies with a low electrodensity, but that, at higher magnification, show the presence of characteristic thin parallel lamellae (Fig. 1).

A549 cells treated with PM_{1.0} at TEM observation showed a round shape. The microvilli, distributed on the cell surface are increased compared to control cells, and these allow the increase of the cell surface. In the cytoplasm, the nucleus is irregularly shaped. Mitochondria, of a round/oval shape, are numerous and equally distributed in the cytoplasm. Low electrodensity granules, probably of a lipidic nature are present in numbers similar to control cells. The cells are characterized by a cytoplasm particularly rich in multilamellar bodies compared to control cells. In addition, these multilamellar bodies show a higher electrodensity probably due to the presence of lysosomal enzymes. In the cytoplasm of cells at 72 h after treatment it is possible to highlight the presence of some vesicles containing small particles. These probably are particles that have been phagocytized/endocytosed by epithelial cells (Fig. 2).

A549 cells treated with ALL show a rounded shape, with a greater number of microvilli distributed on the cell surface compared to control cells. In the cytoplasm, the nucleus is irregularly shaped. Mitochondria are numerous of round/oval shape, equally distributed in the cytoplasm. Granules of low electrodensity, probably of lipidic kind, are present in numbers similar to control cells and to those treated with the particulate.

In these cells, it is possible to highlight a significant increase in multilamellar bodies, as compared with both the control cells and those treated with the particulate only. Even multilamellar bodies with higher electrodensity appear to be increased in number compared both to control cells and to cells treated with the particulate. There are also numerous vesicles containing small particles (probably allergen particles that have been phagocytized). These vesicles are markedly increased compared to cells treated with the particulate (Fig. 3).




Fig. 1. Conventional TEM of A549 cells untreated. M: microvilli; N: nucleus; G: granules; MB: multilamellar bodies; MT: mitochondria.

A549 cells treated with PM_{1.0} + ALL together show a rounded shape. The microvilli are reduced in number compared to cells treated individually with the particulate or the allergen. The nucleus is irregularly shaped. Mitochondria are numerous and round/oval shaped, equally distributed in the cytoplasm. The granules with low electrodensity, probably of lipidic nature are present in greater numbers as compared both with the control cells and those treated with the particulate only or the allergene. There are numerous multilamellar bodies, which are present in greater numbers than control cells and those treated with the particulate only. It is possible to highlight inside the cytoplasm a further

increase of multilamellar bodies with a higher electrodensity and vesicles containing small particles as compared with both the control cells and those treated with the particulate or the allergene (Fig. 4).

4. Discussion

Over the past years several biochemical effects have been described after the treatment of lung cells with several particles and nanoparticles; however few informations are present at today about the morphological changes following these treatments.




Fig. 2. A: untreated Cell; B, C, D: cells treated for 72 h with PM. M: microvilli; N: nucleus; G: granules; MB: multilamellar bodies; V: vesicles with small particles; MT: mitochondria.




Fig. 3. A: untreated Cell; B, C, D: cells treated for 72 h with ALL. M: microvilli; N: nucleus; MB: multilamellar bodies; V: vesicles with small particles; MT: mitochondria.

The principal interest of our study was to investigate if type II lung epithelium cells undergo morphological changes after treatment with sub-micron $PM_{1.0}$ particles ($PM_{1.0}$), *P. officinalis* (ALL), and $PM_{1.0}$ + ALL together. In particular we analyzed the morphological distribution and the quantitative incorporation of this particles using a Transmission Electron Microscope (TEM). We used A549 cells because it is a widely used and well characterized cell culture line which shares characteristics with alveolar epithelial cells.

The morphological results obtained in the study point out that with the concentration used in our model we do not observe any apoptotic event.

A549 cells treated with $PM_{1.0}$ at TEM observation showed that the vesicles containing the particulate matter are much more evident at 72 h compared with 48 h. This increase of vesicles is probably due to the fact that the cells at 48 h have not yet had the time or the way to incorporate the PM.

A549 cells treated with $PM_{1.0}$ and ALL showed an increase of microvilli, distributed on the cell surface compared to control cells, and these allow the enlarge of the cell surface. As previously described this phenomenon is due to an altered cellular lipidic metabolism favoring lipid trafficking to the membrane after exposure to high particle concentrations (Vlahakis et al., 2001). The interaction between particles and membrane interferes with endocytosis and exocytosis leading to an increase in membrane surface. This may protect the epithelial cells against injury and helps to reseal plasma membrane injuries (Borgonovo et al., 2002).

Interestingly the microvilli are reduced in number compared to cells treated individually with the particulate or the allergen.

On the contrary granules with low electrodensity, probably of lipidic nature are present in greater numbers as compared both with the control cells and those treated with the particulate only or the allergen. The hypothesis could be that these granules are lipid reserve and could be used to make multilamellar bodies.

In all cells treated, it is possible to highlight a significant increase in multilamellar bodies, as compared with the control cells. In particular the multilamellar bodies are increased in the treatment with the ALL compared with the cells treated with the PM. The major increase is obtained in the cells treated with $PM_{1.0}$ + ALL together.

The multilamellar bodies are those producing the surfactant that has a function to lower the surface tension at the interface. Probably the increase of multilamellar bodies is used by the cells to increase the production of surfactant as defence against the external agent. Morphologically the multilamellar bodies appear with a higher electrodensity in the cells treated with ALL compared with the one treated with $PM_{1.0}$. This higher electrodensity is due to the presence of vesicles that probably contain lysosomal enzymes. Their increase is probably a defence reaction. The cells react to the allergen phagocytising and destroying it as defensive reaction. We found this kind of defence also in samples treated with $PM_{1.0}$, even if with a minor extent and only at 72 h.

In summary the cells treated appear healthy and not apoptotic. The main effect is the increase of multilamellar bodies and the relative surfactant. Moreover it is possible to observe an increase of lysosomal enzymes which probably are involved in digestion of foreign bodies. Finally it is important to underline also the increase of microvilli that allow the enhancement of the cell surface and the presence of vesicle/vacuoles containing particles, which indicates the ability of epithelial cells to phagocytose foreign bodies.




Fig. 4. A: untreated Cell; B, C, D: cells treated for 72 h PM + ALL. M: microvilli; N: nucleus; G: granules; MB: multilamellar bodies; V: vesicles with small particles; MT: mitochondria.

As previously described, we showed that lung epithelial cells (A549) can also participate in the release of inflammatory cytokines (IL-4, IL-5 and IL-13) in response to challenge with PM_{1.0} and *P. officinalis* together supporting the potential of lung epithelial cells to participate in inflammation and disease. The interaction between diesel particulates and pollen is capable of forming micro compounds that can reach the acinar lung units and activate cells of the immune system. This activation may depend on the interaction between susceptibility genes and environmental factors that trigger the onset of allergic disease (Mazzarella et al., 2012). Also other authors have shown that the fine urban air particulate interferes with epithelial cell physiology by reducing the proliferation rate and increasing the release of proinflammatory cytokines typically secreted by epithelial cells, tumor necrosis factor- α , and IL-6, IL-8 a multifunctional cytokine with immunoregulatory and inflammatory effects (Pozzi et al., 2003; Singh et al., 2007).

The observations reported demonstrate morphological and functional alterations related to the PM_{1.0} and *P. officinalis* and confirm the induction of the inflammatory response in lung cells exposed to the inhalable particles.

Acknowledgments

The study was partially supported by Second University of Naples.

References

Anderson, J.O., Thundiyil, J.G., Stolbach, A., 2011. Clearing the air: a review of the effects of particulate matter air pollution on human health. *J. Med. Toxicol.* 8, 165–166.

- Annesi-Maesano, I., Moreau, D., Caillaud, D., et al., 2007. Residential proximity fine particles related to allergic sensitization and asthma in primary school children. *Respir. Med.* 101, 1721–1729.
- Borgonovo, B., Cocucci, E., Racchetti, G., et al., 2002. Regulated exocytosis: a novel, widely expressed system. *Nat. Cell. Biol.* 4, 955–962.
- Brunekef, B., Holgate, S.T., 2002. Air pollution and health. *Lancet* 360, 1233–1242.
- Catena, E., Mazzarella, G., Peluso, G.F., et al., 1993. Phenotypic features and secretory pattern of alveolar macrophages in atopic asthmatic patients. *Monaldi Arch. Chest Dis.* 48, 6–15.
- Cazzola, M., Matera, M.G., Di Perna, F., et al., 1999. Influence of higher than conventional doses of oxitropium bromide on formoterol-induced bronchodilation in COPD. *Respir. Med.* 93, 909–911.
- Churg, A., Brauer, M., 1997. Human lung parenchyma retains PM_{2.5}. *Am. J. Resp Crit. Care Med.* 155, 2109–2111.
- D'Amato, G., Cecchi, L., D'Amato, M., Liccardi, G., 2010. Urban air pollution and climate changes as environmental risk factors of respiratory allergy: an update. *J. Invest. Allergol. Clin. Immunol.* 20, 95–102.
- Diaz Sanchez, D., Tsien, A., Fleming, J., Saxon, A., 1997. Combined diesel exhaust particulate and ragweed allergen challenge markedly enhances human in vivo nasal ragweed specific IgE and skews cytokine production to a T helper cell 2-type pattern. *J. Immunol.* 158, 2406–2413.
- Donaldson, K., Tran, L., Jimenez, L.A., et al., 2005. Combustion-derived nanoparticles: a review of their toxicology following inhalation exposure. *Part Fibre Toxicol.* 2, 10.
- Gehr, P., Blank, F., Rothen-Rutishauser, B.M., 2006. Fate of inhaled particles after interaction with the lung surface. *Paediatr. Respir. Rev.* 7, 73–75.
- Gilmour, M.I., Jaakkola, M.S., London, S.J., Nel, A.E., Rogers, C.A., 2006. How exposure to environmental tobacco smoke, outdoor air pollutants, and increased pollen burdens influences the incidence of asthma. *Environ. Health Perspect.* 114, 627–633.
- Gualtieri, M., Mantecchia, P., Corvaja, V., et al., 2009. Winter fine particulate matter from Milan induces morphological and functional alterations in human pulmonary epithelial cells (A549). *Toxicol. Lett.* 188, 52–62.
- Knaapen, A.M., Borm, P.J., Albrecht, C., Schins, R.P., 2004. Inhaled particles and lung cancer. Part A. Mechanism. *Int. J. Cancer* 109, 799–809.
- Kreyling, W.G., Semmler-Behnke, M., Möller, W., 2006. Ultrafine particle lung interactions: does size matter? *J. Aerosol. Med.* 19, 74–83.
- Mazzarella, G., Esposito, V., Bianco, A., et al., 2012. Inflammatory effects on human lung epithelial cells after exposure to diesel exhaust micron sub particles (PM_{1.0}) and pollen allergens. *Env. Pollut.* 161, 64–69.

- Mazzarella, G., Ferraraccio, F., Prati, M.V., et al., 2007. Effect of diesel exhaust particles on human lung epithelial cells: an in vitro study. *Respir. Med.* 101, 1155–1162.
- Mazzarella, G., Petillo, O., Margarucci, S., Calabrese, C., Peluso, G., 1998. Role of monocyte/macrophage population in immune response. *Monaldi Arch. Chest Dis.* 53, 92–96.
- Mühlfeld, C., Rothen-Rutishauser, B., Blank, F., et al., 2008. Interactions of nanoparticles with pulmonary structures and cellular responses. *Am. J. Physiol. Lung Cell. Mol. Physiol.* 294, 817–829.
- Pozzi, R., De Bernardis, B., Paoletti, L., Guastadisegni, C., 2003. Inflammatory mediators induced by coarse (PM_{2.5–10}) and fine (PM-2.5) urban air particles in RAW 264.7 cells. *Toxicology* 183, 243–254.
- Singh, S., Shi, T., Duffin, R., et al., 2007. Endocytosis, oxidative stress and IL-8 expression in human lung epithelial cells upon treatment with fine and ultrafine TiO₂: role of the specific surface area and of surface methylation of the particles. *Toxicol. Appl. Pharmacol.* 222, 141–151.
- Suphioglu, C., Singh, M.B., Taylor, P., Knox, R.B., 1992. Mechanism of grass pollen-induced asthma. *Lancet* 339, 569–572.
- Taketomi, E.A., Sopelete, M.C., de Sousa Moreira, P.F., et al., 2006. Pollen allergic disease: pollens and its major allergens. *Braz. J. Otorhinolaryngol.* 72, 569–572.
- Turner, M.C., Krewski, D., Pope 3rd, C.A., et al., 2011. Long-term ambient fine particulate matter air pollution and lung cancer in a large cohort of never-smokers. *Am. J. Respir. Crit. Care Med.* 184, 1374–1381.
- Vlahakis, N.E., Schroeder, M.A., Pagano, R.E., Hubmayr, R.D., 2001. Deformation-induced lipid trafficking in alveolar epithelial cells. *Am. J. Physiol. Lung Cell. Mol. Physiol.* 280, 938–946.