

Threshold and safe ingestion dose among infants sensitized to hen's egg

To the Editor,

Generally, hen's egg is recommended to be introduced in infancy.¹ However, infants who have food allergies (FA) other than egg allergy (EA) or eczema are sometimes found to be sensitized to egg before introduction of egg, and they are often advised to completely avoid egg consumption. As some infants do not have true EA, evaluating whether they can ingest eggs is crucial. To date, no method has been established for initiating egg consumption in sensitized infants. Unlike oral food challenges (OFCs) described in the PRACTALL guidelines, the 2020 Japanese guidelines for food allergy recommend stepwise OFCs for the diagnosis and management of FA.² Stepwise OFCs, which start with a low-dose OFC, and in which the next step would be performed after approximately 3 months if one step was negative, are useful both for diagnosis and to determine the amount of causative food that can be safely consumed. We have previously reported that stepwise OFCs are useful and safe for patients suspected with EA.^{3,4} However, most patients in the study had a history of immediate reaction to eggs and were aged >1 year. There have been no reports regarding stepwise OFCs in infants without a history of previous consumption of egg who were instructed to completely avoid egg consumption owing to sensitization (egg-sensitized infants). This study aimed to evaluate the usefulness of stepwise OFCs in egg-sensitized infants for introducing egg to determine the threshold and safe ingestion dose of egg.

We retrospectively examined egg-sensitized infants who underwent stepwise OFCs between May 2016 and December 2018. Egg-sensitization was found in these infants because of blood tests performed to evaluate eczema or FA other than EA. Stepwise OFCs were performed starting with a low dose (containing 1/25 of a heated whole egg [HE], 250mg of egg protein). After the low-dose OFC, we performed an OFC containing 1/8 of a HE (775–769mg of egg protein, medium-dose OFC) and later 1/2 of a HE (3076–3100mg of egg protein, high-dose OFC) if the results of each previous OFC were negative. After having passed each OFC, the patients were instructed to consume foods at home that contained the same amounts of egg protein as they had consumed in the OFC. Patients who passed the high-dose OFC received an OFC with scrambled egg containing a whole egg (6591.5 mg of egg protein, full-dose OFC) or were instructed to increase egg consumption up to one HE at home. If the result of the low-dose OFC was positive, the patients were advised to completely avoid egg consumption. The patients who failed medium-, high-, or

full-dose OFCs were advised to consume the amount of egg that they could safely ingest (e.g., a seasoned powder, bread or processed meat, and a donut or slice of cake for positive medium-, high-, and full-dose OFC patients, respectively, Figure S1). The challenge foods used in this study are shown in Table S1. Serum-specific immunoglobulin E to egg white (EW-sIgE) and ovomucoid (OVM-sIgE) levels were measured within 3 months of the low-dose OFC. Sensitization to egg was defined as an EW-sIgE level >0.10 kU_A/L.

Of the 148 egg-sensitized infants who underwent low-dose OFC, 61 (41%) were excluded because they did not meet the eligibility criteria (Figure 1). The median OVM-sIgE level of the patients included in this study was significantly higher than that for the excluded ones (Table S2). The median age was 9.9 months, and the median EW-sIgE and OVM-sIgE levels were 18.0 and 0.99 kU_A/L, respectively.

Among the 87 egg-sensitized infants, 12 (14%) failed the low-dose OFC. Nine (12%) of 75 who passed the low-dose OFC failed the medium-dose OFC, and 8 (12%) of 66 who passed the medium-dose OFC failed the high-dose OFC. Of the 58 patients who passed the high-dose OFC, 32 underwent full-dose OFC and 26 were instructed to increase egg consumption at home. Two patients failed the full-dose OFC, and 56 were able to ingest a full dose (Figure 1). Thus, 64% (56/87) of egg-sensitized infants did not have an allergic reaction to egg. Of the 31 patients allergic to egg, the threshold dose was ≤1/25 of a HE for 39% (12/31), >1/25–1/8 for 29% (9/31), >1/8–1/2 for 26% (8/31), and >1/2–1 for 6% (2/31) (Figure 2). Therefore, 14% (12/87) of egg-sensitized infants needed to continue to completely avoid egg consumption. The median time between the low- and full-dose OFC was 10.8 months. The median time between the low-dose OFC and the date when the patients were able to ingest a full dose was 16.1 months. Probability curves based on the results of these OFCs are shown in Figure 3. The probability curves show that >50% of egg-sensitized infants would pass a full-dose OFC even if their EW-sIgE levels exceeded 100kU_A/L. In this study, six patients had EW-sIgE levels over 100kU_A/L, and one of them also had an OVM-sIgE level exceeding 100kU_A/L. Of these six patients, five were able to ingest one scrambled egg. The symptoms and treatment during OFCs are described in Table S3. There were no severe symptoms in the medium- and full-dose OFCs. However, one patient received intramuscular adrenaline during the low-dose OFC because of vomiting and appearance of a pale face. In the high-dose OFC, one patient developed a severe reaction (barking cough).

This study revealed that the rate of positive reactions to stepwise OFCs for egg-sensitized infants was not high, and stepwise OFCs clarified the threshold and safe ingestion dose of eggs in these infants. Majority of egg-sensitized infants were able to ingest one scrambled egg, even if they have relatively high levels of EW-sIgE.

Regarding the threshold dose in egg-sensitized infants, a previous study reported that 33% reacted to a boiled egg white OFC equivalent to one half of an egg.⁵ In our study, the proportion of egg-sensitized infants who reacted to $\leq 1/2$ of a HE was similar (33%, 29/87). However, by performing stepwise OFCs, 59% (17/29) could ingest $\leq 1/8$ of a HE. For patients with EA, avoiding complete elimination of eggs may improve their prognosis^{6,7} and quality of life (QOL).⁸ Therefore, stepwise OFCs appear to be beneficial for egg-sensitized infants.

This study had some limitations. First, OFCs were not double-blind placebo-controlled. However, as most symptoms induced during OFCs were objective, the number of false-positive OFCs was considered to be low. Second, a few patients may have achieved tolerance during intervals between OFCs. However, as only 30% of Japanese EA patients

Key messages

This study showed that only a small number of the infants for whom egg consumption was completely avoided owing to sensitization needed complete avoidance of eggs, and the majority of them were able to ingest a whole egg in the form of scrambled egg. Stepwise oral food challenges starting from a low dose are relatively safe to introduce egg and effective in determining the threshold and safe ingestion dose for such infants.

achieve tolerance by 3 years of age,⁹ the influence of intervals between OFCs is likely minimal. Third, approximately 40% of the patients were excluded because of not undergoing stepwise OFCs, and their median OVM-sIgE level was significantly lower than for the included ones. Therefore, our study may have overestimated the rate of positive reactions to stepwise OFCs for egg-sensitized infants.




FIGURE 1 Enrollment of subjects. Stepwise oral food challenges (low-dose, medium-dose, high-dose, and full-dose) performed. After passing the high-dose OFC, 32 patients underwent full-dose OFC and 26 were instructed to increase egg consumption at home. OFC, oral food challenge

FIGURE 2 Percentage of egg-sensitized patients who had an allergic reaction to egg (A) and the threshold dose of egg in these patients (B)





FIGURE 3 Probability curve for failing the stepwise oral food challenge (OFC) at a given sIgE level against egg white (A) and ovomucoid (B). The probability curves were constructed using results of multinomial logistic regression analysis. OFC, oral food challenge; sIgE, specific IgE

In conclusion, stepwise OFCs appeared to be relatively safe for the introduction of egg in egg-sensitized infants. We found that only a small number of these infants required complete avoidance of eggs and that the majority of them were able to ingest one scrambled egg. We expect that stepwise OFCs will help improve the management of egg-sensitized infants who have completely avoided egg consumption.

AUTHOR CONTRIBUTIONS

Masatoshi Mitomori: Conceptualization (equal); data curation (lead); formal analysis (lead); investigation (lead); methodology (lead); project administration (lead); resources (lead); software (equal); supervision (equal); validation (supporting); visualization (supporting); writing – original draft (lead); writing – review and editing (equal). **Makoto Nishino:** Conceptualization (supporting); data curation (supporting);

formal analysis (supporting); investigation (equal); methodology (supporting); project administration (supporting); resources (supporting); software (supporting); supervision (supporting); validation (supporting); visualization (supporting); writing – original draft (lead); writing – review and editing (equal). **Kyohei Takahashi:** Conceptualization (supporting); data curation (supporting); formal analysis (supporting); investigation (equal); methodology (supporting); project administration (supporting); resources (supporting); software (supporting); supervision (supporting); validation (supporting); visualization (supporting); writing – original draft (lead); writing – review and editing (equal). **Ken-ichi Nagakura:** Conceptualization (supporting); data curation (supporting); formal analysis (supporting); investigation (equal); methodology (supporting); project administration (supporting); resources (supporting); software (supporting); supervision (supporting); validation (supporting); visualization (supporting); writing

– original draft (lead); writing – review and editing (equal). **Kiyotake Ogura:** Conceptualization (supporting); data curation (supporting); formal analysis (supporting); investigation (equal); methodology (supporting); project administration (supporting); resources (supporting); software (supporting); supervision (supporting); validation (supporting); visualization (supporting); writing – original draft (lead); writing – review and editing (equal). **Mari Takei:** Conceptualization (supporting); data curation (supporting); formal analysis (supporting); investigation (equal); methodology (supporting); project administration (supporting); resources (supporting); software (supporting); supervision (supporting); validation (supporting); visualization (supporting); writing – original draft (supporting); writing – review and editing (supporting). **Sakura Sato:** Conceptualization (lead); data curation (supporting); formal analysis (supporting); investigation (equal); methodology (equal); project administration (equal); resources (equal); software (equal); supervision (lead); validation (lead); visualization (lead); writing – original draft (equal); writing – review and editing (lead). **Motohiro Ebisawa:** Conceptualization (lead); data curation (supporting); formal analysis (supporting); investigation (equal); methodology (equal); project administration (equal); resources (equal); software (equal); supervision (lead); validation (lead); visualization (equal); writing – original draft (equal); writing – review and editing (lead).

ACKNOWLEDGMENTS

We would like to thank Editage (www.editage.jp) for their assistance with English Language editing.

KEYWORDS

egg, food allergy, oral food challenge, pediatric, sensitization, stepwise, threshold

FUNDING INFORMATION

This study was supported by funds from Nipponham Foundation for the Future of Food, Japanese Society of Allergology, Clinical Research Support Program and JSPS KAKENHI grant number (16K00899).

CONFLICT OF INTEREST

Motohiro Ebisawa and Sato Sakura has received lecture fees from Mylan EPD. The other authors declare that they have no conflicts of interest.

PEER REVIEW

The peer review history for this article is available at <https://publons.com/publon/10.1111/pai.13830>.

ETHICAL APPROVAL AND TRIAL REGISTRATION STATEMENTS

This study was conducted in accordance with the Declaration of Helsinki and was approved by the ethics committee of Sagami National Hospital (approval number: 2013–20,130,318,017).

Written informed consent was obtained from the guardians of all patients before each oral food challenge.

Masatoshi Mitomori¹
 Noriyuki Yanagida^{1,2} 
 Makoto Nishino¹ 
 Kyohei Takahashi¹ 
 Ken-ichi Nagakura¹ 
 Kiyotake Ogura¹
 Mari Takei²
 Sakura Sato² 
 Motohiro Ebisawa² 

¹Department of Pediatrics, National Hospital Organization, Sagami National Hospital, Kanagawa, Japan

²Clinical Research Center for Allergy and Rheumatology, National Hospital Organization, Sagami National Hospital, Kanagawa, Japan

Correspondence

Noriyuki Yanagida, Clinical Research Center for Allergy and Rheumatology, National Hospital Organization Sagami National Hospital, 18-1, Sakuradai, Minami-ku, Sagami, Kanagawa 252-0392, Japan.

Email: [yana@foodallergy.jp](mailto: yana@foodallergy.jp)

Editor: Alexandra Santos

ORCID

Noriyuki Yanagida  <https://orcid.org/0000-0001-9643-744X>
 Makoto Nishino  <https://orcid.org/0000-0002-5381-8848>
 Kyohei Takahashi  <https://orcid.org/0000-0003-1220-7386>
 Ken-ichi Nagakura  <https://orcid.org/0000-0002-9381-9044>
 Sakura Sato  <https://orcid.org/0000-0003-3674-0759>
 Motohiro Ebisawa  <https://orcid.org/0000-0003-4117-558X>

REFERENCES

1. Halken S, Muraro A, de Silva D, et al. EAACI guideline: preventing the development of food allergy in infants and young children (2020 update). *Pediatr Allergy Immunol*. 2021;32:843-858.
2. Ebisawa M, Ito K, Fujisawa T, Committee for Japanese Pediatric Guideline for food allergy, the Japanese Society of Pediatric Allergy and Clinical Immunology, Japanese Society of Allergology. Japanese guidelines for food allergy 2020. *Allergol Int*. 2020;69:370-386.
3. Yanagida N, Sato S, Takahashi K, et al. Stepwise single-dose oral egg challenge: a multicenter prospective study. *J Allergy Clin Immunol Pract*. 2019;7:716-718.e6.
4. Yanagida N, Minoura T, Kitaoka S, Ebisawa M. A three-level stepwise oral food challenge for egg, milk, and wheat allergy. *J Allergy Clin Immunol Pract*. 2018;6:658-660.e10.
5. Haneda Y, Kando N, Yasui M, et al. Ovomucoids IgE is a better marker than egg white-specific IgE to diagnose boiled egg allergy. *J Allergy Clin Immunol*. 2012;129:1681-1682.

6. Peters RL, Dharmage SC, Gurrin LC, et al. The natural history and clinical predictors of egg allergy in the first 2 years of life: a prospective, population-based cohort study. *J Allergy Clin Immunol*. 2014;133:485-491.
7. Okada Y, Yanagida N, Sato S, Ebisawa M. Heated egg yolk challenge predicts the natural course of hen's egg allergy: a retrospective study. *World Allergy Organ J*. 2016;9:31.
8. Yanagida N, Sato S, Asaumi T, Ogura K, Borres MP, Ebisawa M. Safety and feasibility of heated egg yolk challenge for children with egg allergies. *Pediatr Allergy Immunol*. 2017;28:348-354.
9. Ohtani K, Sato S, Syukuya A, et al. Natural history of immediate-type hen's egg allergy in Japanese children. *Allergol Int*. 2016;65:153-157.

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.