

RESEARCH

Open Access



Having concomitant asthma phenotypes is common and independently relates to poor lung function in NHANES 2007–2012

Rita Amaral^{1,2*} , João A. Fonseca^{1,3,4}, Tiago Jacinto^{1,2,4}, Ana M. Pereira^{1,4}, Andrei Malinovski⁵, Christer Janson⁶ and Kjell Alving⁷

Abstract

Background: Evidence for distinct asthma phenotypes and their overlap is becoming increasingly relevant to identify personalized and targeted therapeutic strategies. In this study, we aimed to describe the overlap of five commonly reported asthma phenotypes in US adults with current asthma and assess its association with asthma outcomes.

Methods: Data from the National Health and Nutrition Examination Surveys (NHANES) 2007–2012 were used ($n = 30,442$). Adults with current asthma were selected. Asthma phenotypes were: B-Eos-high [if blood eosinophils (B-Eos) $\geq 300/\text{mm}^3$]; FeNO-high (FeNO ≥ 35 ppb); B-Eos&FeNO-low (B-Eos $< 150/\text{mm}^3$ and FeNO < 20 ppb); asthma with obesity (AwObesity) (BMI $\geq 30 \text{ kg/m}^2$); and asthma with concurrent COPD. Data were weighted for the US population and analyses were stratified by age (< 40 and ≥ 40 years old).

Results: Of the 18,619 adults included, 1059 (5.6% [95% CI 5.1–5.9]) had current asthma. A substantial overlap was observed both in subjects aged < 40 years (44%) and ≥ 40 years (54%). The more prevalent specific overlaps in both age groups were AwObesity associated with either B-Eos-high (15 and 12%, respectively) or B-Eos&FeNO-low asthma (13 and 11%, respectively). About 14% of the current asthma patients were “non-classified”. Regardless of phenotype classification, having concomitant phenotypes was significantly associated with (adjusted OR, 95% CI) ≥ 2 controller medications (2.03, 1.16–3.57), and FEV₁ $< \text{LLN}$ (3.21, 1.74–5.94), adjusted for confounding variables.

Conclusions: A prevalent overlap of commonly reported asthma phenotypes was observed among asthma patients from the general population, with implications for objective asthma outcomes. A broader approach may be required to better characterize asthma patients and prevent poor asthma outcomes.

Keywords: Asthma, Asthma-related outcomes, Epidemiological study, Overlap, Phenotypes

Background

Profiling asthma phenotypes is becoming increasingly relevant to choose the most appropriate therapeutic strategy for individual patients, and to provide optimal improvement of disease control and quality of life [1, 2].

The predominant pathophysiological mechanism of asthma is type 2-mediated, associated with atopy and eosinophilic inflammation [3, 4]. However, it has been

shown that asthma is a heterogeneous disease that involves other mechanisms that are not so well understood and respond poorly to corticosteroid therapy (e.g. non-type 2-mediated) [3, 5, 6].

There has been a recent rise in the number of studies that try to identify asthma phenotypes based on non-invasive type 2-markers, such as blood eosinophils (B-Eos) count, fraction of exhaled nitric oxide (FeNO), serum IgE, and/or serum periostin [7–10]. Moreover, there appears to be an additive role of biomarkers, such as B-Eos and FeNO, in relation to recent asthma morbidity [7, 11, 12].

*Correspondence: rita.samaral@gmail.com

¹ CINTESIS- Center for Health Technology and Services Research, Faculty of Medicine, University of Porto, Edifício Nascente, Piso 2, Rua Dr. Plácido da Costa, s/n, 4200-450 Porto, Portugal

Full list of author information is available at the end of the article

However, there is little information regarding the appropriate use of these biomarkers in asthma phenotype classification, particularly when a significant overlap occurs. Also, the importance of having concomitant asthma phenotypes for disease outcomes has scarcely been studied in the general population. This information may be useful to identify personalized and targeted therapeutic strategies [13, 14].

Recently, an extensive overlap of asthma phenotypes was described [15]. However, only type 2-high, atopic, and eosinophilic asthma were examined. The extent of overlap with other phenotypes commonly reported in the literature, among adults with asthma from the general population remains unknown. Asthma phenotypes are frequently reported in the literature according to the high levels of systemic and local type 2-markers (B-Eos high and FeNO-high, respectively) [1–4, 16–18]. However, other distinct subgroups of asthma phenotypes are increasingly being reported due to its characteristics of steroid therapy resistance and lack of inflammatory markers: e.g. subjects with asthma without evidence of type 2 inflammation (Th2-low phenotype); obese asthmatic subjects (obesity-related asthma phenotype); and patients with asthma-COPD overlap syndrome [19–22]. Therefore, we hypothesized that if, in general population, occurs a high proportion of overlap of commonly reported asthma phenotypes, there may be a need for improving the definition of asthma phenotypes. Additionally, asthma subjects with multiple phenotypes may have poorer asthma-related outcomes.

The aims of this study were to describe the proportion of overlap of five commonly reported asthma phenotypes: asthma with obesity (AwObesity), asthma with concurrent COPD (AwCOPD), B-Eos-high, FeNO-high and B-Eos&FeNO-low asthma, and to examine the association of their overlap with asthma-related outcomes, using population-based data from the National Health and Nutrition Examination Surveys (NHANES), 2007–2012.

Methods

Study design

The NHANES is a nationally representative survey of the civilian, non-institutionalized U.S. population that uses a complex stratified, multistage probability sampling. Further details on NHANES survey design databases can be found in Additional file 1: Supplementary methods. The National Center for Health Statistics, Ethics Review Board approved NHANES protocol, and all participants gave written informed consent.

Subjects selection

Six survey years (NHANES 2007–2012) were analyzed, resulting in 30,442 individuals of all ages (Fig. 1). We included adults (≥ 18 years-old) with current asthma ($n=1059$), defined by a positive answer to the questions: “Has a doctor ever told you that you have asthma?” together with “Do you still have asthma?”, and either “wheezing/whistling in the chest in the past 12 months” or “asthma attack in the past 12 months.”

Variables


Demographic characteristics, such as age, gender, body mass index (BMI), race/ethnicity, and educational status were analyzed. B-Eos count, FeNO and spirometric measurements, collected at the NHANES Mobile Examination Center were also examined. A detailed description of the procedures can be found elsewhere [23–25]. FeNO and spirometric measurements not fulfilling ATS/ERS recommendations [26, 27] were excluded ($n=653$). After predicted values of basal FEV₁ and FEV₁/FVC were calculated [28], with a correction factor for ethnicity [29], abnormal lung function was defined if either one of them were less than the lower limit of normal (LLN), defined as lower fifth percentile of the reference population [30].

Prescription medications used last month were also analyzed [31]. More details regarding the inclusion of reliever and controller medications for asthma and the definitions of each asthma-related variable included in the analysis (asthma attack, asthma-related emergency department (ED) visit, work/school absenteeism, asthma symptoms, smoking status and rhinitis) are provided in the supplementary material (see Additional file 1: Supplementary methods).

Asthma phenotypes definition

A B-Eos count $\geq 300/\text{mm}^3$ was used to define an B-Eos-high asthma phenotype [32, 33], while FeNO-high was defined as FeNO ≥ 35 ppb [34]. Asthma patients with both B-Eos $< 150/\text{mm}^3$ and FeNO < 20 ppb were categorized as B-Eos&FeNO-low asthma [35]. Additionally, we considered subjects with either B-Eos-high or FeNO-high as having “Type 2-high” asthma.

The AwObesity phenotype was defined by a BMI $\geq 30 \text{ kg/m}^2$ in individuals with current asthma [36]. Finally, the AwCOPD phenotype was considered if participants ≥ 40 years-old had concurrent asthma and COPD, defined by a positive answer to “Has a doctor ever told you that you have chronic bronchitis/emphysema”, with age of diagnosis ≥ 40 years and having self-reported smoking history (being either a current or ex-smoker) [37, 38].



Statistical analysis

In accordance with the NHANES sampling design, the weights for each full sample 2-year mobile examination center were used to obtain weighted percentages adjusted to the US adult population.

Categorical variables were described as frequencies and weighted proportions, and continuous variables were described as median and first and third quartiles (Q1–Q3). Chi square test and Mann–Whitney U-test were used to compare groups. To explore the association of concomitant (having at least 2 concurrent) phenotypes with each asthma-related outcome we performed multivariate logistic regression modelling. Separate models were run using each asthma-related outcome and abnormal lung function as dependent variable and having multiple phenotypes as independent variables. Adjustments were also made for potential confounders: sex, age, race, current smoking and rhinitis. Adjusted odds ratios (aOR)

with 95% confidence intervals (95% CI) were presented, and model fit was assessed using the *svylogitgof* function [39].

According to age (<40 or ≥40 years-old), a four- or five-set Venn-Euler diagram was used to quantify the proportion of individuals with different asthma phenotypes and to illustrate the overlap.

The diagrams were created using R software version 3.2.0 (“VennDiagram”, “venneuler” and “reshape2” packages) and all statistical analyses were performed in Stata version 13.1 (StataCorp, TX, USA), using the *survey* command to account for the complex sampling design and weights in the NHANES. The *MI* command was used to perform sensitivity analysis by multiple-imputation of missing values; however, to create the Venn-Euler diagrams, a listwise deletion for missing data was applied. A *p* value <0.05 was considered statistically significant.

Results

Of the 18,619 adults included in NHANES 2007–2012 datasets, 1059 (5.6% [95% CI 5.1–5.9]) had current asthma (Fig. 1). Of these, 63% were female, and the median (Q1–Q3) age was 48.0 (32.0–62.0) years. After excluding subjects with missing data on the main variables, 634 individuals were included for phenotype classification (Fig. 1). Despite having all information available, 77 patients did not meet the criteria for any of the defined asthma phenotypes and were considered “non-classified”. These were non-obese subjects with asthma who did not meet the criteria for COPD, had B-Eos values ranging between 150 and 300/mm³, and FeNO ranging 20–34 ppb.

Demographic characteristics of adults with current asthma included and excluded from the analysis and patients with single (n=271) and multiple phenotypes (n=286) are described in Table 1.

There is a female predominance in both groups (64 and 66%, respectively). Subjects with multiple phenotypes were older ($p=0.003$), had higher BMI ($p<0.001$), were more often obese ($p<0.001$) and ex-smokers ($p=0.003$), and a higher proportion of patients were treated with inhaled corticosteroids (ICS) ($p=0.01$), than those with only one phenotype. Females were more obese, regardless the number of concomitant asthma phenotypes (data not shown).

Phenotypes and overlap description

The weighted proportions of asthma phenotypes were (in descending order): 49% for AwObesity, 36% for B-Eos-high asthma, 26% for B-Eos&FeNO-low asthma, 18% for FeNO-high asthma, and 8% for AwCOPD (Table 1).

Demographic and clinical characteristics among all 5 asthma phenotypes and the “non-classified” group are described in Table 2.

There is a female predominance among all phenotypes, particularly in the B-Eos&FeNO-low (78%). Subjects with AwCOPD phenotype were the oldest group (median [Q1–Q3]: 61.0 [52.0–69.0] years-old), with the lowest proportion of individuals that had $\geq$ high school and lowest FEV₁/FVC (0.63 [0.50–0.75]), comparing to the other phenotypes.

When categorized by age, <40 (n=227) and $\geq$ 40 years-old (n=330), the most prevalent phenotypes were AwObesity (42 and 53%, respectively) and B-Eos-high asthma (34 and 37%). The less ones were FeNO-high asthma (18 and 19%) and AwCOPD (19% in the older group) (Fig. 2).

The areas of intersection in the four- and five-set Venn-Euler diagrams revealed 5 and 12 overlapping categories, and proportions of 17 and 12% of non-classified asthma subjects, respectively.

In both diagrams, a substantial total overlap was observed: 44% in subjects <40 years-old and 54% in subjects $\geq$ 40 years-old. About 40% of the individuals in both age groups had two concomitant asthma phenotypes, 4% of the younger group had 3 concomitant phenotypes and 13% of the older group had $\geq$ 3 (Table 3 and Fig. 2). Furthermore, 1% of the older subjects had four concomitant asthma phenotypes: AwObesity, AwCOPD, FeNO-high, and B-Eos-high asthma.

The most prevalent overlaps in both groups (<40 and $\geq$ 40 years-old) were AwObesity together with either B-Eos-high (15 and 12%, respectively) or B-Eos&FeNO-low asthma (13 and 11%) (Fig. 2).

Moreover, the proportions of subjects having AwObesity together with other phenotypes were high: 53% for the B-Eos-high phenotype, 48% for AwCOPD, 45% for the B-Eos&FeNO-low, and 44% for the FeNO-high phenotype. Also, the proportion of individuals having AwCOPD together with the B-Eos-high phenotype was high (36%), whereas the proportions were lower for the B-Eos&FeNO-low and the FeNO-high asthma phenotypes (15 and 10%, respectively) (data not shown).

In this population, only 12 and 15% of asthma subjects (<40 and $\geq$ 40 years-old, respectively) with high B-Eos count had a concomitant high FeNO values (Fig. 2). Moreover, the two biomarkers were non-congruent across cut-offs. For example, when comparing groups with B-Eos count <150/mm³ and 150–300/mm³, the proportion of asthma subjects having low FeNO (<20 ppb), was not significantly different (Additional file 2: Table S1).

Associations between asthma-related outcomes and phenotype overlap

A comparison of the clinical characteristics of participants with one, two or three or more asthma phenotypes, stratified by age, is presented in Table 3 and no significant differences were observed in any age groups with regard to asthma attacks, asthma-related ED, $\geq$ 2 asthma symptoms, and use of $\geq$ 1 reliever. In the older group, the proportion of individuals with work/school absenteeism, $\geq$ 2 controller medications and with FEV₁/FVC<LLN was significantly higher in participants with concomitant phenotypes than in those with a single phenotype (Table 3). In both age groups, the proportion of patients with FEV₁<LLN was significantly higher when participants presented multiple phenotypes, as well as they presented lower median FEV₁% predicted values.

When analyzing the asthma-related outcomes in subjects with a single phenotype with those having specific combination of asthma phenotypes, the overall findings were that subjects having multiple phenotypes had significantly higher proportion of using $\geq$ 1 reliever and $\geq$ 2

Table 1 Characteristics of adults with current asthma: included and excluded from phenotype classification, and stratified by single or multiple phenotypes

Demographic characteristics n (wt%)	Included subjects n = 634	Excluded subjects n = 425	<i>p</i> value*	Single phenotype [†] n = 271	Multiple phenotypes [†] n = 286	<i>p</i> value*
Female gender	410 (63)	261 (64)	0.93	174 (64)	192 (66)	0.68
Age (yrs), median (Q1–Q3)	44.0 (31.0–57.0)	48.9 (33.7–68.0)	< 0.001	42.0 (30.0–55.0)	47.5 (34.0–60.0)	0.003
BMI (kg/m ²), median (Q1–Q3)	30.8 (25.4–35.9)	31.4 (24.4–35.7)	0.99	28.7 (24.2–35.0)	33.7 (30.7–39.0)	< 0.001
Obesity status						
Underweight (≤ 18.4 kg/m ²)	2 (0.3)	13 (4)	< 0.001	1 (0.2)	1 (0.5)	0.41
Normal (18.5–24.9 kg/m ²)	142 (25)	98 (27)	0.44	78 (30)	20 (8)	< 0.001
Overweight (25–29.9 kg/m ²)	151 (26)	77 (19)	0.07	80 (32)	38 (15)	< 0.001
Obese (≥ 30 kg/m ²)	339 (49)	187 (49)	0.94	112 (38)	227 (77)	< 0.001
Race and/or ethnicity						
Hispanic	105 (8)	85 (10)	0.30	38 (8)	56 (9)	0.24
Non-Hispanic white	323 (74)	184 (63)	0.005	135 (72)	138 (72)	0.93
Non-Hispanic black	167 (14)	117 (18)	0.03	76 (14)	81 (16)	0.71
Other Race	39 (4)	39 (7)	0.06	22 (6)	11 (3)	0.11
Smoking status						
Current smoker	199 (29)	114 (32)	0.40	89 (31)	87 (27)	0.46
Ex-smoker	163 (29)	98 (23)	0.13	56 (20)	89 (36)	0.003
Non-smoker	272 (43)	154 (45)	0.53	126 (49)	110 (37)	0.02
Education						
$\geq$ High school	478 (84)	225 (69)	< 0.001	205 (86)	209 (81)	0.13
Asthma-related medication [‡]						
Reliever medication**	276 (41)	202 (48)	0.16	113 (37)	132 (47)	0.10
Oral corticosteroids	33 (8)	29 (3)	0.001	12 (3)	15 (4)	0.75
Inhaled corticosteroids [§]	153 (25)	122 (30)	0.14	55 (19)	81 (32)	0.01
Other control medications	53 (9)	63 (14)	0.19	15 (8)	33 (10)	0.50
Asthma phenotype						
AwObesity	339 (49)	–		112 (38)	227 (76)	< 0.001
B-Eos-high	237 (36)	–		61 (22)	176 (62)	< 0.001
B-Eos&FeNO-low	157 (26)	–		74 (30)	83 (30)	0.95
FeNO-high	110 (18)	–		18 (8)	92 (34)	< 0.001
AwCOPD	57 (8)	–		6 (2)	51 (17)	< 0.001
Non-classified ^{††}	77 (14)	–		–	–	

Data presented as absolute numbers and proportions weighted for the U.S. population. *p* values < 0.05 are presented in italic

Yrs years, BMI body mass index, Q1 first quartile, Q3 third quartile, BMI body mass index, AwObesity Asthma with obesity, AwCOPD Asthma with concurrent COPD

* Chi square test or Mann–Whitney U-test was used

[†] Seventy seven subjects included in the “non-classified” group were considered as missing

[‡] Prescribed medication taken in the past 30 days

[§] Alone or in combination with long-acting inhaled β_2 -agonist

^{||} Included long-acting inhaled β_2 -agonist (without corticosteroids), leukotriene inhibitors, and mast cell stabilizers

**Short-acting β_2 -agonist and/or anticholinergic

^{††} Subjects with non-single and non-multiple asthma phenotype

controller medications and had decreased lung function, with the exception of those with the B-Eos&FeNO-low phenotype combined with any of the other phenotypes (Additional file 3: Table S2, and Additional file 4: Table S3).

Moreover, a lower proportion of subjects reporting asthma attacks was observed in subjects with AwObesity and either FeNO-high (26%) or AwCOPD (20%), compared to those with a single phenotype (67%) (Additional file 3: Table S2). Subjects with concomitant AwCOPD and B-Eos&FeNO-low phenotypes had the lowest

Table 2 Demographic and clinical characteristics among all 5 phenotypes and in the “Non-classified” group

Characteristics n (wt%)	AwObesity n = 339	B-Eos-high n = 237	B-Eos&FeNO-low n = 157	FeNO-high n = 110	AwCOPD n = 57	Non-classified ^{††} n = 77
Female gender	366 (67)	196 (55)	138 (78)	78 (52)	71 (58)	44 (56)
Age (yrs), median (Q1–Q3)	48.0 (34.0–59.0)	47.0 (31.0–59.0)	41.0 (27.0–57.0)	45.0 (30.0–54.0)	61.0 (52.0–69.0)	39.0 (28.0–53.0)
BMI (kg/m ²), median (Q1–Q3)	35.4 (32.5–40.4)	30.3 (25.9–36.8)	28.9 (24.2–33.0)	27.6 (24.9–33.4)	30.3 (25.1–35.1)	24.3 (22.8–27.5)
Race and/or ethnicity						
Hispanic	94 (9)	78 (11)	30 (8)	29 (10)	18 (4)	11 (6)
Non-Hispanic white	230 (65)	175 (73)	79 (69)	65 (75)	80 (81)	50 (86)
Non-Hispanic black	177 (21)	74 (12)	62 (17)	38 (13)	27 (11)	10 (6)
Other race	25 (5)	21 (4)	13 (6)	6 (2)	10 (4)	6 (2)
Smoking status						
Current smoker	150 (27)	104 (32)	48 (23)	18 (13)	65 (52)	23 (28)
Ex-smoker	134 (27)	100 (32)	35 (25)	42 (39)	70 (48)	18 (30)
Non-smoker	229 (45)	124 (35)	88 (51)	66 (49)	0 (0)	36 (42)
Education						
≥ High school	358 (79)	227 (80)	125 (81)	102 (87)	73 (59)	64 (91)
Asthma-related medication [‡]						
Reliever medication ^{**}	234 (42)	172 (51)	82 (44)	65 (46)	69 (57)	31 (36)
Oral corticosteroids	36 (5)	17 (5)	4 (2)	10 (7)	15 (7)	2 (2)
Inhaled corticosteroids [§]	144 (27)	103 (32)	36 (20)	36 (33)	60 (47)	17 (19)
Asthma-related outcomes						
Asthma attack	363 (68)	252 (74)	125 (68)	98 (71)	85 (63)	54 (75)
Asthma-related ED	130 (27)	80 (23)	41 (23)	26 (13)	37 (32)	9 (8)
> 2 asthma symptoms	311 (66)	199 (65)	92 (55)	80 (57)	90 (74)	42 (59)
Work/school absenteeism	66 (18)	43 (16)	25 (18)	23 (14)	12 (20)	10 (14)
Lung function						
FEV ₁ % predicted, median (Q1–Q3)	89.0 (75.6–99.2)	84.1 (75.2–95.4)	93.2 (83.8–100.8)	82.7 (75.9–95.4)	74.0 (62.9–90.1)	89.9 (80.5–103.5)
FEV ₁ /FVC, median (Q1–Q3)	0.77 (0.62–0.82)	0.74 (0.65–0.80)	0.79 (0.72–0.83)	0.72 (0.66–0.79)	0.63 (0.50–0.75)	0.76 (0.69–0.82)

Data presented as absolute numbers and proportions weighted for the U.S. population

Yrs years, BMI body mass index, Q1 first quartile, Q3 third quartile, BMI body mass index, AwObesity Asthma with obesity, AwCOPD Asthma with concurrent (COPD), ED emergency-department, FEV₁ Forced expiratory volume in 1 s, FEV₁/FVC forced expiratory volume in 1 s and functional vital capacity ratio, LLN lower limit of normality

^{††} Subjects with non-single and non-multiple asthma phenotype

[‡] Prescribed medication taken in the past 30 days

^{**} Short-acting β_2 -agonist and/or anticholinergic

[§] Alone or in combination with long-acting inhaled β_2 -agonist


proportion of ≥ 2 asthma symptoms (20%), but had the highest proportion of using ≥ 1 reliever medication (84%) as well as having FEV₁ < LLN (71%).

In multivariate regression analysis, adjusting for co-variables, having multiple phenotypes was significantly associated with using ≥ 2 controller medications (aOR, 95% CI 2.03, 1.16–3.57), and having reduced FEV₁ (3.21, 1.73–5.94) (Table 4). However, no associations were seen with asthma attacks, asthma-related ED, ≥ 2 asthma symptoms, work/school absenteeism, use of reliever medication or FEV₁/FVC < LLN (Additional file 5: Table S4).

Furthermore, subjects aged ≥ 40 years-old, had significantly higher odds of using ≥ 2 controller medications

and having FEV₁ < LLN predicted, compared to those < 40 years-old, adjusted for covariates (Table 4). Being a current smoker was significantly associated with using ≥ 1 reliever medication (1.95, 1.35–2.83) and with reduced lung function: FEV₁ < LLN predicted (2.01, 1.21–3.33) and FEV₁/FVC < LLN (2.02, 1.16–3.51) and not associated with any other asthma-related outcomes (Table 4 and Additional file 5: Table S4).

The association between having concomitant phenotypes and using multiple controller medications was consistent when considering oral corticosteroids (OCS) separated from other controller medications (1.87, 1.09–3.21) (data not shown).



We also analyzed the potential bias of controller medications in the phenotype classification, particularly in the B-Eos-high and FeNO-high phenotypes (Additional file 6: Fig. S1). No significant differences in asthma-related treatment were found between the phenotypes, with exception for a higher proportion of patients treated with ICS within the FeNO-high and B-Eos-high phenotypes compared to those with B-Eos&FeNO-low phenotype ($p=0.03$). When restricting to subjects with a single asthma phenotype no significant differences were found.

Moreover, sensitivity analyses showed that the proportion of total overlap (weighted 53%), and the associations between having multiple phenotypes and asthma outcomes were similar when imputing all missing values (data not shown). The goodness-of-fit test revealed adequate fitting for all regression models, except when using $FEV_1/FVC < LLN$ as dependent variable (Additional file 5: Table S4) and no statistically significant interactions between co-variables were observed.

Discussion

We report a substantial overlap of commonly reported asthma phenotypes among adults with current asthma in a large population sample, with almost half of them having two or more concomitant phenotypes. Furthermore, having multiple asthma phenotypes, regardless of their classification, was associated with poorer asthma outcomes, particularly the use of more controller medication and reduced lung function.

These findings illustrate the complexity and unique features of the concomitant asthma phenotypes when categorizing asthma in adults, using only the “classical”

(hypothesis-driven) approach, based on measures readily available in the clinic (such as non-invasive biomarkers and medical records).

Hypothesis-driven asthma phenotypes are usually based on single dimensions of the disease, such as clinical symptoms, triggers, pathology or patterns of airway obstruction [16–18, 40–42]. However, evidence has shown that this approach is highly heterogeneous, as it depends on the a priori assumptions and target population [43–45]. Also, it is of note that the 77 subjects with asthma that could not be classified as having any of the studied phenotypes, supports the fact that there is a considerable number of asthma patients whose clinical phenotype is not easily classified (e.g. asthmatics with irreversible airflow obstruction, patients with similar airways symptoms but with different pattern of airway inflammation), suggesting the presence of sub-phenotypes [1, 22, 44, 45].

In an attempt to explore the pathophysiology of specific asthma subgroups, and help stratify patients for targeted therapies, data-driven or unsupervised approaches (such as k-means, hierarchical clustering, partition-around-medoids methods or latent class analysis) are being applied in airways disease to identify “novel” accurate and distinct phenotypes, taking into account the heterogeneity and multidimensional characteristics of the disease [8, 46–52].

Our study results seem to be in line with the view of those that argue for a combination of both hypothesis- and data-driven approaches as a way forward to progress our knowledge on asthma endotypes and clinical phenotypes in an iterative way [52–54]. The data-driven

Table 3 Distribution and comparisons of the asthma-related outcomes among asthma phenotypes, stratified by age

	Total n (wt%)	Asthma attack	Asthma-related ED	≥ 2 asthma symptoms	Work/school absenteeism	Asthma medication		Lung function		
						≥ 1 reliever medication [†]	≥ 2 controller medication	FEV ₁ < LLN	FEV ₁ % predicted [§]	FEV ₁ /FVC < LLN
<40 yrs										
1 phenotype	118 (56)	85 (70)	27 (20)	59 (56)	15 (21)	55 (43)	17 (11)	10 (7)	95.6 (90.2–102.1)	26 (20)
2 phenotypes	97 (40)	69 (75)	26 (32)	57 (73)	12 (9)	41 (40)	8 (9)	19 (23)	91.8 (81.3–99.2)	25 (29)
3 phenotypes	12 (4)	10 (73)	5 (40)	8 (72)	4 (25)	9 (68)	1 (4)	3 (36)	81.9 (75.3–84.6)	4 (29)
<i>p</i> value*										
1 versus 2		0.43	0.10	0.052	0.04	0.72	0.64	0.006	0.07	0.25
2 versus 3		0.92	0.55	0.96	0.07	0.15	0.46	0.43	0.009	0.98
1 versus 3		0.86	0.11	0.39	0.73	0.20	0.34	0.01	<0.001	0.48
≥40 yrs										
1 phenotype	153 (46)	104 (65)	35 (23)	76 (62)	20 (15)	58 (33)	26 (17)	29 (20)	91.2 (81.6–99.2)	34 (26)
2 phenotypes	136 (41)	92 (69)	28 (21)	70 (55)	12 (9)	60 (46)	43 (36)	31 (37)	80.4 (70.0–91.7)	27 (32)
≥3 pheno- types	41 (13)	26 (72)	4 (9)	30 (70)	8 (40)	22 (60)	16 (40)	12 (46)	74.0 (63.0–85.8)	16 (53)
<i>p</i> value*										
1 versus 2		0.56	0.82	0.38	0.41	0.12	0.02	0.01	0.007	0.46
2 versus ≥3		0.80	0.27	0.30	0.002	0.24	0.67	0.47	0.22	0.09
1 versus ≥3		0.50	0.21	0.54	0.03	0.053	0.02	0.01	0.006	0.02

Data presented as absolute numbers and proportions weighted for the U.S. population. *p* values < 0.05 are presented in italic. The 77 subjects included in the “non-classified” group were considered as missing

ED emergency-department, FEV₁ forced expiratory volume in 1 s, FEV₁/FVC forced expiratory volume in 1 s and functional vital capacity ratio, LLN lower limit of normality, Q1 first quartile; Q3: third quartile

* Chi square test or Mann–Whitney U-test was used

[†] Short-acting β₂-agonist or/and anticholinergic

[§] Presented as median (Q1–Q3)

Table 4 Regression models with significant associations between having multiple asthma phenotypes and asthma-related outcomes, adjusted for co-variables

	≥ 2 controller medications		FEV ₁ < LLN	
	aOR	95% CI	aOR	95% CI
Multiple versus single phenotype	2.03	1.16–3.57	3.21	1.74–5.94
Female gender	1.39	0.77–2.50	1.51	0.81–2.81
Age ≥ 40 yrs	3.01	1.52–5.95	2.55	1.29–5.03
Caucasian versus others	1.38	0.86–2.23	1.37	0.78–2.42
Current smoker versus non-/ex-smokers	1.02	0.52–2.02	2.01	1.21–3.33
Rhinitis	1.08	0.57–2.16	0.94	0.54–1.63
Goodness-of-fit test				
χ ² (p value)	0.86 (0.56)		0.80 (0.61)	

Multivariate logistic regression models adjusted for gender, age, race, current smoking and rhinitis. The aOR values with $p < 0.05$ are presented in italic

FEV₁, forced expiratory volume in 1 s, LLN lower limit of normal, CI confidence interval, aOR adjusted odds ratio, χ² Chi square goodness-of-fit

phenotypes studies obtained some of the phenotypes already defined by hypothesis-driven approaches (e.g. obese, non-eosinophilic asthmatics [8]; persistent airway inflammation [46]; low type-2 inflammation [49]; fixed obstructive, non-eosinophilic and neutrophilic [50]), but, importantly, they identified other phenotypes that differ by certain characteristics: clinical parameters [8, 47–49], clinical response to treatment [46, 52], and airway inflammation [49, 51]. Therefore, further studies are required to compare and validate the asthma phenotypes obtained using different unsupervised methods.

The high overlap of asthma phenotypes seen in this study was similar to the findings of Tran et al. [15], who used datasets from previous NHANES surveys to evaluate the overlap of asthma phenotypes. However, Tran et al. study focused on allergic asthma phenotypes, based on IgE levels, and was therefore limited to the 2005–2006 survey that lacks data on FeNO. We provided a broader analysis of phenotypes that included not only the eosinophil-based phenotype (associated more closely with IL-5-driven) and the one based on FeNO values (mostly dependent on IL-4/IL-13-driven) [12], but also other phenotypes not defined by biomarkers and in a much larger dataset.

In this study, we extended previous observations [7, 11] suggesting that FeNO and B-Eos count partially reflect different inflammatory pathways, representing a local and a systemic type 2-marker, respectively. We observed that only 12–15% of asthma subjects with high B-Eos count had concomitant high FeNO levels, in this population. Also, a similar proportion of subjects with multiple

phenotypes was obtained when considering the “Type 2-high” phenotype, supporting the view that these two biomarkers are not interchangeable and that the use of both biomarkers in combination may allow for better targeted and personalized treatment for at least certain subsets of asthma patients [7, 10, 11].

The more prevalent combinations of phenotypes observed in this study were AwObesity together with either B-Eos-high or B-Eos&FeNO-low phenotypes. This supports the view that obesity-related asthma, despite often suggested to be a separate asthma phenotype associated with non-eosinophilic airway inflammation [9, 55, 56], may also be associated to eosinophilic inflammation.

Given the high prevalence in the US population, in this sample, obesity is likely to be a comorbidity, rather than the primary reason for asthma [57]; however, we defined the AwObesity phenotype as a separate group, since the interdependence on inflammatory markers to targeting different asthma therapies makes essential the accurate characterization of inflammation in obese asthmatic subjects [19, 58]. In addition, the relevance of defining the AwObesity phenotype is supported by the data as the weighted proportion of overlap is similar when excluding AwObesity or B-Eos-high phenotype from the analysis (data not shown). Moreover, the weighted proportion of subjects with “non-classified” asthma doubled when excluding the AwObesity phenotype (increasing to 32% in the < 40 years-old and 29% in ≥ 40 years old).

Having multiple asthma phenotypes was more common in older subjects and in non- and former- smokers; whether this is due to a general increase of comorbidities with age [58–60] and/or an interaction with environmental factors [61] cannot be specifically addressed by this study design. Nevertheless, when interpreting these results one should bear in mind that AwCOPD is associated to older age and prior/current smoking while FeNO increases with age and decreases with smoking [62, 63].

Interestingly, subjects with a higher number of concomitant asthma phenotypes presented reduced lung function and this association remained when controlling for potential confounders by multiple regression analysis. This shows that having multiple phenotypes is independently associated with reduced lung function, suggesting a cumulative effect of different disease processes.

Moreover, patients having several commonly reported asthma phenotypes had higher odds of using more controller medications, supporting the view that these patients are those with more complex disease and higher asthma morbidity [9, 15]. This also suggests that these asthma patients have an inadequate response to prescribed therapies, since lung function was reduced, and that they may represent a group of patients with the need for add-on treatment. However, the choice of specific

treatments, such as biological therapies, will be more difficult considering the complexity introduced by having multiple phenotypes.

The lack of significant associations between multiple phenotypes and the other asthma-related outcomes may be difficult to understand. A possible explanation could be that the prescribed medication is effective against asthma symptoms and attacks but less effective against (subclinical) processes that cause long-term reduction in lung function. However, the results could also be related to data collection methods, as lung function measurement and the way medication use was ascertained, were less dependent on patient recall than the self-reported variables that were used for the outcomes with null results in the present study. Further studies should be done, adding the quantitative assessment of asthma attacks/asthma-related ED visits, and also including the age of asthma onset, that could have an influence on asthma-related outcomes.

This study has several limitations. First, because of its cross-sectional design, it was not possible to evaluate interactions between phenotypes over time in patients with concomitant phenotypes, nor was it possible to determine which phenotype occurred first. Second, although there were differences between the included and excluded groups in the variables age, BMI, non-Hispanic white/black subjects, having finished high school and OCS use, the majority were not used for phenotypic classification and did not affect the outcomes, as shown in sensitivity analysis. Third, as our asthma and COPD definitions were based on self-reported diagnosis, rather than relying on lung function tests, the acquired information is subject to recall bias and misclassification. However, these definitions have been commonly used in NHANES reports [7, 11, 15] and have proven to be reasonably reliable [64, 65]. Moreover, we have used the most frequent combination of questions seen in epidemiological studies [65, 66], and we also included questions on recent wheeze and/or asthma attacks, which should reduce the risk of including individuals without true disease. Also, we stratified the analysis by age (at 40 years), as used in other COPD studies [37, 67], in order to improve the clinical value into the interpretation of phenotyping data, as the overlap among asthma phenotypes will be different among those less than age 40 and those older than 40 (with higher possibility of having COPD) [37]. Fourth, the lack of other biomarkers in the present NHANES years, prevented the analysis of other asthma phenotypes and the use of alternative definitions. However, we analyzed biomarkers of type 2-inflammation in both blood and exhaled air that previously have been shown to independently relate to asthma morbidity [7, 11]. Fifth, as there is no consensual definition of

biomarker-defined asthma phenotypes, we based our definitions on cut-offs used in previous studies to discriminate patients in single asthma phenotypes [32–35], rather than on any reported specificity or sensitivity for predicting asthma morbidity or response to therapeutics [68, 69]. For high probability of eosinophilic inflammation, the cut-off value for FeNO has been suggested to be >50 ppb for adults [70]. However, we chose a FeNO cut-off of 35 ppb, based on the mean baseline FeNO levels of patients included in randomized controlled trials of anti-IL-13 treatment [33, 69]. In spite of using this lower cut-off, 77 subjects with current asthma could not be classified as having any of the predefined phenotypes, indicating the need for better, and probably personalized, cut-offs for biomarkers in asthma.

Furthermore, we could not demonstrate a clear effect of ongoing controller medication in the phenotypic classification in our data. No significant associations were observed, probably at least partially explained with the exclusion of the participants with ICS/OCS use <48 h prior to the exhaled NO measurements. Also, contrary to the expected, we observed a higher proportion of patients treated with ICS/OCS within both B-Eos-high and FeNO-high phenotypes than the B-Eos&FeNO-low phenotype. A plausible explanation is that subjects with ongoing inflammation have more clear asthma, with more symptoms, and, thus, a higher need of treatment. B-Eos&FeNO-low asthma is a heterogeneous group with less need of controller treatment, and because the treatment is ineffective, it may be that medication use and even prescription has been stopped.

Finally, even though we did not specifically analyze the overlap of asthma phenotypes in patients with severe asthma, the significant association between presenting more than one phenotype and being treated with multiple asthma controller medications suggests higher asthma severity in this subset [71]. Also, we did not consider individual environmental factors, such as air pollution and/or indoor allergens, that could influence asthma phenotypes. Further studies describing the overlap in patients with severe asthma and studies examining asthma patients exposed to different environmental factors, such as subjects who live in cities versus in rural areas are needed.

This study indicates that the overlap of commonly reported asthma phenotypes is observed also in non-selected asthma patients from the general population. Our findings highlight the importance of classifying asthma patients with regard to applicable phenotypes, rather than using a single asthma phenotype, to enable the development of adequate targeted strategies to avoid lung function impairment. However, further data is required, such as that from higher order analysis, using

data-mining methods possibly combined with those that rely on predefined hypotheses. This synergy is expected to improve the knowledge on asthma phenotypes and, ultimately, to lead to more personalized treatment strategies [53, 54].

Conclusions

In conclusion, a prevalent overlap of commonly reported phenotypes was observed in asthma patients identified from the general population. Subjects classified as having multiple phenotypes used more controller medications and had reduced lung function. Thus, the complexity and unique features of concomitant asthma phenotypes may require a broader data analysis approach, based on a combination of clinical information and biomarkers resulting in better characterization of patients. This could lead to better asthma outcomes, particularly preserved lung function.

Additional files

Additional file 1. Supplementary Methods.

Additional file 2: Table S1. Distribution and comparisons between the FeNO and B-Eos cut-offs used in this study, among individuals with current asthma.

Additional file 3: Table S2. Weighted percentages and comparisons of asthma-related outcomes among subjects with a single asthma phenotype versus: non-classified, and specific combinations of asthma phenotypes.

Additional file 4: Table S3. Weighted percentages and comparisons of asthma-related outcomes among subjects with a single asthma phenotype versus: non-classified, and specific combinations of asthma phenotypes.

Additional file 5: Table S4. Multivariable logistic regression models between each asthma-related outcome and having multiple asthma phenotypes, adjusted for co-variables.

Additional file 6: Fig. S1. Proportions (weighted to the US population) of subjects taking asthma controller medications stratified into the different phenotypes, among all participants included for asthma phenotype classification (left) and only in those with a single phenotype (right). P-values <0.05 were indicated. NA: Non-applicable (not possible to determine because some participants had both B-Eos-high and FeNO-high asthma phenotypes).

Abbreviations

aOR: adjusted odds ratios; ATS/ERS: American Thoracic Society/European Respiratory Society; AwCOPD: asthma with concurrent COPD; AwObesity: asthma with obesity; B-Eos: blood eosinophils; BMI: body mass index; COPD: chronic obstructive pulmonary disease; CI: confidence intervals; ED: emergency department; FeNO: fraction of exhaled nitric oxide; FEV₁: forced expiratory volume in 1 s; FEV₁/FVC: ratio of forced expiratory volume in 1 s to forced vital capacity; ICS: inhaled corticosteroids; IgE: immunoglobulin E; IL: interleukin; LLN: lower limit of normal; NHANES: National Health and Nutrition Examination Survey (USA); OCS: oral corticosteroids; Q1: first quartile; Q3: third quartile; Th2: T helper cell type 2.

Authors' contributions

RA, JAF, AM and KA contributed to study conception and design, analysis and interpretation of data, writing and revising the article. TJ, AMP and CJ

contributed to data interpretation, writing and revising the article. All authors read and approved the final version of the manuscript.

Author details

¹ CINTESIS- Center for Health Technology and Services Research, Faculty of Medicine, University of Porto, Edifício Nascente, Piso 2, Rua Dr. Plácido da Costa, s/n, 4200-450 Porto, Portugal. ² Department of Cardiovascular and Respiratory Sciences, Porto Health School, Porto, Portugal. ³ MEDCIDS- Department of Community Medicine, Information, and Health Sciences, Faculty of Medicine, University of Porto, Porto, Portugal. ⁴ Department of Allergy, Instituto & Hospital CUF, Porto, Portugal. ⁵ Department of Medical Sciences: Clinical Physiology, Uppsala University, Uppsala, Sweden. ⁶ Department of Medical Sciences: Respiratory Medicine and Allergology, Uppsala University, Uppsala, Sweden. ⁷ Department of Women's and Children's Health: Paediatric Research, Uppsala University, Uppsala, Sweden.

Acknowledgements

RA is supported by Fundação para a Ciência e Tecnologia (PhD grant – Ref. PD/BD/113659/2015), Portugal.

Competing interests

The authors declare that they have no competing interests.

Availability of data

The datasets used and analyzed during this study are available in the NHANES website: <https://www.cdc.gov/nchs/nhanes/index.htm>.

Consent for publication

Not applicable.

Ethics approval and consent to participate

The NHANES survey operates under the approval of the National Center for Health Statistics Research Ethics Review Board, Protocols #2005-06, and #2011-17 (www.cdc.gov/nchs/nhanes/irba98.htm). All of the NHANES data meet the circumstances described in Policy #39, Research Using Publicly Available Data-sets (Secondary Analysis) for use without application to Institutional Review Board. Moreover, all study participants provided written informed consent.

Funding

Project "NORTE-01-0145-FEDER-000016" is financed by the North Portugal Regional Operational Programme (NORTE 2020), under the PORTUGAL 2020 Partnership Agreement, and through the European Regional Development Fund (ERDF).

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 12 December 2017 Accepted: 26 March 2018

Published online: 04 May 2018

References

1. Froidure A, Mouthuy J, Durham SR, Chanez P, Sibille Y, Pilette C. Asthma phenotypes and IgE responses. *Eur Respir J*. 2016;47:304–19.
2. Corren J. Asthma phenotypes and endotypes: an evolving paradigm for classification. *Discov Med*. 2013;15(83):243–9.
3. Woodruff PG, Modrek B, Choy DF, Jia G, Abbas AR, Ellwanger A, Koth LL, Arron JR, Fahy JV. T-helper type 2-driven inflammation defines major sub-phenotypes of asthma. *Am J Respir Crit Care Med*. 2009;180(5):388–95.
4. Robinson D, Humbert M, Buhl R, Cruz AC, Inoue H, Korom S, Hanania NA, Nair P. Revisiting Type 2-high and Type 2-low airway inflammation in asthma: current knowledge and therapeutic implications. *Clin Exp Allergy*. 2017;47(2):161–75.
5. Hastie AT, Moore WC, Li H, Rector BM, Ortega VE, Pascual RM, Peters SP, Meyers DA, Bleecker ER, National Heart, Lung, and Blood Institute's Severe Asthma Research Program. Biomarker surrogates do not accurately

- predict sputum eosinophil and neutrophil percentages in asthmatic subjects. *J Allergy Clin Immunol*. 2013;132(1):72.e12–80.e12.
6. Green RH, Brightling CE, Woltmann G, Parker D, Wardlaw AJ, Pavord ID. Analysis of induced sputum in adults with asthma: identification of subgroup with isolated sputum neutrophilia and poor response to inhaled corticosteroids. *Thorax*. 2002;57(10):875–9.
 7. Malinovschi A, Fonseca JA, Jacinto T, Alving K, Janson C. Exhaled nitric oxide levels and blood eosinophil counts independently associate with wheeze and asthma events in National Health and Nutrition Examination Survey subjects. *J Allergy Clin Immunol*. 2013;132(4):821.e5–27.e5.
 8. Jia G, Erickson RW, Choy DF, Mosesova S, Wu LC, Solberg OD, et al. Periostin is a systemic biomarker of eosinophilic airway inflammation in asthmatic patients. *J Allergy Clin Immunol*. 2012;130(3):647.e10–54.e10.
 9. Haldar P, Pavord ID, Shaw DE, Berry MA, Thomas M, Brightling CE, et al. Cluster analysis and clinical asthma phenotypes. *Am J Respir Crit Care Med*. 2008;178(3):218–24.
 10. Jia G, Erickson RW, Choy DF, Mosesova S, Wu LC, Solberg OD, Shikotra A, Carter R, Audusseau S, Hamid Q, et al. Identification of airway mucosal type 2 inflammation by using clinical biomarkers in asthmatic patients. *J Allergy Clin Immunol*. 2017;140(3):710–9.
 11. Malinovschi A, Janson C, Borres M, Alving K. Simultaneously increased fraction of exhaled nitric oxide levels and blood eosinophil counts relate to increased asthma morbidity. *J Allergy Clin Immunol*. 2016;138(5):1301.e2–308.e2.
 12. Katial RK, Bensch GW, Busse WW, Chipps BE, Denson JL, Gerber AN, Jacobs JS, Kraft M, Martin RJ, Nair P, et al. Changing paradigms in the treatment of severe asthma: the role of biologic therapies. *J Allergy Clin Immunol Pract*. 2017;5(2):S1–14.
 13. Chung KF, Adcock IM. How variability in clinical phenotypes should guide research into disease mechanisms in asthma. *Ann Am Thorac Soc*. 2013;10(Suppl):S109–17.
 14. Merritt F. Blood eosinophils: the Holy Grail for asthma phenotyping? *Ann Allergy Asthma Immunol*. 2016;116(2):90–1.
 15. Tran TN, Zeiger RS, Peters SP, Colice G, Newbold P, Goldman M, Chipps BE. Overlap of atopic, eosinophilic, and TH2-high asthma phenotypes in a general population with current asthma. *Ann Allergy Asthma Immunol*. 2016;116(1):37–42.
 16. Borish L. The immunology of asthma: asthma phenotypes and their implications for personalized treatment. *Ann Allergy Asthma Immunol*. 2016;117(2):108–14.
 17. Hekking PPW, Bel EH. Developing and emerging clinical asthma phenotypes. *J Allergy Clin Immunol Pract*. 2014;2(6):671–80.
 18. Wenzel SE. Asthma phenotypes: the evolution from clinical to molecular approaches. *Nat Med*. 2012;18(5):716–25.
 19. Gibeon D, Batuwita K, Osmond M, Heaney LG, Brightling CE, Niven R, et al. Obesity-associated severe asthma represents a distinct clinical phenotype: analysis of the British Thoracic Society Difficult Asthma Registry Patient cohort according to BMI. *Chest*. 2013;143(2):406–14.
 20. Kobayashi S, Hanagama M, Yamada S, Ishida M, Yanai M. Inflammatory biomarkers in asthma-COPD overlap syndrome. *Int J Chronic Obstr Pulm Dis*. 2016;11:2117–23.
 21. Carr TF, Zeki AA, Kraft M. Eosinophilic and noneosinophilic asthma. *Am J Respir Crit Care Med*. 2018;197(1):22–37.
 22. Christenson SA, Steiling K, van den Berge M, Hijazi K, Hiemstra PS, Postma DS, Lenburg ME, Spira A, Woodruff PG, et al. Asthma-COPD overlap. Clinical relevance of genomic signatures of type 2 inflammation in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med*. 2015;191(7):758–66.
 23. Complete Blood Count. http://www.cdc.gov/nchs/data/nhanes/nhanes_11_12/cbc_g_met_he.pdf. Accessed 17 June 2017.
 24. Respiratory Health ENO Procedures Manual. www.cdc.gov/nchs/data/nhanes/nhanes_11_12/Respiratory_Health_ENO_Procedures_Manual.pdf. Accessed 26 May 2017.
 25. Respiratory Health Spirometry Procedures Manual. https://www.cdc.gov/nchs/data/nhanes/nhanes_11_12/spirometry_procedures_manual.pdf. Accessed 6 June 2017.
 26. ATS/ERS. ATS/ERS Recommendations for Standardized Procedures for the Online and Offline Measurement of Exhaled Lower Respiratory Nitric Oxide and Nasal Nitric Oxide, 2005. *Am J Respir Crit Care Med*. 2005;171(8):912–30.
 27. Miller MR. Standardisation of spirometry. *Eur Respir J*. 2005;26(2):319–38.
 28. Hankinson JL, Odencrantz JR, Fedan KB. Spirometric reference values from a sample of the general U.S. population. *Am J Respir Crit Care Med*. 1999;159(1):179–87.
 29. Hankinson JL, Kawut SM, Shahar E, Smith LJ, Stukovsky KH, Barr RG. Performance of American Thoracic Society-recommended spirometry reference values in a multiethnic sample of adults. *Chest*. 2010;137(1):138–45.
 30. Pellegrino R, Viegi G, Brusasco V, Crapo RO, Burgos F, Casaburi R, Coates A, van der Grinten CP, Gustafsson P, Hankinson J, et al. Interpretative strategies for lung function tests. *Eur Respir J*. 2005;26(5):948–68.
 31. Prescription Medications - Drug Information. http://www.cdc.gov/Nchs/Nhanes/1999-2000/RXQ_DRUG.htm. Accessed 6 June 2017.
 32. Castro M, Zangrilli J, Wechsler ME, Bateman ED, Brusselle GG, Bardin P, Murphy K, Maspero JF, O'Brien C, Korn S. Reslizumab for inadequately controlled asthma with elevated blood eosinophil counts: results from two multicentre, parallel, double-blind, randomised, placebo-controlled, phase 3 trials. *Lancet Respir Med*. 2015;3:355–66.
 33. Wenzel S, Ford L, Pearlman D, Spector S, Sher L, Skobieranda F, Wang L, Kirkesseli S, Rocklin R, Bock B, et al. Dupilumab in persistent asthma with elevated eosinophil levels. *N Engl J Med*. 2013;368(26):2455–66.
 34. Dweik RA, Sorkness RL, Wenzel S, Hammel J, Curran-Everett D, Comhair SA, Bleecker E, Busse W, Calhoun WJ, Castro M, et al. Use of exhaled nitric oxide measurement to identify a reactive, at-risk phenotype among patients with asthma. *Am J Respir Crit Care Med*. 2010;181(10):1033–41.
 35. McGrath KW, Icitovic N, Boushey HA, Lazarus SC, Sutherland ER, Chinchilli VM, Fahy JV. Asthma Clinical Research Network of the National Heart, Lung, and Blood Institute. A large subgroup of mild-to-moderate asthma is persistently noneosinophilic. *Am J Respir Crit Care Med*. 2012;185(6):612–9.
 36. Flegal KM, Carroll MD, Johnson CL. Overweight and obesity in the United States: prevalence and trends, 1960–1994. *Int J Obes Relat Metab Disord*. 1998;22(1):39–47.
 37. Buist AS, McBurnie MA, Vollmer WM, Gillespie S, Burney P, Mannino DM, et al. International variation in the prevalence of COPD (The BOLD Study): a population-based prevalence study. *Lancet*. 2007;370(9589):741–50.
 38. Diagnosis of Diseases of Chronic Airflow Limitation: Asthma, COPD and Asthma-COPD Overlap Syndrome (ACOS). <http://goldcopd.org/asthma-copd-asthma-copd-overlap-syndrome/>. Accessed 3 Jan 2017.
 39. Archer KJ, Lemeshow S. Goodness-of-fit test for a logistic regression model fitted using survey sample data. *Stata J*. 2006;6(1):97–105.
 40. Hirano T, Matsunaga K. Late-onset asthma: current perspectives. *J Asthma and Allergy*. 2018;11:19–27.
 41. Wenzel SE. Asthma: defining of the persistent adult phenotypes. *Lancet*. 2006;368:804–13.
 42. Vonk JM, Jongepier H, Panhuysen CIM, Schouten JP, Bleecker ER, Postma DS. Risk factors associated with the presence of irreversible airflow limitation and reduced transfer coefficient in patients with asthma after 26 years of follow up. *Thorax*. 2003;58(4):322–7.
 43. Bousquet J, Anto JM, Sterk PJ, Adcock IM, Chung KF, Roca J, et al. Systems medicine and integrated care to combat chronic noncommunicable diseases. *Genome Med*. 2011;3:43.
 44. Agusti A, Bel E, Thomas M, Vogelmeier C, Brusselle G, Holgate S, Humbert M, Jones P, Gibson PG, Vestbo J, et al. Treatable traits: toward precision medicine of chronic airway diseases. *Eur Respir J*. 2016;47(2):410–9.
 45. Zedan MM, Laimon WN, Osman AM, Zedan MM. Clinical asthma phenotyping: a trial for bridging gaps in asthma management. *World J Clin Pediatr*. 2015;4(2):13–8.
 46. Wu W, Bleecker E, Moore W, Busse WW, Castro M, Chung KF, Calhoun WJ, Erzurum S, Gaston B, Israel E, et al. Unsupervised phenotyping of Severe Asthma Research Program participants using expanded lung data. *J Allergy Clin Immunol*. 2014;133(5):1280–8.
 47. Moore WC, Meyers DA, Wenzel SE, Teague WG, Li H, Li X, D'Agostino R Jr, Castro M, Curran-Everett D, Fitzpatrick AM, et al. Identification of asthma phenotypes using cluster analysis in the Severe Asthma Research Program. *Am J Respir Crit Care Med*. 2010;181(4):315–23.
 48. Siroux V, Basagaña X, Boudier A, Pin I, Garcia-Aymerich J, Vesin A, Slama R, Jarvis D, Anto JM, Kauffmann F, Sunyer J. Identifying adult asthma phenotypes using a clustering approach. *Eur Respir J*. 2011;38(2):310–7.
 49. Leflaudeux D, de Meulder B, Loza MJ, Pepper N, Rowe A, Baribaud F, Bansal AT, Lutter R, Sousa AR, Corfield J, et al. U-BIOPRED clinical adult asthma clusters linked to a subset of sputum omics. *J Allergy Clin Immunol*. 2017;139:1797–807.

50. Loza MJ, Adcock I, Auffray C, Chung KF, Djukanovic R, Sterk PJ, Susulic VS, Barnathan ES, Baribaud F, Silkoff PE, et al. Longitudinally stable, clinically defined clusters of patients with asthma independently identified in the ADEPT and U-BIOPRED asthma studies. *Ann Am Thorac Soc*. 2016;13(Suppl 1):S102–3.
51. Loza MJ, Djukanovic R, Chung KF, Horowitz D, Ma K, Branigan P, Barnathan ES, Susulic VS, Silkoff PE, Sterk PJ, et al. Validated and longitudinally stable asthma phenotypes based on cluster analysis of the ADEPT study. *Respir Res*. 2016;17(1):165.
52. Clemmer GL, Wu AC, Rosner B, McGeachie MJ, Litonjua AA, Tantisira KG, Weiss ST. Measuring the corticosteroid responsiveness endophenotype in asthmatic patients. *J Allergy Clin Immunol*. 2015;136(2):274.e8–81.e8.
53. Belgrave D, Henderson J, Simpson A, Buchan I, Bishop C, Custovic A. Disaggregating asthma: big investigation vs. big data. *J Allergy Clin Immunol*. 2016;139(2):400–7.
54. Bousquet J, Anto JM, Akdis M, Auffray C, Keil T, Momas I, Postma DS, Valenta R, Wickman M, Cambon-Thomsen A, et al. Paving the way of systems biology and precision medicine in allergic diseases: the MeDALL success story. *Allergy*. 2016;71(11):1513–25.
55. Leiria LOS, Martins MA, Saad MJA. Obesity and asthma: beyond TH2 inflammation. *Metabolism*. 2015;64(2):172–81.
56. Bates JHT, Poynter ME, Frodella CM, Peters U, Dixon AE, Suratt BT. Pathophysiology to phenotype in the asthma of obesity. *Ann Am Thorac Soc*. 2017;14(Supplement 5):S395–8.
57. Gonzalez-Barcala FJ, Pertega S, Perez-Castro T, Sampedro M, Sanchez-Lastres J, San-Jose-Gonzalez MA, Bamonde L, Garnelo L, Valdés-Cuadrado L, Moure JD, et al. Obesity and asthma: an association modified by age. *Allergol Immunopathol (Madr)*. 2013;41(3):176–80.
58. Amelink M, De Nijs SB, De Groot JC, Van Tilburg PMB, Van Spiegel PI, Krouwels FH, et al. Three phenotypes of adult-onset asthma. *Allergy*. 2013;68(5):674–80.
59. Ledford DK, Lockey RF. Asthma and comorbidities. *Curr Opin Allergy Clin Immunol*. 2013;13(1):78–86.
60. Yanez A, Cho S-H, Soriano JB, Rosenwasser LJ, Rodrigo GJ, Rabe KF, Peters S, Niimi A, Ledford DK, Katial R, et al. Asthma in the elderly: what we know and what we have yet to know. *World Allergy Organ J*. 2014;7(1):8.
61. Miller RL, Ho SM. Environmental epigenetics and asthma: current concepts and call for studies. *Am J Respir Crit Care Med*. 2008;177(6):567–73.
62. Jacinto T, Malinovschi A, Janson C, Fonseca J, Alving K. Evolution of exhaled nitric oxide levels throughout development and aging of healthy humans. *J Breath Res*. 2015;9(3):36005.
63. Jacinto T, Malinovschi A, Janson C, Fonseca J, Alving K. Differential effect of cigarette smoke exposure on exhaled nitric oxide and blood eosinophils in healthy and asthmatic individuals. *J Breath Res*. 2017;11(3):036006.
64. Halldin CN, Doney BC, Hnizdo E. Changes in prevalence of chronic obstructive pulmonary disease and asthma in the US population and associated risk factors. *Chron Respir Dis*. 2015;12(1):47–60.
65. Sá-Sousa A, Jacinto T, Azevedo LF, Morais-Almeida M, Robalo-Cordeiro C, Bugalho-Almeida A, Bousquet J, Fonseca JA. Operational definitions of asthma in recent epidemiological studies are inconsistent. *Clin Transl Allergy*. 2014;4:24.
66. de Marco R, Cerveri I, Bugiani M, Ferreri M, Verlato G. An undetected burden of asthma in Italy: the relationship between clinical and epidemiological diagnosis of asthma. *Eur Respir J*. 1998;11(3):599–605.
67. Menezes AMB, Perez-Padilla R, Jardim JRB, Muiño A, Lopez MV, Valdivia G, et al. Chronic obstructive pulmonary disease in five Latin American cities (the PLATINO study): a prevalence study. *Lancet*. 2005;366(9500):1875–81.
68. Hanania NA, Noonan M, Corren J, Korenblat P, Zheng Y, Fischer SK, Cheu M, Putnam WS, Murray E, Scheerens H, et al. Lebrikizumab in moderate-to-severe asthma: pooled data from two randomised placebo-controlled studies. *Thorax*. 2015;70(8):748–56.
69. Corren J, Lemanske RF, Hanania NA, Korenblat PE, Parsey MV, Arron JR, Harris JM, Scheerens H, Wu LC, Su Z, et al. Lebrikizumab treatment in adults with asthma. *N Engl J Med*. 2011;365(12):1088–98.
70. Dweik RA, Boggs PB, Erzurum SC, Irvin CG, Leigh MW, Lundberg JO, Olin AC, Plummer AL, Taylor DR; American Thoracic Society Committee on Interpretation of Exhaled Nitric Oxide Levels (FENO) for Clinical Applications. An official ATS clinical practice guideline: interpretation of exhaled nitric oxide levels (FENO) for clinical applications. *Am J Respir Crit Care Med*. 2011;184(5):602–15.
71. Global Strategy for Asthma Management and Prevention 2016. <http://ginasthma.org/2016-gina-report-global-strategy-for-asthma-management-and-prevention/>. Accessed 12 May 2017.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit



ONLINE SUPPLEMENT

Having concomitant asthma phenotypes is common and independently relates to poor lung function in NHANES 2007-2012

Rita Amaral, MSc.^{1,2}, João A. Fonseca, M.D., Ph.D.^{1,3,4}, Tiago Jacinto, Ph.D.^{1,2,4}, Ana M. Pereira, M.D.^{1,4}, Andrei Malinowski, M.D., Ph.D.⁵, Christer Janson, M.D., Ph.D.⁶, Kjell Alving, M.D., Ph.D.⁷

¹CINTESIS- Center for Health Technology and Services Research, Faculty of Medicine, University of Porto, Portugal;

²Dept. of Cardiovascular and Respiratory Sciences, Porto Health School, Porto, Portugal

³MEDCIDS- Dept. of Community Medicine, Information, and Health Sciences: Faculty of Medicine, University of Porto, Portugal;

⁴Dept. of Allergy: Instituto & Hospital CUF, Porto, Portugal;

⁵Dept. of Medical Sciences: Clinical Physiology, Uppsala University, Uppsala, Sweden;

⁶Dept. of Medical Sciences: Respiratory Medicine and Allergology, Uppsala University, Uppsala, Sweden;

⁷Dept. of Women's and Children's Health: Paediatric Research, Uppsala University, Uppsala, Sweden

SUPPLEMENTARY METHODS

Study design

The National Health and Nutrition Examination Survey (NHANES) is a program of studies designed to assess the health and nutritional status of adults and children in the United States. The survey is unique in that it combines interviews and physical examinations.

The NHANES program began in the early 1960s and has been conducted as a series of surveys focusing on different population groups or health topics. The survey examines a nationally representative sample of about 5,000 persons each year. These persons are located in counties across the country, 15 of which are visited each year, to produce reliable statistics, NHANES over-samples persons 60 and older, African Americans, and Hispanics.

The NHANES interview includes demographic, socioeconomic, dietary, and health-related questions. The examination component consists of medical, dental, and physiological measurements, as well as laboratory tests administered by highly trained medical personnel.

All participants visit the physician. Dietary interviews and body measurements are included for everyone. All but the very young have a blood sample taken and will have a dental screening.

Health interviews are conducted in respondents' homes. Health measurements are performed in specially-designed and equipped mobile centers, which travel to locations throughout the country. An advanced computer system using high-end servers, desktop PCs, and wide-area networking collect and process all of the NHANES data. Touch-sensitive computer screens let respondents enter their own responses to certain sensitive questions in complete privacy.

Further details on survey design and respective survey questionnaires can be found in http://www.cdc.gov/nchs/nhanes/about_nhanes.htm.

Variables

Demographic characteristics, such as age, gender, body mass index (BMI), race/ethnicity, and educational status were analyzed:

- *Body mass index (BMI)* was calculated and classified based on the WHO definition (S1): underweight ($\text{BMI} \leq 18.4 \text{ kg/m}^2$); normal ($18.5\text{--}24.9 \text{ kg/m}^2$), overweight ($25\text{--}29.9 \text{ kg/m}^2$), and obese ($\geq 30 \text{ kg/m}^2$);
- *Educational status* was divided in: less than high school ($< \text{high school}$) and completion or greater than high school ($\geq \text{high school}$).

Current smoking was considered if participants had a positive answer to both questions: “Have you smoked ≥ 100 cigarettes during lifetime” and “Do you now smoke cigarettes?”. If participants answered positively to the first question but negatively to the second one, they were considered ex-smokers.

Blood eosinophils (B-Eos) were part of the complete blood counts and assessed on a Beckman Coulter MAXM® instrument (Beckman Coulter, Fullerton, Calif). Blood samples were collected at the NHANES Mobile Examination Center. A detailed description of the laboratory method used can be found elsewhere (S2).

FeNO measurements were also performed at the mobile examination center using the analyzer NIOX MINO® (Aerocrine, Solna, Sweden) (S3) and valid measurements were defined in accordance with ATS/ERS guidelines (S4). *Spirometric* procedures followed the ATS/ERS recommendations (S5), and a minimum of 3 acceptable and reproducible measurements were obtained (S6). After predicted values of basal FEV1 and FEV1/FVC were calculated according to Hankinson et al (S7) with an ethnicity correction (S8), abnormal lung function was defined according to the lower limit of normal (LLN), which classifies subjects with spirometry values below the lower fifth percentile of an aged-matched healthy reference group as abnormal (S9). FeNO and spirometric measurements not fulfilling ATS/ERS recommendations (S4,S5) were excluded (n=653).

Self-reported asthma attacks and asthma-related emergency department (ED) visits (Yes/No), in the past 12 months were analyzed.

Work/school absenteeism was defined as having at least one day lost at work/school due to wheezing (Yes/No).

Asthma symptoms were evaluated with the following questions regarding the last 12 months: “Had wheezing/whistling in your chest?”; “Had disturbed sleep due to wheezing?”; “Had dry hard cough at night, not associated with a cold for at least 14 days in a row?”; “Had wheezing during/after exercise?”; and “Had limited activity due to wheezing?”.

Self-reported rhinitis was defined by an affirmative answer to “During the past 12 months, have you had an episode of hay fever?”.

Use of reliever/rescue medication for asthma was considered if the participant used short-acting β_2 -agonist, anticholinergic, or inhaled corticosteroids (ICS)/formoterol.

Controller medication included: ICS; leukotriene modifiers; long-acting inhaled β_2 -agonist (LABA) and ICS combination; ICS/LABA associated with methylxanthine, cromoglycate, and/or oral corticosteroids. Details on the prescription medication data collection in NHANES 1999–2012 are available elsewhere (S10).

Statistical analysis

A multivariate logistic regression was performed in order to explore the association of concomitant (having at least 2 concurrent) phenotypes with each asthma-related outcomes. Separate models were run using each asthma-related outcome (asthma attack, asthma-related ED visit, ≥ 2 asthma symptoms, work/school absenteeism, rescue medication, and ≥ 2 controller medication and abnormal lung function) as dependent variable and having multiple phenotypes as independent variables.

Model fit was assessed using the *svylogitgof* function for complex survey data (S11).

To create the Venn-Euler diagram, we excluded subjects with incomplete information for at least one of the following: self-reported medical conditions, BMI, smoking history, and FeNO or B-Eos count (listwise deletion of missing data) (S12).

REFERENCES OF SUPPLEMENTARY METHODS

- S1. WHO. Physical status: the use and interpretation of anthropometry. Report of a WHO Expert Committee. *World Health Organization technical report series* 1995;1–452.
- S2. CDC; National Health and Nutrition Examination 2011-2012. Complete Blood Count [updated 2013; accessed 2017 Aug]. Available from:
http://www.cdc.gov/nchs/data/nhanes/nhanes_11_12/cbc_g_met_he.pdf
- S3. CDC; National Health and Nutrition Examination Survey. Respiratory Health ENO Procedures Manual [updated 2011; accessed 2017 Aug]. Available from:
www.cdc.gov/nchs/data/nhanes/nhanes_11_12/Respiratory_Health_ENO_Procedures_Manual.pdf
- S4. ATS/ERS. ATS/ERS Recommendations for Standardized Procedures for the Online and Offline Measurement of Exhaled Lower Respiratory Nitric Oxide and Nasal Nitric Oxide. *Am J Respir Crit Care Med* 2005;**171**(8):912–3
- S5. Miller MR. Standardisation of spirometry. *Eur Respir J* 2005;**26**(2):319–38.
- S6. CDC; National Health and Nutrition Examination Survey. Respiratory Health Spirometry Procedures Manual [updated 2011; accessed 2017 Aug]. Available from:
https://www.cdc.gov/nchs/data/nhanes/nhanes_11_12/spirometry_procedures_manual.pdf
- S7. Hankinson JL, Odencrantz JR, Fedan KB. Spirometric Reference Values from a Sample of the General U.S. Population. *Am J Respir Crit Care Med* 1999;**159**(1):179–87.
- S8. Hankinson JL, Kawut SM, Shahar E, Smith LJ, Stukovsky KH, Barr RG. Performance of American Thoracic Society-Recommended Spirometry Reference Values in a Multiethnic Sample of Adults. *Chest* 2010;**137**(1):138–45.
- S9. Pellegrino R, Viegi G, Brusasco V, Crapo RO, Burgos F, Casaburi R, et al. Interpretative strategies for lung function tests. *Eur Respir J* 2005;**26**(5):948–68.

- S10. CDC; National Health and Nutrition Examination Survey 1988-2012. Prescription Medications - Drug Information [updated 2014; accessed 2017 Aug]. Available from: http://wwwn.cdc.gov/Nchs/Nhanes/1999-2000/RXQ_DRUG.htm
- S11. Archer K, Lemeshow S. Goodness-of-fit test for a logistic regression model fitted using survey sample data. *Stata J* 2006;**6**:97–105.
- S12. Peugh JL, Enders CK. Missing data in educational research: A review of reporting practices and suggestions for improvement. *Review of Educational Research* 2004;**74**:525-55.

Additional file 2: Table S1. Distribution and comparisons between the FeNO and B-Eos cut-offs used in this study, among individuals with current asthma

		B-Eos n (weighed %)			p-value		
Total n (weighed %)		Class I <150/mm ³	Class II 150-300/mm ³	Class III ≥300/mm ³	I vs II	II vs III	I vs III
Current asthma (n= 1,059)		332 (37)	276 (27)	348 (36)			
FeNO	<20ppb	501 (65)	184 (73)	141 (50)	0.12	<0.001	<0.001
	≥20ppb	273 (35)	60 (27)	130 (50)			
	<35ppb	636 (82)	229 (94)	184 (65)	0.02	<0.001	<0.001
	≥35ppb	138 (18)	15 (6)	87 (35)			

FeNO: Fraction of exhaled nitric oxide; B-Eos: blood eosinophils.

P-values <0.05 are presented.

Additional file 3: Table S2. Weighted percentages and comparisons of asthma-related outcomes among subjects with a single asthma phenotype versus: non-classified, and specific combinations of asthma phenotypes.

Weighted %	Total	Asthma attack	Asthma-related ED	≥2 asthma symptoms	Work/school absenteeism	Asthma medication		Lung function		
						≥1 reliever medication	≥2 controller medication	FEV ₁ <LLN	FEV ₁ % predicted§	FEV ₁ /FVC <LLN
Single phenotypes*	43	67	21	59	17	38	14	14	94 (86-97)	23
AwObesity	38	61	27	59	15	31	14	18	92 (84-104)	16
B-Eos-high	22	81	29	70	19	47	15	17	95 (84-99)	27
FeNO-high	8	80	6	31	3	33	28	11	94 (87-96)	55
B-Eos&FeNO-low	30	64	14	55	20	46	9	7	93 (88-102)	20
AwCOPD	2	24	20	93	61	6	23	30	97 (57-97)	30
Non-classified†	14	75	8	59	14	36	15	25	90 (80-103)	23
Multiple phenotypes‡	43	72	23	64	14	49	27	34	83	34

									(74-95)	
AwObesity + others	48	71	23	65	16	48	25	30	81 (74-92)	26
B-Eos-high + others	19	75	20	65	14	51	31	39	80 (69-91)	42
FeNO-high + others	10	68	16	66	17	51	30	37	81 (74-92)	38
B-Eos&FeNO-low + others	9	78	33	59	18	42	18	25	90 (74-100)	21
AwCOPD + others	5	61	22	61	20	66	38	50	74 (63-86)	63
Specific combinations of phenotypes										
B-Eos-high + AwObesity	8	77	25	62	8	46	27	32	85 (77-95)	32
AwObesity+ B- Eos&FeNO-low	7	79	30	64	13	35	16	18	94 (77-101)	9
B-Eos-high + FeNO- high	5	73	19	66	6	44	35	41	80 (69-90)	51
B-Eos-high + FeNO- high + AwObesity	3	80	16	65	40	67	22	33	79 (75-86)	30
FeNO-high + AwObesity	2	26	8	67	12	47	27	32	93 (74-96)	9

B-Eos-high + AwCOPD	1	71	14	73	25	69	4	22	110 (90-110)	43
AwObesity + AwCOPD	1	20	19	67	0	71	12	14	85 (83-85)	14
AwCOPD+ B- Eos&FeNO-low	1	76	44	20	12	84	44	71	66 (63-77)	85

AwObesity: Asthma with obesity; AwCOPD: Asthma with COPD; ED: Emergency-department; FEV₁: Forced expiratory volume in 1 second; FEV₁/FVC: Forced expiratory volume in 1 second and functional vital capacity ratio; LLN: Lower limit of normality.

Values are presented as weighted percentages and significant associations (p<0.05) between multiple and single phenotypes are presented in bold.

*Subjects having only one of the 5 asthma phenotypes: AwObesity, B-Eos-high, FeNO-high, B-Eos&FeNO-low or AwCOPD. † Subjects with non-single and non-multiple phenotypes. ‡Subjects having at least one of the other asthma phenotypes. § Presented as median (Q1-Q3).

Additional file 4: Table S3. Weighted percentages and comparisons of asthma-related outcomes among subjects with a single asthma phenotype versus: non-classified, and specific combinations of asthma phenotypes.

Weighted %	Total	Asthma attack	Asthma-related ED	≥2 asthma symptoms	Work/school absenteeism	Asthma medication		Lung function		
						≥1 reliever medication	≥2 controller medication	FEV ₁ <LLN	FEV ₁ % predicted§	FEV ₁ /FVC <LLN
Single phenotypes *	49	68	22	59	15	40	18	18	92 (82-99)	28
AwObesity	31	61	27	59	15	29	15	18	92 (84-111)	16
Type 2-high†	43	77	23	59	11	46	25	25	90 (80-98)	41
B-Eos&FeNO-low	24	64	14	55	20	45	9	7	93 (88-102)	20
AwCOPD	2	24	20	93	61	11	24	20	97 (57-97)	30
Non-classified‡	13	75	8	59	14	36	15	25	90 (80-103)	23
Multiple phenotypes 	38	71	24	65	19	51	27	32	85 (74-96)	30
AwObesity+ others	26	71	23	67	20	49	26	30	85 (75-96)	26
Type 2-high+ others	19	70	20	68	21	54	32	37	82	36

									(74-93)	
B-Eos&FeNO-low+ others	9	78	33	59	18	42	18	25	90 (74-100)	21
AwCOPD+ others	6	64	23	66	18	66	43	50	74 (63-86)	63
Specific combinations of phenotypes										
Type 2-high+ AwObesity	15	70	22	66	22	52	25	32	83 (76-94)	28
Type 2-high+ AwCOPD	2	74	24	86	11	69	47	18	90 (81-110)	52

AwObesity: Asthma with obesity; AwCOPD: Asthma with COPD; ED: Emergency-department; FEV₁: Forced expiratory volume in 1 second; FEV₁/FVC: Forced expiratory volume in 1 second and functional vital capacity ratio; LLN: Lower limit of normality.

Values are presented as weighted percentages and significant associations (p<0.05) between multiple and single phenotypes are presented in bold.

*Subjects with only one of the 4 asthma phenotypes: AwObesity, Type 2-high, B-Eos&FeNO-low or AwCOPD. †Type 2-high (B-Eos-high or FeNO-high), ‡ subjects with non-single and non-multiple phenotypes. ||Subjects having at least one of the other asthma phenotypes. § Presented as median (Q1-Q3)

Additional file 5: Table S4. Multivariable logistic regression models between each asthma-related outcome and having multiple asthma phenotypes, adjusted for co-variables.

	Asthma attack aOR (95% CI)	Asthma-related ED aOR (95% CI)	≥2 asthma symptoms aOR (95% CI)	Work/school absenteeism aOR (95% CI)	≥1 reliever medication* aOR (95% CI)	FEV ₁ /FVC <LLN aOR (95% CI)
Multiple vs. single phenotype	1.27 (0.78-2.06)	1.17 (0.74-1.86)	1.26 (0.76-2.11)	0.79 (0.37-1.68)	1.55 (0.97-2.49)	1.74 (0.94-3.24)
Female	1.34 (0.87-2.07)	2.05 (1.08-3.90)	1.96 (1.16-3.31)	1.35 (0.66-2.79)	0.88 (0.53-1.49)	0.53 (0.31-0.90)
Age ≥ 40 yrs	0.80 (0.50-1.27)	0.91 (0.59-1.41)	0.94 (0.55-1.60)	1.09 (0.60-2.00)	1.08 (0.67-1.73)	1.50 (0.90-2.49)
Caucasian vs. other	0.89 (0.54-1.46)	0.38 (0.23-0.64)	0.72 (0.46-1.15)	0.76 (0.39-1.50)	1.03 (0.65-1.63)	0.92 (1.16-3.51)
Current smoker vs. non-/ex-smokers	0.89 (0.55-1.43)	1.60 (0.94-2.72)	1.22 (0.69-2.14)	0.82 (0.37-1.83)	1.95 (1.35-2.83)	2.02 (1.16-3.51)
Rhinitis	0.96 (0.59-1.57)	0.77 (0.38-1.56)	0.88 (0.50-1.52)	0.70 (0.32-1.53)	0.66 (0.42-1.04)	1.04 (0.62-1.75)
Goodness-of-fit test						
χ ² (p-value)	0.59 (0.80)	0.59 (0.81)	1.33(0.22)	1.27 (0.25)	1.14 (0.33)	3.05 (0.002)

CI: confidence interval; aOR: adjusted odds ratio; ED: emergency-department; FEV₁/FVC: Forced expiratory volume in one second and forced vital capacity ratio; LLN: Lower Limit of Normal; χ²: Chi-square.

All asthma-related outcomes were analyzed separately and treated as dependent variables. The aOR values with p<0.05 are presented in bold.

*Short-acting β₂-agonist and/or anticholinergic.




Figure S1: Proportions (weighted to the US population) of subjects taking asthma controller medications stratified into the different phenotypes, among all participants included for asthma phenotype classification (left) and only in those with a single phenotype (right). P-values <0.05 were indicated. NA: Non-applicable (not possible to determine because some participants had both B-Eos-high and FeNO-high asthma phenotypes).