



COMMUNICATION

Structural Basis for the Activation of Muscle Contraction by Troponin and Tropomyosin

William Lehman¹, Agnieszka Galińska-Rakoczy¹, Victoria Hatch^{1*},
Larry S. Tobacman² and Roger Craig³

¹*Department of Physiology and Biophysics, Boston University School of Medicine, 72 East Concord Street, Boston, MA 02118, USA*

²*Departments of Medicine and Physiology and Biophysics, University of Illinois at Chicago, 840 South Wood Street, Chicago, IL 60612, USA*

³*Department of Cell Biology, University of Massachusetts Medical School, 55 Lake Avenue North, Worcester, MA 01655, USA*

Received 11 February 2009;
received in revised form
23 March 2009;
accepted 25 March 2009
Available online
31 March 2009

The molecular regulation of striated muscle contraction couples the binding and dissociation of Ca^{2+} on troponin (Tn) to the movement of tropomyosin on actin filaments. In turn, this process exposes or blocks myosin binding sites on actin, thereby controlling myosin crossbridge dynamics and consequently muscle contraction. Using 3D electron microscopy, we recently provided structural evidence that a C-terminal extension of TnI is anchored on actin at low Ca^{2+} and competes with tropomyosin for a common site to drive tropomyosin to the B-state location, a constrained, relaxing position on actin that inhibits myosin-crossbridge association. Here, we show that release of this constraint at high Ca^{2+} allows a second segment of troponin, probably representing parts of TnT or the troponin core domain, to promote tropomyosin movement on actin to the Ca^{2+} -induced C-state location. With tropomyosin stabilized in this position, myosin binding interactions can begin. Tropomyosin appears to oscillate to a higher degree between respective B- and C-state positions on troponin-free filaments than on fully regulated filaments, suggesting that tropomyosin positioning in both states is troponin-dependent. By biasing tropomyosin to either of these two positions, troponin appears to have two distinct structural functions; in relaxed muscles at low Ca^{2+} , troponin operates as an inhibitor, while in activated muscles at high Ca^{2+} , it acts as a promoter to initiate contraction.

© 2009 Elsevier Ltd. All rights reserved.

Edited by J. Karn

Keywords: actin; troponin; tropomyosin; calcium; electron microscopy

Contraction in all muscles results from the relative sliding of thick and thin filaments. The process is driven by the myosin-crossbridge motors projecting from thick filaments and interacting cyclically with actin subunits on the thin filament molecular track. In skeletal and cardiac muscles, contraction is switched on and off by changes in sarcoplasmic free Ca^{2+} concentration, and by the corresponding binding and dissociation of Ca^{2+} from the troponin

complex, located on thin filaments. Crossbridge dynamics is a function of the myosin ATPase, which itself is activated by myosin-crossbridge binding to actin. Thus, any modulation of actin-activation would, in turn, serve to regulate contraction. Indeed, at low Ca^{2+} , actin–myosin interaction is inhibited because myosin binding sites on actin become inaccessible. It is generally recognized that this inhibition occurs when Ca^{2+} -free troponin impinges on elongated tropomyosin to then block myosin binding.^{1,2} In contrast, the involvement of troponin in muscle activation is less well characterized. The crossbridge cycle may turn on simply because the troponin constraint is released at high Ca^{2+} and tropomyosin defaults to an unstrained, non-blocking position on actin. Alternatively, Ca^{2+} -saturated troponin may have a more dynamic role and facilitate regulatory switching by actively promoting tropo-

*Corresponding author. E-mail address: wlehman@bu.edu.

Abbreviations used: TnI, the inhibitory subunit of troponin; cTerm-TnI, the C-terminal 80 amino acid domain of TnI that links to actin at low Ca^{2+} ; TnC, the Ca^{2+} -sensor of troponin that releases inhibition; TnT, the element linking troponin to tropomyosin; B-state, the blocked state; C-state, the closed state; M-state, the open state.

myosin movement away from the blocking position. We have investigated these possibilities using electron microscopy and 3D reconstruction of thin filaments, and report here on troponin's structural influence over tropomyosin at high Ca^{2+} .

Tropomyosin is a ~40 nm long "coiled-coil" α -helical protein, which lies along the long-pitch double helical array of actin monomers on thin filaments.³⁻⁵ Tropomyosin molecules associate together in an end-to-end fashion,⁶ and form a continuous strand, with each tropomyosin spanning seven successive actin molecules. This arrangement is possible because tropomyosin possesses a series of seven quasi-repeating motifs designed to bind to neighboring actin monomers along filaments.⁷⁻⁹ In turn, troponin, consisting of three subunits (TnT, TnI, and TnC), binds to tropomyosin at specific points along the tropomyosin molecule;¹⁰⁻¹³ troponin complexes therefore assume the 40 nm periodicity of tropomyosin on thin filaments. Troponin is thought to have multiple, compartmentalized functions, with each subunit having a particular role in binding tropomyosin or Ca^{2+} or in inhibiting actomyosin ATPase.¹⁴ TnC is well characterized and functions as the Ca^{2+} receptor. After binding Ca^{2+} , it neutralizes the inhibition of actomyosin ATPase imposed by TnI (the inhibitory subunit). TnT, a fairly long asymmetric molecule (~19 nm),¹⁵ links the entire troponin complex to tropomyosin.^{14,16,17} The N-terminal "tail" of TnT binds alongside tropomyosin on thin filaments, bridging the head-to-tail joint between adjacent tropomyosin molecules. However, the C-terminal part of TnT converges on TnC and TnI, and interacts with them, forming the core domain of troponin.^{18,19} Other than providing a scaffold for TnI and TnC, the actions of TnT near or within the core domain are obscure. The view of TnT solely as a structural intermediary between troponin subunits and tropomyosin may be an oversimplification. In this regard, biochemical approaches have shown that in concert with tropomyosin and the rest of the troponin complex, TnT enhances actomyosin ATPase at high Ca^{2+} ,^{2,20,21} suggesting that it and the troponin core domain have more than just a permissive role during activation. Thus, troponin subunits may have a dual role in thin filament regulation, with TnI being linked to inhibition at low Ca^{2+} , and TnT and the rest of the complex linked to activation at high Ca^{2+} .²²

In the complete absence of other factors, tropomyosin is thought to be able to oscillate laterally over a narrow region of the flat surface of actin.²³⁻²⁸ It is generally assumed, although not explicitly established, that this intrinsic ability to shift azimuthally around actin filaments at low energy cost is inherent to and necessary for the thin filament regulatory mechanism.²³⁻²⁵ Presumably, tropomyosin location becomes biased towards specific regulatory positions on actin^{25,29} in the presence of troponin and/or myosin, and depending on levels of Ca^{2+} binding to troponin and myosin binding to actin. At low Ca^{2+} , tropomyosin localizes over the outer domain of actin (on actin subdomains 1 and 2,

covering myosin binding sites; the B-state position). At high Ca^{2+} , tropomyosin moves to actin's inner domain (to the edge of subdomains 3 and 4, exposing most but not all of the myosin binding site; the C-state position), and after myosin binding it moves further onto the inner domain (exposing the entire myosin binding site; the M-state position).²⁹ Thus, the Ca^{2+} -induced movement of tropomyosin is thought to increase the probability of myosin binding, and the resulting myosin interaction leads to a further tropomyosin shift and full activation of the thin filament.²⁹

We showed recently that the C-terminal end of TnI (cTerm-TnI) drives tropomyosin to the B-state blocking position on the actin outer domain at low Ca^{2+} .^{30,31} Here, we have extended these studies to characterize how troponin influences tropomyosin in the presence of Ca^{2+} . Our results demonstrate that a troponin extension, likely to involve TnT and/or parts of the troponin core domain complex, promotes tropomyosin movement away from the blocking position and stabilizes it in the C-state position. Our work therefore supports the view that troponin exhibits dual function, inhibiting actin-myosin interaction at low Ca^{2+} and facilitating interaction at high Ca^{2+} . Our results suggest further that troponin transforms tropomyosin's fundamentally ambiguous position on striated muscle actin by dampening tropomyosin's oscillatory behavior, while actively promoting its movement to B- or C-state configurations.

3D reconstruction of Ca^{2+} -treated thin filaments

Thin filaments were reconstituted from F-actin, cardiac troponin and tropomyosin under conditions known to saturate the filaments with the regulatory proteins.^{25,26} Filaments were negatively stained,^{25,26} and EM images of the reconstituted thin filaments showed characteristic double-helical arrays of actin monomers, tropomyosin strands, and troponin densities repeating with a 40 nm periodicity (Fig. 1). The visually apparent, in-register binding of troponin on each helical strand of F-actin at 40 nm intervals has been directly quantified by analysis of class averaged 2D projections of these same filaments,³² corroborating that troponin-tropomyosin binds with the same structural arrangement and molar stoichiometry displayed by native thin filaments isolated directly from muscle. Troponin densities were absent from actin-tropomyosin controls (Fig. 1).

Helical reconstructions of Ca^{2+} -treated filaments showed actin subunits and densities that were attributable to tropomyosin, as observed earlier (Fig. 2c and d). As expected, the longitudinally continuous tropomyosin strands were well defined and localized over the outer part of the inner domain of actin. This position is distinctly different from that assumed by tropomyosin on the actin outer domain in reconstructions of EGTA-treated filaments (Fig. 2e and f; cf. Refs. 25 and 26). In an




Fig. 1. Electron micrographs of negatively stained filaments. (a) F-actin-troponin-tropomyosin; (b) F-actin-tropomyosin. Note that the presence of troponin (arrowheads), distributed with a 40 nm periodicity in a increases the maximum width of thin filaments; also note the obliquely oriented tropomyosin strands in a and b (arrows). Filaments are shown with their pointed ends facing up; polarity was determined with alignment tools as described.³⁶ The scale bar represents 50 nm. *Preparation of proteins:* F-actin, bovine cardiac troponin and tropomyosin were purified as described.³⁷ SDS-PAGE indicated that the cardiac tropomyosin was >90% $\alpha\alpha$ -isoform and the remainder was $\alpha\beta$ -isoform; it was presumed to be 20–30% phosphorylated, as is tropomyosin in the adult cardiac tissue of other mammals.^{38,39} The cardiac troponin as isolated was monophosphorylated (not biphosphorylated as occurs after adrenergic stimulation). *EM and image processing:* Earlier, we showed that a moderate excess of troponin-tropomyosin relative to actin is needed to saturate filaments at the low concentrations of actin required for successful EM work.^{25,26} Filaments were thus prepared by mixing a twofold excess of tropomyosin or troponin-tropomyosin (6 μ M) with F-actin (20 μ M) to optimize binding, in 100 mM NaCl, 3 mM $MgCl_2$, 1 mM NaN_3 , 0.2 mM EGTA, 1 mM dithiothreitol, 5 mM sodium phosphate/5 mM Pipes buffer (pH 7.0) at

25°C.^{25,26} The mixture was diluted 20-fold, applied to carbon-coated grids and negatively stained with 1% (w/v) uranyl acetate.^{25,26} Filaments from two preparations of F-actin-troponin-tropomyosin and three preparations of F-actin-tropomyosin were recorded and processed in these studies. EM was done with a Philips CM120 instrument at a magnification of 60,000 $\times$ under low-dose conditions (~ 12 e $^-$ /Å). Helical reconstruction³⁶ was performed as described.^{25,26} Helical reconstruction was particularly useful for identification of the relatively low-density troponin signal observed, as the method takes advantage of layer line indexing information to filter out noise, which otherwise would have obscured the troponin density observed. In addition, robust statistical programs accompany the helical reconstruction package, allowing the significance of weak densities to be assessed by Student's *t*-test.^{40,41} Real space reconstruction methods,⁴² treating filament segments as single particles, were used to confirm the results obtained by helical reconstruction and to attempt to sort filament segments according to their tropomyosin positional modes, as described.²⁶

effort to identify material derived from troponin, the threshold density cutoff was reduced to 1.5–3.0 standard deviation units above the mean density (normally >5.0 σ is used). Additional densities, not detected previously, were then evident in the maps of high Ca^{2+} filaments, but were absent at low Ca^{2+} at the same or considerably lower threshold density levels. The densities emerged from the center of actin subdomain 1 and traversed the face of the domain in a path that approached and then abutted the C-state tropomyosin (Fig. 2d); i.e. they bridged sites on actin normally occupied by B-state tropomyosin at low Ca^{2+} . As these densities were not

seen in actin-tropomyosin controls (Fig. 2b), we attribute them to troponin. As they were not seen in reconstructions of low Ca^{2+} -treated troponin-tropomyosin filaments (Fig. 2e) either, we ascribe them to a part of troponin that is visualized only in the Ca^{2+} -saturated conformation.

Densities specific to the high Ca^{2+} filaments are statistically significant

Helical reconstruction is an averaging technique that treats each successive actin molecule along thin filaments with its associated proteins as equivalent




Fig. 2. Surface views of thin filament reconstructions showing the position of the high- Ca^{2+} troponin extension. Reconstructions of: (a) F-actin (cyan, subdomains numbered on one actin subunit); and (b) F-actin-tropomyosin (green) controls; (c-e) F-actin-troponin-tropomyosin: (c and d) high Ca^{2+} (gold); (e), low Ca^{2+} (maroon). Tropomyosin positions are indicated by arrows. High Ca^{2+} reconstructions are displayed at two density cutoff thresholds; (c) 5σ and (d) 2.5σ greater than the mean density. Note the presence of extra densities in d emerging from the centers of subdomain 1, which traverse the face of the domain to about C-state tropomyosin (open arrowheads). Note also that these densities are absent from F-actin-tropomyosin controls (b) and low- Ca^{2+} F-actin-troponin-tropomyosin (e) (each displayed at 2.5σ). (f) Reconstructions in (d) and (e) superimposed for comparison. Note that the respective tropomyosin strands occupy different positions, with tropomyosin from the high- and low- Ca^{2+} maps in C- and B-states. Also note that Ca^{2+} -specific extra density attributed to troponin (open arrows) crosses the B-state tropomyosin position. All of the reconstructions of thin filaments obtained were aligned to each other and are shown with filament pointed ends facing up. Each map shows densities with amplitudes that are at a threshold of 2.5σ greater than the mean density (with the exception of the map in (c)).

units. Thus well-ordered troponin domains present only on every seventh actin along thin filaments will distribute as if they derived from every single actin and would be expected to appear so in reconstructions. The distribution process dilutes the troponin

signal, which becomes comparatively low in amplitude relative to actin and tropomyosin. In this case, the amplitude of each extra density observed on every actin molecule in the reconstructions of Ca^{2+} -treated filaments at best may represent one-seventh




Fig. 3. Statistical significance of densities contributing to reconstructions of Ca^{2+} -treated F-actin-troponin-tropomyosin. (a) A z-section of the reconstruction shown in Fig. 2d (gold). Note the actin subdomains (numbered) and tropomyosin positions (indicated by arrows and labeled Tm) on the inner domain of actin over subdomains 3 and 4. Note also the extra density emerging from subdomain 1 and connecting to tropomyosin (indicated by open arrowhead; the outer edge of the extra density is highlighted in cyan). (b) Densities associated with the z-section in (a) that have a confidence level of $>99\%$ (pink). (c) The z-section in (a) superimposed on the statistical map in (b) showing that all densities in (a) including the extra density attributable to troponin, have a high level of statistical significance.

that of the original troponin mass. Nonetheless, the troponin densities observed in the reconstructions were quite robust and displayed high statistical significance at greater than the 99% confidence level (Fig. 3).

The position of striated muscle tropomyosin is indeterminate in the absence of troponin

In order to assess the impact of troponin on the structure of thin filaments further, we examined the structural interactions of tropomyosin on troponin-free actin, particularly since biochemical studies suggest that cardiac tropomyosin binding is weakened when troponin is absent.²⁰ Helical reconstruction of 500–1000 nm stretches of actin–tropomyosin (free of troponin) revealed that the mean location of tropomyosin was midway between the B- and C-state configurations, possibly reflecting an assemblage of filaments with variable tropomyosin position (Fig. 4a and b). Inspection of reconstructions of the individual filaments contributing to the average map confirmed this possibility. About half of the filament reconstructions analyzed could be clearly identified by eye (as well as by fitting protocols) as either being distinctly B- or C-state examples of tropomyosin association. Dividing these two filament classes into separate data sets and averaging each set independently yielded two distinctly different reconstructions, where C- and B-state modes were each indistinguishable from those displayed by troponin–tropomyosin filaments at corresponding

high or low Ca^{2+} (Fig. 4c and d; cf. Refs. 25–27). This demonstrates that approximately half of the filaments possessed long stretches that, on average, were in one or another well-defined regulatory state. Hence, a mixed population of B- and C-state conformations does coexist in these cardiac filament preparations, explaining, in part, the diffuse localization of cardiac muscle tropomyosin in averaged data.²⁶ The more ambiguous positioning of tropomyosin, seemingly between B- and C-state positions on the remainder of the filaments, may reflect a lack of correlation between the tropomyosin position on the two opposing actin helical strands and/or a disordering of tropomyosin position resulting from a dynamic fluctuation between states.

Previous single-particle reconstructions of 40 nm long segments of actin–cardiac tropomyosin filaments suggested a similar positional ambiguity for tropomyosin.²⁶ Attempts were made to classify segments from the above set of filaments showing poorly localized tropomyosin. Cross-correlation against B-, C- or intermediate state models did not succeed in sorting the data into demonstrably separate categories, and the reconstructions generated showed no unique tropomyosin positional state. Thus, single-particle methods and helical analysis both identified a set of filaments characterized by an apparent local disordering of cardiac tropomyosin on actin in the absence of troponin.

By contrast, marked variability in tropomyosin position has not been observed for preparations of filaments composed of cardiac troponin–




Fig. 4. Reconstructions of troponin-free actin–tropomyosin. (a) A z-section of the reconstruction shown in Fig. 2b of cardiac tropomyosin bound to F-actin (no troponin). Note that tropomyosin (arrows) occupies a position midway between the inner and outer domains of actin, i.e. between actin subdomains 1 and 3 on one side of the filament and between subdomains 2 and 4 on the other side. (b) Tropomyosin densities determined by subtracting densities in the F-actin reconstruction from those of the map in a and then superimposing them on the corresponding z-section of F-actin in order to highlight the position of tropomyosin. (c and d) The filaments contributing to the average density map in a were sorted according to their best alignment to B- or C-state models. (c) Reconstruction of the sorted filaments

that were in the C-state mode; note tropomyosin densities on the actin inner domain (i.e. associated with actin subdomains 3 and 4). (d) Reconstruction of the sorted filaments that were in the B-state mode, with tropomyosin densities on the outer domain (i.e. associated with subdomains 1 and 2). Comparable results were obtained when the skeletal muscle isoform of tropomyosin was examined on troponin-free F-actin, viz. an average position midway between B- and C-state, but here individual filament reconstructions showed a slightly greater bias for the C-state mode than found with cardiac tropomyosin.

tropomyosin; without exception, helical reconstruction showed that, on average, tropomyosin along all single filaments examined at low Ca^{2+} could be readily classified as belonging to the B-state mode and analogously tropomyosin along all filaments in high Ca^{2+} to the C-state configuration. Similarly, single-particle analysis of short segments of these filaments indicated that while localized oscillation between positional modes appears to occur, tropomyosin occupied characteristic regulatory state positions along roughly 80 % of filament lengths.²⁶

Different tropomyosin isoforms interact differently on F-actin

The precision with which tropomyosin is positioned is not simply a matter of troponin being present or absent, but rather depends also on the isoform of tropomyosin examined. For example, polymorphic positioning of tropomyosin was not observed here or previously^{25,33} on thin filaments containing smooth muscle tropomyosin (without troponin), where a single mode of binding interaction was detected. In fact, the variance associated with densities connecting cardiac tropomyosin to actin is higher than that for the smooth muscle tropomyosin connection, particularly near subdomain 1 (Fig. 5).

The location of tropomyosin on actin is defined by weak electrostatic contacts.⁸ In turn, both the collective strength of these weak interactions and the stiffness of the tropomyosin strand on actin will influence tropomyosin's responsiveness to mechanical or chemical perturbations. Maytum *et al.* likened the behavior of different tropomyosin isoforms to variably vibrating guitar strings being held taut or not, where the additional displacement of the guitar strings by a pick, would be the counterpart of tro-

ponin or other proteins disturbing preset average equilibrium positions.²⁸ Thus, in a case in which tropomyosin is well localized on actin, for example in smooth muscle filaments,²⁵ the tropomyosin in effect is taut and well positioned by electrostatics. Non-muscle tropomyosins, also well localized on actin, could be similarly described.^{25,28,34} A high degree of chemomechanical specificity may be required in these cases of troponin-free filaments in order to maximally stabilize tropomyosin on actin. In contrast, we show here that the position of cardiac striated muscle tropomyosin on actin is less well defined, as if the isoform were held more loosely on actin, at least in the absence of troponin (cf. Ref. 21). Given this extra degree of plasticity, limited as it might be to yielding azimuthal oscillations amounting to between 15 Å and 25 Å,²⁷ the behavior of tropomyosin on actin filaments may be more easily fine-tuned by troponin.

We demonstrated earlier that the mechanochemical equilibrium balance of tropomyosin that determines its positional state is easily perturbed by small changes in electrostatic interactions between actin and tropomyosin.²⁵ We showed also that the content of $\alpha\alpha$ -, $\beta\beta$ -, or $\alpha\beta$ -tropomyosin isoforms may strongly influence the equilibrium balance of tropomyosin on actin.²⁵ Less well characterized variation in the end-to-end contacts between successive tropomyosins along tropomyosin and their degree of phosphorylation may also influence tropomyosin interactions with actin. Whether such subtleties affect tropomyosin positioning *in vivo* and are physiologically meaningful is not known.

Troponin promotes tropomyosin movement on actin

We indicated recently that at low Ca^{2+} , a C-terminal domain of TnI (cTerm-TnI) binds to actin,




Fig. 5. Comparison of cardiac and smooth muscle tropomyosin positions on actin. z-Sections of reconstructions of F-actin combined with cardiac muscle tropomyosin (a) and aortic smooth muscle tropomyosin (b). Note the closer association of the smooth muscle tropomyosin density with the actin outer domain (compare white arrows in (a) and (b)). (c) Maps plotting the variance associated with the contributing density points^{40,41} in the reconstruction in a show that a high variance (red) occurs over the site where tropomyosin normally contacts actin in the B-state. (d) In contrast, corresponding variance maps of actin – smooth muscle tropomyosin reconstructions show low variance at this site and only comparably high variance (red) at the very edges of the tropomyosin density.




Fig. 6. A cartoon representation of the organization of the thin filament at high Ca^{2+} . As previously,³¹ the troponin core domain complexes on either side of F-actin are depicted as W-shaped TnIT structures supporting dumbbell-shaped TnC;^{18,19} actin, gray; tropomyosin, salmon; TnI, cyan; TnC, red; TnT, yellow. At high Ca^{2+} , cTerm-TnI dissociates from actin and binds to the N-terminal lobe of TnC. We argue that this releases chemomechanical constraints on troponin and tropomyosin. Troponin then unfurls, here depicted for simplicity as TnT propelling tropomyosin to the C-state position. Other related depictions of troponin on thin filaments can be found in the literature.^{18,19,43}

displaces tropomyosin from the C-state and stabilizes it in the B-state position.³¹ This would account for the unambiguous position of tropomyosin in low Ca^{2+} -treated thin filaments. We suggest here that at high Ca^{2+} , C-terminal domains of TnT or parts of the troponin core domain complex move over the B-state binding sites, promote tropomyosin movement and then stabilize tropomyosin in the C-state, thus accounting for tropomyosin's positional fidelity in Ca^{2+} -treated filaments. Hence, the presence of troponin appears to diminish the positional promiscuity of cardiac tropomyosin at both low and high Ca^{2+} . We also indicated earlier that the N-terminal TnT tail domain stabilizes tropomyosin in a B-state position.³⁵ Thus, the TnT tail alone cannot promote the transition of tropomyosin to the C-state at high Ca^{2+} . Clearly, additional parts of troponin, whose domain structures are yet to be identified, are required for inducing tropomyosin movement to the C-state.

We have suggested that the blocked, B-state is brought about and muscle relaxation ensues at low Ca^{2+} because tropomyosin is wedged in an inhibitory position between the cTerm-TnI regulatory domain on one side, and TnT and the troponin core domain on the other.³¹ This configuration may compress the C-terminal end of TnT against the troponin core domain complex. At low Ca^{2+} , the core domain may also become constricted and thus more limited in its binding to targets on actin. It follows that once cTerm-TnI dissociates from actin at high Ca^{2+} , such strain will be released, thrusting tropomyosin toward the C-state position (Fig. 6). The extra density seen in high Ca^{2+} reconstructions is likely to represent an obliquely oriented segment of troponin that associates closely with actin and drives this process. Comparable densities are not resolved from tropomyosin in low Ca^{2+} reconstructions, possibly because they may be compressed

linearly against the elongated protein or become disordered by a radial displacement from actin.

Our results support the hypothesis that troponin is intimately involved in both the inhibition and the activation of muscle contraction. We propose that the counterpunching action of the mobile parts of troponin determines the positions of tropomyosin at low and high Ca^{2+} , thereby regulating the thin filament. We argue further that the inherent positional ambiguity of striated muscle tropomyosin on actin is an adaptation attuned to troponin function. While we have not defined how the entire troponin complex reconfigures during muscle activation and inhibition, our results indicate that key structural domains of troponin bring about striking steric effects that control actin-activation of myosin ATPase and consequently contraction.

Acknowledgements

We thank Ms Karen Moore (Moore Design) for artwork in Fig. 6. This research was supported by grants from the National Institutes of Health to W.L. (HL36153, HL86655), L.S.T. (HL38834, HL63774), and R.C. (AR34711, RR08426).

References

1. Gordon, A. M., Homsher, E. & Regnier, M. (2000). Regulation of contraction in striated muscle. *Physiol. Rev.* **80**, 853–924.
2. Tobacman, L. S. (1996). Thin filament-mediated regulation of cardiac contraction. *Annu. Rev. Physiol.* **58**, 447–481.
3. Moore, P. B., Huxley, H. E. & DeRosier, D. J. (1970). Three-dimensional reconstruction of F-actin, thin

- filaments and decorated thin filaments. *J. Mol. Biol.* **50**, 279–292.
4. O'Brien, E. J., Bennett, P. M. & Hanson, J. (1971). Optical diffraction studies of myofibrillar structure. *Phil. Trans. Roy. Soc. B*, **261**, 201–208.
 5. Spudich, J. A., Huxley, H. E. & Finch, J. T. (1972). The regulation of skeletal muscle contraction. II. Structural studies of the interaction of the tropomyosin-troponin complex with actin. *J. Mol. Biol.* **72**, 619–632.
 6. Greenfield, N. J., Huang, Y. J., Swapna, G. V., Bhattacharya, A., Rapp, B., Singh, A. *et al.* (2006). Solution NMR structure of the junction between tropomyosin molecules: implications for actin binding and regulation. *J. Mol. Biol.* **364**, 80–96.
 7. McLachlan, A. D. & Stewart, M. (1976). The 14-fold periodicity in α -tropomyosin and the interaction with actin. *J. Mol. Biol.* **103**, 271–298.
 8. Lorenz, M., Poole, K. J. V., Popp, D., Rosenbaum, G. & Holmes, K. C. (1995). An atomic model of the unregulated thin filament obtained by X-ray fiber diffraction on oriented actin-tropomyosin gels. *J. Mol. Biol.* **246**, 108–119.
 9. Hitchcock-DeGregori, S. E. & An, Y. (1996). Integral repeats and continuous coiled coil are required for binding of striated muscle tropomyosin to the regulated actin filament. *J. Biol. Chem.* **271**, 3600–3603.
 10. Ohtsuki, I., Masaki, T., Nonomura, Y. & Ebashi, S. (1967). Periodic distribution of troponin along the thin filament. *J. Biochem. (Tokyo)*, **61**, 817–819.
 11. Ebashi, S. & Endo, M. (1968). Calcium ion and muscle contraction. *Prog. Biophys. Mol. Biol.* **28**, 123–183.
 12. Cohen, I. & Cohen, C. (1972). A tropomyosin-like protein from human platelets. *J. Mol. Biol.* **68**, 383–387.
 13. Ohtsuki, I. (1974). Localization of troponin in thin filaments and in tropomyosin paracrystals. *J. Biochem. (Tokyo)*, **75**, 753–765.
 14. Greaser, M. & Gergely, J. (1971). Reconstitution of troponin activity from three protein components. *J. Biol. Chem.* **246**, 4226–4233.
 15. Flicker, P. F., Phillips, G. N. & Cohen, C. (1982). Troponin and its interactions with tropomyosin: an electron microscope study. *J. Mol. Biol.* **162**, 495–501.
 16. Hitchcock, S. E., Huxley, H. E. & Szent-Györgyi, A. G. (1973). Calcium sensitive binding of troponin to actin-tropomyosin: a two-site model for troponin action. *J. Mol. Biol.* **80**, 825–836.
 17. Potter, J. D. & Gergely, J. (1974). Troponin, tropomyosin and actin interactions in the Ca^{2+} regulation of muscle contraction. *Biochemistry*, **13**, 2697–2703.
 18. Takeda, S., Yamashita, A., Maeda, K. & Maeda, Y. (2003). Structure of the core domain of human cardiac troponin in the Ca^{2+} -saturated form. *Nature*, **424**, 35–41.
 19. Vinogradova, M. V., Stone, D. B., Malanina, G. G., Karatzaferi, C., Cooke, R., Mendelson, R. A. *et al.* (2005). Ca^{2+} -regulated structural changes in troponin. *Proc. Natl Acad. Sci. USA*, **102**, 5038–5043.
 20. Daniya, R., Butters, C. A. & Tobacman, L. S. (1994). Equilibrium linkage analysis of cardiac thin filament assembly. Implications for the regulation of muscle contraction. *J. Biol. Chem.* **269**, 29457–29461.
 21. Potter, J. D., Sheng, Z., Pan, B.-S. & Zhao, J. (1995). A direct regulatory role for troponin T and a dual role for troponin C in the Ca^{2+} regulation of muscle contraction. *J. Biol. Chem.* **270**, 2557–2562.
 22. Gomes, A. V., Potter, J. D. & Szczesna-Cordary, D. (2004). Molecular and cellular aspects of troponin cardiomyopathies. *Ann. N.Y. Acad. Sci.* **1015**, 214–224.
 23. McKillop, D. F. A. & Geeves, M. A. (1993). Regulation of the interaction between actin and myosin subfragment-1: Evidence for three states of the thin filament. *Biophys. J.* **65**, 693–701.
 24. Lehrer, S. S. & Geeves, M. A. (1998). The muscle thin filament as a classical cooperative/allosteric regulatory system. *J. Mol. Biol.* **277**, 1081–1089.
 25. Lehman, W., Hatch, V., Korman, V., Rosol, M., Thomas, L., Maytum, R. & *et al.* (2000). Tropomyosin and actin isoforms modulate the localization of tropomyosin strands on actin filaments. *J. Mol. Biol.* **302**, 593–606.
 26. Pirani, A., Xu, C., Hatch, V., Craig, R., Tobacman, L. S. & Lehman, W. (2005). Single particle analysis of relaxed and activated muscle thin filaments. *J. Mol. Biol.* **346**, 761–772.
 27. Poole, K. J., Lorenz, M., Evans, G., Rosenbaum, G., Pirani, A., Tobacman, L. S. *et al.* (2006). A comparison of muscle thin filament models obtained from electron microscopy reconstructions and low-angle X-ray fibre diagrams from non-overlap muscle. *J. Struct. Biol.* **155**, 273–284.
 28. Maytum, R., Hatch, V., Konrad, M., Lehman, W. & Geeves, M. A. (2008). Ultra short yeast tropomyosins show novel myosin regulation. *J. Biol. Chem.* **283**, 1902–1910.
 29. Vibert, P., Craig, R. & Lehman, W. (1997). Steric-model for activation of muscle thin filaments. *J. Mol. Biol.* **266**, 8–14.
 30. Pirani, A., Vinogradova, M. V., Curmi, P. M. G., King, W. A., Fletterick, R. J., Craig, R. *et al.* (2006). An atomic model of the thin filament in the relaxed and Ca^{2+} -activated states. *J. Mol. Biol.* **357**, 707–717.
 31. Galińska-Rakoczy, A., Engel, P., Xu, C., Jung, H., Craig, R., Tobacman, L. S. & Lehman, W. (2008). Structural basis for the regulation of muscle contraction by troponin and tropomyosin. *J. Mol. Biol.* **379**, 929–935.
 32. Paul, D., Lehman, W., Pirani, A., Craig, R., Tobacman, L. S., Squire, J. M. & Morris, E. P. (2009). Reference free single particle analysis of reconstituted thin filaments. *Proceedings of the Biophysical Society 53rd Annual Meeting*. Biophys. J. **96**, 376a (Cell Press, Cambridge, MA).
 33. Hodgkinson, J. L., Marston, S. B., Craig, R., Vibert, P. & Lehman, W. (1997). Three-dimensional image reconstruction of reconstituted smooth muscle thin filaments: effects of caldesmon. *Biophys. J.* **72**, 2398–2404.
 34. Skoumpla, K., Coulton, A. T., Lehman, W., Geeves, M. A. & Mulvihill, D. P. (2007). Acetylation regulates tropomyosin function in the fission yeast *Schizosaccharomyces pombe*. *J. Cell Sci.* **120**, 1635–1645.
 35. Tobacman, L. S., Nihli, M., Butters, C., Heller, M., Hatch, V., Craig, R. *et al.* (2002). The troponin tail domain promotes a conformational state of the thin filament that suppresses myosin activity. *J. Biol. Chem.* **277**, 636–27642.
 36. Owen, C. & DeRosier, D. J. (1993). A 13-Å map of the actin-scrutin filament from the *Limulus* acrosomal process. *J. Cell Biol.* **123**, 337–344.
 37. Landis, C. A., Bobkova, A., Homsher, E. & Tobacman, L. S. (1997). The active state of the thin filament is destabilized by an internal deletion in tropomyosin. *J. Biol. Chem.* **272**, 14051–14056.
 38. Heeley, D. H., Moir, A. J. G. & Perry, S. V. (1982). Phosphorylation of tropomyosin during development in mammalian striated muscle. *FEBS Lett.* **146**, 115–118.

39. Heeley, D. H., Watson, D. H., Mak, A. S., Dubord, P. & Smillie, L. B. (1989). Effect of phosphorylation on the interaction and functional properties of rabbit striated muscle alpha alpha-tropomyosin. *J. Biol. Chem.* **264**, 2424–2430.
40. Milligan, R. A. & Flicker, P. F. (1987). Structural relationships of actin, myosin, and tropomyosin revealed by cryo-electron microscopy. *J. Cell Biol.* **105**, 29–39.
41. Trachtenberg, S. & DeRosier, D. J. (1987). Three-dimensional structure of frozen hydrated flagellar filament of *Salmonella typhimurium*. *J. Mol. Biol.* **195**, 581–601.
42. Egelman, E. H. (2000). A robust algorithm for the reconstruction of helical filaments using single-particle methods. *Ultramicroscopy*, **85**, 225–234.
43. Schutt, C. E. (1987). Muscle regulation. Movement on the Aufbaubahn. *Nature*, **325**, 757–758.