


Journal Pre-proof



Development and Initial Validation of the Asthma Severity Scoring System (ASSESS)

Anne M. Fitzpatrick, Ph.D., Stanley J. Szefler, M.D., David T. Mauger, Ph.D., Brenda R. Phillips, M.S., Loren C. Denlinger, M.D., Ph.D., Wendy C. Moore, M.D., Ronald L. Sorkness, Ph.D., Sally E. Wenzel, M.D., Peter J. Gergen, M.D., M.P.H., Eugene R. Bleecker, M.D., Mario Castro, M.D., Serpil C. Erzurum, M.D., John V. Fahy, M.D., Benjamin M. Gaston, M.D., Elliot Israel, M.D., Bruce D. Levy, M.D., Deborah A. Meyers, Ph.D., W.Gerald Teague, M.D., Leonard B. Bacharier, M.D., Ngoc P. Ly, M.D., Wanda Phipatanakul, M.D., M.Sc., Kristie R. Ross, M.D., Joe Zein, M.D., Nizar N. Jarjour, M.D.

PII: S0091-6749(19)31254-0

DOI: <https://doi.org/10.1016/j.jaci.2019.09.018>

Reference: YMAI 14196

To appear in: *Journal of Allergy and Clinical Immunology*

Received Date: 7 January 2019

Revised Date: 9 September 2019


Accepted Date: 16 September 2019

Please cite this article as: Fitzpatrick AM, Szefler SJ, Mauger DT, Phillips BR, Denlinger LC, Moore WC, Sorkness RL, Wenzel SE, Gergen PJ, Bleecker ER, Castro M, Erzurum SC, Fahy JV, Gaston BM, Israel E, Levy BD, Meyers DA, Teague WG, Bacharier LB, Ly NP, Phipatanakul W, Ross KR, Zein J, Jarjour NN, Development and Initial Validation of the Asthma Severity Scoring System (ASSESS), *Journal of Allergy and Clinical Immunology* (2019), doi: <https://doi.org/10.1016/j.jaci.2019.09.018>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Development and Initial validation of a tool for assessing asthma treatment response



Development and Initial Validation of the Asthma Severity Scoring System (ASSESS)

Anne M. Fitzpatrick, Ph.D.^{1,2}
Stanley J. Szeffler, M.D.^{3,4}
David T. Mauger, Ph.D.⁵
Brenda R. Phillips, M.S.⁵
Loren C. Denlinger, M.D., Ph.D.⁶
Wendy C. Moore, M.D.⁷
Ronald L. Sorkness, Ph.D.⁶
Sally E. Wenzel, M.D.⁸
Peter J. Gergen, M.D., M.P.H.⁹
Eugene R. Bleeker, M.D.¹⁰
Mario Castro, M.D.¹¹
Serpil C. Erzurum, M.D.¹²
John V. Fahy, M.D.¹³
Benjamin M. Gaston, M.D.¹⁴
Elliot Israel, M.D.¹⁵
Bruce D. Levy, M.D.¹⁵
Deborah A. Meyers, Ph.D.¹⁰
W. Gerald Teague, M.D.¹⁶
Leonard B. Bacharier, M.D.¹⁷
Ngoc P. Ly, M.D.¹⁸
Wanda Phipatanakul, M.D., M.Sc.¹⁹
Kristie R. Ross, M.D.¹⁴
Joe Zein, M.D.¹²
Nizar N. Jarjour, M.D.⁶

Author affiliations:

¹ Emory University, Department of Pediatrics, Atlanta, Georgia

² Children's Healthcare of Atlanta, Atlanta, Georgia

³ Children's Hospital Colorado, Aurora, Colorado

⁴ University of Colorado School of Medicine, Department of Pediatrics, Aurora, Colorado

⁵ The Pennsylvania State University, Department of Public Health Sciences, Hershey, Pennsylvania

⁶ University of Wisconsin, Department of Medicine, Madison, Wisconsin

⁷ Wake Forest University, Department of Internal Medicine, Winston-Salem, North Carolina

⁸ University of Pittsburgh, Department of Medicine, Pittsburgh, Pennsylvania

⁹ National Institute of Allergy and Infectious Diseases, Bethesda, Maryland

¹⁰ University of Arizona, College of Medicine, Tucson, Arizona

¹¹ Washington University, Department of Internal Medicine, St. Louis, Missouri

¹² Cleveland Clinic, Lerner Research Institute, Cleveland, Ohio

¹³ University of California San Francisco, Department of Medicine, San Francisco, California

¹⁴ Case Western Reserve University Department of Pediatrics, Cleveland, Ohio

¹⁵ Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts

¹⁶ University of Virginia Department of Pediatrics, Charlottesville, Virginia

¹⁷ Washington University, Department of Pediatrics, St. Louis, Missouri

¹⁸ University of California San Francisco, Department of Pediatrics, San Francisco, California

¹⁹ Boston Children's Hospital, Division of Immunology, Boston, Massachusetts

Please address correspondence to:

Anne M. Fitzpatrick, Ph.D.

Emory University Department of Pediatrics

2015 Uppergate Drive

Atlanta, Georgia 30322

Telephone: 404-727-9112

Facsimile: 404-712-0920

Email: anne.fitzpatrick@emory.edu

Supported by NHLBI to the Severe Asthma Research Program (SARP):

U10 HL109086

U10 HL109146

U10 HL109152

U10 HL109164

U10 HL109168

U10 HL109172

U10 HL109250

U10 HL109257

In addition, this program is supported through the following National Institutes of Health National Center for Advancing Translational Sciences awards:

UL1 TR001420 (Wake Forest University)

UL1 TR000427 (University of Wisconsin)

UL1 TR001102 (Harvard University)

UL1 TR002378 (Emory University)

Disclosures:

Anne M. Fitzpatrick:

Nothing to disclose.

Stanley J. Szeffler:

Dr. Szeffler reports consultancy fees from Merck, Boehringer-Ingelheim, Genentech, GlaxoSmithKline, Aerocrine, Novartis, Astra Zeneca, Daiichi Sankyo, Roche, Sanofi, Regeneron and Teva; grants from GlaxoSmithKline, outside the submitted work.

David T. Mauger:

Dr. Mauger reports non-financial support from Merck, non-financial support from Boehringer-Ingelheim, non-financial support from GSK, non-financial support from TEVA, non-financial support from Vifor, outside the submitted work.

Brenda R. Phillips:

Nothing to disclose.

Loren C. Denlinger:

Dr. Denlinger reports personal fees from AstraZeneca, personal fees from Sanofi, personal fees from GSK, outside the submitted work.

Wendy C. Moore:

Dr. Moore reports consultancy fees from AstraZeneca, Sanofi, and GlaxoSmithKline; and is the principal investigator in multicenter clinical trials with sponsors Astrazeneca, GSK, Pearl Therapeutics, and Sanofi.

Ronald L. Sorkness:

Nothing to disclose.

Sally E. Wenzel:

Dr. Wenzel reports grants and personal fees from AstraZeneca, grants from Boehringer-Ingelheim, grants from GSK, grants from Novartis, grants and personal fees from Sanofi, personal fees from Pieris, personal fees from Up to Date, outside the submitted work; In addition, Dr. Wenzel has a patent null pending.

Peter J. Gergen:

Nothing to disclose.

Eugene R. Bleecker:

Dr. Bleecker reports other from ERB has undertaken clinical trials through his employer, Wake Forest School of Medicine and University of Arizona, for AstraZeneca, MedImmune, Boehringer Ingelheim, Genentech, Johnson and Johnson (Janssen), Novartis, Regeneron, and Sanofi Genzyme, personal fees from ERB has also served as a paid consultant for AstraZeneca, MedImmune, Boehringer Ingelheim, Glaxo Smith Kline, Novartis, Regeneron, and Sanofi Genzyme, outside the submitted work.

Mario Castro:

Dr. Castro reports personal fees from Astra-Zeneca, Aviragen, Boehringer-Ingelheim, Boston Scientific, Elsevier, Genentech, 4D Pharma, Mallinckrodt, Neutronic, Nuaira, Teva, Theravance and VIDA; grants from Boehringer-Ingelheim, Chiesi, Genentech, Novartis, Sanofi-Aventis, Vectura; all outside the submitted work.

Serpil C. Erzurum:

Dr. Erzurum reports serving as the Chair of the American Board of Internal Medicine Pulmonary Disease Board, outside the submitted work.

John V. Fahy:

Dr. Fahy reports consultancy fees from Boehringer Ingelheim, Pieris, Entrinsic Health Solutions, Sanofi Genzyme; and is a named inventor on three patents, outside the submitted work.

Benjamin M. Gaston:

Dr. Gaston reports minority ownership in Respiratory Research, Inc., and intellectual property regarding the treatment of severe asthma.

Elliot Israel:

Dr. Israel reports personal fees from AstraZeneca, personal fees from Novartis, personal fees from Philips Respironics, personal fees from Regeneron Pharmaceuticals, personal fees and other from TEVA Specialty Pharmaceuticals, grants from Genentech, non-financial support from Boehringer Ingelheim, non-financial support from GlaxoSmithKline, non-financial support from Merck, non-financial support from Sunovion, non-financial support from TEVA, grants from Sanofi, personal fees from Bird Rock Bio, personal fees from Nuvelution Pharmaceuticals, personal fees from Vitaeris, Inc, grants from Boehringer Ingelheim, non-financial support from TEVA Specialty Pharmaceuticals, personal fees from Sanofi Genzyme, personal fees from Merck, personal fees from Entrinsic Health Solutions, personal fees from GlaxoSmithKline,

other from Vorso Corp., personal fees from Pneuma Respiratory , personal fees from 4D Pharma, outside the submitted work.

Bruce D. Levy:

Nothing to disclose.

Deborah A. Meyers:

Nothing to disclose.

W. Gerald Teague:

Dr. Teague reports salary support from the University of Virginia Ivy Foundation (Endowed Chair); Grant support from Panera Bread, TEVA Respiratory, Astra Zeneca and Sanofi/Regeneron; Advisory Boards for Sanofi/Regeneron, TEVA Respiratory, GSK, Genentech, Aviragen; Speaker Bureaus with personal fees from Genentech/Novartis, TEVA Respiratory (QVAR); and Writing committees for the American College of Allergy and Immunology, outside the submitted work.

Kristie R. Ross:

Dr. Ross reports grants from National Institutes of Health, during the conduct of the study; grants from Ohio Department of Jobs and Family Services, non-financial support from Sunovion, non-financial support from Merck, non-financial support from BI, grants and non-financial support from TEVA, non-financial support from GSK, grants from Astra Zeneca, grants from Roche, outside the submitted work.

Leonard B. Bacharier:

Dr. Bacharier reports personal fees from GlaxoSmithKline, personal fees from Genentech/Novartis, personal fees from Merck, personal fees from DBV Technologies, personal fees from Teva, personal fees from Boehringer Ingelheim, personal fees from Sanofi/Regeneron, personal fees from Vectura, personal fees from Circassia, personal fees from AstraZeneca, outside the submitted work.

Ngoc Ly:

Dr. Ly reports grants from Vertex 2017, personal fees from Gilead 2017, outside the submitted work.

Wanda Phipatanakul:

Nothing to disclose.

Kristie R. Ross:

Nothing to disclose.

Joe Zein:

Dr. Zein reports research support from the Cleveland Clinic.

Nizar N. Jarjour:

Dr. Jarjour received honorarium from Astra Zeneca and Boehringer Ingelheim for consultation unrelated to the submitted work. Dr. Jarjour also served on the ABIM Pulmonary Disease Test Committee, outside the submitted work.

ClinicalTrials.Gov registration number: NCT01606826

Word count = 3,809

Abstract (word count = 250)**Background**

Tools for quantification of asthma severity are limited.

Objective

To develop a continuous measure of asthma severity, the Asthma Severity Scoring System (ASSESS), for adolescents and adults incorporating domains of asthma control, lung function, medications, and exacerbations.

Methods

Baseline and 36-month longitudinal data from participants in Phase 3 of the Severe Asthma Research Program (SARP, NCT01606826) were utilized. Scale properties, responsiveness, and a minimal important difference (MID) were determined. External replication was performed in participants enrolled in SARP Phase 1/2. Utility of ASSESS for detecting treatment response was explored in participants undergoing corticosteroid responsiveness testing with intramuscular triamcinolone and participants receiving biologics.

Results

ASSESS scores ranged from 0 to 20 (8.78 ± 3.9 ; higher scores reflect worse severity) and differed between 5 phenotypic groups. Measurement properties were acceptable. ASSESS was responsive to changes in quality of life with a MID of 2, with good specificity for outcomes of asthma improvement and worsening, but poor sensitivity. Replication analyses yielded similar results, with a 2-point decrease (improvement) associated with improvements in quality of life. Participants with ≥ 2 point decrease (improvement) in ASSESS scores also had greater improvement in lung function and asthma control after triamcinolone, but these differences were

limited to phenotypic clusters 3, 4 and 5. Participants treated with biologics also had ≥ 2 point decrease (improvement) in ASSESS scores overall.

Conclusions

The ASSESS tool is an objective measure that may be useful in epidemiologic and clinical research studies for quantification of treatment response in individual patients and in phenotypic groups. However, validation studies are warranted.

Clinical implications (word count = 31)

The Asthma Severity Scoring System (ASSESS) incorporates four asthma domains and may be useful in research settings for the quantification of treatment response in individual patients and in phenotypic groups.

Capsule summary (word count = 34)

Objective tools for quantifying asthma severity are limited. The Asthma Severity Scoring System (ASSESS) assesses 4 domains of asthma and may be useful for quantifying treatment responses in research settings.

Key words

Asthma control

Asthma severity classification

Severe asthma

Psychometric testing

Tool development

228 **Abbreviations**

229	ACQ	Asthma Control Questionnaire
230	ACT	Asthma Control Test
231	ASSESS	Asthma Severity Scoring System
232	ATS	American Thoracic Society
233	AQLQ	Asthma Quality of Life Questionnaire
234	CASI	Composite Asthma Severity Index
235	ERS	European Respiratory Society
236	FEV ₁	Forced expiratory volume in one second
237	FVC	Forced vital capacity
238	GINA	Global Initiative for Asthma
239	ICS	Inhaled corticosteroid
240	LABA	Long-acting beta-agonist
241	MID	Minimally important difference
242	ROC	Receiver operating curve
243	SARP	Severe Asthma Research Program

Introduction

Asthma severity is a unique construct, independent of asthma control, that encompasses a range of asthma patients from mild to severe.^{1,2} Whereas asthma “control” refers to the extent to which symptoms and other features of asthma are present in patients, asthma “severity” reflects the level of treatment required to control symptoms and exacerbations.² Like asthma control, asthma severity is not a static feature and can change over the course of months or years in response to seasonal variations, infections and other exposures.³⁻⁵ However, while there are many standardized tools available for the assessment of asthma control in both clinical and research settings, tools for quantification of asthma severity are not widely available.

The National Institute of Allergy and Infectious Disease-sponsored Inner-City Asthma Consortium recently developed a Composite Asthma Severity Index (**CASI**) which quantifies asthma severity through domains of impairment, risk, and medication requirements, but that tool was developed in a population of children (and adolescents) and lacks details on systemic corticosteroids, emerging controller medications such as tiotropium, and biologic therapies that are currently utilized for the treatment of severe asthma.² Recognizing that severe asthma is a multidimensional construct, we aimed to develop a continuous measure of asthma severity for adolescents and adults that: 1) incorporates fundamental aspects of asthma control, lung function, current controller medications, and recent exacerbations in the assessment of the construct without extensive questioning, 2) is sensitive to short-term temporal changes in asthma and its management, 3) does not require prior knowledge of the patient, and 4) could be utilized by future investigators as well as clinicians in real-world practice settings. The development, measurement properties, and validation of this tool, named the “Asthma Severity Scoring System” (**ASSESS**), are reviewed below.

Methods

Participants age 12 and older enrolled in Phase 3 of the National Heart, Lung and Blood Institute's Severe Asthma Research Program (**SARP**) between November 2012 and February 2015 with at least one year of follow-up data were included in the analysis (**Figure E1**). Eligibility criteria for SARP3 included a physician diagnosis of asthma and either $\geq 12\%$ reversibility in the forced expiratory volume in one second (**FEV₁**) after bronchodilator administration or airway hyperresponsiveness to methacholine, evidenced by a provocative concentration of methacholine ≤ 16 mg/mL. Thirteen participants with FEV₁ values below 50% of predicted who could not undergo methacholine challenge due to concerns for safety were also enrolled at the discretion of the principal investigator. Current smoking, smoking history >10 pack years if ≥ 30 years of age or >5 pack years if <30 years of age, premature birth before 35 weeks gestation, and other chronic airway disorders such as aspiration or vocal cord dysfunction were criteria for exclusion. Permission to proceed with SARP studies was granted by the Institutional Review Board of each institution and an independent Data Safety and Monitoring Board. The study was also registered at ClinicalTrials.gov (NCT01606826). Informed written consent and assent (if less than 18 years) were obtained from all participants prior to enrollment in these studies.

SARP design and procedures. Details of the study design have been reported previously.^{6,7} Participants completed a baseline characterization visit during which systemic triamcinolone acetonide was administered, a follow-up visit at 18 ± 3 days to assess triamcinolone response, and follow-up characterization visits at 12, 24 and 36 months (± 90 days). Visits were postponed if an asthma exacerbation treated with systemic corticosteroids or a respiratory infection treated with antibiotics was reported within the preceding four or two

weeks, respectively. Telephone calls were completed every 6 months (± 60 days) to assess for asthma control and healthcare utilization. Exacerbations were defined according to a consensus report⁸ as a worsening of asthma necessitating treatment with systemic corticosteroids. Severe asthma was defined using a modification of the European Respiratory Society (**ERS**)/American Thoracic Society (**ATS**) definition⁹ that required the presence of: 1) asthma requiring Global Initiative for Asthma (**GINA**) step 4-5 treatment with high doses of inhaled corticosteroids (**ICS**) and other controller medications, or 2) systemic corticosteroid treatment for $\geq 50\%$ of the previous year to achieve or maintain asthma control. Management by an asthma specialist was not a criterion for severe asthma in SARP3. Other SARP3 characterization procedures including assignment of participants to 5 phenotypic asthma clusters previously identified in Phase 1 and Phase 2 of SARP¹⁰ are detailed in the online repository.

Tool development. ASSESS is an adaptation of the CASI,¹¹ which was developed using the Delphi method and tested with data from children enrolled in a clinical trials network (**Table 1**). Details of the ASSESS tool development are provided in the online repository and in **Tables E1-E3** and **Figure E2**.

Scale properties of the Asthma Severity Scoring System. Internal consistency of the ASSESS tool was evaluated with Cronbach's alpha. Test-retest reliability was evaluated with intraclass correlation using two-way random models with the following format:

$$(\text{Mean square}_{\text{rows}} - \text{Mean square}_{\text{error}}) / \text{Mean square}_{\text{rows}},$$

which is the preferred approach for evaluation of rater-based clinical assessments that are designed for routine use by any clinician.¹² Given that there is no available gold-standard measure for asthma severity in adults, validity could not be determined with traditional approaches. Instead, baseline ASSESS scores were compared in participants treated with ICS

who reported 3 or more exacerbations in the previous 12 months, and participants with a hospitalization or intensive care unit admission in the previous year using t-tests. ASSESS scores were also compared in participants who did and did not satisfy the modified dichotomous ERS/ATS definition of severe asthma using Point-Biserial correlation analysis.

Responsiveness of the Asthma Severity Scoring System. Changes in ASSESS scores were compared to changes in Asthma Control Test (**ACT**) score, absolute change in FEV₁ percent predicted value (calculated as FEV₁ % predicted_{follow-up visit} - FEV₁ % predicted_{prior visit}), and the change in the number of asthma controller medications using Pearson correlational analyses. Responsiveness of the ASSESS tool to changes in the Asthma Quality of Life Questionnaire (**AQLQ**) total score and individual AQLQ domains (symptoms, activities, emotions, environment) was assessed using Pearson correlational analyses. For comparison purposes, the ability of the modified dichotomous ERS/ATS definition of severe asthma to discriminate these same outcomes was also assessed with t-tests.

Mean changes and Minimal Important Difference. Mean changes in ASSESS scores were further calculated for groups of participants meeting pre-specified cut-points as follows: 1) ACT score, 3 point increase¹³; 2) FEV₁ absolute change, 10% increase;^{7, 14, 15} 3) controller medications, 2 medication decrease; and 4) AQLQ, 0.5 point increase.¹⁶ Mean differences between groups were compared with t-tests.

The Minimal Important Difference (**MID**) of the ASSESS tool was determined at baseline and 12, 24 and 36 months with a distribution-based approach, which utilizes the statistical properties of the scale.¹⁷ Data from each corresponding time point were used to determine the standard deviation (**SD**) and standard error of measurement (**SEM**). The MID was defined by both $0.5 \times SD$ ¹⁸ and $1 \times SEM$.¹⁹ The SEM was calculated as follows:

$$SEM = SD_{ASSESS} \times \sqrt{1 - Reliability_{ASSESS}},$$

Where reliability corresponds to the Cronbach α value. To aid in clinical interpretation of MID results, receiver operating characteristic (**ROC**) analyses were performed with one-year differences in ASSESS scores as the predictor and outcomes of asthma improvement (≥ 1 fewer exacerbations, ≥ 1 fewer controller medications, ≥ 0.5 point increase in AQLQ¹⁶) and asthma worsening (≥ 1 more exacerbation, ≥ 1 more controller medication, ≥ 0.5 point decrease in AQLQ¹⁶, hospitalization). Analyses were performed in a merged dataset of repeated measures from 12, 24 and 36 months (N=1686 observations).

External replication. External replication was performed on a sample of 100 participants age ≥ 12 years enrolled in phase 1 and 2 of SARP program between 2003 and 2010 with baseline and 6 month follow-up data. None of these participants were enrolled in SARP3. Methods for participant characterization in the SARP 1 and 2 program have been published previously.²⁰

Utility for detection of treatment response. Utility of the ASSESS tool in detecting treatment responses was assessed in participants in SARP3 who underwent corticosteroid response testing with intramuscular triamcinolone. Changes in Asthma Control Questionnaire (**ACQ**) scores and FEV₁ % predicted values before and after triamcinolone receipt were compared in participants with a ≥ 2 point improvement (i.e., decrease) on the ASSESS tool versus participants with ≤ 2 point improvement. Exploratory analyses also compared ASSESS scores in participants in whom biologic therapy was initiated or discontinued.

Data analysis. Data were analyzed with IBM SPSS Statistics (Version 24.0). For all analyses, an alpha level of 0.05 was considered the threshold for statistical significance. Given the exploratory nature of the analyses, no adjustments were made for multiple comparisons.

Results

Five hundred ninety eight participants were initially selected for inclusion; however, 562 participants provided at least 12 months of follow up data and were included in the analysis. Features of these participants are shown in **Table 2**.

ASSESS score distribution. In the total sample of SARP3 participants, ASSESS scores ranged from 0 to 20 (mean $\pm$ standard deviation, 8.78 ± 3.98 ; 95% confidence interval (CI) for the mean, 8.45 – 9.11). Responses to the individual items of the ASSESS tool were varied, but some clustering was noted in the pattern of responses (**Figure E3**). For example, one group of participants with high ASSESS scores had poor asthma control yet normal lung function and few exacerbations, whereas another group of participants with high ASSESS scores had poor asthma control with significant airflow obstruction and frequent exacerbations, consistent with the heterogeneity of asthma previously reported in the SARP3 program.²¹

The distribution of ASSESS scores according to the modified dichotomous ERS/ATS definition of severe asthma and SARP Phase 1 and 2 phenotypic cluster assignment (Cluster 1: well controlled early-onset atopic asthma; Cluster 2: early-onset atopic asthma with increased medication requirements; Cluster 3: late-onset, non-atopic obese asthma with moderately reduced lung function and frequent exacerbations; Cluster 4: severe airflow obstruction with bronchodilator reversibility and frequent exacerbations despite multiple controller medications; Cluster 5: severe airflow obstruction with less bronchodilator reversibility and frequent exacerbations despite multiple controller medications) is shown in **Figure 1**. There were no significant differences in ASSESS scores between adolescents 12-17 years (8.1 ± 3.4) and adults ≥ 18 years (8.9 ± 4.0).

Reliability measures. Given the multi-dimensional (versus unidimensional) nature of the ASSESS tool, internal consistency of the item questions was expected to be less than 0.7.²² The Cronbach α value was 0.639 for the entire sample, 0.468 for adolescents 12-17 years, and 0.662 for adults ≥ 18 years, which reflects an expected lack of concordance between symptoms, lung function, medication use and exacerbations in all asthma patients. However, the intraclass correlation coefficients between baseline and 12 months, 12-24 months, and 24-36 months were 0.764, 0.768, and 0.813 for the entire sample, respectively (**Figure E4**) and suggested “good” test-retest reliability according to the conventions of Koo et al.¹² Intraclass correlation coefficients were also similar between age groups (age 12-17: 0.717, 0.841, 0.732; age ≥ 18 years: 0.768, 0.766, 0.816). Therefore, further analyses were not stratified by age.

Construct validity. At baseline, ASSESS scores were higher in participants who reported 3 or more exacerbations, hospitalization or intensive care unit admission for asthma in the previous 12 months (**Figure 2**). Agreement between ASSESS scores and the modified dichotomous ERS/ATS definition of severe asthma was 0.639 at baseline and 0.635, 0.533, and 0.532 at 12, 24, and 36 months, respectively.

Responsiveness. By design, the ASSESS tool was sensitive to changes in ACT score, FEV₁ % predicted, and the number of asthma controller medications between 0-12 months, 12-24 months, and 24-36 months (**Figure E5**). The ASSESS tool was also responsive to changes in the AQLQ score (total score and each domain), as shown in **Figure 3**. In contrast, the modified dichotomous ERS/ATS definition of severe asthma (also assessed at baseline, 12 months and 24 months) did not discriminate changes in these same outcomes (**Figure E6**).

Mean differences and Minimal Important Difference. Mean differences in ASSESS scores between pre-specified participant groups are shown in **Table 3** and were approximately 2

overall. Calculation of the ASSESS MID, defined as the smallest difference in the ASSESS score that represents a meaningful change, produced an MID of approximately 2 using a distribution-based approach (for the $0.5 \times \text{SD}$ calculation, range 1.83-1.96; for the $1 \times \text{SEM}$ calculation, range 2.39 – 2.56) (**Table E4**). Clinical interpretation of the calculated ASSESS score MID was further guided by ROC analyses with ASSESS scores as the predictor and outcomes of asthma improvement and worsening reflected by changes in the number of exacerbations, number of controller medications, and AQLQ score and hospitalization occurrence. A MID of 2 had high specificity for outcome detection (80.3 – 85.5% for asthma improvement and 88.0 – 91.4% for asthma worsening), but poor sensitivity (45.8 – 46.5% and 30.0 – 37.3% for improvement and worsening, respectively) (**Table E5**). A MID of 1 increased sensitivity for asthma improvement (~62%), but only marginally increased sensitivity for asthma worsening (~46%).

External replication. Features of the external replication sample are shown in **Table E6** and were similar to those of participants in SARP3. Mean ASSESS scores at baseline and the follow-up visit were 8.04 ± 4.61 (95% CI, 6.63 – 8.36) and 7.00 ± 4.42 (95% CI, 5.54 – 7.12). Changes in ASSESS scores were also linearly associated with changes in AQLQ scores (**Figure 4A-E**). Patients who achieved a clinically meaningful improvement in quality of life (≥ 0.5 point improvement) also had significant reductions in ASSESS score, although not all participants met the MID of 2 for the ASSESS instrument (**Figure 4F-J**).

Utility for detection of treatment response. Overall, ASSESS scores were significantly lower after the triamcinolone intervention ($p < 0.001$, **Figure 5A**). Participants with ≥ 2 point improvement (i.e., decrease) in ASSESS scores after triamcinolone also had greater mean improvements in FEV_1 % predicted and ACQ scores at 18 ± 3 days after triamcinolone administration (**Figure 5B-D**). However, these differences were limited to participants with a

modified dichotomous ERS/ATS definition of severe asthma (**Figure E7, Table E7**) and participants assigned to phenotypic clusters 3, 4 and 5 (**Figure E8**). Further exploratory analyses of patients with severe asthma in whom biologic therapy was initiated (N = 22) also demonstrated higher baseline ASSESS scores than severe patients not treated with biologics and improvement by approximately 2 points (-1.95 ± 3.27) after biologic initiation (**Figure E9**).

Discussion

Objective measures of asthma severity are limited, despite the emphasis of asthma severity by treatment guidelines for over a decade.^{2, 23} The relative explosion of new product development for asthma in recent years now necessitates objective measures for distinguishing patients who might benefit from new treatments such as biologics as well as patients who do not respond to those therapies.²⁴ Advantages of objective versus subjective patient assessments are clearly demonstrated in the asthma literature. For example, other studies of asthma control and asthma functional status have shown that concordance between objective and judgment-based assessments is often poor and may even worsen in patients receiving higher Step-based therapy.²⁵⁻²⁸ Asthma severity determination is also discordant between physicians in both primary care²⁹ and specialist³⁰ settings. Objective measures such as ASSESS have potential to mitigate the rising costs of asthma severity both in the United States^{31, 32} and elsewhere^{33, 34} through comprehensive assessment of treatment responses and ultimately, elimination of unnecessary treatments in patients who do not benefit.

ASSESS was developed in response to the unmet need for a severity scoring system in adolescent and adult patients with asthma.³⁵ Despite the multidimensional nature of the ASSESS tool, it has acceptable measurement properties including test-retest reliability and responsiveness

to changes in asthma-related quality of life in heterogeneous asthma populations. Our exploratory analyses also suggest that a reduction (i.e., improvement) of ASSESS scores by at least 2 points may also aid with quantification of response to severe asthma treatments in individual patients and in some phenotypic subgroups. While single outcome variables (such as FEV₁ or exacerbation rate) have been used previously in many clinical trials, such narrowly focused analyses may miss the complexity of this disease manifestation and the effects of selected outcome variables on each other. We feel that ASSESS can overcome the limitation of single response variable by offering an integrated numerical score that is sensitive to changes in individual patients. The ASSESS tool is also the first to include biologics and anticholinergics in the assignment of medication treatment step.

Other tools have been proposed for determining asthma severity, but these have limitations in adult asthma populations. The CASI tool was derived from a population of children enrolled in clinical trials¹¹ and the distribution of its components (i.e., lung function measures) differ in adults. The CASI tool also does not address medications that are commonly used in adult asthma populations.¹¹ A separate continuous score of asthma severity developed for adults has also been proposed for epidemiological studies,³⁶ but this score does not adequately capture the constructs of severity as recognized by treatment guidelines and focuses only on symptom frequency and pharmacologic treatment. The coding of the variables in that study (i.e., “yes” or “no” for most symptoms and “none,” “GINA step 1,” or “≥GINA step 2” for medications) also does not distinguish more severe patients with multiple controller medications.³⁶

Nonetheless, there are limitations to our approach. Most importantly, there is no gold standard measure of asthma severity for the purpose of comparison. We instead evaluated

performance of the ASSESS score against its own component parts, so it is not surprising that associations were seen. We also were unable to directly compare ASSESS scores to CASI scores. The CASI tool requires that daytime and nighttime symptoms be assessed by self-report using a 2-week recall period prior to the patient encounter, and this information was not available in SARP. However, given the extensive utilization of the ACT tool in clinical practices across the world, our intent was to develop a tool that could utilize information captured by the ACT. An advantage of this approach is that the ACT tool uses a 4-week recall period and collects symptom information as well as information on self-rated asthma control and impact of asthma on functioning. Nonetheless, it is also recognized that despite the 4-week recall window, ACT scores may more strongly reflect symptoms in the immediate 2 weeks.³⁷ We were also limited in our assessment of exacerbation rates at the baseline visit, which impacted tool development, and assessment of treatment responsiveness given the design of the SARP3 study. For example, post-triamcinolone ASSESS scores were calculated from data obtained at the 6-month phone call, with the exception of FEV₁, which was obtained at the post-triamcinolone follow-up visit 18 ± 3 days after triamcinolone administration. This long interval between triamcinolone administration and follow-up may not have been optimal for detection of asthma control responses. Indeed, other studies have shown that peak symptom effects with systemic triamcinolone are evident at 7-10 days.³⁸

Although the Cronbach alpha value associated with the ASSESS tool are lower than those associated with other tools, asthma “severity” is not a unidimensional construct and therefore struct cut-points of 0.70 for “acceptability” are not appropriate. Indeed, Brunner et al.²² argue that is worth re-thinking recommended benchmark values for Cronbach alpha in the context of multi-dimensional measures, because scale scores below these values may still contain

valuable information and may still have predictive power. That fact that the Cronbach alpha was less than 0.7 was therefore expected and reflects a lack of concordance between symptoms, lung function, medication use and exacerbations in all patients.

The ASSESS score also does not take into account medication adherence and could be limited if prescribing practices are not aligned with treatment recommendations. Indeed, over-prescription of systemic corticosteroids and under-prescription (or under-utilization) of ICS and other asthma controller medications have been reported in the literature and may be particularly problematic in certain patient populations.³⁹⁻⁴¹ The ASSESS tool also focuses on current medications and does not account for treatment duration or the timing of medication initiation (for example, the previous week or 12 months earlier). Generalization to other asthma populations is also cautioned since the ethnic distribution of the SARP3 population was limited and participant recruitment occurred predominately at large academic medical centers across the United States.

The lack of a gold standard measure of asthma severity also poses a challenge for calculation of the MID of the ASSESS tool. The two broad methods of estimating MID, the distribution method and the anchor-based methods, are conceptually different.¹⁷ Whereas the distribution method uses the statistical properties of the score distribution, the anchor-based method uses a direct response from the patient on another tool that measures the same construct to evaluate the meaning of a particular degree of change.¹⁷ Anchor-based methods often yield more conservative estimates of MID compared with the distribution method⁴² and thus the clinical meaningfulness of our calculated MID should be interpreted with caution until further studies are performed. We did compare changes in ASSESS scores to changes in patient-reported quality of life, and found the ASSESS tool to be responsive in this regard. Patients achieving a

MID of 0.5 point improvement¹⁶ on the AQLQ instrument also had improvement (i.e., decrease) of ~2 points on the ASSESS tool, which was similar to our calculated MID value of ~2. The specificity of a 2-point MID was also good, but the sensitivity for outcome detection (either asthma improvement or asthma worsening) was poor. Although a separate analysis of the CASI tool found that a MID of 1 point or greater may be appropriate for future studies,⁴³ a one-point change in ASSESS could occur spuriously (for example, from a 1-point increase in FEV₁ from 69 to 70% predicted, or a 1-point increase in ACT score from 10 to 11) and have little clinical meaningfulness. Recognizing that asthma severity is a multi-dimensional construct and not a unidimensional measure, an MID of 2 for ASSESS is also more likely to capture changes in more than one severity dimension.

Despite these limitations, there are a number of strengths. A major strength of the present study is the comprehensive characterization of enrolled participants and enrichment of the study population with severe asthma as defined by ERS/ATS criteria, which has a prevalence rate of less than 5% in the entire asthma population.⁴⁴ Another strength is the availability of longitudinal data of the majority of the participants, allowing for testing this tool in the same group of subjects over time. The multi-center design of SARP is another strength since asthma features and related outcomes can be subject to wide geographic variability.^{45, 46} The SARP study also utilized crude measures of socioeconomic status such as education and income and had adequate representation across socioeconomic brackets.

In conclusion, the ASSESS tool, which assesses 4 domains of asthma severity, has acceptable measurement properties and is responsive to changes in asthma quality of life with a MID of approximately 2. ASSESS explains significant variance in selected asthma outcomes and may be useful in epidemiologic and research studies to quantify treatment response in individual

patients and in phenotypic subgroups. However, additional validation studies are needed. We caution against the use of ASSESS in clinical practice until such studies are undertaken, to avoid inappropriate changes in asthma treatment.

Acknowledgments

The authors would like to acknowledge Dr. William W. Busse for his input regarding the tool development and for his review of the manuscript.

548 **Table 1.** Asthma Severity Scoring System (ASSESS) for age 12 and older.

549

Original CASI tool (reference ¹¹)		ASSESS tool	
Dimension of severity (weight)		Dimension of severity (weight)	
Days with symptoms and albuterol in the last 2 weeks (15%)		ACT composite score (30%)	
0 – 3 days	___ 0 points	ACT score 23-25	___ 0 points
4 – 9 days	___ 1 point	ACT score 20-22	___ 1 point
10 – 13 days	___ 2 points	ACT score 17-19	___ 2 points
14 days	___ 3 points	ACT score 14-16	___ 3 points
Nights with symptoms and albuterol in the last 2 weeks (15%)		ACT score 11-13	___ 4 points
0 – 1 nights	___ 0 points	ACT score 8-10	___ 5 points
2 nights	___ 1 point	ACT score 5-7	___ 6 points
3 – 4 nights	___ 2 points		
5 – 14 nights	___ 3 points		
Current pre-bronchodilator lung function (15%)		Current pre-bronchodilator lung function (15%)	
FEV ₁ ≥85% predicted	___ 0 points	FEV ₁ ≥80% predicted	___ 0 points
FEV ₁ =80-84% predicted	___ 1 point	FEV ₁ =70-80% predicted	___ 1 point
FEV ₁ =70-79% predicted	___ 2 points	FEV ₁ =60-70% predicted	___ 2 points
FEV ₁ <70% predicted	___ 3 points	FEV ₁ <60% predicted	___ 3 points
Current medications (25%)		Current medications (25%)	
Controller medications:		Controller medications:	
No treatment	___ 0 points	No treatment	___ 0 points
Albuterol	___ 1 point	Step 1: Albuterol only	___ 1 point
Low-dose ICS (or LTRA)	___ 2 points	Step 2: Low-dose ICS only	___ 2 points
Low-dose ICS + LABA	___ 3 points	or LTRA only	
Medium-dose ICS + LABA	___ 4 points	Step 3: Low-dose ICS plus 1 or more controllers	___ 3 points
High-dose ICS	___ 5 points	or medium-dose ICS only	
		or high-dose ICS only	
		Step 4: Medium-dose ICS + 1 or more controllers	___ 4 points
		or high-dose ICS + 1 or more controllers	

		Step 5: High-dose ICS + 2 or more controllers	___ 5 points
		Other medications:	
		Daily or alternative-day oral corticosteroids or monthly corticosteroid injections	___ 1 point
		Current biologic	___ 1 point
Asthma exacerbations (last 2 months)		Asthma exacerbations (last 6 months) ¹	
Prednisone burst	___ 2 points	Prednisone burst	___ 2 points
Prednisone burst + hospitalization	___ 4 points	Prednisone burst + hospitalization	___ 4 points
Maximum possible score	18	Maximum possible score	20

ACT = Asthma Control Test, CASI = Composite Asthma Severity Index, ICS = Inhaled corticosteroid, LABA = Long-acting-beta agonist, LTRA = leukotriene receptor antagonist

¹ For the baseline SARP visit, exacerbations were assessed over the preceding 12 months. Follow-up SARP visits (at 12, 24 and 36 months) assessed exacerbations over the preceding 6 months

Table 2. Features of the participants, shown as mean $\pm$ SEM or number of participants (%).

Groups are not significantly different.

Feature	N = 562
Age (years)	44.1 $\pm$ 0.7
Age < 18 years	65 (11.6)
Age $\geq$ 18 years	497 (88.4)
Asthma duration (years)	28.5 $\pm$ 0.7
Male	202 (35.9)
Hispanic ethnicity	24 (4.3)
Race	
White	350 (62.3)
Black	142 (25.3)
More than one race	49 (8.7)
Other	21 (3.7)
Severe asthma (modified dichotomous ERS/ATS definition)	328 (58.4)
Highest household educational attainment	
Did not complete high school	10 (1.8)
High school diploma	57 (10.1)
Some college or technical training	115 (20.5)
Associate degree	87 (15.5)
Bachelor's degree	289 (51.4)
Decline to answer	4 (0.7)
Highest annual household income	
Less than \$25,000	104 (18.5)
\$25,000 to \$49,999	114 (20.3)
\$50,000 to \$99,999	159 (28.3)
\$100,000 or more	109 (19.4)
Decline to answer	76 (13.5)
Saw an asthma specialist (previous year) ¹	306 (54.4)
Asthma healthcare utilization (previous year)	
Unscheduled physician visit	250 (44.5)
Emergency department	143 (25.4)
Hospitalization	70 (12.5)

Intubation for asthma (ever in lifetime)	35 (6.2)
Daily asthma medications	
Inhaled corticosteroid	489 (87.0)
Long-acting beta-agonist	439 (78.1)
Long-acting anti-muscarinic	52 (9.3)
Leukotriene receptor antagonist	226 (40.2)
Theophylline	23 (4.1)
Systemic corticosteroids	91 (16.2)
Biologic	43 (7.7)
Asthma Control Test Score	17.2 ± 0.2
23-25	66 (11.8)
20-22	153 (27.3)
17-19	116 (20.7)
14-16	98 (17.5)
11-13	65 (11.6)
8-10	42 (7.5)
5-7	20 (3.6)
Baseline lung function (% predicted value)	
FVC	86.7 ± 0.8
FEV ₁	74.2 ± 0.9

¹ Severe asthma (modified dichotomous ERS/ATS definition), n = 236 (72.0%), non-severe asthma, n = 70 (29.9%)

Table 3. Mean difference in ASSESS scores between visits. Negative values indicate a lower ASSESS score at the follow up visit.

Interval	Asthma outcome	n (%) with outcome	Change in ASSESS score		Mean difference in ASSESS score (95% CI)	p-value
			Participants without outcome mean (SD)	Participants with outcome mean (SD)		
Baseline – 12 months	ACT score, ≥ 3 point increase	169 (28.3)	-0.20 ± 2.23	-3.15 ± 2.34	-2.95 (-3.36, -2.54)	< 0.001
	FEV ₁ , $\geq 10\%$ absolute increase	84 (15.1)	-0.77 ± 2.49	-2.86 ± 2.77	-2.08 (-2.67, -1.50)	< 0.001
	Medication, decrease of ≥ 2	35 (6.3)	-0.90 ± 2.50	-3.80 ± 3.11	-2.90 (-3.77, -2.02)	< 0.001
	AQLQ score, ≥ 0.5 point increase	159 (26.6)	-0.44 ± 2.36	-2.73 ± 2.62	-2.29 (-2.74, -1.84)	< 0.001
12-24 months	ACT score, ≥ 3 point increase	111 (18.6)	0.41 ± 2.12	-2.46 ± 2.72	-2.88 (-3.35, -2.40)	< 0.001
	FEV ₁ , $\geq 10\%$ absolute increase	60 (11.8)	-0.01 ± 2.45	-1.72 ± 2.79	-1.71 (-2.38, -1.04)	< 0.001
	Medication, decrease of ≥ 2	5 (1.0)	-0.17 ± 2.52	-4.00 ± 3.16	-3.82 (-6.06, -1.60)	0.001
	AQLQ score, ≥ 0.5 point increase	117 (19.6)	0.21 ± 2.30	-1.62 ± 2.85	-1.83 (-2.33, -1.32)	< 0.001
24-36 months	ACT score, ≥ 3 point increase	97 (16.2)	0.21 ± 2.00	-2.25 ± 2.17	-2.46 (-2.92, -2.00)	< 0.001
	FEV ₁ , $\geq 10\%$ absolute increase	59 (12.7)	-0.02 ± 2.08	-2.18 ± 2.60	-2.16 (-2.75, -1.57)	< 0.001
	Medication, decrease of ≥ 2	11 (2.3)	-0.27 ± 2.23	-1.45 ± 3.35	-1.18 (-2.53, 0.17)	0.087
	AQLQ score, ≥ 0.5 point increase	101 (16.9)	0.12 ± 2.12	-1.48 ± 2.41	-1.50 (-1.98, -1.01)	< 0.001

Figure Legends

Figure 1. (word count = 47)

Distribution of baseline and longitudinal Asthma Severity Scoring System (ASSESS) scores in all participants, participants according to the modified dichotomous European Respiratory Society/American Thoracic Society definition of severe asthma, and participants according to phenotypic cluster assignment. Data in panels A and B reflect the mean $\pm$ SEM.

Figure 2. (word count = 33)

Asthma Severity Scoring System (ASSESS) scores in participants with 3 or more exacerbations hospitalization, or intensive care unit admission in the previous year. Boxplot whiskers correspond to the 5th and 95th percentiles. **p<0.001

Figure 3. (word count = 35)

Responsiveness of the Asthma Severity Scoring System (ASSESS) score to changes in the Asthma Quality of Life Questionnaire (AQLQ) score between baseline and 12 months, 12 and 24 months, and 24 and 36 months. *p<0.001

Figure 4. (word count = 59)

Associations between changes in Asthma Severity Scoring System (ASSESS) and Asthma Quality of Life Questionnaire (AQLQ) scores in the replication cohort. Figures on the right are stratified by changes in AQLQ scores. Boxplot whiskers and dashed lines correspond to the 5th

and 95th percentiles and the minimal important difference of -2 for ASSESS improvement. *p < 0.05, **p<0.01

Figure 5. (word count = 60)





Radar plots depicting changes in individual domains of the Asthma Severity Scoring System (ASSESS) at baseline and after intramuscular triamcinolone. Spikes correspond to individual participants responses. Panels B-D depict changes (mean $\pm$ SEM) in FEV₁, ACQ7, and ACQ6 scores after triamcinolone in participants in whom ASSESS scores improved (≥ 2 point decrease), did not change, and worsened (≥ 2 point increase). **p<0.01




References


1. Taylor DR, Bateman ED, Boulet LP, Boushey HA, Busse WW, Casale TB, et al. A new perspective on concepts of asthma severity and control. *Eur Respir J* 2008; 32:545-54.
2. Global Initiative for Asthma. Global Strategy for Asthma Management and Prevention, 2018. Available from www.ginasthma.org. Last accessed November 1, 2018.
3. Newby C, Heaney LG, Menzies-Gow A, Niven RM, Mansur A, Bucknall C, et al. Statistical cluster analysis of the British Thoracic Society Severe refractory Asthma Registry: clinical outcomes and phenotype stability. *PLoS One* 2014; 9:e102987.
4. Silkoff PE, Laviolette M, Singh D, FitzGerald JM, Kelsen S, Backer V, et al. Longitudinal stability of asthma characteristics and biomarkers from the Airways Disease Endotyping for Personalized Therapeutics (ADEPT) study. *Respir Res* 2016; 17:43.
5. Zaihra T, Walsh CJ, Ahmed S, Fugere C, Hamid QA, Olivenstein R, et al. Phenotyping of difficult asthma using longitudinal physiological and biomarker measurements reveals significant differences in stability between clusters. *BMC Pulm Med* 2016; 16:74.
6. Teague WG, Phillips BR, Fahy JV, Wenzel SE, Fitzpatrick AM, Moore WC, et al. Baseline Features of the Severe Asthma Research Program (SARP III) Cohort: Differences with Age. *J Allergy Clin Immunol Pract* 2018; 6:545-54 e4.
7. Phipatanakul W, Mauger DT, Sorkness RL, Gaffin JM, Holguin F, Woodruff PG, et al. Effects of Age and Disease Severity on Systemic Corticosteroid Responses in Asthma. *Am J Respir Crit Care Med* 2017; 195:1439-48.
8. Fuhlbrigge A, Peden D, Apter AJ, Boushey HA, Camargo CA, Jr., Gern J, et al. Asthma outcomes: exacerbations. *J Allergy Clin Immunol* 2012; 129:S34-48.
9. Chung KF, Wenzel SE, Brozek JL, Bush A, Castro M, Sterk PJ, et al. International ERS/ATS guidelines on definition, evaluation and treatment of severe asthma. *Eur Respir J* 2014; 43:343-73.
10. Moore WC, Meyers DA, Wenzel SE, Teague WG, Li H, Li X, et al. Identification of asthma phenotypes using cluster analysis in the Severe Asthma Research Program. *Am J Respir Crit Care Med* 2010; 181:315-23.
11. Wildfire JJ, Gergen PJ, Sorkness CA, Mitchell HE, Calatroni A, Kattan M, et al. Development and validation of the Composite Asthma Severity Index--an outcome measure for use in children and adolescents. *J Allergy Clin Immunol* 2012; 129:694-701.
12. Koo TK, Li MY. A Guideline of Selecting and Reporting Intraclass Correlation Coefficients for Reliability Research. *J Chiropr Med* 2016; 15:155-63.
13. Schatz M, Kosinski M, Yarlas AS, Hanlon J, Watson ME, Jhingran P. The minimally important difference of the Asthma Control Test. *J Allergy Clin Immunol* 2009; 124:719-23 e1.
14. Brand PL, Quanjer PH, Postma DS, Kerstjens HA, Koeter GH, Dekhuijzen PN, et al. Interpretation of bronchodilator response in patients with obstructive airways disease. The Dutch Chronic Non-Specific Lung Disease (CNSLD) Study Group. *Thorax* 1992; 47:429-36.
15. Ward H, Cooper BG, Miller MR. Improved criterion for assessing lung function reversibility. *Chest* 2015; 148:877-86.
16. Juniper EF, Guyatt GH, Willan A, Griffith LE. Determining a minimal important change in a disease-specific Quality of Life Questionnaire. *J Clin Epidemiol* 1994; 47:81-7.











17. Norman GR, Sridhar FG, Guyatt GH, Walter SD. Relation of distribution- and anchor-based approaches in interpretation of changes in health-related quality of life. *Med Care* 2001; 39:1039-47.
18. Norman GR, Sloan JA, Wyrwich KW. Interpretation of changes in health-related quality of life: the remarkable universality of half a standard deviation. *Med Care* 2003; 41:582-92.
19. Wyrwich KW, Tierney WM, Wolinsky FD. Further evidence supporting an SEM-based criterion for identifying meaningful intra-individual changes in health-related quality of life. *J Clin Epidemiol* 1999; 52:861-73.
20. Moore WC, Bleecker ER, Curran-Everett D, Erzurum SC, Ameredes BT, Bacharier L, et al. Characterization of the severe asthma phenotype by the National Heart, Lung, and Blood Institute's Severe Asthma Research Program. *J Allergy Clin Immunol* 2007; 119:405-13.
21. Denlinger LC, Phillips BR, Ramratnam S, Ross K, Bhakta NR, Cardet JC, et al. Inflammatory and Comorbid Features of Patients with Severe Asthma and Frequent Exacerbations. *Am J Respir Crit Care Med* 2017; 195:302-13.
22. Brunner M, SUB H-M. Analyzing the reliability of multidimensional measures: An example from intelligence research. *Educ Psychol Meas* 2005; 65:227-40.
23. National Asthma E, Prevention P. Expert Panel Report 3 (EPR-3): Guidelines for the Diagnosis and Management of Asthma-Summary Report 2007. *J Allergy Clin Immunol* 2007; 120:S94-138.
24. Tabatabaian F, Ledford DK, Casale TB. Biologic and New Therapies in Asthma. *Immunol Allergy Clin North Am* 2017; 37:329-43.
25. Crespo-Lessmann A, Plaza V, Gonzalez-Barcala FJ, Fernandez-Sanchez T, Sastre J. Concordance of opinions between patients and physicians and their relationship with symptomatic control and future risk in patients with moderate-severe asthma. *BMJ Open Respir Res* 2017; 4:e000189.
26. Urrutia I, Plaza V, Pascual S, Cisneros C, Entrenas LM, Luengo MT, et al. Asthma control and concordance of opinions between patients and pulmonologists. *J Asthma* 2013; 50:877-83.
27. Menzies-Gow A, Chiu G. Perceptions of asthma control in the United Kingdom: a cross-sectional study comparing patient and healthcare professionals' perceptions of asthma control with validated ACT scores. *NPJ Prim Care Respir Med* 2017; 27:48.
28. Juniper EF, Chauhan A, Neville E, Chatterjee A, Svensson K, Mork AC, et al. Clinicians tend to overestimate improvements in asthma control: an unexpected observation. *Prim Care Respir J* 2004; 13:181-4.
29. Gillis RME, van Litsenburg W, van Balkom RH, Muris JW, Smeenk FW. The contribution of an asthma diagnostic consultation service in obtaining an accurate asthma diagnosis for primary care patients: results of a real-life study. *NPJ Prim Care Respir Med* 2017; 27:35.
30. Braido F, Baiardini I, Alleri P, Bacci E, Barbetta C, Bellocchia M, et al. Asthma management in a specialist setting: Results of an Italian Respiratory Society survey. *Pulm Pharmacol Ther* 2017; 44:83-7.
31. Sullivan PW, Campbell JD, Ghushchyan VH, Globe G. Outcomes before and after treatment escalation to Global Initiative for Asthma steps 4 and 5 in severe asthma. *Ann Allergy Asthma Immunol* 2015; 114:462-9.





32. Sullivan PW, Campbell JD, Ghushchyan VH, Globe G, Lange J, Woolley JM. Characterizing the severe asthma population in the United States: claims-based analysis of three treatment cohorts in the year prior to treatment escalation. *J Asthma* 2015; 52:669-80.
33. Nordon C, Grimaldi-Bensouda L, Pribil C, Nachbaur G, Amzal B, Thabut G, et al. Clinical and economic burden of severe asthma: A French cohort study. *Respir Med* 2018; 144:42-9.
34. Barry LE, Sweeney J, O'Neill C, Price D, Heaney LG. The cost of systemic corticosteroid-induced morbidity in severe asthma: a health economic analysis. *Respir Res* 2017; 18:129.
35. Szeffler SJ. Asthma across the lifespan: Time for a paradigm shift. *J Allergy Clin Immunol* 2018; 142:773-80.
36. Calciano L, Corsico AG, Pirina P, Trucco G, Jarvis D, Janson C, et al. Assessment of asthma severity in adults with ever asthma: A continuous score. *PLoS One* 2017; 12:e0177538.
37. Okupa AY, Sorkness CA, Mauger DT, Jackson DJ, Lemanske RF, Jr. Daily diaries vs retrospective questionnaires to assess asthma control and therapeutic responses in asthma clinical trials: is participant burden worth the effort? *Chest* 2013; 143:993-9.
38. Fitzpatrick AM, Stephenson ST, Brown MR, Nguyen K, Douglas S, Brown LAS. Systemic Corticosteroid Responses in Children with Severe Asthma: Phenotypic and Endotypic Features. *J Allergy Clin Immunol Pract* 2017; 5:410-9 e4.
39. Farber HJ, Silveira EA, Vicere DR, Kothari VD, Giardino AP. Oral Corticosteroid Prescribing for Children With Asthma in a Medicaid Managed Care Program. *Pediatrics* 2017; 139.
40. Patel M, Pilcher J, Reddel HK, Qi V, Mackey B, Tranquilino T, et al. Predictors of severe exacerbations, poor asthma control, and beta-agonist overuse for patients with asthma. *J Allergy Clin Immunol Pract* 2014; 2:751-8.
41. Sweeney J, Patterson CC, O'Neill S, O'Neill C, Plant G, Lynch V, et al. Inappropriate prescribing of combination inhalers in Northern Ireland: retrospective cross-sectional cohort study of prescribing practice in primary care. *Prim Care Respir J* 2014; 23:74-8.
42. Jayadevappa R, Cook R, Chhatre S. Minimal important difference to infer changes in health-related quality of life-a systematic review. *J Clin Epidemiol* 2017; 89:188-98.
43. Krouse RZ, Sorkness CA, Wildfire JJ, Calatroni A, Gruchalla R, Hershey GKK, et al. Minimally important differences and risk levels for the Composite Asthma Severity Index. *J Allergy Clin Immunol* 2017; 139:1052-5.
44. Israel E, Reddel HK. Severe and Difficult-to-Treat Asthma in Adults. *N Engl J Med* 2017; 377:965-76.
45. Malhotra K, Baltrus P, Zhang S, McRoy L, Immergluck LC, Rust G. Geographic and racial variation in asthma prevalence and emergency department use among Medicaid-enrolled children in 14 southern states. *J Asthma* 2014; 51:913-21.
46. Garcia E, Serban N, Swann J, Fitzpatrick A. The effect of geographic access on severe health outcomes for pediatric asthma. *J Allergy Clin Immunol* 2015; 136:610-8.











Development and Initial Validation of the Asthma Severity Scoring System (ASSESS)

Online Data Repository

Journal Pre-proof

Methods

SARP3 recruitment procedures. The Severe Asthma Research Program (**SARP**) cohort was assembled in response to a National Institutes of Health mandate (RFA HL-11-018, June 11, 2010) requesting applications for clinical centers of establish a severe asthma research program that outlined specific recruiting methods. Clinical partnerships and a Data Coordinating Center were named and a broad recruiting process was initiated targeting a sample enriched in severe asthma (60%) and including 30% minorities, 50% women, and 25% children. Study coordinators at each clinical site screened potential enrollees with both severe and non-severe asthma by various methods including program notices on the ClinicalTrials.gov web site, data queries of patients with asthma receiving care at the respective clinical sites, research databases of patients available for clinical trials, social media, advertisement in local media, and archives of participants who had previously participated in research studies.

SARP3 participant characterization methods. Medical history questionnaires assessed self-reported healthcare utilization, medications, and demographic features and exposures such as tobacco smoke. Severe asthma was defined using a modification of the European Respiratory Society (**ERS**) and American Thoracic Society (**ATS**).¹ Asthma control over the past 4 weeks was assessed with the Asthma Control Test (**ACT**).^{2,3} Global asthma-related quality of life over the preceding two weeks was assessed with the Asthma Quality of Life Questionnaire (**AQLQ**).⁴ Spirometry (KoKo® PDS, Ferraris, Louisville, CO, donated by nSpire Health, Longmont, CO) was performed according to technical standards⁵ with centralized over-reading for accuracy. Participants withheld short-acting and long-acting bronchodilator medication for a minimum of 4 or 12 hours, respectively. Forced vital capacity (**FVC**), and forced expiratory volume in one

second (FEV_1) were interpreted according to Global Lung Function Initiative population reference equations⁶ and expressed as a percentage of the predicted value.

Triamcinolone response testing. At the end of the baseline characterization visits, participants received a single dose of intramuscular triamcinolone (40 mg for age ≥ 18 and 1 mg/kg up to 40 mg maximum for age ≤ 17) administered deep in the gluteal muscle.

Triamcinolone response was assessed at a follow-up visit (18 ± 3 days) where spirometry was repeated. Asthma control over the past week was also assessed prior to and following receipt of intramuscular triamcinolone with the 6- and 7-item Asthma Control Questionnaire (**ACQ**).^{7, 8}

Asthma phenotypic cluster assignment. At the baseline visit, participants in SARP3 were assigned to 5 phenotypic asthma clusters previously identified in Phase 1 and Phase 2 of SARP: Cluster 1, early-onset atopic asthma with normal lung function and few controller medications; Cluster 2, early-onset atopic asthma with preserved lung function but increased medication requirements; Cluster 3, late-onset, non-atopic obese asthma with moderate reductions in FEV_1 and frequent exacerbations; Cluster 4, severe airflow obstruction with bronchodilator reversibility and frequent exacerbations despite multiple controller medications; and Cluster 5, severe airflow obstruction with less bronchodilator reversibility and frequent exacerbations despite multiple controller medications.⁹ Phenotypic cluster assignment was accomplished by adding one additional participant to the original dataset of N=726 participants in SARP Phase 1 and Phase 2, and then repeating the Ward's minimum-variance hierarchical clustering method with Ward's linkage on 11 of the strongest discriminatory variables for cluster assignment in the previous report, to minimize the within-cluster sum of squares or to maximize the between-cluster sum of squares.⁹ The 11 variables used for cluster assignment included pre-bronchodilator FEV_1 (% predicted) and FVC (% predicted), absolute change in FEV_1 %

predicted value after bronchodilator administration, age of asthma onset, asthma duration, sex, frequency of beta-agonist use, and dose of inhaled corticosteroids.

Tool development. Scoring for each dimension of the Asthma Severity Scoring System (**ASSESS**) tool was based on consensus of a working group of SARP3 investigators with expertise in pulmonary/critical care medicine (L.C.D., W.C.M., N.N.J.), physiology (R.L.S.), nursing (A.M.F), and biostatistics (D.T.M., B.R.P.) Three external experts in allergy/immunology familiar with the CASI (S.J.S., P.J.G., W.W.B.) also provided input on the tool development. For the medication dimension, participants were assigned to a treatment step developed by consensus, based on the 2018 Global Initiative for Asthma (**GINA**) treatment guidelines (**Table E1**).¹⁰ Thresholds for low, medium and high-dose ICS treatment were adapted from the GINA guidelines for estimated clinical comparability of drug products, with minor modification for United States formulations (**Table E2**).

Two versions of the ASSESS tool were compared. The first version contained separate dimensions for daytime symptoms (weighted at 15%) and nighttime symptoms (weighted at 15%) over the previous 4 weeks. The second version replaced the daytime and nighttime symptom dimensions with the total ACT score (weighted at 30%). Strong agreement was noted between versions ($r = 0.980$), but measurement properties were enhanced with the ACT score modification (inter-item correlation range for daytime and nighttime symptoms, 0.190 – 0.511; for ACT score, 0.591 – 0.737) (**Table E3**). Therefore, the modification with total ACT score was retained for further testing. Principal components analysis of the four dimensions of the retained ASSESS tool identified two components that explained 69.4% of the total variance. The first component contained medications (i.e., Treatment Step), exacerbations and ACT score

(component loadings: 0.749, 0.724, and 0.712, respectively) and the second component contained FEV₁ % predicted (component loading: 0.776) (**Figure E1**).

Visualization of ASSESS item-response clustering. Visualization of responses to individual items of the ASSESS tool was accomplished with a clustered heat map (i.e., double dendrogram) depicting hierarchical clustering in two directions, by ASSESS item response and patient (NCSS software, version 12). Item responses were transformed to a 0-1 range to eliminate imbalances between individual item scoring and weighting.

Utility of ASSESS for detection of treatment response. Post-triamcinolone ASSESS scores were calculated from data obtained at the 6-month phone call, with the exception of FEV₁, which was obtained at the post-triamcinolone follow-up visit 18 ± 3 days after triamcinolone administration. Participants with a ≥ 2 point improvement (i.e., decrease) on the ASSESS tool were compared to participants with no change and participants with ≥ 2 point worsening (i.e., increase) on the ASSESS tool for the following outcomes: 1) absolute change in FEV₁ %predicted value, 2) change in ACQ-6 score, and 3) ACQ-7 between baseline and the post-triamcinolone visit at 18 ± 3 days.

Table E1. Stepwise approach to asthma treatment, developed from consensus opinion of the SARP investigators. Modifications are shown as footnotes.

	Preferred controller (Daily treatment) ¹	Other controller options
Step 1	N/A ²	N/A
Step 2	Low-dose ICS	LTRA Theophylline
Step 3	Low-dose ICS + LABA	Medium or high-dose ICS Low-dose ICS + LTRA Low-dose ICS + Theophylline <i>Low-dose ICS + LABA + LTRA</i>
Step 4	Medium or high dose ICS + LABA	Add tiotropium Medium or high-dose ICS + LTRA Medium or high-dose ICS + Theophylline <i>Medium-dose ICS + LAMA</i> <i>Medium-dose ICS + LABA + LTRA</i> <i>Medium-dose ICS + LABA + LAMA</i> <i>Medium-dose ICS + LABA + LAMA + LTRA</i> <i>High-dose ICS + LAMA³</i> <i>High-dose ICS + LTRA³</i>
Step 5	High dose ICS + LABA + add-on treatment (i.e., tiotropium)	Add low-dose oral corticosteroids <i>High-dose ICS + LAMA + LTRA</i>

¹ For participants using ICS as needed, ICS use ≥ 21 days/month was considered daily ICS use.

² As-needed ICS without LTRA considered Step 1

Table E2. Inhaled corticosteroid dose thresholds (for age 12 and older) used for treatment step assignment. Thresholds were developed from consensus opinion of the SARP investigators. Modifications are shown as footnotes. Values represent estimated clinical comparability and are not necessarily bioequivalent.

	ICS dose threshold ¹		
	Low	Medium	High
Beclomethasone dipropionate (HFA)	100-200	> 200-400	> 400
Budesonide (DPI)	200-400	> 400-800	> 800
Budesonide (nebules)	1000	2000	4000
Budesonide (HFA)	80 up to 320	320 up to 640	≥ 640
Ciclesonide (HFA)	80-160	> 160-320	> 320
Fluticasone furoate (DPI)	100	N/A	200
Fluticasone propionate (DPI)	100-250	> 250-500	> 500
Fluticasone propionate (HFA)	100-250	> 250-500	> 500
Mometasone furoate (DPI or HFA)	110-220	> 220-440	> 440

¹ For participants with two ICS products, ICS dose threshold assignments were made as follows:
 Low-dose primary ICS + low-dose secondary ICS = medium dose ICS
 Medium-dose primary ICS + low or medium-dose secondary ICS = high dose ICS
 If either the primary or secondary ICS is high-dose = high-dose ICS

DPI = dry powder inhaler, GINA = Global Initiative for Asthma, HFA = hydrofluoroalkane,

Table E3. Item-total statistics for an initial version of ASSESS tool with separate dimensions for daytime and nighttime symptoms, compared to the final version of the ASSESS tool, which replaces daytime and nighttime symptoms with Asthma Control Test (ACT) total scores.

Tool version	Cronbach's α	Item	Corrected Item-Total Correlation	Cronbach's α if Item Deleted
Separate dimensions for daytime and nighttime symptoms	0.669	Daytime symptoms	0.407	0.627
		Nighttime symptoms	0.498	0.589
		Lung function (FEV ₁)	0.357	0.645
		Controller medications	0.448	0.609
		Exacerbations	0.357	0.645
Single symptom dimension with ACT total score	0.726	ACT score	0.931	0.569
		Lung function (FEV ₁)	0.498	0.716
		Controller medications	0.671	0.635
		Exacerbations	0.616	0.666

Table E4. Minimal Important Differences (MIDs) in ASSESS scores determined by the distribution approach.

Timepoint	Standard deviation (SD)	Cronbach α	MID (0.5*SD)	MID (1*SEM)
Baseline	3.98	0.639	1.96	2.39
12 months	3.65	0.590	1.83	2.51
24 months	3.74	0.590	1.87	2.51
36 months	3.65	0.574	1.83	2.56
All timepoints merged	3.84	0.605	1.92	2.46

SEM = standard error of measurement

Table E5. Receiver operative curve (ROC) analyses with change in ASSESS score as a predictor for selected asthma outcomes in a merged dataset of repeated measures at 12, 24 and 36 months (N=1686 observations). A negative value for the change in ASSESS score indicates an improvement in asthma severity, whereas a positive value reflects a worsening of asthma severity.

Asthma outcome at 12 months ¹	Number (%) with outcome	Cut-point for ASSESS score	Sensitivity (%)	Specificity (%)	Area under the curve (95% CI)	p-value
Asthma improvement						
≥1 fewer exacerbation(s) in the past year	449 (26.6)	-3	32.1	92.3	0.73 (0.703, 0.760)	< 0.001
		-2	46.5	85.5		
		-1	62.6	72.8		
≥1 fewer controller medication(s) for asthma	231 (13.7)	-3	33.1	88.8	0.687 (0.649, 0.726)	< 0.001
		-2	45.9	80.3		
		-1	61.5	69.6		
≥ 0.5 point increase in Asthma Quality of Life Questionnaire (AQLQ) total score	370 (21.9)	-3	32.2	92.5	0.732 (0.702, 0.763)	< 0.001
		-2	45.8	85.4		
		-1	62.4	72.8		
Asthma worsening						
≥1 more exacerbation(s) in the past year	281 (16.7)	3	18.0	95.6	0.688 (0.652, 0.724)	< 0.001
		2	30.4	91.1		
		1	46.1	82.0		
≥1 more controller medication(s) for asthma	203 (12.0)	3	20.0	95.1	0.681 (0.638, 0.724)	< 0.001
		2	31.5	90.2		
		1	47.1	80.8		

≥ 0.5 point decrease in Asthma Quality of Life Questionnaire (AQLQ) total score	290 (17.2)	3	19.5	96.2	0.705 (0.670, 0.739)	< .001
		2	30.0	91.4		
		1	45.7	82.4		
Hospitalization for asthma	59 (3.5)	3	27.1	93.8	0.708 (0.643, 0.774)	< 0.001
		2	37.3	88.0		
		1	47.5	77.6		

CI = confidence interval

¹ Compared to the prior visit

Table E6. Features of the participants in the external replication cohort, shown as mean $\pm$ SEM or number of participants (%).

Feature	N = 100
Age (years)	35.1 $\pm$ 1.5
Age 12-18 years	7 (7.0)
Age $\geq$ 18 years	93 (93.0)
Asthma duration (years)	21.9 $\pm$ 1.3
Severe asthma (modified dichotomous ERS/ATS definition)	29 (29.0)
Male	27 (27.0)
Hispanic ethnicity	6 (6.0)
Race	
White	60 (60.0)
Black	33 (33.0)
More than one race	3 (3.0)
Other	4 (4.0)
Highest household educational attainment	
Did not complete high school	3 (3.0)
High school diploma	12 (12.0)
Some college or technical training	30 (30.0)
College degree	45 (45.0)
Decline to answer	10 (10.0)
Asthma healthcare utilization (previous year)	
Unscheduled physician visit	64 (64.0)
Emergency department	28 (28.0)
Hospitalization	9 (9.0)
Intubation for asthma (ever in lifetime)	10 (10.0)
Daily asthma medications	
Inhaled corticosteroid	70 (70.0)
Long-acting beta-agonist	53 (53.0)
Leukotriene receptor antagonist	29 (29.0)
Theophylline	4 (4.0)
Systemic corticosteroids	13 (13.0)
Biologic (omalizumab)	3 (3.0)

Asthma Control Test score	17.1 ± 0.4
Lung function	
FVC % predicted	91.7 ± 1.8
FEV ₁ % predicted	81.9 ± 2.1

Table E7. Differences (mean $\pm$ SEM) in outcomes in participant subgroups before and 18 $\pm$ 3 days after triamcinolone administration. * $p \leq 0.05$, ** $p \leq 0.01$ vs. ASSESS score improvement

	ASSESS score improvement (≥ 2 point decrease)	No change in ASSESS score	ASSESS score worsening (≥ 2 point increase)
Absolute change in FEV ₁ % predicted (%)			
Non-severe asthma ¹	2.40 $\pm$ 1.03	1.53 $\pm$ 0.71	0.66 $\pm$ 1.60
Severe asthma ²	6.69 $\pm$ 0.81	0.45 $\pm$ 0.87**	-1.65 $\pm$ 1.72**
Cluster 1 ³	2.61 $\pm$ 3.90	-0.04 $\pm$ 2.13	1.40 $\pm$ 3.61
Cluster 2 ⁴	3.76 $\pm$ 1.03	-2.55 $\pm$ 1.74	-5.15 $\pm$ 5.33
Cluster 3 ⁵	7.35 $\pm$ 2.02	3.09 $\pm$ 2.33*	0.17 $\pm$ 2.07**
Cluster 4 ⁶	7.61 $\pm$ 1.95	2.49 $\pm$ 0.95**	1.30 $\pm$ 2.83
Cluster 5 ⁷	9.41 $\pm$ 1.60	1.26 $\pm$ 1.98**	-2.05 $\pm$ 2.88
Change in ACQ7 score			
Non-severe asthma	-0.36 $\pm$ 0.10	-0.31 $\pm$ 0.06	-0.25 $\pm$ 0.13
Severe asthma	-0.67 $\pm$ 0.70	-0.10 $\pm$ 0.07**	0.01 $\pm$ 0.12**
Cluster 1	-1.16 $\pm$ 0.34	-0.48 $\pm$ 0.17	-0.57 $\pm$ 0.29
Cluster 2	-0.66 $\pm$ 0.13	-0.11 $\pm$ 0.11	0.04 $\pm$ 0.37
Cluster 3	-0.50 $\pm$ 0.15	-0.34 $\pm$ 0.21	-0.27 $\pm$ 0.64*
Cluster 4	-0.60 $\pm$ 0.12	-0.08 $\pm$ 0.11**	-0.29 $\pm$ 0.00
Cluster 5	-0.67 $\pm$ 0.14	0.23 $\pm$ 0.13**	0.46 $\pm$ 0.37**
Change in ACQ6 score			
Non-severe asthma	-0.34 $\pm$ 0.11	-0.33 $\pm$ 0.07	-0.27 $\pm$ 0.14
Severe asthma	-0.66 $\pm$ 0.08	-0.10 $\pm$ 0.08**	-0.04 $\pm$ 0.26**
Cluster 1	-0.67 $\pm$ 0.22	-0.35 $\pm$ 0.12	-0.29 $\pm$ 0.23
Cluster 2	-0.40 $\pm$ 0.11	-0.28 $\pm$ 0.08	-0.30 $\pm$ 0.19
Cluster 3	-0.54 $\pm$ 0.15	-0.32 $\pm$ 0.16	-0.11 $\pm$ 0.39
Cluster 4	-0.65 $\pm$ 0.13	-0.18 $\pm$ 0.10**	-0.28 $\pm$ 0.15
Cluster 5	-0.69 $\pm$ 0.15	0.22 $\pm$ 0.14**	0.28 $\pm$ 0.45*

¹ Non-severe asthma: Improvement, n = 84; no change, n = 108, worsened, n = 34

² Severe asthma: Improvement, n = 166; no change, n = 115, worsened, n = 27

³ Cluster 1: Improvement, n = 28; no change, n = 26, worsened, n = 7

⁴ Cluster 2: Improvement, n = 88; no change, n = 94, worsened, n = 28

⁵ Cluster 3: Improvement, n = 37; no change, n = 33, worsened, n = 14

⁶ Cluster 4: Improvement, n = 39; no change, n = 41, worsened, n = 3

⁷ Cluster 5: Improvement, n = 59; no change, n = 29, worsened, n = 9

Figure Legends

Figure E1. (word count = 11)

Inclusion of participants. SARP = Severe Asthma Research Program.

Figure E2 (word count = 25)

Scree plot depicting the fraction of total variance (shown as Eigenvalues) in the Asthma Severity Scoring System (ASSESS) data, as explained by each principal component.

Figure E3. (word count = 53)

Clustered heat map (i.e., double dendrogram) depicting patterns of responses for each item (i.e., domain) in the Asthma Severity Scoring System (ASSESS) tool. Items were transformed to a 0-1 range prior to clustering. Blue reflect a lower response value to the item whereas orange and red reflect higher response values to the item.

Figure E4. (word count = 22)

Intraclass correlations for the Asthma Severity Scoring System (ASSESS) tool between baseline and 12 months, 12 months and 24 months, and 24 and 36 months.

Figure E5. (word count = 33)

Associations between changes in Asthma Severity Scoring System (ASSESS) scores and changes in Asthma Control Test (ACT) scores, FEV₁ (measured as absolute change in percent predicted value), and controller medications across 36 months.

Figure E6. (word count = 57)

Association between the modified dichotomous European Respiratory Society/American Thoracic Society definition of severe asthma (assessed at baseline, 12 months and 24 months) and changes in Asthma Control Test (ACT) score, FEV₁ (measured as absolute change), controller medications and Asthma Quality of Life Questionnaire (AQLQ) score over the following year. Boxplot whiskers correspond to 5th and 95th percentiles.

Figure E7. (word count = 40)

Radar plots depicting changes in individual domains of the Asthma Severity Scoring System (ASSESS) at baseline and after intramuscular triamcinolone, stratified by the modified dichotomous European Respiratory Society/American Thoracic Society definition of severe asthma. Spikes correspond to individual participant responses.

Figure E8. (word count = 39)

Radar plots depicting changes in individual domains of the Asthma Severity Scoring System (ASSESS) at baseline and after intramuscular triamcinolone, stratified by phenotypic cluster assignment. Spikes correspond to individual participant responses.


Figure E9. (word count = 57)


(A) Baseline Asthma Severity Scoring System (ASSESS) score and (B) change in ASSESS score over 12 months between patients in whom biologic therapy was initiated, compared with patients

on stable biologic therapy and patients not treated with biologics. The dashed line at -2 corresponds to the meaningful improvement (i.e., reduction of 2 points) of the ASSESS score.


References


1. Chung KF, Wenzel SE, Brozek JL, Bush A, Castro M, Sterk PJ, et al. International ERS/ATS guidelines on definition, evaluation and treatment of severe asthma. *Eur Respir J* 2014; 43:343-73.
2. Nathan RA, Sorkness CA, Kosinski M, Schatz M, Li JT, Marcus P, et al. Development of the asthma control test: a survey for assessing asthma control. *J Allergy Clin Immunol* 2004; 113:59-65.
3. Schatz M, Zeiger RS, Drane A, Harden K, Cibildak A, Oosterman JE, et al. Reliability and predictive validity of the Asthma Control Test administered by telephone calls using speech recognition technology. *J Allergy Clin Immunol* 2007; 119:336-43.
4. Juniper EF, Guyatt GH, Ferrie PJ, Griffith LE. Measuring quality of life in asthma. *Am Rev Respir Dis* 1993; 147:832-8.
5. Miller MR, Hankinson J, Brusasco V, Burgos F, Casaburi R, Coates A, et al. Standardisation of spirometry. *Eur Respir J* 2005; 26:319-38.
6. Quanjer PH, Stanojevic S, Cole TJ, Baur X, Hall GL, Culver BH, et al. Multi-ethnic reference values for spirometry for the 3-95-yr age range: the global lung function 2012 equations. *Eur Respir J* 2012; 40:1324-43.
7. Juniper EF, O'Byrne PM, Guyatt GH, Ferrie PJ, King DR. Development and validation of a questionnaire to measure asthma control. *Eur Respir J* 1999; 14:902-7.
8. Juniper EF, Svensson K, Mork AC, Stahl E. Measurement properties and interpretation of three shortened versions of the asthma control questionnaire. *Respir Med* 2005; 99:553-8.
9. Moore WC, Meyers DA, Wenzel SE, Teague WG, Li H, Li X, et al. Identification of asthma phenotypes using cluster analysis in the Severe Asthma Research Program. *Am J Respir Crit Care Med* 2010; 181:315-23.
10. Global Initiative for Asthma. Global Strategy for Asthma Management and Prevention, 2018. Available from www.ginasthma.org. Last accessed November 1, 2018.






Clustered Heat Map




















0-12 months

12-24 months


24-36 months




Non-severe asthma
Baseline
ASSESS score = 5.77




Non-severe asthma
Post-triamcinolone
ASSESS score = 5.08



Severe asthma
Baseline
ASSESS score = 10.95



Severe asthma
Post-triamcinolone
ASSESS score = 9.01




- ACT domain
- FEV₁ domain
- Medication domain
- Exacerbation domain

Baseline


Cluster 1

ASSESS
score
= 6.33




Cluster 2

ASSESS
score
= 7.07




Cluster 3

ASSESS
score
= 8.84




Cluster 4

ASSESS
score
= 10.41



Cluster 5

ASSESS
score
= 12.37




ACT domain
FEV₁ domain
Medication
domain
Exacerbation
domain

Post-triamcinolone


Cluster 1

ASSESS
score
= 5.24




Cluster 2

ASSESS
score
= 6.12




Cluster 3

ASSESS
score
= 7.79




Cluster 4

ASSESS
score
= 8.48




Cluster 5

ASSESS
Score
= 10.21



A.



B.

