

REVIEW

Hypersensitivities following allergen antigen recognition by unconventional T cells

Marcela de Lima Moreira¹  | Michael N. T. Souter^{1,2}  | Zhenjun Chen¹  |
Liye Loh^{1,3}  | James McCluskey¹  | Daniel G. Pellicci⁴  | Sidonia B. G. Eckle¹ 

¹Department of Microbiology and Immunology, Peter Doherty Institute for Infection and Immunity, University of Melbourne, Melbourne, Vic., Australia

²Australian Research Council Centre of Excellence in Advanced Molecular Imaging, University of Melbourne, Parkville, Vic., Australia

³Department of Immunology and Microbiology, School of Medicine, University of Colorado Anschutz Medical Campus, Aurora, CO, USA

⁴Murdoch Children's Research Institute, Parkville, Vic., Australia

Correspondence

Sidonia B. G. Eckle, Department of Microbiology and Immunology, Peter Doherty Institute for Infection and Immunity, University of Melbourne, Melbourne, Vic., Australia.
Email: seckle@unimelb.edu.au

Funding information

CSL limited, Grant/Award Number: CSL Centenary Fellowship; Australian Government, Department of Education, Grant/Award Number: Australian Postgraduate Award; Australian Research Council, Grant/Award Number: DECRA (DE170100407); National Health and Medical Research Council, Grant/Award Number: NHMRC program grant (1113293); Allergy and Immunology Foundation of Australasia (AIFA), Grant/Award Number: 2019 AIFA grant

Abstract

Conventional T cells recognise protein-derived antigens in the context of major histocompatibility complex (MHC) class Ia and class II molecules and provide anti-microbial and anti-tumour immunity. Conventional T cells have also been implicated in type IV (also termed delayed-type or T cell-mediated) hypersensitivity reactions in response to protein-derived allergen antigens. In addition to conventional T cells, subsets of unconventional T cells exist, which recognise non-protein antigens in the context of monomorphic MHC class I-like molecules. These include T cells that are restricted to the cluster of differentiation 1 (CD1) family members, known as CD1-restricted T cells, and mucosal-associated invariant T cells (MAIT cells) that are restricted to the MHC-related protein 1 (MR1). Compared with conventional T cells, much less is known about the immune functions of unconventional T cells and their role in hypersensitivities. Here, we review allergen antigen presentation by MHC-I-like molecules, their recognition by unconventional T cells, and the potential role of unconventional T cells in hypersensitivities. We also speculate on possible scenarios of allergen antigen presentation by MHC-I-like molecules to unconventional T cells, the hallmarks of such responses, and the expected frequencies of hypersensitivities within the human population.

KEYWORDS

antigen, CD1, MAIT cells, MR1, NKT cells

de Lima Moreira and Souter are joint first authors.

© 2020 EAACI and John Wiley and Sons A/S. Published by John Wiley and Sons Ltd.

1 | INTRODUCTION

1.1 | Type IV hypersensitivities

According to the type of immune response, four broad subtypes of hypersensitivities can be distinguished based on the traditional classification by Gell and Coombs.¹ Type I-III involve immunoglobulin responses whilst type IV is T cell-mediated (also termed delayed-type or T cell-mediated hypersensitivity).¹ Within type IV hypersensitivities, four categories exist, where type IVa is a CD4⁺ T helper (Th) 1 lymphocyte-mediated reaction with activation of macrophages; type IVb is CD4⁺ Th2 lymphocyte mediated with eosinophilic involvement; type IVc is cytotoxic CD8⁺ T lymphocyte mediated with involvement of perforin-granzyme B in apoptosis; type IVd is T cell-driven neutrophilic inflammation.² Most studies on type IV hypersensitivity¹ have focused on protein allergen-derived peptide antigens (Ags) presented by classical major histocompatibility complex class I or class II (MHC-I or MHC-II) molecules. In contrast,

little is known about hypersensitivities caused by non-peptide Ags,³ presented by MHC-I-like molecules, the restriction elements of various subsets of unconventional T cells. Discovered ~30 years ago,⁴⁻⁸ unconventional T cells remain an emerging field of research. Despite their higher frequencies and broad tissue distributions,^{9,10} much less is known about the role and function of unconventional T cell subsets in disease and at steady state as compared to conventional T cells. Here, we provide an overview on possible concepts and review the current knowledge on how MHC-I-like presented allergen Ags cause hypersensitivities.

1.2 | Ag presentation by MHC-I-like molecules and their recognition by unconventional T cells

Peptide Ags, presented by classical MHC-I (also termed MHC-Ia) and MHC-II molecules, are recognised by conventional T cells. In




FIGURE 1 Overview of antigen presentation by MHC-I and MHC-I-like molecules and their recognition by T cells. Row 1 displays top-views onto the antigen cleft based on crystal structures of HLA-A*02:01 in complex with the cytomegalovirus pp65-derived peptide antigen NLV⁴⁹⁵⁻⁵⁰³ (PDB ID: 2X4R¹²⁷), as a representative of an MHC-Ia molecule; MR1 in complex with the bacterial/fungal small molecule metabolite antigen 5-(2-oxopropylideneamino)-6-D-ribitylaminoouracil (5-OP-RU) derived from a biosynthetic precursor to riboflavin (PDB ID: 4NQC¹⁶); CD1a in complex with the *Mycobacterium tuberculosis* lipid antigen Dideoxymycobactin (DDM) (PDB ID: IXZ0¹²⁸); and CD1d in complex with the marine sponge-derived lipid antigen α-galactosylceramide (α-GalCer) (PDB ID: 2PO6¹²⁹) or the endogenous lipid antigen sulfatide (PDB ID: 4MQ7³²). Row 2 shows chemical structures of the relevant antigens, followed by the name and approximate frequency of the relevant MHC-restricted T-cell type in row 3.⁹ The frequency of CD1d-restricted γδ T cells, most of which recognise endogenous lipid antigens and sulfatides, is estimated to be 0.05%-3.5% of CD1d-α-GalCer reactive cells.³¹ Row 4 shows schematics of the antigen presentation by antigen-presenting cells (APCs) and their recognition by T-cell receptors (TCRs) expressed by T cells. In each case, the antigen type, TCR usage and effector function molecules are highlighted.^{9,16,18,31,37,43,49,130-139} Row 5 includes phenotypic markers commonly used to identify these cells by flow cytometry^{131,140-143}

contrast, non-peptide Ags are recognised by distinct subsets of unconventional T cells, restricted by a number of Ag-presenting molecules that are homologues of classical MHC-I molecules, namely MHC-I-like molecules⁹ (Figure 1). Both conventional and unconventional T cells can express $\alpha\beta$ T-cell receptors (TCRs), whereas unconventional T cells can alternatively express a $\gamma\delta$ TCR.⁹ Some $\gamma\delta$ TCR⁺ unconventional T cells can recognise MHC-I-like molecules, whilst others are not restricted by MHC molecules^{9,11} (Figure 1). In contrast to MHC-Ia and MHC-II molecules which are highly polymorphic and thus present diverse Ags and vary significantly from one individual to the next, MHC-I-like molecules are typically monomorphic.⁹ In line with the monomorphic nature of the MHC-I-like molecules, where to date Ag diversity appears limited, unconventional T cells unlike conventional T cells can express limited TCR diversity, often featuring clonally expanded and similar, but nonidentical, TCR sequences (intradonor conservation).¹² Furthermore, similar TCRs can be found in nearly all individuals (interdonor conservation), allowing for public as compared to private immune responses at the population level.¹² In the following, we describe each set of unconventional T cells. Notably, the characteristics of unconventional T cells, including the breadth of Ags recognised, are an active area of research.

The MHC-I-like molecule MHC-related protein 1 (MR1) is expressed widely amongst nucleated cells and possibly in all tissues based on mRNA.¹³ MR1 presents small molecule metabolite Ags, derived from folic acid (non-stimulating)^{14,15} and a biosynthetic precursor to microbial riboflavin (stimulating), with 5-(2-oxopropylideneamino)-6-D-ribitylaminouracil (5-OP-RU) being the most potent Ag.¹⁶ MR1 is the restriction element of $\alpha\beta$ TCR⁺ mucosal-associated invariant T cells (MAIT cells), which are found in most tissues and represent up to 10% of peripheral blood T cells in humans¹⁷ (Figure 1), and 0.5–2% of T cells in a broad range of tissues in naïve C57BL/6 and BALB/c mice^{18,19} (Table 1). MAIT cells express a semi-invariant TCR α chain, composed of the variable region 1-2 and joining region 33 in humans (TRAV1-2-TRAJ33 and TRAV1-TRAJ33 in mice) (Figure 1, Table 1), where in humans TRAJ12 and TRAJ20 are also commonly incorporated but less frequently than TRAJ33.²⁰ Although the repertoire of TCR β chains of MAIT cells appears skewed, being dominated by TRBV6 and TRBV20 in humans (Figure 1) and TRBV19 and TRBV13 in mice (Table 1), there is appreciable variation in TCR β chain usage, especially within the hypervariable loop of the TCR β -chain, the complementarity-determining region 3 β (CDR3 β) loop.²⁰ After MR1-mediated stimulation via the TCR, human MAIT cells produce diverse cytokines, including interleukin 2 (IL-2), interferon- γ (IFN γ),

tumour necrosis factor (TNF), IL-17A¹⁷ and IL-13 during chronic stimulation²¹ and can be cytotoxic^{22,23} (Figure 1).

Cluster of differentiation 1 (CD1) molecules, CD1a, CD1b, CD1c and CD1d, present lipid Ags, with the unique size and architecture of each CD1 cleft, facilitating the capture and presentation of shared but predominantly distinct lipid Ags.^{12,24,25} As a whole, CD1a-, CD1b- and CD1c-restricted T cells range from 0.1% to 10% of T cells in human blood.⁹ CD1a, which amongst the CD1 molecules features the smallest Ag-binding cleft and shallow pockets, presents glycolipids as well as “headless” lipids and lipo-peptides such as dideoxymycobactin (DDM) to $\alpha\beta$ TCR⁺ CD1a-restricted T cells⁹ (Figure 1). CD1a-reactive cells are particularly abundant in blood and also in the skin where they produce IL-22 (Figure 1); and CD1a is expressed on dendritic cells and in high levels on Langerhans cells.¹² The TCR usage of CD1a-restricted T cells is largely unexplored, but many of these T cells appear autoreactive and current literature suggests that these cells participate in allergic responses,⁹ which will be discussed in further detail below.

Other CD1-restricted T cells that play a role in allergic reactions include CD1d-restricted natural killer T (NKT) cells (Figure 1). CD1d is expressed in many tissues such as the kidney, pancreas, breast and conjunctiva of the eye.²⁶ CD1d is broadly distributed on many haemopoietic cell types, including monocytes, macrophages, dendritic cells, B cells and non-haemopoietic cells such as epithelial cells of the gastrointestinal tract,²⁷ hepatocytes and keratinocytes.²⁸ CD1d is the restriction element of three subsets of NKT cells: (a) $\alpha\beta$ TCR⁺ type I NKT cells, (b) $\alpha\beta$ TCR⁺ type II NKT cells and (c) CD1d-restricted $\gamma\delta$ TCR⁺ T cells. Type I NKT cells recognise the marine sponge-derived glycolipid α -galactosylceramide (α -GalCer) and other α -linked glycolipids (Figure 1). Type I NKT cells make up ~0.1% of T cells in peripheral blood of humans (Figure 1), ~1% of T cells in most tissues of mice (Table 1) and up to 50% of all T cells in mouse liver.⁹ Type I NKT cells express an invariant TCR α chain (TRAV10-TRAJ18 in humans [Figure 1]; TRAV11-TRAJ18 in mice [Table 1]) paired with a limited array of TCR β chains and hence are also referred to as “invariant NKT cells” or “iNKT cells.” Type I NKT cells encompass distinct functional subsets that resemble Th1, Th2 and Th17 cells and predominantly express IFN γ , IL-4 and IL-17, respectively⁹ (Figure 1).

$\alpha\beta$ TCR⁺ type II NKT cells recognise various lipid Ags presented by CD1d, but not α -GalCer. Type II NKT cells are recognised as T cells that have much greater TCR diversity, compared with type I NKT cells and are hence also referred to as “diverse NKT cells” (Figure 1). Although type II NKT cells are less-well studied, these cells are thought to outnumber type I NKT cells in humans²⁹ and

TABLE 1 Unconventional T cell frequency and TCR usage in mice

MHC molecules	MR1	CD1d	
T cell subset	MAIT cell	Type I NKT cell	Type II NKT cell
Cell frequency	0.5%–2%	1%	Not well defined
TCR usage	TRAV1 (V α 19)- TRAJ33 (J α 33) TRBV19 (V β 6.1)/13 (V β 8)	TRAV11 (V α 14)- TRAJ18 (J α 18) TRBV13 (V β 8)/29 (V β 7)/1 (V β 2)	Predominantly TRAV9(V α 3)/7 (V α 1)- TRAJ7/9 (J α 7/9) TRBV13-3 (V β 8.1)/26 (V β 3.1)-J β 2.7

some studies suggest that type I and type II NKT cells have opposing roles, reviewed in.³⁰ A subset of $\gamma\delta$ TCR⁺ T cells in humans (predominantly TRDV1⁺ $\gamma\delta$ T cells) can also recognise CD1d- α -GalCer,³¹ whilst other $\gamma\delta$ T cells recognise the endogenous lipid Ag sulfatide.³² $\gamma\delta$ TCR⁺ NKT cells are also found in mice at a frequency of <1% in liver and spleen (PMID: 11592065). Some murine $\gamma\delta$ TCR⁺ cells recognise cardiolipin, a major phospholipid in bacterial cell walls, in the context of CD1d (ref: PMID: 21389252). It is unclear if CD1d-restricted type II NKT cells and $\gamma\delta$ T cells contribute to hypersensitivity; therefore, this review will mainly focus on the role of type I NKT cells in these aberrant immune responses.

In the following, we highlight some of the key features of unconventional T-cell responses as compared to those by conventional T cells. Precursor frequencies of Ag-specific unconventional T cell subsets in immune tissues are generally much higher ($\sim 1\text{--}10 \times 10^3$ per million of human T cells) than those of conventional T cells (1–10 per million of human T cells)⁹ (Figure 1). Furthermore, in non-lymphoid tissues, at the site of infection, unconventional T cells are often present in relatively high frequencies at steady state.⁹ Naïve conventional T cells require Ag contact in secondary lymphoid organs, a process termed priming, which leads to activation followed by clonal T cell expansion and differentiation over the course of 3–7 days.³³ In contrast, driven by the upregulation of the master transcription factor promyelocytic leukaemia zinc finger (PLZF) during thymic development,^{34–36} MAIT cells and type I NKT cells (and possibly other unconventional T cells) acquire a “preprimed”

state, that is somewhere in between the states of a naïve and effector-memory conventional T cell. Whilst this state is not fully characterised, both unconventional T cell subsets express markers broadly consistent with conventional effector-memory T cells (human MAIT cells: CD45RA⁺CD45RO⁺CD95^{hi}CD62L^{lo},^{36–38} human type I NKT cells: CD45RA^{dim}CD45RO⁺CD62L^{–/lo39}). In particular for MAIT cells, this preprimed state evolves further following thymic egress, when MAIT cells expand in the periphery,^{36,38} probably in response to commensal flora. In mice, type I NKT cell expansion, lineage commitment and acquisition of a preprimed state occur during thymic development and are independent of exogenous Ag exposure.^{40,41} However, for most NKT cells, the upregulation of NK1.1 and further maturation occurs in the periphery.^{41,42} Similarly, in humans, the peripheral environment contributes to both maturation and expansion of type I NKT cells.⁴³ Consistent with their “preprimed” state, whilst present in low frequencies in naïve mice,^{18,19} upon Ag exposure MAIT cells rapidly expand to large numbers⁴⁴ and produce cytokines.¹⁸ Human MAIT cells also rapidly produce cytokines upon Ag recognition.^{17,45} It has further been shown in mice that the MAIT cell effector response involves the formation of a long-lived population with memory-like recall properties and characterised by a polarised and more potent immune response upon restimulation.^{44,46} Similarly, mature type I NKT cells expand and produce cytokines rapidly in response to Ag^{47–50}; however, unlike MAIT cells, these cells contract to a pre-stimulation frequency over subsequent days,⁵¹ indicating that priming does not lead to memory formation. The development of




FIGURE 2 Possible scenarios of allergen antigen presentation by MHC-I-like molecules. A schematic of the four possible scenarios of allergen antigen display by MHC-I-like molecules in comparison with microbial/endogenous antigen presentation (left column). (i) The allergen antigen replaces the microbial/endogenous antigen. (ii) The allergen antigen and a microbial or endogenous antigen are simultaneously presented, but as distinct entities. The repertoire of microbial or endogenous antigens may be identical to that presented in the absence of the allergen antigen or distinct. (iii) The allergen antigen is directly conjugated to an antigen (iii), selected from the microbial or endogenous Ag repertoire that is either identical or distinct to that presented in the absence of the allergen antigen. (iv) The allergen might “act on” endogenous material, eliciting presentation of endogenous antigens (in the absence of allergen antigen and referred to as neoantigens), that would not be presented in steady state or that are normally presented at very low levels

other CD1-restricted T cells is not well understood, although some studies suggest CD1a-, CD1b- and CD1c-restricted T cells may exit the thymus as naïve T cells (CD45RA⁺CD45RO⁻) and follow a similar pathway as conventional T cells with regards to the following: priming⁵²; clonal expansion⁵³; and memory formation.⁴³

2 | SPECULATION ON POSSIBLE SCENARIOS OF ALLERGEN Ag PRESENTATION BY MHC-I-LIKE MOLECULES AND T-CELL RECOGNITION

Drawing on a limited number of published examples of allergen Ag presentation by MHC-I-like molecules as well as known concepts of allergen Ag presentation by classical MHC molecules, the following different scenarios of allergen Ag presentation by MHC-I-like molecules can be envisaged: (i) The allergen Ag may displace the microbial or endogenous Ag in the Ag-binding cleft (Figure 2i). This has been demonstrated for MR1 presentation of drugs, drug-metabolites and drug-like molecules⁵⁴ as well as for CD1a presentation of the poison ivy allergen-derived antigen urushiol⁵² and farnesol present in cosmetics and perfumes⁵⁵ (described in detail below). (ii) The allergen Ag and a microbial or endogenous Ag are simultaneously presented, but as distinct entities (Figure 2ii). This scenario would be similar to the altered repertoire concept for delayed-type hypersensitivity (DTH), involving conventional T cell-mediated, drug-specific recall responses⁵⁶: an altered repertoire of endogenous peptides was simultaneously presented with abacavir by the MHC-Ia molecule HLA-B*57:01 and carbamazepine by the MHC-Ia molecule HLA-B*15:02.⁵⁷

(iii) In a yet different scenario, the allergen Ag might be directly conjugated to an Ag. Again, the allergen Ag may be conjugated to an Ag selected from the microbial or endogenous Ag repertoire that is either identical or altered to that presented in the absence of the allergen Ag (Figure 2iii). This scenario would be similar to the hapten concept for DTH, in which a drug covalently bound to a peptide, a drug-haptenated peptide, is presented by the MHC-Ia molecule.^{56,58} If not the parent drug itself but a metabolite of the parent drug is bound to the peptide this is referred to the prohaptens concept.⁵⁶

Whilst scenarios (ii) and (iii) have not been described yet for MR1 or CD1, they may be possible: the Ag-binding cleft of MR1 has sufficient plasticity and versatility within the A'-pocket (equivalent to the MHC-Ia pocket that binds the N-terminal peptide residue) to accommodate diverse chemical scaffolds.⁵⁴ In addition, Ags or parts of Ags might be accommodated in the F'-pocket of MR1 (equivalent to the MHC-Ia pocket that binds the C-terminal residue of peptides).²⁰ Similarly, the Ag-binding pockets of the various CD1 molecules are larger in volume than the Ag-binding clefts of MHC and MR1 and are capable of binding multiple lipid species simultaneously, as well as accommodating lipids of greater volume than the Ag-binding pocket itself.¹²

(iv) In a last possible scenario, the allergen might elicit presentation of endogenous Ags (in the absence of allergen Ag and referred

TABLE 2 Hypersensitivities related to allergen recognition by unconventional T cells

Allergen category	Cell type	Potential allergy/ allergen
Environmental	CD1a-restricted T cells	Bee/wasp ^{59,65}
		House dust mites ⁶⁶
		Urushiol ⁵²
		Cypress pollen ⁷¹
	CD1d-restricted T cells	Cypress pollen ⁷¹
		NKT cells
	MAIT cells	Olive pollen ⁷²
		Ragweed pollen ⁷³
		Asthma ^{60,78-83,126}
Asthma ^{85-86,127}		
Food	NKT cells	Brazil nut ⁸⁹
		Cow's milk ⁹⁰⁻⁹⁴
	γδ T cells	Gluten ⁹⁷⁻¹⁰⁰
		Peanut ¹⁰¹
		Metal
Chromium ¹¹⁵		
Drug	Type II NKT cells	PPBF ¹¹⁶
	MAIT cells	Diclofenac ⁵⁴

to as neoantigens), that would not be presented at steady state or that are presented at very low levels at steady state (Figure 2iv). For instance, allergen-derived enzymes create neoantigens for presentation by CD1a⁵⁹ (see below). Or, as it has been speculated, allergens cause inflammation and dysregulation in the mucosa leading to the release or synthesis of endogenous Ags for Ag display.⁶⁰

For any of the given allergen Ag presentation scenarios described above, different scenarios of T-cell recognition can be extrapolated from published examples of T-cell recognition of allergen Ags presented by MHC-I-like as well as classical MHC molecules. (i) There might be an overlap in the T-cell clones that recognise the microbial or endogenous Ags as well as the allergen Ags presented by MHC-I-like molecules, that is they cross-react with allergen Ags. The entire repertoire of clonally distributed TCRs or only a subset of TCRs might cross-react with the allergen Ag. For example, only a subset of 5-OP-RU-specific MAIT TCRs cross-reacted with metabolites of the drug diclofenac and responding MAIT cells reacted to different diclofenac metabolites.⁵⁴ (ii) Alternatively, a new, distinct repertoire of TCRs might cross-react with the allergen Ag, as seen for recognition of peptides co-presented with abacavir and carbamazepine.⁵⁷ (iii) Whilst there are currently no examples of T-cell cross-reactivities between classical MHC and MHC-I-like molecules, or amongst different MHC-I-like molecules, these types of cross-reactivity might occur: MR1-reactive MAIT-like cells, atypical MAIT cells and MR1T cells have been described, some of which do not express the invariant MAIT TCR.^{61,62} These cells do not possess the same characteristics as invariant MAIT cells and some of these are likely conventional T cells that cross-react with MR1.⁶¹ Interestingly, in MAIT TCR transgenic mice that lack MR1, a significant population

of "MAIT-like" T cells develops, apparently selected by MHC-Ia molecules or CD1d.⁶³ Similarly, in *Mus musculus castaneus* (CAST mice), modified to lack MR1, a small population of TRAV1-TRAJ33 expressing cells was identified that was selected by other MHC molecules, possibly MHC-II molecules.⁶⁴

3 | RECOGNITION OF ALLERGEN Ags BY UNCONVENTIONAL T CELLS

In the following, we provide an overview on the current stage of the literature on environmental, food, metal and drug allergen Ag recognition by unconventional T cells, summarised in Table 2.

3.1 | Environmental allergens

CD1a-restricted T-cell responses to lipid Ags derived from environmental allergens such as bee and wasp venom,^{59,65} aerosolised

extracts from house dust mites (HDM)⁶⁶ and tree saps⁵² have been described and may contribute to skin-related hypersensitivity reactions in predisposed individuals. During a bee/wasp sting, phospholipase (PLA) enzymes present within venom were introduced in the skin as allergenic products. PLA catalysed the cleavage of host phospholipids into antigenic neolipids, including lysophosphatidylcholine (LPC), that were presented by CD1a (Figure 3A) and potentially stimulated skin-derived CD1a-restricted T cells.⁵⁹ Similarly, a skin-topical CD1a-restricted T-cell response to HDM extract as allergen has been attributed to both HDM- and host-derived PLA activities.^{66,67} CD1a-restricted T cells reactive to bee venom and HDM were more frequent in the peripheral blood of hypersensitive individuals than non-hypersensitive controls,^{65,66} suggesting that CD1a-restricted T cells may contribute to the hypersensitivity response. In support of this, the frequency of IFN γ -producing CD1a-restricted T cells increased significantly in bee sting hypersensitive patients during desensitisation therapy.⁶⁵ In addition, HDM-responsive CD1a-restricted T cells were also responsive to bee-derived PLA.⁶⁶ These data align closely with another study that identified an increased




FIGURE 3 Allergen antigen presentation by CD1a and MR1. (A) Chemical structures of the *Mycobacterium tuberculosis*-derived antigen Dideoxymycobactin (DDM), the bee/wasp venom allergen-derived antigen Lysophosphatidylcholine (LPC), the poison ivy allergen-derived antigen urushiol (C15:2) and the allergen antigen farnesol contained in cosmetics/perfumes and their presentation by CD1a (PDB IDs: 1XZ0,¹²⁸ 4X6E,⁶⁹ 5JIA,⁵² 6NUX⁵⁵), displaying top-views onto the CD1a-antigen complexes. (B) Chemical structures of the bacterial/fungal riboflavin biosynthesis derived antigen 5-(2-oxopropylideneamino)-6-D-ribitylamouracil (5-OP-RU), the drug-like small molecule antigen 3-formyl-salicylic acid (3-F-SA) and the diclofenac drug metabolite antigen 5-hydroxy-diclofenac (5-OH-DCF) and their presentation by MR1 (PDB IDs: 4NQC,¹⁶ 5U6Q,⁵⁴ 5U72⁵⁴), displaying top-views onto the MR1-antigen complexes

frequency of autoreactive CD1a-restricted T cells in psoriasis patients compared with controls.⁶⁸ Similarly, stimulatory neolipids were generated by host-derived PLA activity and in this case, localised to the dermis of psoriasiform lesions.⁶⁸ Thus, the aberrant activity of host-derived PLAs, triggered on an environmental and/or genetic basis, appears to be a shared factor in CD1a-auto- and allergen-reactivity.

Direct recognition of allergen-derived lipids has also been established for CD1a-restricted T cells in response to the poison ivy-derived lipid, urushiol.⁵² In sensitised mice, urushiol treatment caused an inflammatory response that was greater in CD1a-transgenic mice compared with CD1a-deficient wild-type mice and was characterised by IL-17 and IL-22 production by clonally expanded (utilising TCR β -chain TRBV1 or TRBV2 gene segments) CD4⁺ T cells.⁵² Similarly, in hypersensitive donors, urushiol displayed a greater frequency of IL-17- and IL-22-producing CD1a-responsive T cells than control donors.⁵² Structural determination of the dominant antigenic species of urushiol (C15:2), presented by CD1a, revealed that the lipid was buried deep within the cleft of CD1a, positioned such that the lipid was only minimally exposed for TCR recognition,⁵² perhaps indicating a minor role for urushiol in overall TCR binding to CD1a (Figure 3A). Structural analysis of an autoreactive TCR (isolated from an autoimmune patient) bound to CD1a presenting endogenous lipids revealed that small, solvent protected CD1a-binding lipids that did not appreciably alter the conformation of CD1a were permissive of TCR recognition.⁶⁹ In contrast, lipids with longer acyl chains or a bulky head group disrupted this interaction.⁶⁹ Importantly, the molecular contacts occurred exclusively between TCR and CD1a protein itself, indicating that the lipid Ag was not surveyed by the TCR directly⁶⁹ and suggestive of a model of TCR-CD1a recognition described as "absence of interference".⁷⁰ It is therefore possible that these CD1a-autoreactive T cells may respond to both CD1a-urushiol and other allergenic small lipids that do not disrupt the TCR-CD1a interface.

In support of this hypothesis, several small hydrophobic compounds, derived from cosmetics and perfumes were recently shown to stimulate T cells in a CD1a-dependent manner.⁵⁵ This included farnesol, related to farnesyl pyrophosphate, the biosynthetic precursor of the human skin oil squalene. Farnesol was able to stimulate polyclonal T cells *in vitro* when presented by CD1a in some donors.⁵⁵ Structural analysis revealed farnesol occupied only 36% of the CD1a cleft (Figure 3A) and was likely protected entirely from TCR surveillance.⁵⁵ Despite its small size, farnesol was able to displace larger, amphipathic self-lipids from CD1a, thus restoring "absence of interference" conducive to TCR recognition.⁵⁵

CD1a-restricted T cells and CD1d-restricted NKT cells have also been suggested to play a role in airway hypersensitivity reactions to plant pollens and other aerosolised allergens. In a study examining the cypress pollen, CD1a and CD1d molecules expressed by human pulmonary DCs were able to directly capture the pollen particles.⁷¹ Analysis of the lipids extracted from the cypress pollen revealed a range of phospholipid species, of which phosphatidylcholine (PC) and phosphatidylethanolamine (PE) were demonstrated to be

antigenic, and stimulated T-cell clones derived from pollen hypersensitive donors in a CD1-dependent manner.⁷¹ Similarly, olive pollen extract upregulated CD1d expression on human monocytes and macrophages, and NKT cells were able to lyse Ag-presenting cells (APCs) cultured with olive pollen lipid extract,⁷² suggesting that pollen-derived lipids may be directly recognised by CD1-restricted T cells. In a mouse model, intranasal administration of the allergen ragweed (RW) pollen in mice deficient of NKT cells reduced the typical hypersensitive response, measured by pulmonary mucus production, serum IgE and circulating eosinophils,⁷³ suggesting that NKT cells are involved in the hypersensitive response to RW pollen. Interestingly, wild-type mice treated with α -GalCer prior to sensitisation with RW further exacerbated the hypersensitive response compared with non-treated mice.⁷³ Some inroads have been made to identify key factors responsible for regulating NKT cell pathogenicity during hypersensitive responses. Recently, the histone methyltransferase enhancer of zeste homolog 2 (EZH2) was shown to be important in the differentiation of pathogenic NKT cells in an airway hyperresponsiveness (AHR) mouse model.⁷⁴ Conditional deletion of EZH2 in CD4⁺ T cells induced spontaneous AHR that relied predominantly on NKT cell IL-4 secretion, which was significantly increased in these animals compared with wild-type mice.⁷⁴ In humans, expression of EZH2 was shown to be reduced in blood Th1 and Th2 cells of allergic rhinitis (AR) patients compared with controls and EZH2 expression was negatively correlated with serum IL-17A in AR patients challenged with HDM.⁷⁵ Thus, EZH2 appears to be important for the epigenetic control of T helper^{74,75} and NKT cell differentiation⁷⁶ in allergic responses. Similarly, NKT cells were shown to be sensitive to the anti-inflammatory effects of the histamine receptor 2 (H₂R) in AHR mouse models.⁷⁷ Pharmacological activation of H₂R attenuated NKT cell lung accumulation and the inflammatory response in AHR, whereas inhibition or genetic deletion of H₂R significantly enhanced NKT cell pathology, suggesting that histamine signalling may be a key modulator of NKT cells during an allergic response.⁷⁷

Many studies have examined the role of NKT cells in the development of allergic asthma using an ovalbumin (OVA)-induced mouse model. In sensitised mice, challenge with OVA-induced AHR characterised by airway inflammation, leucocyte infiltration and increased serum IgE.^{60,78} Amongst studies that are reviewed elsewhere,⁷⁹⁻⁸¹ the consensus is that whilst OVA-induced AHR is not directly mediated by NKT cells, the disease is significantly attenuated in the absence of NKT cells, such as in CD1d^{-/-} and J α 18^{-/-} mice as well as in mice treated with anti-CD1d antibodies to deplete NKT cells.^{60,78,82} In spite of these findings, a subset of NKT cells has been described that appears to suppress OVA-induced AHR.⁸³ This suppressive subset of NKT cells was CD38⁺ and CD4/CD8 co-receptor deficient and could be expanded in neonatal mice after infection with influenza A virus or upon stimulation with an α -GalCer analogue.⁸³ In the absence of an exogenously administered lipid Ag in the OVA-induced AHR model, some have speculated that NKT cells may become exposed to antigenic endogenous lipids in the allergen-induced inflammatory mucosal environment.⁶⁰

The role of MAIT cells in asthma is unclear; however, clinical evidence suggests MAIT cells can be both protective and pathogenic in asthma. In asthmatic adults, disease severity correlated with a lower frequency of MAIT cells in lung biopsies, blood and sputum samples.⁸⁴ A potential protective role of MAIT cells in asthma was also observed in a population of infants, whereby lower frequencies of circulating MAIT cells in 1-year-old children correlated with the later development of asthma at age 7.⁸⁵ However, higher frequencies of circulating IL-17A producing MAIT cells were associated with severe asthma compared with non-severe asthma, suggesting a pathogenic role for this MAIT cell subset.⁸⁶ The observed differential roles of MAIT cells in asthma might be related to age, ethnicity and genetic differences, as well as to the exposure of distinct sets of environmental Ags,^{87,88} which may include unknown MAIT cell Ags.

3.2 | Food allergens

NKT cells have been implicated in the recognition of food-derived allergens, including from Brazil nuts,⁸⁹ cow milk⁹⁰⁻⁹² and other mammal milks.⁹³ For the study of Brazil nut hypersensitivity, a mouse model of allergic disease has been established.⁸⁹ In this model, the purified Brazil nut protein, Ber e 1, acted as a sensitising Ag in conjunction with a fractionated Brazil nut lipid extract to stimulate a Th2-related antibody response.⁸⁹ NKT cell-deficient mice displayed reduced antibody responses including those specific to Ber e 1, compared with wild-type mice, suggesting a role for NKT cells in the generation of Ber e 1-specific antibodies, possibly involving the secretion of IL-4.⁸⁹ Analysis of the lipid fraction revealed a mixture of neutral and polar lipids, including CD1d-binding phospholipids,¹² of which phosphatidylethanolamine (PE) and phosphatidylinositol (PI) species were most abundant.⁸⁹ In nut hypersensitive donors, Brazil nut lipid extract-responsive CD161⁺ T cells were enriched for NKT cells.⁸⁹ Thus, NKT cells may contribute to the hypersensitivity reaction to Brazil nuts in allergic patients, specifically by responding to CD1d-presented lipids derived from the allergen.

A number of studies have assessed the reactivity of NKT cells to common mammalian lipids found in dairy products, most notably cow's milk,^{90,91} with a focus on two candidates for inducing NKT cell antigenicity: sphingomyelin (SM)^{91,94} and β -glucosylceramide (β -GluCer).^{90,93} In one study, NKT cells were found to be significantly less frequent in the peripheral blood of children with a milk allergy compared with non-allergic children and tended to produce more IL-13 in response to a milk-derived SM, suggesting a skewed functional response to milk SM by NKT cells in hypersensitive children.⁹¹ SM was shown to be recognised in the context of CD1d by a modest subset of NKT cells in the peripheral blood of non-allergic donors and was a less potent Ag than α -GalCer.⁹¹ Similarly, in a cohort of children diagnosed with eosinophilia oesophagitis (EoE), an IgE-mediated and food-related atopic disease, NKT cells were less frequent in the peripheral blood of children with active compared to controlled disease or in healthy donors.⁹⁴ Further, a greater frequency of NKT cells was reported in oesophageal biopsies from children with active

compared to controlled disease,⁹⁴ indicative of NKT cell recruitment to the oesophagus during active disease. In all children, NKT cells proliferated in response to milk SM, yet significantly more NKT cells from active EoE produced IL-4 and IL-13 than control children.⁹⁴ Therefore, NKT cells in two separate cohorts appeared responsive to milk SM, with some evidence of CD1d-dependent recognition.

In terms of cow's milk β -GluCer, it initially appeared that mouse NKT cells were stimulated by β -GluCer in a CD1d-dependent manner, and two relevant β -GluCer species were identified as candidate Ags (C12:0 and C24:1).⁹⁰ However, digestion of mouse and human milk lipid extracts with an enzyme that cleaves β -linked GluCer did not abolish the broad reactivity to the milk lipids displayed by NKT cells, suggesting that the stimulatory response was not caused by β -GluCer, rather, a naturally occurring α -linked lipid present in the milk lipid extract. In a further analysis, the minor lipid species type 2 α -linked monohexosylceramide was identified from cow's milk which was recognised strongly by a murine type I NKT TCR when bound to CD1d, similarly to α -GalCer bound CD1d.^{92,95}

Food hypersensitivity has also been associated with $\gamma\delta$ T-cell recruitment, despite an incomplete understanding of the underlying immunological mechanisms. Biopsies of inflamed gut tissues showed increased infiltration of $\gamma\delta$ T cells, predominantly V δ 1, in the terminal ileum and duodenum associated with allergic reactions, when compared to non-allergic and chronic inflammatory diseases.⁹⁶⁻⁹⁹ Also, increased numbers of intraepithelial $\gamma\delta$ T cells have been suggested as a biomarker for the diagnosis of coeliac disease when histology is inconclusive.^{98,100}

However, in a mouse model of food allergy, the allergic sensitisation induced by co-administration of cholera toxin and peanut antigens was followed by a decrease of number and proportion of $\gamma\delta$ T cells in the intestine.¹¹ In the same study, the functional depletion of $\gamma\delta$ T cells, through co-administration of anti- $\gamma\delta$ TCR blocking antibody during the sensitisation, led to a higher production of Th2 cytokines and peanut-specific IgE.¹¹ Whilst this suggested a protective role for $\gamma\delta$ T cells in allergic sensitisation in this model, the involvement of the $\gamma\delta$ TCR in the activation of the regulatory response is not clear.¹¹

Given that MAIT cell Ags are small molecule metabolites, it is tantalising to speculate that MAIT cells might also recognise small molecule food metabolites and this way causes food hypersensitivities or intolerances. Food-derived flavonoids, for example, represent likely MR1 Ag candidates due to their chemical structures resembling known MAIT cell Ags. Indeed, dietary isoflavone intake has been associated not only with anti-inflammatory, but also pro-inflammatory effects in the gastrointestinal tract.¹⁰²⁻¹⁰⁴

3.3 | Metal allergens

Nickel (Ni)-induced allergic contact dermatitis (ACD) is the most common metal allergy in humans¹⁵ and invokes both innate and adaptive immune responses that are not fully understood.¹⁶ During sensitisation, Ni metal ions bind to a histidine-rich motif of TLR4 and trigger an inflammatory response.¹⁷ In mice, TLR4 lacks

the Ni ion-binding site.¹⁷ However, Ni hypersensitivity in mouse models can be induced by co-administration of Ni with a classical TLR4 agonist such as lipopolysaccharide (LPS).¹⁷ In mice sensitised with Ni and LPS, subsequent footpad challenge with a Ni solution caused a sustained inflammatory response as well as accumulation of T cells in the footpad epithelial basal layer.¹⁸ Analysis of the TCR repertoire from the footpads of challenged mice showed a bias in NKT TCR gene segment usage (TRAV11 and TRBV13-2) and a clonotypic invariant NKT TCR CDR3 α chain,¹⁸ suggestive of NKT cell accumulation in the inflamed footpad. In another mouse model of Ni hypersensitivity, sensitised NKT cell-deficient mice displayed significantly increased ear swelling upon Ni challenge, compared with wild-type mice.¹⁹ The difference in swelling was reduced 96 hours after challenge, suggesting that NKT cells may contribute early in the immune response.¹⁹ Interestingly, when mice were treated with α -GalCer at the same time as receiving Ni challenge, ear swelling induced by Ni was significantly reduced at 24 hours.¹⁹ Together, these data suggest NKT cells may dampen the Ni hypersensitive response, particularly after direct activation with potent Ag.

Further, MAIT cells might be implicated in human Ni hypersensitivity, where preferential activation of human $\alpha\beta$ CD8⁺ T cells expressing a selected TCR-V β repertoire, including TRBV6 and TRBV20, has been observed¹¹⁰ which is typical for MAIT cells; the TCR-V α repertoire was not assessed in this study. In this regard, two Ni-reactive CD8⁺ T-cell clones isolated from sensitised patients showed Ni-specific activation independently of MHC-Ia, MHC-II, or CD1d. Earlier studies have demonstrated the potential of these clones to proliferate and to mediate specific cytotoxicity of different human cell lines in a TCR-dependent way in the presence of Ni-sensitised APCs.^{111,112}

MHC-Ia-independent activation of CD8⁺ T cells in response to gold-sensitised APCs is likely related to NKT cell activation.¹¹³ Indeed, in biopsies of skin lesions, NKT cell infiltration was suspected based on increased levels of CD161 and CD1d expression.¹¹⁴ In addition, in a mouse model of chromium hypersensitivity, NKT cells accumulated in inflamed skin.¹¹⁵ Together, these results underline the key role that NKT cells play in metal allergy pathogenesis.

3.4 | Drug allergens

A study by Van Rhijn and colleagues¹¹⁶ identified a type II NKT cell clone that in the context of CD1d could recognise a non-lipid molecule called phenyl 2,2,4,6,7-pentamethyldihydrobenzofuran-5-sulphonate (PPBF), which resembles sulpha drugs that induce hypersensitivity in some individuals.¹¹⁶ Data suggested that PPBF bound in or near the CD1d Ag cleft.¹¹⁶ Precisely how PPBF binds to CD1d and is recognised by the type II NKT cell clone remains unclear.

The capacity of MAIT cells to recognise small molecules has prompted the proposition that MAIT cells are involved in drug hypersensitivities.⁵⁴ Following multiple parallel *in silico* screens of 6000 in-house organic compounds and 1216 drugs (approved by the US Food and Drug Administration), 183 candidate molecules were

identified. Of those, 81 were subjected to cellular assays assessing MAIT cell activation and MR1 binding. A quarter of the tested drugs, drug-metabolites and drug-like molecules were able to bind to MR1 and/or activate MAIT TCR reporter cell lines.⁵⁴ 3-formyl-salicylic acid, a synthetic analogue of salicylate (aspirin) which strongly bound MR1 (Figure 3B) but did not activate human or mouse MAIT cells, competitively inhibited MAIT cell activation by 5-OP-RU, demonstrating the capacity of drug-like small molecules to modulate MAIT cell function.⁵⁴ Consequently, it was demonstrated that MR1 can capture chemically diverse scaffolds. These observations indicate that some drugs and drug-like molecules affect MAIT cell function.⁵⁴

Further, MAIT TCR reporter cell lines responded to diclofenac at a concentration that can be achieved in patients after an oral dose.^{54,117} The activation of MAIT cells was attributed to diclofenac metabolites, specifically to 4- and 5-hydroxy-diclofenac (4-OH-DCF, 5-OH-DCF) (Figure 3B). Not all 5-OP-RU-specific MAIT TCR reporter cell lines responded to 4-OH-DCF and 5-OH-DCF, suggesting the involvement of specific subsets of MAIT cells in their respective recognition.⁵⁴ Hypersensitivity to diclofenac has been attributed to 5-OH-DCF in mouse model studies.¹¹⁸ Moreover, the cytotoxic activity of diclofenac-activated T cells against sensitised hepatocytes was only partially MHC-Ia-dependent,¹¹⁹ in line with a potential role for other T cells such as MAIT cells in diclofenac hypersensitivity. Whilst clinical studies are needed, it is possible that drugs and drug-metabolites modulate MAIT cell function, causing potentially drug hypersensitivities.

4 | SPECULATION ON THE CLINICAL IMPACT AND HALLMARKS OF UNCONVENTIONAL T CELL-MEDIATED HYPERSENSITIVITIES

4.1 | Speculations on frequencies of hypersensitivities at the population level, involving allergen Ags presented by MHC-I-like Ag-presenting molecules

Given the donor-unrestricted nature of MHC-I-like molecules and unconventional T cell subsets,¹² a given allergen Ag would be predicted to be presented and recognised by most/all donors. Consequently, whilst not enumerated yet, hypersensitivities directed to MHC-I-like molecules should be more frequent than those directed to classical MHC molecules. Unconventional T cell allergen Ag recognition could however be donor restricted, such as in the following cases:

- A donor-specific microbial or endogenous Ag is presented simultaneously, as per allergen Ag presentation scenarios (ii) and (iii).
- The allergen Ag recognition is mediated by conventional T cells which are donor restricted, as per allergen Ag T cell recognition scenario (iii).
- Relevant private, donor-specific unconventional T cell repertoires exist.

The latter is exemplified by microbial GMM-reactive CD1b-restricted T cells that in many donors comprise a public TCR repertoire (eg GEM T cells) as well as private TCR repertoire.^{53,120,121} Further, it is unclear which other immunological mechanisms are in place to prevent or encourage reactions to allergens presented by MHC-I-like molecules. Some unconventional T cell responses appear to correlate with genetic polymorphisms established for allergic disease, such as the *Filaggrin-null* mutation that is associated with atopic dermatitis severity.⁶⁶ This also resonates with conventional T cell-mediated hypersensitivities. For instance, in the case of abacavir hypersensitivity, ~ 40% of abacavir treated individuals that are HLA-B*57:01⁺ tolerate abacavir.¹²² So, whilst HLA-B*57:01 is necessary for abacavir hypersensitivity, it is not sufficient to allow for hypersensitivity reactions to occur. In summary, whilst unconventional T cells are donor-unrestricted, genetic predispositions likely play a major role in the involvement of unconventional T cells in allergic disease. Notably, in the case of drug hypersensitivities, actual frequencies of hypersensitive individuals could be masked as presumably drugs that cause a strong allergic response in most individuals would not pass clinical testing towards drug approval.

4.2 | Expected hallmarks of hypersensitivities involving T-cell responses to allergen Ags presented by MHC-I-like molecules

In the context of a type IV hypersensitivity, priming of conventional T cells occurs in the lymph nodes upon the first allergen exposure, which is often referred to as Ag sensitisation. Following a second allergen exposure, effector responses are detected with a delay of 24–72 hours, a T-cell response termed DTH.¹²³ In contrast, naïve (or preprimed) MAIT cells and type I NKT cells are present at high frequencies at steady state in most, if not all tissues.⁹ Additionally, MHC-I-like molecules are broadly expressed across tissues.^{13,23,124} Therefore, one might speculate that MAIT cells and type I NKT cells are poised to encounter allergen Ag directly at these sites. Moreover, given the preprimed nature of these cells, priming might not be needed to the same extent, resulting in a more rapid effector response.

Given the unique chemical properties of the Ag-binding clefts of the various MHC-I-like molecules, different classes of non-protein molecules represent allergen Ag candidates for each MHC-I-like molecule. For example, small molecules including pyrimidines, phenols/anilines, enones, aromatic aldehydes, aromatic carboxylates, quinones, flavones, and isoflavones (150–400 Da) can be presented by MR1,²⁰ whilst a diverse repertoire of endogenous lipids can bind mutually to the different CD1 molecules.^{12,24,125} The class of allergen Ag in turn is recognised by a subset of T cells restricted by the relevant MHC-I-like molecule, thus eliciting a hypersensitivity reaction, in line with the functional capacity of the relevant T-cell subset. Thus, a range of type IV hypersensitivity responses

Box 1 Future research perspectives (open areas for future research)

In the last decade, there have been significant advances, demonstrating allergen Ag display of lipids by CD1 and small molecules by MR1. Our emerging knowledge of the function of unconventional T cells and development of better reagents will facilitate allergen Ag discoveries and delineation of the underlying mechanisms governing hypersensitivity responses. More studies into the clinical relevance, especially in the case of potential MAIT cell-mediated hypersensitivities to small molecules, are needed. Could some of the suspected T cell-mediated hypersensitivities including those to antibiotics, for which mechanisms are unclear, be mediated by CD1- or MR1-restricted T cells? For example, metal hypersensitivities have often been described as TCR-dependent but classical MHC-independent, thus not excluding MHC-I-like molecules as targets. T-cell immunotherapies with allergen Ags relevant to MHC-I-like molecules would likely apply to the genetically diverse population. Like conventional T cell-based immunotherapies, they would lack IgE-binding capacity so that the risk of adverse reactions would be low3.

can be envisaged, similar to the existing subgrouping for conventional T cells (type IVa-d).² Type IV hypersensitivities can also be accompanied by IgE production, as described for iNKT cells in the OVA-induced AHR mouse model.^{60,78} Future research perspectives are discussed in Box 1.

ACKNOWLEDGMENTS

M. M., L. L. and S. B. G. E. would like to acknowledge research funding in the form of the 2019 AIFA (Allergy and Immunology Foundation of Australasia) research grant. AIFA is a charity initiative of ASCIA (Australasian Society of Clinical Immunology and Allergy), the not for profit peak professional body for clinical immunology and allergy in Australasia (www.allergyimmunology.org.au/about-aifa). MNTS is supported by an Australian Postgraduate Award, from the Australian Government, Department of Education. JM acknowledges funding from an NHMRC program grant (1113293). DGP is supported by a CSL Centenary Fellowship from CSL limited. SBGE is the recipient of a Discovery Early Career Research Award (DECRA) (DE170100407) from the Australian Research Council (ARC).

CONFLICT OF INTEREST

James McCluskey, Zhenjun Chen and Sidonia BG Eckle are inventors on patents describing MR1 antigens and MR1 tetramers. The other authors have no conflict of interest in relation to this work.

ORCID

Marcela de Lima Moreira  <https://orcid.org/0000-0003-3477-0727>

Michael N. T. Souter  <https://orcid.org/0000-0002-1459-7923>

Zhenjun Chen  <https://orcid.org/0000-0003-1205-683X>

Liyen Loh  <https://orcid.org/0000-0002-1676-0579>

James McCluskey  <https://orcid.org/0000-0002-8597-815X>

Daniel G. Pellicci  <https://orcid.org/0000-0001-5305-2398>

Sidonia B. G. Eckle  <https://orcid.org/0000-0003-0766-171X>

REFERENCES

- Gell PGH, Coombs RRA. *Clinical Aspects of Immunology*. 24–25, Broad Street. Oxford: Blackwell Scientific Publications Ltd.; 1963.
- Uzzaman A, Cho SH. Chapter 28: Classification of hypersensitivity reactions. *Allergy asthma Proc*. 2012;33:96–99.
- Woodfolk JA. T-cell responses to allergens. *J Allergy Clin Immunol*. 2007;119(2):280–294.
- Imai K, Kanno M, Kimoto H, Shigemoto K, Yamamoto S, Taniguchi M. Sequence and expression of transcripts of the T-cell antigen receptor α -chain gene in a functional, antigen-specific suppressor-T-cell hybridoma. *Proc Natl Acad Sci USA*. 1986;83(22):8708–8712.
- Fowlkes BJ, Kruisbeek AM, Ton-That H, et al. A novel population of T-cell receptor $\alpha\beta$ -bearing thymocytes which predominantly expresses a single V β gene family. *Nature*. 1987;329 (6136):251–254.
- Porcelli S, Yockey CE, Brenner MB, Balk SP. Analysis of T cell antigen receptor (TCR) expression by human peripheral blood CD4–8– $\alpha\beta$ T cells demonstrates preferential use of several V β genes and an invariant TCR α chain. *J Exp Med*. 1993;178(1):1–16.
- Budd RC, Miescher GC, Howe RC, Lees RK, Bron C, Robson MH. Developmentally regulated expression of T cell receptor β chain variable domains in immature thymocytes. *J Exp Med*. 1987;166(2):577–582.
- Bendelac A, Schwartz RH. CD4+ and CD8+ T cells acquire specific lymphokine secretion potentials during thymic maturation. *Nature*. 1991;353(6339):68–71.
- Godfrey DI, Uldrich AP, Mccluskey J, Rossjohn J, Moody DB. The burgeoning family of unconventional T cells. *Nat Immunol*. 2015;16(11):1114–1123.
- D'Souza MP, Adams E, Altman JD, et al. Casting a wider net: Immunosurveillance by nonclassical MHC molecules. *PLoS Pathog*. 2019;15(2):e1007567.
- Le Nours J, Gherardin NA, Ramarathinam SH, et al. A class of gd T cell receptors recognize the underside of the antigen-presenting molecule MR1. *Science*. 2019;66(6472):1522–1527.
- Van Rhijn I, Moody DB. Donor unrestricted T cells: a shared human T cell response. *J Immunol*. 2015;195(5):1927–1932.
- Lamichhane R, Ussher JE. Expression and trafficking of MR1. *Immunology*. 2017;151(3):270–279.
- Eckle SBG, Birkinshaw RW, Kostenko L, et al. A molecular basis underpinning the T cell receptor heterogeneity of mucosal-associated invariant T cells. *J Exp Med*. 2014;211(8):1585–1600.
- Kjer-Nielsen L, Corbett AJ, Chen Z, et al. An overview on the identification of MAIT cell antigens. *Immunol Cell Biol*. 2018;96(6):573–587.
- Corbett AJ, Eckle SBG, Birkinshaw RW, et al. T-cell activation by transitory neo-antigens derived from distinct microbial pathways. *Nature*. 2014;509(7500):361–365.
- Gherardin NA, Souter MNT, Koay HF, et al. Human blood MAIT cell subsets defined using MR1 tetramers. *Immunol Cell Biol*. 2018;96(5):507–525.
- Rahimpour A, Koay HF, Enders A, et al. Identification of phenotypically and functionally heterogeneous mouse mucosal-associated invariant T cells using MR1 tetramers. *J Exp Med*. 2015;212(7):1095–1108.
- Chen Z, Wang H, D'Souza C, et al. Mucosal-associated invariant T-cell activation and accumulation after in vivo infection depends on microbial riboflavin synthesis and co-stimulatory signals. *Mucosal Immunol*. 2017;10(1):58–68.
- Eckle SBG, Corbett AJ, Keller AN, et al. Recognition of Vitamin B precursors and byproducts by mucosal associated invariant T cells. *J Biol Chem*. 2015;290(51):30204–30211.
- Kelly J, Minoda Y, Meredith T, et al. Chronically stimulated human MAIT cells are unexpectedly potent IL-13 producers. *Immunol Cell Biol*. 2019;97(8):689–699.
- Le Bourhis L, Dusseaux M, Bohineust A, et al. MAIT cells detect and efficiently lyse bacterially-infected epithelial cells. *PLoS Pathog*. 2013;9(10):e1003681.
- Kurioka A, Ussher JE, Cosgrove C, et al. MAIT cells are licensed through granzyme exchange to kill bacterially sensitized targets. *Mucosal Immunol*. 2015;8(2):429–440.
- Shamshiev A, Gober HJ, Donda A, Mazorra Z, Mori L, De Libero G. Presentation of the same glycolipid by different CD1 molecules. *J Exp Med*. 2002;195(8):1013–1021.
- Chancellor A, Gadola SD, Mansour S. The versatility of the CD1 lipid antigen presentation pathway. *Immunology*. 2018;154(2):196–203.
- Canchis PW, Bhan AK, Landau SB, et al. Tissue distribution of the non-polymorphic major histocompatibility complex class I-like molecule, CD1d. *Immunology*. 1993;80(4):561–565.
- Blumberg RS, Terhorst C, Bleicher P, et al. Expression of a non-polymorphic MHC class I-like molecule, CD1D, by human intestinal epithelial cells. *J Immunol*. 1991;147(8):2518–2524.
- Bonish B, Jullien D, Dutronc Y, et al. Overexpression of CD1d by Keratinocytes in psoriasis and CD1d-dependent IFN- γ production by NK-T Cells. *J Immunol*. 2000;165(7):4076–4085.
- Exley MA, Tahir SMA, Cheng O, et al. Cutting edge: a major fraction of human bone marrow lymphocytes are Th2-Like CD1d-reactive T cells that can suppress mixed lymphocyte responses. *J Immunol*. 2001;167(10):5531–5534.
- Pellicci DG, Uldrich AP. Unappreciated diversity within the pool of CD1d-restricted T cells. *Semin Cell Dev Biol*. 2018;84:42–47.
- Uldrich AP, Le Nours J, Pellicci DG, et al. CD1d-lipid antigen recognition by the $\gamma\delta$ TCR. *Nat Immunol*. 2013;14(11):1137–1145.
- Luoma AM, Castro CD, Mayassi T, et al. Crystal structure of V δ 1T cell receptor in complex with CD1d-sulfatide shows mhc-like recognition of a self-lipid by human $\gamma\delta$ T cells. *Immunity*. 2013;147(8):2518–2524.
- Obst R. The timing of T cell priming and cycling. *Front Immunol*. 2015;6:563.
- Kovalovsky D, Uche OU, Eladad S, et al. The BTB-zinc finger transcriptional regulator PLZF controls the development of invariant natural killer T cell effector functions. *Nat Immunol*. 2008;9(9):1055–1064.
- Savage AK, Constantinides MG, Han J, et al. The transcription factor PLZF directs the effector program of the NKT cell lineage. *Immunity*. 2008;29(3):391–403.
- Martin E, Treiner E, Duban L, et al. Stepwise development of mait cells in mouse and human. *PLoS Biol*. 2009;7:e1000054.
- Dusseaux M, Martin E, Serriari N, et al. Human MAIT cells are xenobiotic-resistant, tissue-targeted, CD161 hi IL-17-secreting T cells. *Blood*. 2011;117(4):1250–1259.
- Koay HF, Gherardin NA, Enders A, et al. A three-stage intrathymic development pathway for the mucosal-associated invariant T cell lineage. *Nat Immunol*. 2016;17(11):1300–1311.

39. Van Der Vliet HJJ, Nishi N, De Gruijl TD, et al. Human natural killer T cells acquire a memory-activated phenotype before birth. *Blood*. 2000;95(7):2440-2442.
40. Gapin L, Matsuda JL, Surh CD, Kronenberg M. NKT cells derive from double-positive thymocytes that are positively selected by CD1d. *Nat Immunol*. 2001;2(10):971-978.
41. Wei DG, Lee H, Park SH, et al. Expansion and long-range differentiation of the NKT cell lineage in mice expressing CD1d exclusively on cortical thymocytes. *J Exp Med*. 2005;202(2):239-248.
42. McNab FW, Berzins SP, Pellicci DG, et al. The influence of CD1d in postselection NKT cell maturation and homeostasis. *J Immunol*. 2005;175(6):3762-3768.
43. De Lalla C, Lepore M, Piccolo FM, et al. High-frequency and adaptive-like dynamics of human CD1 self-reactive T cells. *Eur J Immunol*. 2011;41(3):602-610.
44. Wang H, Kjer-Nielsen L, Shi M, et al. IL-23 costimulates antigen-specific MAIT cell activation and enables vaccination against bacterial infection. *Sci Immunol*. 2019;4(41):pii: eaaw0402.
45. Kurioka A, Jahun AS, Hannaway RF, et al. Shared and distinct phenotypes and functions of human cD161+ V α 7.2+ T cell subsets. *Front Immunol*. 2017;8:1031.
46. Wang H, D'Souza C, Lim XY, et al. MAIT cells protect against pulmonary *Legionella longbeachae* infection. *Nat Commun*. 2018;9(1):3350.
47. Gumperz JE, Miyake S, Yamamura T, Brenner MB. Functionally distinct subsets of CD1d-restricted natural killer T cells revealed by CD1d tetramer staining. *J Exp Med*. 2002;195(5):625-636.
48. Lee PT, Benlagha K, Teyton L, Bendelac A. Distinct functional lineages of human V α 24 natural killer T cells. *J Exp Med*. 2002;195(5):637-641.
49. Matsuda JL, Naidenko OV, Gapin L, et al. Tracking the response of natural killer T cells to a glycolipid antigen using CD1d tetramers. *J Exp Med*. 2000;192(5):741-754.
50. Coquet JM, Chakravarti S, Kyriassoudis K, et al. Diverse cytokine production by NKT cell subsets and identification of an IL-17-producing CD4-NK1.1- NKT cell population. *Proc Natl Acad Sci USA*. 2008;105(32):11287-11292.
51. Crowe NY, Uldrich AP, Kyriassoudis K, et al. Glycolipid antigen drives rapid expansion and sustained cytokine production by NK T cells. *J Immunol*. 2003;171(8):4020-4027.
52. Kim JH, Hu Y, Yongqing T, et al. CD1a on Langerhans cells controls inflammatory skin disease. *Nat Immunol*. 2016;17(10):1159-1166.
53. DeWitt WS, Yu KKQ, Wilburn DB, et al. A diverse lipid antigen-specific TCR repertoire is clonally expanded during active tuberculosis. *J Immunol*. 2018;201(3):888-896.
54. Keller AN, Eckle SBG, Xu W, et al. Drugs and drug-like molecules can modulate the function of mucosal-associated invariant T cells. *Nat Immunol*. 2017;18(4):402-411.
55. Nicolai S, Wegrecki M, Cheng TY, et al. Human T cell response to CD1a and contact dermatitis allergens in botanical extracts and commercial skin care products. *Sci Immunol*. 2020;5(43):pii: eaax5430.
56. Bharadwaj M, Illing P, Theodossis A, Purcell AW, Rossjohn J, McCluskey J. Drug hypersensitivity and human leukocyte antigens of the major histocompatibility complex. *Annu Rev Pharmacol Toxicol*. 2012;52(1):401-431.
57. Illing PT, Vivian JP, Dudek NL, et al. Immune self-reactivity triggered by drug-modified HLA-peptide repertoire. *Nature*. 2012;486(7404):554-558.
58. Pichler WJ, Beeler A, Keller M, et al. Pharmacological interaction of drugs with immune receptors: The p-i concept. *Allergol Int*. 2006;55(1):17-25.
59. Bourgeois EA, Subramaniam S, Cheng TY, et al. Bee venom processes human skin lipids for presentation by CD1a. *J Exp Med*. 2015;212(2):149-163.
60. Akbari O, Stock P, Meyer E, et al. Essential role of NKT cells producing IL-4 and IL-13 in the development of allergen-induced airway hyperreactivity. *Nat Med*. 2003;9(5):582-588.
61. Godfrey DI, Koay H-F, McCluskey J, Gherardin NA. The biology and functional importance of MAIT cells. *Nat Immunol*. 2019;20(9):1110-1128.
62. Loh L, Gherardin NA, Sant S, et al. Human mucosal-associated invariant T cells in older individuals display expanded TCR $\alpha\beta$ clonotypes with potent antimicrobial responses. *J Immunol*. 2020;204(5):1119-1133.
63. Sakala IG, Kjer-Nielsen L, Eickhoff CS, et al. Functional heterogeneity and antimycobacterial effects of mouse mucosal-associated invariant T cells specific for riboflavin metabolites. *J Immunol*. 2015;195(2):587-601.
64. Legoux F, Gilet J, Procopio E, Echasserieu K, Bernardeau K, Lantz O. Molecular mechanisms of lineage decisions in metabolite-specific T cells. *Nat Immunol*. 2019;20(9):1244-1255.
65. Subramaniam S, Aslam A, Misbah SA, et al. Elevated and cross-responsive CD1a-reactive T cells in bee and wasp venom allergic individuals. *Eur J Immunol*. 2016;46(1):242-252.
66. Jarrett R, Salio M, Lloyd-Lavery A, et al. Filaggrin inhibits generation of CD1a neolipid antigens by house dust mite-derived phospholipase. *Sci Transl Med*. 2016;8(325):325ra18-325ra18.
67. Hardman CS, Chen YL, Salimi M, et al. CD1a presentation of endogenous antigens by group 2 innate lymphoid cells. *Sci Immunol*. 2017;2(18):pii: eaan5918.
68. Cheung KL, Jarrett R, Subramaniam S, et al. Psoriatic T cells recognize neolipid antigens generated by mast cell phospholipase delivered by exosomes and presented by CD1a. *J Exp Med*. 2016;213(11):2399-2412.
69. Birkinshaw RW, Pellicci DG, Cheng TY, et al. $\alpha\beta$ T cell antigen receptor recognition of CD1a presenting self lipid ligands. *Nat Immunol*. 2015;16(3):258-266.
70. Cotton RN, Shahine A, Rossjohn J, Moody DB. Lipids hide or step aside for CD1-autoreactive T cell receptors. *Curr Opin Immunol*. 2018;52:93-99.
71. Agea E, Russano A, Bistoni O, et al. Human CD1-restricted T cell recognition of lipids from pollens. *J Exp Med*. 2005;202(2):295-308.
72. Gracia BA, López Relanó J, Revilla A, et al. Human invariant natural killer T cells respond to antigen-presenting cells exposed to lipids from *Olea europaea* pollen. *Int Arch Allergy Immunol*. 2017;173(1):12-22.
73. Bilenki L, Yang J, Fan Y, Wang S, Yang X. Natural killer T cells contribute to airway eosinophilic inflammation induced by ragweed through enhanced IL-4 and eotaxin production. *Eur J Immunol*. 2004;34(2):345-354.
74. Tumes D, Hirahara K, Papadopoulos M, et al. Ezh2 controls development of natural killer T cells, which cause spontaneous asthma-like pathology. *J Allergy Clin Immunol*. 2019;144(2):549-560. e10.
75. Hou TY, Chen MR, Chou YC, Kan PC, Tsai YT, Cha TL. Impact of enhancer of zeste homolog 2 on T helper cell-mediated allergic rhinitis. *Front Immunol*. 2017;8:790.
76. Tumes DJ, Onodera A, Suzuki A, et al. The polycomb protein Ezh2 regulates differentiation and plasticity of CD4+ T helper Type 1 and type 2 cells. *Immunity*. 2013;39(5):819-832.
77. Ferstl R, Frei R, Barcik W, et al. Histamine receptor 2 modifies iNKT cell activity within the inflamed lung. *Allergy Eur J Allergy Clin Immunol*. 2017;72(12):1925-1935.
78. Lisbonne M, Diem S, de Castro KA, et al. Cutting edge: invariant V α 14 NKT cells are required for allergen-induced airway inflammation and hyperreactivity in an experimental asthma model. *J Immunol*. 2003;171(4):1637-1641.
79. Meyer EH, DeKruyff RH, Umetsu DT. T Cells and NKT cells in the pathogenesis of asthma. *Annu Rev Med*. 2008;59(1):281-292.

80. Iwamura C, Nakayama T. Role of NKT cells in allergic asthma. *Curr Opin Immunol*. 2010;22(6):807-813.
81. Iwamura C, Nakayama T. Role of CD1d- and MR1-restricted T cells in asthma. *Front Immunol*. 2018;9:1942.
82. Kim JO, Kim DH, Chang WS, et al. Asthma is induced by intranasal coadministration of allergen and natural killer T-cell ligand in a mouse model. *J Allergy Clin Immunol*. 2004;114(6):1332-1338.
83. Chuang YT, Leung K, Chang YJ, et al. A natural killer T-cell subset that protects against airway hyperreactivity. *J Allergy Clin Immunol*. 2019;143(2):565-576.e7.
84. Hinks TSC, Zhou X, Staples KJ, et al. Innate and adaptive T cells in asthmatic patients: relationship to severity and disease mechanisms. *J Allergy Clin Immunol*. 2015;136(2):323-333.
85. Chandra S, Wingender G, Greenbaum JA, et al. Development of asthma in inner-city children: possible roles of MAIT cells and variation in the home environment. *J Immunol*. 2018;200(6):1995-2003.
86. Lezmi G, Abou Taam R, Dietrich C, Chatenoud L, de Blic J, Leite-de-Moraes M. Circulating IL-17-producing mucosal-associated invariant T cells (MAIT) are associated with symptoms in children with asthma. *Clin Immunol*. 2018;188:7-11.
87. Lezmi G, Leite-de-Moraes MC. Comment on "development of asthma in inner-city children: possible roles of MAIT Cells and variation in the home environment". *J Immunol*. 2018;200(10):3317.
88. Chandra S, Wingender G, Greenbaum JA, Kronenberg M. Response to comment on "development of asthma in inner-city children: possible roles of MAIT cells and variation in the home environment". *J Immunol*. 2018;200(10):3317-3318.
89. Mirotti L, Florsheim E, Rundqvist L, et al. Lipids are required for the development of Brazil nut allergy: the role of mouse and human iNKT cells. *Allergy*. 2013;68(1):74-83.
90. Brennan PJ, Titituri RVV, Brigl M, et al. Invariant natural killer T cells recognize lipid self antigen induced by microbial danger signals. *Nat Immunol*. 2011;12(12):1202-1211.
91. Jyonouchi S, Abraham V, Orange JS, et al. Invariant natural killer T cells from children with versus without food allergy exhibit differential responsiveness to milk-derived sphingomyelin. *J Allergy Clin Immunol*. 2011;128(1):102-109.e13.
92. Brennan PJ, Cheng TY, Pellicci DG, et al. Structural determination of lipid antigens captured at the CD1d-T-cell receptor interface. *Proc Natl Acad Sci USA*. 2017;114(31):8348-8353.
93. Brennan PJ, Titituri RVV, Heiss C, et al. Activation of iNKT cells by a distinct constituent of the endogenous glucosylceramide fraction. *Proc Natl Acad Sci USA*. 2014;111(37):13433-13438.
94. Jyonouchi S, Smith CL, Saretta F, et al. Invariant natural killer T cells in children with eosinophilic esophagitis. *Clin Exp Allergy*. 2014;44(1):58-68.
95. Sidobre S, Naidenko OV, Sim B-C, Gascoigne NRJ, Garcia KC, Kronenberg M. The V α 14 NKT Cell TCR exhibits high-affinity binding to a Glycolipid/CD1d complex. *J Immunol*. 2002;169(3):1340-1348.
96. Mansueto P, Iacono G, Seidita A, D'Alcamo A, Sprini D, Carroccio A. Review article: intestinal lymphoid nodular hyperplasia in children - the relationship to food hypersensitivity. *Aliment Pharmacol Ther*. 2012;35(9):1000-1009.
97. Kokkonen J, Holm K, Karttunen TJ, Maki M. Children with untreated food allergy express a relative increment in the density of duodenal $\gamma\delta$ + T cells. *Scand J Gastroenterol*. 2000;35(11):1137-1142.
98. Järvinen TT, Kaukinen K, Laurila K, et al. Intraepithelial lymphocytes in celiac disease. *Am J Gastroenterol*. 2003;98(6):1332-1337.
99. Dunne MR, Elliott L, Hussey S, et al. Persistent changes in circulating and intestinal $\gamma\delta$ T cell subsets, invariant natural killer T cells and mucosal-associated invariant T cells in children and adults with coeliac disease. *PLoS ONE*. 2013;8(10):e76008.
100. Nijeboer P, Van Gils T, Reijm M, et al. Gamma-delta T lymphocytes in the diagnostic approach of coeliac disease. *J Clin Gastroenterol*. 2019;53(5):e208-e213.
101. Bol-Schoenmakers M, Marcondes Rezende M, Bleumink R, et al. Regulation by intestinal $\gamma\delta$ T cells during establishment of food allergic sensitization in mice. *Allergy*. 2011;66(3):331-340.
102. Portman MA. Kawasaki disease and soy: potential role for isoflavone interaction with Fc γ receptors. *Pediatr Res*. 2013;73(2):130-134.
103. Maleki SJ, Crespo JF, Cabanillas B. Anti-inflammatory effects of flavonoids. *Food Chem*. 2019;299:125124.
104. Yu J, Bi X, Yu B, Chen D. Isoflavones: anti-inflammatory benefit and possible caveats. *Nutrients*. 2016;8(6):361.
105. Schram SE, Warshaw EM. Genetics of nickel allergic contact dermatitis. *Dermatitis*. 2007;18(3):125-133.
106. Martin SF. New concepts in cutaneous allergy. *Contact Dermatitis*. 2015;72(1):2-10.
107. Schmidt M, Raghavan B, Müller V, et al. Crucial role for human toll-like receptor 4 in the development of contact allergy to nickel. *Nat Immunol*. 2010;11(9):814-819.
108. Eguchi T, Kumagai K, Kobayashi H, et al. Accumulation of invariant NKT cells into inflamed skin in a novel murine model of nickel allergy. *Cell Immunol*. 2013;284(1-2):163-171.
109. Okuno H, Satoh M, Takeuchi E, et al. Inhibitory function of NKT cells during early induction phase of nickel allergy. *Immunobiology*. 2016;221(7):833-838.
110. Werfel T, Hentschel M, Kapp A, Renz H. Dichotomy of blood- and skin-derived IL-4-producing allergen-specific T cells and restricted V β repertoire in nickel-mediated contact dermatitis. *J Immunol*. 1997;35(11):1137-1142.
111. Moulon C, Wild D, Dormoy A, Weltzien HU. MHC-dependent and -independent activation of human nickel-specific CD8+ cytotoxic T cells from allergic donors. *J Invest Dermatol*. 1998;111(3):360-366.
112. Moulon C, Choleva Y, Thierse HJ, Wild D, Weltzien HU. T cell receptor transfection shows non-HLA-restricted recognition of nickel by CD8+ human T cells to be mediated by $\alpha\beta$ T cell receptors. *J Invest Dermatol*. 2003;121(3):496-501.
113. Hashizume H, Seo N, Ito T, Takigawa M, Yagi H. Promiscuous interaction between gold-specific T cells and APCs in gold allergy. *J Immunol*. 2008;181(11):8096-8102.
114. Gober MD, Fischelevich R, Zhao Y, Unutmaz D, Gaspari AA. Human natural killer T cells infiltrate into the skin at elicitation sites of allergic contact dermatitis. *J Invest Dermatol*. 2008;128(6):1460-1469.
115. Shigematsu H, Kumagai K, Kobayashi H, et al. Accumulation of metal-specific T cells in inflamed skin in a novel murine model of chromium-induced allergic contact dermatitis. *PLoS ONE*. 2014;9(1):e85983.
116. Van Rhijn I, Young DC, Im JS, et al. CD1d-restricted T cell activation by nonlipidic small molecules. *Proc Natl Acad Sci USA*. 2004;101(37):13578-13583.
117. Davies NM, Andersen KE. Clinical pharmacokinetics of diclofenac. Therapeutic insights and pitfalls. *Clin Pharmacokinet*. 1997;33(3):184-213.
118. Naisbitt DJ, Sanderson LS, Meng X, Stachulski AV, Clarke SE, Park BK. Investigation of the immunogenicity of diclofenac and diclofenac metabolites. *Toxicol Lett*. 2007;168(1):45-50.
119. Kretz-Rommel A, Boelsterli UA. Cytotoxic activity of T cells and non-T cells from diclofenac-immunized mice against cultured syngeneic hepatocytes exposed to diclofenac. *Hepatology*. 1995;22(1):213-222.
120. Van Rhijn I, Kasmar A, De Jong A, et al. A conserved human T cell population targets mycobacterial antigens presented by CD1b. *Nat Immunol*. 2013;14(7):706-713.
121. Van Rhijn I, Gherardin NA, Kasmar A, et al. TCR bias and affinity define two compartments of the CD1b-glycolipid-specific T cell repertoire. *J Immunol*. 2014;192(9):4054-4060.

122. Mallal S, Phillips E, Carosi G, et al. HLA-B*5701 screening for hypersensitivity to abacavir. *N Engl J Med*. 2008;5(3):297-300.
123. Black CA. Delayed type hypersensitivity: current theories with an historic perspective. *Dermatol Online J*. 1999;5(1):7.
124. Brigl M, Brenner MB. CD1: antigen presentation and T cell function. *Annu Rev Immunol*. 2004;22(1):817-890.
125. Bai L, Picard D, Anderson B, et al. The majority of CD1d-sulfatide-specific T cells in human blood use a semiinvariant V δ 1 TCR. *Eur J Immunol*. 2012;42(9):2505-2510.
126. Huang S, Martin E, Kim S, et al. MR1 antigen presentation to mucosal-associated invariant T cells was highly conserved in evolution. *Proc Natl Acad Sci USA*. 2009;106(20):8290-8295.
127. Celie PHN, Toebes M, Rodenko B, Ovaa H, Perrakis A, Schumacher TNM. UV-induced ligand exchange in MHC class I protein crystals. *J Am Chem Soc*. 2009;131(34):12298-12304.
128. Zajonc DM, Crispin MDM, Bowden TA, et al. Molecular mechanism of lipopeptide presentation by CD1a. *Immunity*. 2005;22(2):209-219.
129. Borg NA, Wun KS, Kjer-Nielsen L, et al. CD1d-lipid-antigen recognition by the semi-invariant NKT T-cell receptor. *Nature*. 2007;448(7149):44-49.
130. Chien Y, Meyer C, Bonneville M. $\gamma\delta$ T cells: first line of defense and beyond. *Annu Rev Immunol*. 2014;32:121-155.
131. Vantourout P, Hayday A. Six-of-the-best: unique contributions of $\gamma\delta$ T cells to immunology. *Nat Rev Immunol*. 2013;13(2):88-100.
132. Berzins SP, Cochrane AD, Pellicci DG, Smyth MJ, Godfrey DI. Limited correlation between human thymus and blood NKT cell content revealed by an ontogeny study of paired tissue samples. *Eur J Immunol*. 2005;35(5):1399-1407.
133. Arrenberg P, Halder R, Dai Y, Maricic I, Kumar V. Oligoclonality and innate-like features in the TCR repertoire of type II NKT cells reactive to a β -linked self-glycolipid. *Proc Natl Acad Sci USA*. 2010;107(24):10984-10989.
134. Mangan BA, Dunne MR, O'Reilly VP, et al. Cutting edge: CD1d restriction and Th1/Th2/Th17 cytokine secretion by human V δ 3 T cells. *J Immunol*. 2013;191(1):30-34.
135. Mittrücker HW, Visekruna A, Huber M. Heterogeneity in the differentiation and function of CD8⁺ T cells. *Arch Immunol Ther Exp (Warsz)*. 2014;62(6):449-458.
136. Rouxel O, Da Silva J, Beaudoin L, et al. Cytotoxic and regulatory roles of mucosal-associated invariant T cells in type 1 diabetes. *Nat Immunol*. 2017;18(12):1321-1331.
137. Rossjohn J, Pellicci DG, Patel O, Gapin L, Godfrey DI. Recognition of CD1d-restricted antigens by natural killer T cells. *Nat Rev Immunol*. 2012;12(12):845-857.
138. Salio M, Silk JD, Yvonne Jones E, Cerundolo V. Biology of CD1- and MR1-restricted T cells. *Annu Rev Immunol*. 2014;32(1):323-366.
139. De Jong A, Peña-Cruz V, Cheng TY, Clark RA, Van Rhijn I, Moody DB. CD1a-autoreactive T cells are a normal component of the human $\alpha\beta$ T cell repertoire. *Nat Immunol*. 2010;11(12):1102-1109.
140. Souter MNT, Loh L, Li S, et al. Characterization of human mucosal-associated invariant T (MAIT) Cells. *Curr Protoc Immunol*. 2019;127(1):e90.
141. Gilleron M, Stenger S, Mazorra Z, et al. Diacylated sulfoglycolipids are novel mycobacterial antigens stimulating CD1-restricted T cells during infection with mycobacterium tuberculosis. *J Exp Med*. 2004;199(5):649-659.
142. Liao CM, Zimmer MI, Wang CR. The functions of type I and Type II natural killer T cells in inflammatory bowel diseases. *Inflamm Bowel Dis*. 2013;19(6):1330-1338.
143. Kasmar AG, Van Rhijn I, Magalhaes KG, et al. Cutting edge: CD1a tetramers and dextramers identify human lipopeptide-specific T cells ex vivo. *J Immunol*. 2013;191(9):4499-4503.

How to cite this article: de Lima Moreira M, Souter MNT, Chen Z, et al. Hypersensitivities following allergen antigen recognition by unconventional T cells. *Allergy*. 2020;00:1-14. <https://doi.org/10.1111/all.14279>