



Occupation and risk of lymphoma: a multicentre prospective cohort study (EPIC)

David Neasham, Ahlem Sifi, Kaspar Rene Nielsen, et al.

Occup Environ Med published online September 30, 2010

doi: 10.1136/oem.2009.048173

Updated information and services can be found at:

<http://oem.bmj.com/content/early/2010/09/30/oem.2009.048173.full.html>

These include:

Data Supplement

"Web Only Data"

<http://oem.bmj.com/content/suppl/2010/10/06/oem.2009.048173.DC1.html>

References

This article cites 21 articles, 12 of which can be accessed free at:

<http://oem.bmj.com/content/early/2010/09/30/oem.2009.048173.full.html#ref-list-1>

P<P

Published online September 30, 2010 in advance of the print journal.

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Notes

Advance online articles have been peer reviewed and accepted for publication but have not yet appeared in the paper journal (edited, typeset versions may be posted when available prior to final publication). Advance online articles are citable and establish publication priority; they are indexed by PubMed from initial publication. Citations to Advance online articles must include the digital object identifier (DOIs) and date of initial publication.

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://journals.bmj.com/cgi/ep>

Occupation and risk of lymphoma: a multicentre prospective cohort study (EPIC)

David Neasham,¹ Ahlem Sifi,¹ Kaspar Rene Nielsen,² Kim Overvad,³ Ole Raaschou-Nielsen,⁴ Anne Tjønneland,⁴ Aurelio Barricarte,⁵ Carlos A González,⁶ Carmen Navarro,^{7,8} Laudina Rodriguez Suarez,⁹ Ruth C Travis,¹⁰ Tim Key,¹⁰ Jakob Linseisen,¹¹ Rudolf Kaaks,¹¹ Paolo Crosignani,¹² Franco Berrino,¹³ Stefano Rosso,¹⁴ Amalia Mattiello,¹⁵ R C H Vermeulen,¹⁶ H Bas Bueno-de-Mesquita,¹⁷ Göran Berglund,¹⁸ Jonas Manjer,¹⁹ Sophia Zackrisson,²⁰ Goran Hallmans,²¹ Beatrice Malmer,²¹ Sheila Bingham,²² Kay Tee Khaw,²³ Manuela M Bergmann,²⁴ Heiner Boeing,²⁴ Antonia Trichopoulou,²⁵ Giovanna Masala,²⁶ Rosario Tumino,²⁷ Eiliv Lund,²⁸ Nadia Slimani,²⁹ Pietro Ferrari,²⁹ Paolo Boffetta,³⁰ Paolo Vineis,^{1,31} Elio Riboli¹

► Additional tables are published online only. To view these files please visit the journal online (<http://oem.bmj.com>).

For numbered affiliations see end of article.

Correspondence to

Department of Epidemiology and Public Health, Imperial College London, Department of Epidemiology and Public Health, Imperial College Faculty of Medicine Building, Norfolk Place, London W2 1PG, UK; david.neasham@googlemail.com

Received 13 May 2009

Revised 16 February 2010

Accepted 24 February 2010

ABSTRACT

Objectives Evidence suggests that certain occupations and related exposures may increase the risk of malignant lymphoma. Farming, printing and paper industry, wood processing, meat handling and processing, welding, shoe and leather manufacturing and teaching profession are among the categories that have been implicated in previous studies. The relationship between occupation and malignant lymphoma has been investigated in a large European prospective study.

Methods We investigated occupational risks for lymphomas in the European Prospective Investigation into Cancer and Nutrition (EPIC). The mean follow-up time for 348 555 subjects was 9 years (SD: 2 years). The analysis was based on 866 and 48 newly diagnosed cases of non-Hodgkin's lymphoma (NHL) and Hodgkin's lymphoma (HL). These were identified in the EPIC subcohorts with occupational data. Data on 52 occupations were collected through standardised questionnaires. Cox proportional hazard models were used to explore the association between occupation and risk of malignant lymphoma.

Results The following occupations were positively associated with malignant NHL after adjustment for study centre, age, sex, socioeconomic status (SES), smoking and alcohol: butchers (HR=1.53, 95% CI 1.05 to 2.48, including multiple myeloma/plasmacytoma; HR=1.30, 95% CI 1.00 to 2.66, excluding multiple myeloma/plasmacytoma) and car repair workers (HR=1.50, 95% CI 1.01 to 2.00, including multiple myeloma/plasmacytoma; HR=1.51, 95% CI 1.01 to 2.31, excluding multiple myeloma/plasmacytoma). HL was associated with gasoline station occupation (HR=4.59, 95% CI 1.08 to 19.6).

Conclusion The findings in this current study of a higher risk of NHL among car repair workers and butchers and a higher risk of HL among gasoline station workers suggest a possible role from occupationally related exposures, such as solvents and zoonotic viruses, as risk factors for malignant lymphoma.

INTRODUCTION

Lymphomas are a heterogeneous group of diseases caused by a proliferation of malignant lymphocytes

What this paper adds

- Established risk factors for lymphomas are those associated with downregulation of immune function such as genetically determined or acquired immune deficiencies.
- Known risk factors explain only a small proportion of lymphomas. Evidence suggests other factors such as occupations and related exposures may increase risk of disease.
- This study found that certain occupations (butchers and car repair workers) were associated with non-Hodgkin's lymphoma. Gasoline station work was associated with Hodgkin's lymphoma.
- Our findings point to the role of occupationally related exposures (solvents and zoonotic viruses) as risk factors for malignant lymphoma.

that accumulate in lymph nodes. They are subdivided into Hodgkin's lymphoma (HL) and non-Hodgkin's lymphoma (NHL), where HLs are B-cell malignancies distinguishable by the presence of Reed–Sternberg (RS) cells, and NHLs are of either T- or B-cell origin.¹ NHL accounts for 90% of all lymphomas and HL for the remaining 10%.

Lymphomas account for over 3% of cancers occurring worldwide.¹ The aetiology of these diseases remains largely unknown. The best established risk factors are those associated with downregulation of immune function such as genetically determined or acquired immune deficiencies, including HIV infection or iatrogenically induced immune suppression after transplantation.² Known risk factors explain only a small proportion of lymphomas. Evidence suggests that other factors such as certain occupations and related exposures may increase risk of disease.^{3–12}

Very few studies examining occupation and lymphoma have used a prospective design.

Short report

Retrospective (case control) studies can be affected by recall bias and poor comparability of cases and controls. The present investigation evaluates the relationship between job title and lymphoma in a large prospective cohort (EPIC).

MATERIALS AND METHODS

Study population, outcomes and data collection

EPIC is a European multicentre prospective cohort study investigating the aetiology of cancers and other chronic diseases in relation to diet and lifestyle. Recruitment of subjects took place between 1992 and 2000 in 23 centres located in 10 European countries. The cohort includes participants of both genders, mostly in the age range of 35–70 at recruitment. Centres in France (n=61 996), Oxford (n=42 518), and Utrecht (n=15 174) did not collect job questionnaire information, therefore the current cohort size was 348 555 subjects (table 1). Recruitment was aimed at the general population or subgroups (such as blood donors or women attending breast cancer screening facilities). Identification of cancer cases was based on population cancer registries in seven of the participating countries (Denmark, Italy, The Netherlands, Norway, Spain, Sweden and the UK). The other two countries (Germany and Greece) used a combination of methods including health insurance records, cancer and pathology registries, and active follow-up through study subjects and their next of kin. Information was updated in 2006 (end of follow-up).

Diagnosis of lymphoma cases in EPIC was based on the 2nd revision of the International Classification of Diseases for Oncology (ICD-O-2). All lymphoma cases were subsequently reclassified according to the recently published WHO classification of haematopoietic and lymphoid tissues based on ICD-O-3. This was accomplished applying a conversion program available

at the SEER webpage (<http://seer.cancer.gov/tools/conversion/ICD02-3manual.pdf>). NHL and multiple myeloma/plasmacytoma cases were grouped together (termed 'NHL'); HL cases were grouped separately (termed 'HL').¹³ We further evaluated NHL excluding multiple myeloma/plasmacytoma.

Exposure information was collected via standardised dietary intake questionnaires and non-dietary questionnaires on lifestyle variables. Occupational status was self-reported and obtained using a standardised questionnaire in which participants were asked if they had ever worked in up to 52 jobs, selected a priori for being previously linked to risk of developing cancer. Data on educational attainment (a proxy for socioeconomic status), alcohol consumption and smoking status were obtained. Current smoker/drinker refers to status at baseline. Further details can be found elsewhere.¹⁴

Statistical analyses

All statistical analyses were performed using STATA 9 (Stata-corp). Cox proportional hazard regression models were used to examine the association between job category and lymphoma. Time at entry was defined as age at recruitment and exit time was age when individuals were: diagnosed with lymphoma, died, or censored at end of follow-up, whichever came first. Analyses were stratified by centre to control for the differences in follow-up procedures and questionnaire design. Multivariate analyses were further adjusted for gender, education level (proxy for socioeconomic status), alcohol consumption (g/day) and smoking status at baseline.

RESULTS

Mean follow-up time for 348 555 subjects was 9 years (SD: 2 years). Unadjusted NHL and HL incidence rates were 27.7

Table 1 Characteristics of EPIC cohort population, NHL (including and excluding multiple myeloma/plasmacytoma (MM)) and HL separately

Descriptive data	Whole cohort (N = 348555)	NHL (N = 866)	NHL excluding MM (N = 676)	HL (N = 48)
Age at recruitment (%)				
Mean age, years (SD)	51.7 (9.7)	56.4 (8.4)	56.6 (7.8)	50.9 (11.2)
<35	16854 (4.9)	2 (0.2)	2 (0.3)	5 (10.4)
35–44	64930 (18.6)	70 (8.1)	56 (8.3)	8 (16.7)
45–54	133990 (38.4)	254 (29.3)	207 (30.6)	12 (25.0)
55–65	110837 (31.8)	413 (47.7)	309 (45.7)	19 (39.6)
65+	21944 (6.3)	127 (14.7)	102 (15.1)	4 (8.3)
Age*				
Mean age, years (SD)	60.6 (9.8)	61.5 (8.7)	61.7 (8.4)	55.6 (11.8)
Gender, n (%)				
Men	106813 (30.6)	427 (49.3)	330 (48.8)	29 (60.4)
Women	241742 (69.4)	439 (50.7)	346 (51.2)	19 (39.6)
School degree, n (%)				
None	18621 (5.3)	42 (4.8)	34 (5.0)	2 (4.1)
Primary	82940 (23.8)	270 (31.2)	197 (29.1)	18 (37.5)
Technical	75301 (21.6)	243 (28.1)	194 (28.7)	14 (29.2)
Secondary	80692 (23.2)	133 (15.4)	107 (15.8)	1 (2.1)
University	91001 (26.1)	178 (20.5)	144 (21.4)	13 (27.1)
Smoking habits, n (%)				
Never smoked	181384 (52.0)	372 (43.0)	290 (42.9)	12 (25.0)
Former smoker	92743 (26.6)	305 (35.2)	239 (35.4)	17 (35.4)
Current smoker	74428 (21.4)	189 (21.8)	147 (21.7)	19 (39.6)
Alcohol drinking history, n (%)				
Never drinkers	27406 (7.9)	48 (5.5)	37 (5.5)	2 (4.2)
Former drinkers	16733 (4.8)	62 (7.2)	50 (7.4)	3 (6.2)
Drinkers at recruitment only	36785 (10.5)	59 (6.8)	47 (6.9)	2 (4.2)
Lifetime drinkers	267631 (76.8)	697 (80.5)	542 (80.2)	41 (85.4)

*Age at exit from study based on: lymphoma diagnosis, death, or censoring at end of follow-up, whichever came first.
EPIC, European Prospective Investigation into Cancer and Nutrition; HL, Hodgkin's lymphoma; NHL, non-Hodgkin's lymphoma.

cases per 100 000 person-years (95% CI 25.9 to 29.6) and 1.5 cases per 100 000 person-years (95% CI 1.1 to 2.0), respectively, based on 866 and 48 newly diagnosed cases of NHL and HL. These incidence rates are comparable with other data sources.¹⁵ After adjustment for centre, age, sex, educational status, smoking and alcohol consumption, butchers (including slaughterhouse and meat manufacturing workers) and car repair workers had significantly elevated risk of NHL (table 2) while gasoline station workers had elevated risk of HL. Excluding cases of multiple myeloma/plasmacytoma did not change our findings (apart from a significantly increased risk for chemical laboratory work: adjusted HR=3.21 (1.18–8.72)). This result and results from all 52 job categories can be found in the supplementary table.

An analysis by NHL histological subtypes was attempted, but absolute numbers in most occupations were too small for subgroup analysis. Overall there were 117 diffuse large cell lymphomas, 117 follicular lymphomas, 187 chronic lymphocytic leukaemias, 190 multiple myelomas/plasmacytomas, 97 B-cell

lymphomas NOS and 158 other specified non-Hodgkin's lymphomas.

DISCUSSION

We found that butchers, car repair and gasoline station workers were at increased risk of developing lymphoma compared with all other groups (findings consistent with previous studies). Car repair employees and gasoline station workers are known to be exposed to high levels of solvents such as benzene (and benzene-containing petroleum products),¹⁶ and a growing body of evidence exists suggesting an association between occupational benzene exposure and lymphoma, especially NHL.^{8 17 18} Milham analysed death certificates in Washington State and found that occupational groups with petrol or fuel exposure had excess risk of haematopoietic cancers, including lymphoma.¹⁹ The main hypothesis concerning the mechanism of action of solvents in lymphomagenesis has been immunotoxicity. This mechanism has scientific plausibility since increased incidence of lymphoma (in particular, NHL) is well documented among subjects with

Table 2 Unadjusted and adjusted analysis (Cox proportional hazards model stratified by centre). HR and 95% CI by selected job category

Outcome and job title	Person-years	Cases, n (%)	Unadjusted HR (95% CI)	Adjusted HR (95% CI)*
NHL				
Butchers				
No	1678150	646 (97.4)	1.00 (-)	1.00 (-)
Yes	24356	17 (2.6)	1.81 (1.12 to 2.93)	1.53 (1.05 to 2.48)
Gasoline station workers				
No	1688100	652 (98.3)	1.00 (-)	1.00 (-)
Yes	14293	11 (1.7)	1.99 (1.10 to 3.62)	1.32 (0.73 to 2.41)
Car repair workers				
No	1427447	587 (95.4)	1.00 (-)	1.00 (-)
Yes	38997	28 (4.6)	1.74 (1.19 to 2.54)	1.50 (1.01 to 2.00)
NHL excluding MM/plasmacytoma				
Butchers				
No	1678150	509 (97.9)	1.00 (-)	1.00 (-)
Yes	24356	11 (2.1)	1.49 (1.02 to 2.89)	1.30 (1.00 to 2.66)
Gasoline station workers				
No	1688100	514 (98.8)	1.00 (-)	1.00 (-)
Yes	14293	6 (1.2)	1.38 (0.80 to 3.08)	1.19 (0.64 to 2.72)
Car repair workers				
No	1427447	462 (95.3)	1.00 (-)	1.00 (-)
Yes	38997	23 (4.7)	1.82 (1.20 to 2.77)	1.51 (1.01 to 2.31)
MM/plasmacytoma				
Butchers				
No	1678150	137	1.00 (-)	1.00 (-)
Yes	24356	6	3.02 (1.33 to 6.83)	2.34 (1.03 to 5.35)
Gasoline station workers				
No	1688100	138	1.00 (-)	1.00 (-)
Yes	14293	5	4.28 (1.77 to 10.55)	2.61 (1.06 to 6.43)
Car repair workers				
No	1427447	125	1.00 (-)	1.00 (-)
Yes	38997	5	1.46 (0.59 to 3.54)	0.93 (0.38 to 2.30)
HL				
Butchers				
No	1678150	37 (94.9)	1.00 (-)	1.00 (-)
Yes	24356	2 (5.1)	3.72 (0.90 to 15.4)	2.89 (0.68 to 12.2)
Gasoline station workers				
No	1688100	37 (94.9)	1.00 (-)	1.00 (-)
Yes	14293	2 (5.1)	6.38 (1.53 to 26.3)	4.59 (1.08 to 19.6)
Car repair workers				
No	1427447	35 (94.6)	1.00 (-)	1.00 (-)
Yes	38997	2 (5.4)	2.09 (0.51 to 8.80)	1.50 (0.35 to 6.37)

*Multivariate analysis stratified by centre and adjusted for gender, education level, alcohol consumption and smoking status at baseline.

HL, Hodgkin's lymphoma; MM, multiple myeloma; NHL, non-Hodgkin's lymphoma.

Short report

primary immunodeficiency (eg, ataxia telangiectasia), acquired immunodeficiency (HIV), and autoimmune diseases (eg, rheumatoid arthritis, lupus erythematosus and Sjögren syndrome).²⁰ A recent literature review by the Agency for Toxic Substances and Diseases Registry²¹ found sufficient consistent evidence to support an association between benzene exposure and adverse events related to functional immunity; in particular, benzene seems to determine an immunosuppressive response which is mediated by a depressive effect on T and B lymphocytes.^{22 23} In a study of workers in the oil industry, Biro *et al*²⁴ also found that solvents such as benzene activate peripheral lymphocytes and cause changes to the prevalence of CD25+/CD4+ T lymphocytes. Lan *et al*²⁵ showed that benzene haematotoxicity effects were present even at very low exposure levels (<1 ppm). Such levels are consistent with those found in the car repair industry.¹⁶

The increased risk of lymphoma associated with butchers has been noted in some studies^{5 9 10} but not others.^{11 12} Since our study found no association with farming occupations, the results possibly implicate a role for exposures specifically related to meat handling and processing rather than rearing livestock. 't Mannetje *et al*⁹ reported a twofold increase in risk among meat workers, consistent with elevated NHL risk in earlier New Zealand studies among abattoir workers. The increased risk may be due to zoonotic viruses or antigenic stimulation through chronic exposure to animal protein. Most attention has centred on animal oncogenic viruses such as bovine leukaemia virus (BLV).²⁵ However, to date, in vivo studies have not provided evidence that BLV increases NHL risk in humans,²⁶ though the possibility that other animal retroviruses may have an aetiological role cannot be ruled out entirely.

We found no consistent association between other job categories, for example, those with known solvent exposure such as painting, chemical and rubber industries, printing or leather industry. This could be for several reasons: the study has low case numbers to detect effects (particularly for HL); insufficient follow-up time for accrual of NHL or HL cases (latency between first exposure and cancer diagnosis can be much longer than 9 years for some environmental agents); exposure levels may be differentially lower in well-regulated industrial environments where levels of solvents are monitored compared to settings in which little or no monitoring takes place; job categories associated with solvent exposure will also inevitably have a wide range of exposure levels. The job categories where we found an effect (particularly car repair) are largely in the more unregulated environment, with known high levels of solvent exposure.¹⁶ However, we had only limited job exposure information obtained from self-reported questionnaires and other job categories (such as painters) which might also be seen as less regulated in terms of exposure, showed no increased risk.

Our study is limited by small case numbers of lymphoma (particularly HL cases) and no available information on length of exposure which means that possible dose response effects were not assessed. EPIC also contains a higher proportion of women; effect estimates could be diluted as exposure levels may be lower in women compared with men for particular job categories. Multiple comparisons may give rise to 2–3 false positive significant associations for the analysis of up to 52 job categories, though our study is strengthened by having good a priori hypotheses and plausibility. In addition, the study design is prospective, with no recall bias, and has 100% follow-up. Overall, our findings should be corroborated by further research with more detailed exposure information.

Author affiliations:

- ¹Department of Epidemiology and Public Health, Imperial College London, UK
- ²Department of Clinical Immunology, Aarhus University Hospital, Aalborg, Denmark
- ³Department of Clinical Epidemiology, Aarhus University Hospital, Aalborg, Denmark
- ⁴Institute of Cancer Epidemiology, Danish Cancer Society, Copenhagen, Denmark
- ⁵Public Health Institute of Navarra, Pamplona, CIBERESP, Spain
- ⁶Unit of Nutrition, Environment and Cancer, Catalan Institute of Oncology, Barcelona, Spain
- ⁷Department of Epidemiology, Murcia, Spain
- ⁸CIBER Epidemiología y Salud Pública (CIBERESP), Spain
- ⁹Consejería de Salud y Servicios Sanitarios Principado de Asturias, Spain
- ¹⁰Cancer Epidemiology Unit, University of Oxford, Oxford, UK
- ¹¹German Cancer Research Centre, Division of Cancer Epidemiology, Heidelberg, Germany
- ¹²Unità di Epidemiologia Ambientale e Registro Tumori Fondazione IRCCS Istituto Nazionale dei Tumori Via Venezian, Milano, Italy
- ¹³Etiologic Epidemiology Unit, Fondazione IRCCS Istituto Nazionale dei Tumori Via Venezian, Milano, Italy
- ¹⁴CPO - Registro Tumori Piemonte via San Francesco da Paola 31, Torino, Italy
- ¹⁵Department of Clinical and Experimental Medicine Federico II University, Naples, Italy
- ¹⁶Julius Center for Health Sciences and Primary Care, University Medical Center Utrecht, The Netherlands
- ¹⁷Centre for Nutrition and Health, National Institute of Public Health and the Environment, Bilthoven, The Netherlands
- ¹⁸Department of Clinical Sciences, Malmö University Hospital, Malmö, Sweden
- ¹⁹Department of Surgery, Malmö University Hospital, Malmö, Sweden
- ²⁰Department of Clinical Sciences in Malmö, Medical Radiology, Lund University, Sweden
- ²¹Units of Nutritional Research and Oncology, University of Umeå, Sweden
- ²²MRC Centre for Nutritional Epidemiology in Cancer Prevention and Survival, Department of Public Health and Primary Care, University of Cambridge, UK
- ²³Clinical Gerontology, Department of Public Health and Primary Care, University of Cambridge, UK
- ²⁴Department of Epidemiology, German Institute of Human Nutrition, Potsdam, Germany
- ²⁵Department of Hygiene and Epidemiology, School of Medicine, University of Athens, Greece
- ²⁶Molecular and Nutritional Epidemiology Unit, ISPO - Scientific Institute of Tuscany, Florence, Italy
- ²⁷Cancer Registry, Azienda Ospedaliera "Civile MP Arezzo", Ragusa, Italy
- ²⁸Institute of Community Medicine, University of Tromsø, Tromsø, Norway
- ²⁹Nutrition and Hormones Group, International Agency for Research on Cancer, Lyon, France
- ³⁰The Tisch Cancer Institute Mount Sinai School of Medicine New York, NY, USA
- ³¹ISI Foundation, Torino, Italy

Funding Europe Against Cancer Program of the European Commission (SANCO), Deutsche Krebshilfe Deutsches Krebsforschungszentrum German Federal Ministry of Education and Research, Danish Cancer Society Health Research Fund (FIS) of the Spanish Ministry of Health Spanish Regional Governments of Andalucía, Asturias, Basque Country, Murcia and Navarra Grant ISCIII, Red de Centros CESP, C03/09, Spain Cancer Research, United Kingdom Stroke Association, United Kingdom British Heart Foundation, United Kingdom Department of Health, United Kingdom Food Standards Agency, United Kingdom Wellcome Trust, United Kingdom Greek Ministry of Education and Ministry of Health Italian Association for Research on Cancer (AIRC), Italian National Research Council Dutch Ministry of Public Health, Welfare and Sports, World Cancer Research Fund, Swedish Cancer Society, Swedish Scientific Council Regional Government of Skåne, Sweden Norwegian Cancer Society Research Council of Norway. Other funders: Wellcome Trust.

Competing interests None.

Patient consent Obtained.

Ethics approval This study was conducted with the approval of the IARC ethical committee and all the local IRBs.

Contributors DN and AS equally contributed to the manuscript.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

1. Willett EV, Roman E. Epidemiology. In: Marcus R, Sweetenham JW, Williams ME, eds. *Lymphoma: pathology, diagnosis and treatment*. Cambridge, UK: Cambridge University Press, 2007:3–11.
2. Becker N, de Sanjose S, Nieters A, *et al*. Birth order, allergies and lymphoma risk: results of the European collaborative research project Epilymph. *Leuk Res* 2007;**31**:1365–72.

3. **Scherr PA**, Hutchison GB, Neiman RS. Non-Hodgkin's lymphoma and occupational exposure. *Cancer Res* 1992;**52**:5503–9.
4. **Persson B**, Fredriksson M, Olsen K, *et al*. Some occupational exposures as risk factors for malignant lymphomas. *Cancer* 1993;**72**:1773–8.
5. **Metayer C**, Johnson ES, Rice JC. Nested case-control study of tumors of the hemopoietic and lymphatic systems among workers in the meat industry. *Am J Epidemiol* 1998;**147**:727–38.
6. **Mester B**, Nieters A, Deeg E, *et al*. Occupation and malignant lymphoma: a population based case control study in Germany. *Occup Environ Med* 2006;**63**:17–26.
7. **Miligi L**, Costantini AS, Benvenuti A, *et al*. Occupational exposure to solvents and the risk of lymphomas. *Epidemiology* 2006;**17**:552–61.
8. **Vineis P**, Miligi L, Seniori C. Exposure to solvents and risk of non-Hodgkin lymphoma: clues on putative mechanisms. *Cancer Epidemiol Biomarkers Prev* 2007;**16**:381–4.
9. **'t Mannetje A**, Dryson E, Walls C, *et al*. High risk occupations for non-Hodgkin's lymphoma in New Zealand: case-control study. *Occup Environ Med* 2008;**65**:354–63.
10. **Johnson ES**, Fischman HR, Matanowski GM, *et al*. Cancer mortality among white males in the meat industry. *J Occup Med* 1986;**28**:23–32.
11. **Boffetta P**, de Vocht F. Occupation and the risk of non-Hodgkin's lymphoma. *Cancer Epidemiol Biomarkers Prev* 2007;**16**:369–72.
12. **Moore T**, Brennan P, Becker N, *et al*. Occupational exposure to meat and the risk of lymphoma: a multi-center case control study from Europe. *Int J Cancer* 2007;**121**:2761–6.
13. **Jaffe ES**, Harris NL, Stein H, *et al*, eds. *Pathology and genetics of tumours of haematopoietic and lymphoid tissue*. Lyon: IARC Press, 2001.
14. **Riboli E**, Hunt KJ, Slimani N, *et al*. European prospective investigation into cancer and nutrition (EPIC): study populations and data collection. *Public Health Nutr* 2002;**5**:1113–24.
15. **Cancer Research UK**. <http://info.cancerresearchuk.org/> (accessed Jan 2010).
16. **Hunting KL**, Longbottom H, Kalavar SS, *et al*. Haematopoietic cancer mortality among vehicle mechanics. *Occup Environ Med* 1995;**52**:673–8.
17. **Smith MT**, Jones RM, Smith AH. Benzene exposure and risk of non-Hodgkin lymphoma. *Cancer Epidemiol Biomarkers Prev* 2007;**16**:385–91.
18. **Steinmaus C**, Smith AH, Jones RM, *et al*. Meta-analysis of benzene exposure and non-Hodgkin Lymphoma: biases could mask an important association. *Occup Environ Med* 2008;**65**:371–8.
19. **Milham S Jr**. *Occupational mortality in Washington state 1950–79*. Cincinnati, OH: NIOSH, DSHEFS, 1983. (NTIS No. PB84-199769).
20. **Smedby KE**, Baecklund E, Askling J. Malignant lymphomas in autoimmunity and inflammation: a review of risks, risk factors, and lymphoma characteristics. *Cancer Epidemiol Biomarkers Prev* 2006;**15**:2069–77. Review.
21. **Agency for Toxic Substances and Disease Registry (ATSDR) U.S. Department of Health and Human Services, Public Health Services**. Toxicological profile for benzene (Draft for public comment), <http://www.atsdr.cdc.gov/toxprofiles/tp3.html> 2007 (accessed Jan 2010).
22. **Gist GL**, Burg JR. Benzene: a review of the literature from a Health Effects Perspective. *Toxicol Ind Health* 1997;**13**:661–714.
23. **Lan Q**, Zhang L, Li G, *et al*. Hematotoxicity in workers exposed to low levels of benzene. *Science* 2004;**306**:1774–6.
24. **Biro A**, Pallinger E, Major J, *et al*. Lymphocytes phenotype analysis and chromosome aberration frequency of workers occupationally exposed to styrene, benzene, polycyclic aromatic hydrocarbons or mixed solvents. *Immunol Lett* 2002;**81**:133–40.
25. **Miller JM**, Miller LD, Olsen C, *et al*. Virus like particles in phytohemagglutinin simulated cultures with reference to bovine lymphosarcoma. *J Natl Cancer Inst* 1969;**43**:1297–305.
26. **Bender AP**, Robison LL, Kashmiri SV, *et al*. No involvement of bovine leukemia virus in childhood acute lymphoblastic leukemia and non-Hodgkin's lymphoma. *Cancer Res* 1988;**48**:2919–22.

SUPPLEMENTARY TABLE: Person-years of exposure and adjusted Hazards ratio (HR) + 95% confidence intervals (CI) by job category.

Job title	Person-years	NHL (%)	NHL-MM (%)	HL (%)	NHL: Adjusted HR (95% CI)*	NHL-MM: Adjusted HR (95% CI)*	HL: Adjusted HR (95% CI)*
Butchers							
No	1,678,150	646 (97.4)	509 (97.9)	37 (94.9)	1.00 (-)	1.00 (-)	1.00 (-)
Yes	24,356	17 (2.6)	11 (2.1)	2 (5.1)	1.53 (1.05-2.48)	1.30 (1.00-2.66)	2.89 (0.68-12.2)
Gasoline station workers							
No	1,688,100	652 (98.3)	514 (98.8)	37 (94.9)	1.00 (-)	1.00 (-)	1.00 (-)
Yes	14,293	11 (1.7)	6 (1.2)	2 (5.1)	1.32 (0.73-2.41)	1.19 (0.64-2.72)	4.59 (1.08-19.6)
Car repair workers							
No	1,427,447	587 (95.4)	462 (95.3)	35 (94.6)	1.00 (-)	1.00 (-)	1.00 (-)
Yes	38,997	28 (4.6)	23 (4.7)	2 (5.4)	1.50 (1.01-2.00)	1.51 (1.01-2.31)	1.50 (0.35-6.37)
Chemical laboratory work							
No	35,001	11 (40.7)	7 (33.3)	1 (50.0)	1.00 (-)	1.00 (-)	1.00 (-)
Yes	29,242	16 (59.3)	14 (66.7)	1 (50.0)	2.21 (0.95-5.10)	3.21 (1.18-8.72)	0.97 (0.05-18.0)
Asbestos insulation workers							
No	1,693,583	654 (98.6)	514 (98.8)	38 (97.4)	1.00 (-)	1.00 (-)	1.00 (-)
Yes	8,758	9 (1.4)	6 (1.2)	1 (2.6)	1.54 (0.79-2.99)	1.35 (0.60-3.03)	3.07 (0.41-23.2)
Worked in breeding							
No	1,591,090	622 (93.7)	488 (93.7)	35 (89.7)	1.00 (-)	1.00 (-)	1.00 (-)
Yes	111,853	42 (6.3)	33 (6.3)	4 (10.3)	0.87 (0.63-1.20)	0.89 (0.62-1.28)	1.57 (0.53-4.61)
Worked in farming							
No	1,431,754	564 (85.6)	441 (84.5)	30 (76.9)	1.00 (-)	1.00 (-)	1.00 (-)
Yes	261,784	95 (14.4)	75 (14.5)	9 (23.1)	1.00 (0.80-1.27)	1.05 (0.81-1.37)	1.72 (0.77-3.88)
Worked in mines/ quarries							
No	1,451,023	607 (98.9)	479 (98.8)	37 (100.0)	1.00 (-)	1.00 (-)	1.00 (-)
Yes	15,535	7 (1.1)	6 (1.2)	0 (0.0)	0.96 (0.45-2.04)	1.07 (0.48-2.42)	N/A
Worked in foundries							
No	1,441,049	605 (98.7)	477 (98.8)	37 (100.0)	1.00 (-)	1.00 (-)	1.00 (-)
Yes	24,092	8 (1.3)	6 (1.2)	0 (0.0)	0.68 (0.33-1.36)	0.66 (0.29-1.48)	N/A
Worked in electroplating							
No	1,456,436	609 (99.4)	481 (99.4)	37 (100.0)	1.00 (-)	1.00 (-)	1.00 (-)
Yes	9,365	4 (0.6)	3 (0.6)	0 (0.0)	0.77 (0.29-2.06)	0.76 (0.24-2.38)	N/A
Worked in chemical industry							

No	1,633,369	635 (96.1)	498 (96.0)	36 (94.7)	1.00 (-)	1.00 (-)	1.00 (-)
Yes	64,744	26 (3.9)	21 (4.0)	2 (5.3)	0.98 (0.66-1.45)	1.02 (0.66-1.58)	1.13 (0.27-4.71)
Worked in rubber industry							
No	1,677,864	658 (99.2)	515 (99.0)	39 (100.0)	1.00 (-)	1.00 (-)	1.00 (-)
Yes	23,757	5 (0.8)	5 (1.0)	0 (0.0)	0.52 (0.21-1.25)	0.68 (0.28-1.64)	N/A
Worked in textile industry							
No	1,641,674	648 (97.7)	506 (97.3)	36 (94.7)	1.00 (-)	1.00 (-)	1.00 (-)
Yes	58,677	15 (2.3)	14 (2.7)	2 (5.3)	0.72 (0.43-1.20)	0.87 (0.51-1.48)	1.72 (0.40-7.30)
Worked in leather and tanning industry							
No	1,696,937	658 (99.4)	516 (99.4)	39 (100.0)	1.00 (-)	1.00 (-)	1.00 (-)
Yes	5,146	4 (0.6)	3 (0.6)	0 (0.0)	1.68 (0.63-4.49)	1.64 (0.53-5.11)	N/A
Worked in shoes or other leather goods							
No	1,674,098	649 (98.2)	513 (99.0)	38 (97.4)	1.00 (-)	1.00 (-)	1.00 (-)
Yes	27,227	12 (1.8)	5 (1.0)	1 (2.56)	0.87 (0.49-1.55)	0.47 (0.19-1.13)	1.59 (0.21-11.9)
Worked in woodworking							
No	1,641,036	631 (95.3)	496 (95.6)	36 (92.3)	1.00 (-)	1.00 (-)	1.00 (-)
Yes	59,417	31 (4.7)	23 (4.4)	3 (7.7)	0.95 (0.66-1.38)	0.93 (0.61-1.42)	1.56 (0.47-5.19)
Worked in metal working							
No	1,259,257	527 (87.2)	417 (87.6)	27 (75.0)	1.00 (-)	1.00 (-)	1.00 (-)
Yes	194,469	77 (12.8)	59 (12.4)	9 (25.0)	0.76 (0.59-0.98)	0.76 (0.57-1.02)	1.48 (0.66-3.36)
Worked in shipyards							
No	1,445,935	607 (98.7)	479 (98.8)	37 (100.0)	1.00 (-)	1.00 (-)	1.00 (-)
Yes	20,655	8 (1.3)	6 (1.2)	0 (0.0)	0.59 (0.29-1.20)	0.58 (0.26-1.30)	N/A
Worked in electrical							
No	1,608,335	617 (93.1)	482 (92.7)	37 (94.9)	1.00 (-)	1.00 (-)	1.00 (-)
Yes	94,101	46 (6.9)	38 (7.3)	2 (5.1)	1.03 (0.76-1.40)	1.12 (0.80-1.57)	0.75 (0.18-3.15)
Worked in glass industry							
No	1,693,502	658 (99.4)	517 (99.6)	38 (97.4)	1.00 (-)	1.00 (-)	1.00 (-)
Yes	8,584	4 (0.6)	2 (0.4)	1 (2.6)	0.91 (0.34-2.43)	0.60 (1.50-2.39)	3.74 (0.50-27.5)
Worked in printing							
No	1,660,087	641 (96.8)	501 (96.5)	37 (94.9)	1.00 (-)	1.00 (-)	1.00 (-)
Yes	41,972	21 (3.2)	18 (3.5)	2 (5.1)	0.94 (0.61-1.46)	1.04 (0.65-1.67)	1.61 (0.38-6.79)
Worked in construction							

No	1,357,304	561 (91.7)	438 (90.9)	33 (89.2)	1.00 (-)	1.00 (-)	1.00 (-)
Yes	106,877	51 (8.3)	44 (9.1)	4 (10.8)	0.90 (0.67-1.21)	1.05 (0.76-1.45)	1.08 (0.37-3.16)
Worked in transportation							
No	1,608,577	611 (93.3)	482 (93.8)	34 (89.5)	1.00 (-)	1.00 (-)	1.00 (-)
Yes	85,566	44 (6.7)	32 (6.2)	4 (10.5)	1.00 (0.74-1.38)	0.97 (0.67-1.41)	1.37 (0.46-4.02)
Worked in nuclear industry							
No	1,274,387	420 (99.8)	322 (99.7)	27 (100.0)	1.00 (-)	1.00 (-)	1.00 (-)
Yes	4,816	1 (0.2)	1 (0.3)	0 (0.0)	0.46 (0.06-3.30)	0.60 (0.08-4.32)	N/A
Worked in paper or pulp production							
No	1,681,552	650 (98.0)	511 (98.3)	38 (97.4)	1.00 (-)	1.00 (-)	1.00 (-)
Yes	20,891	13 (2.0)	9 (1.7)	1 (2.6)	1.33 (0.76-2.30)	1.17 (0.60-2.27)	1.62 (0.22-11.99)
Worked in asbestos-cement production							
No	1,698,187	509 (99.8)	509 (99.8)	39 (100.0)	1.00 (-)	1.00 (-)	1.00 (-)
Yes	4,097	1 (0.2)	1 (0.2)	0 (0.0)	0.40 (0.06-2.84)	0.53 (0.07-3.74)	N/A
Worked in cement production							
No	1,692,849	661 (99.7)	514 (98.8)	38 (97.4)	1.00 (-)	1.00 (-)	1.00 (-)
Yes	9,480	2 (0.3)	6 (1.2)	1 (2.6)	0.39 (0.10-1.58)	0.26 (0.04-1.86)	2.94 (0.40-21.8)
Worked in ceramics production							
No	1,692,160	661 (99.7)	518 (99.6)	39 (100.0)	1.00 (-)	1.00 (-)	1.00 (-)
Yes	10,297	2 (0.3)	2 (0.4)	0 (0.0)	0.45 (0.11-1.80)	0.58 (0.14-2.31)	N/A
Worked as painter							
No	1,438,603	601 (97.7)	473 (97.5)	37 (100.0)	1.00 (-)	1.00 (-)	1.00 (-)
Yes	27,823	14 (2.3)	12 (2.5)	0 (0.0)	0.84 (0.50-1.44)	0.97 (0.54-1.73)	N/A
Worked as welder							
No	1,245,524	557 (96.0)	438 (95.8)	34 (97.1)	1.00 (-)	1.00 (-)	1.00 (-)
Yes	43,450	23 (4.0)	19 (4.2)	1 (2.9)	0.88 (0.58-1.35)	0.98 (0.62-1.57)	0.55 (0.07-4.13)
Worked as hairdresser							
No	1,677,116	659 (99.4)	517 (99.4)	39 (100.0)	1.00 (-)	1.00 (-)	1.00 (-)
Yes	25,317	4 (0.6)	3 (0.6)	0 (0.0)	0.42 (0.16-1.14)	0.40 (0.13-1.26)	N/A
Worked as bar tender							
No	831,198	293 (96.7)	229 (96.2)	17 (89.5)	1.00 (-)	1.00 (-)	1.00 (-)
Yes	30,363	10 (3.3)	9 (3.8)	2(10.5)	0.78 (0.41-1.46)	0.90 (0.46-1.75)	2.90 (0.66-12.8)

Worked as restaurant worker							
No	1,627,158	636 (95.9)	502 (96.5)	37 (94.9)	1.00 (-)	1.00 (-)	1.00 (-)
Yes	75,478	27 (4.1)	18 (3.5)	2 (5.1)	0.91 (0.62-1.34)	0.78 (0.48-1.24)	1.16 (0.28-4.87)
Worked in medical and health services							
No	1,557,127	606 (91.5)	470 (90.6)	37 (94.9)	1.00 (-)	1.00 (-)	1.00 (-)
Yes	144,554	56 (8.5)	49 (9.4)	2 (5.1)	1.13 (0.85-1.50)	1.24 (0.91-1.68)	0.72 (0.17-3.12)
Worked in textile/dying							
No	51,299	15 (100.0)	14 (100.0)	2 (100.0)	1.00 (-)	1.00 (-)	1.00 (-)
Yes	4,166	0 (0.0)	0 (0.0)	0 (0.0)	N/A	N/A	N/A
Worked in fibre preparation or weaving							
No	39,328	11 (73.3)	11 (78.6)	1 (33.3)	1.00 (-)	1.00 (-)	1.00 (-)
Yes	18,107	4 (26.7)	3 (21.4)	2 (66.6)	1.03 (0.32-3.33)	0.77 (0.21-2.84)	5.00 (0.40-62.1)
Worked in turning/metal engineering							
No	85,987	29 (38.2)	21 (36.2)	6 (66.6)	1.00 (-)	1.00 (-)	1.00 (-)
Yes	105,091	47 (61.8)	37 (63.8)	3 (33.3)	1.09 (0.67-1.80)	1.18 (0.66-2.09)	0.60 (0.12-2.90)
Worked in welding shop							
No	113,837	39 (51.3)	28 (49.1)	6 (66.6)	1.00 (-)	1.00 (-)	1.00 (-)
Yes	80,321	37 (48.7)	29 (50.9)	3 (33.3)	1.16 (0.72-1.88)	1.29 (0.74-2.25)	1.02 (0.23-4.48)
Worked in painting shop							
No	169,287	76 (92.7)	58 (93.6)	7 (77.8)	1.00 (-)	1.00 (-)	1.00 (-)
Yes	26,823	6 (7.3)	4 (6.4)	2 (22.2)	0.40 (0.17-1.15)	0.34 (0.12-0.94)	3.59 (0.64-20.0)
Worked as truck driver							
No	38,119	17 (38.6)	15 (46.9)	1 (25.0)	1.00 (-)	1.00 (-)	1.00 (-)
Yes	46,007	27 (61.4)	17 (53.1)	3 (75.0)	1.18 (0.62-2.26)	0.83 (0.39-1.74)	1.98 (0.18-21.6)
Worked as bus/taxi driver							
No	56,518	30 (62.5)	23 (65.7)	3 (60.0)	1.00 (-)	1.00 (-)	1.00 (-)
Yes	31,865	18 (37.5)	12 (34.3)	2 (40.0)	1.05 (0.55-2.00)	0.80 (0.37-1.75)	1.75 (0.27-11.3)
Worked in furniture production							
No	33,770	18 (58.1)	14 (60.9)	1 (33.3)	1.00 (-)	1.00 (-)	1.00 (-)
Yes	25,211	13 (41.9)	9 (39.1)	2 (66.6)	0.99 (0.47-2.07)	0.86 (0.36-2.04)	3.78 (0.32-44.4)
Worked in steel mill							

No	13,825	5 (62.5)	3 (50.0)	0 (0.0)	1.00 (-)	1.00 (-)	1.00 (-)
Yes	9,704	3 (37.5)	3 (50.0)	0 (0.0)	0.82 (0.16-4.11)	1.20 (0.20-7.03)	N/A
Worked in special alloys							
No	17,205	6 (75.0)	4 (66.7)	0 (0.0)	1.00 (-)	1.00 (-)	1.00 (-)
Yes	6,579	2 (25.0)	2 (33.3)	0 (0.0)	0.56 (0.08-3.87)	0.69 (0.09-5.29)	N/A
Worked as roofer							
No	82,928	40 (80.0)	34 (79.1)	2 (50.0)	1.00 (-)	1.00 (-)	1.00 (-)
Yes	18,735	10 (20.0)	9 (20.9)	2 (50.0)	0.96 (0.47-1.99)	1.01 (0.47-2.19)	3.88 (0.44-33.7)
Worked as asphalt worker							
No	89,961	44 (88.0)	38 (88.4)	3 (75.0)	1.00 (-)	1.00 (-)	1.00 (-)
Yes	11,855	6 (12.0)	5 (11.6)	1 (25.0)	0.93 (0.39-2.22)	0.89 (0.34-2.30)	2.02 (0.19-21.0)
Worked as demolition worker							
No	78,995	44 (86.3)	38 (88.4)	4 (100.0)	1.00 (-)	1.00 (-)	1.00 (-)
Yes	23,156	7 (13.7)	5 (11.6)	0 (0.0)	0.51 (0.22-1.14)	0.61 (0.27-1.39)	N/A
Worked in refinery							
No	55,892	22 (84.6)	18 (85.7)	1 (50.0)	1.00 (-)	1.00 (-)	1.00 (-)
Yes	6,676	4 (15.4)	3 (14.3)	1 (50.0)	1.17 (0.39-3.50)	0.97 (0.28-3.38)	10.4 (0.50-217)
Worked in dye production							
No	58,448	23 (85.2)	18 (85.7)	2 (100.0)	1.00 (-)	1.00 (-)	1.00 (-)
Yes	4,240	4 (14.8)	3 (14.3)	0 (0.0)	1.82 (0.61-5.49)	1.92 (0.55-6.69)	N/A
Worked with pesticides							
No	128,506	47 (50.5)	37 (50.7)	7 (77.8)	1.00 (-)	1.00 (-)	1.00 (-)
Yes	116,299	46 (49.5)	36 (49.3)	2 (22.2)	1.00 (0.66-1.52)	1.02 (0.63-1.64)	0.31 (0.06-1.57)
Worked in agriculture only before 1960							
No	78,770	18 (43.9)	16 (45.7)	2 (40.0)	1.00 (-)	1.00 (-)	1.00 (-)
Yes	81,540	23 (56.1)	19 (54.3)	3 (60.0)	1.09 (0.50-2.36)	1.02 (0.43-2.39)	2.55 (0.34-18.7)
Worked in agriculture also after 1960							
No	40,769	15 (36.6)	13 (37.1)	3 (60.0)	1.00 (-)	1.00 (-)	1.00 (-)
Yes	128,917	26 (63.4)	22 (62.9)	2 (40.0)	0.57 (0.28-1.19)	0.57 (0.26-1.24)	0.30 (0.04-2.25)

* Multivariate analysis stratified by centre and adjusted for gender, education level, alcohol consumption and smoking status at baseline.

Abbreviations: N/A = not applicable (no cases)