

The prevalence of non-allergic rhinitis phenotypes in the general population: A cross-sectional study

Klementina S. Avdeeva¹  | Wytske J. Fokkens¹  | Christine L. Segboer² | Sietze Reitsma¹

¹Department of Otorhinolaryngology, Amsterdam UMC, Location Academic Medical Centre, Amsterdam, The Netherlands

²Department of Otorhinolaryngology, Dijklander Ziekenhuis, Hoorn, The Netherlands

Correspondence

Klementina S. Avdeeva, Department of Otorhinolaryngology, Amsterdam UMC, Location Academic Medical Centre, Amsterdam, The Netherlands.
Email: k.avdeeva@amsterdamumc.nl

Abstract

Background: Non-allergic rhinitis (NAR) can be subdivided into several phenotypes: rhinorrhea of the elderly, rhinitis medicamentosa, smokers', occupational, hormonal, drug-induced, gustatory, and idiopathic rhinitis. There are two pathophysiological endotypes of NAR: inflammatory and neurogenic. Phenotypes may serve as an indicator of an underlying endotype and, therefore, help to guide the treatment. The prevalence of each phenotype in the general population is currently unknown.

Methodology/Principal: Cross-sectional questionnaire-based study in the general population of the Netherlands.

Results: The prevalence of chronic rhinitis in the general population was 40% ($N = 558$, of those, 65% had NAR and 28% AR, in 7% allergy status is unknown). Individuals with NAR ($N = 363$) had significantly more complaints in October–February. Those with AR ($N = 159$) had significantly more complaints in April–August. The most common NAR phenotypes were idiopathic (39%) and rhinitis medicamentosa (14%), followed by occupational (8%), smokers' (6%), hormonal (4%), gustatory (4%), and rhinorrhea of the elderly (4%). The least prevalent phenotype was drug induced (1%). Nineteen percent of the NAR group could not be classified into any of the phenotypes.

Conclusions: This is the first study to describe the prevalences of NAR phenotypes in the general population. AR and NAR have a distinct seasonality pattern with NAR being more prevalent in autumn/winter and AR in spring/summer. Our data on the prevalence of phenotypes may help clinicians to anticipate the type of patients at their clinic and help guide a tailored treatment approach. The high prevalence of rhinitis medicamentosa is alarming, since this is a potentially preventable phenotype.


KEYWORDS

endotype, epidemiology, non-allergic rhinitis, phenotype, prevalence

Abbreviations: AR, Allergic rhinitis; CR, Chronic rhinitis; CRSwNP, Chronic rhinosinusitis with nasal polyps; DR, Drug-induced rhinitis; HR, Hormonal rhinitis; IR, Idiopathic rhinitis; LAR, Local allergic rhinitis; MWU, Mann-Whitney U; NAR, Non-allergic rhinitis; NARES, Non-allergic rhinitis with eosinophilia syndrome; ND, Nasal decongestant; NERD, NSAID-exacerbated respiratory disease; NSAID, Non-steroid anti-inflammatory drug; OCP, Oral contraceptive pill; OR, Occupational rhinitis; PDE5, Phosphodiesterase 5; RM, Rhinitis medicamentosa; RoE, Rhinorrhea of the elderly; SR, Smokers rhinitis; URTI, Upper respiratory tract infection; VAS, Visual analogue scale; WHO, World health organization.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

© 2022 The Authors. *Allergy* published by European Academy of Allergy and Clinical Immunology and John Wiley & Sons Ltd.



GRAPHICAL ABSTRACT

The prevalence of NAR and AR in general population: 27% and 12%. AR symptoms get worse in spring/ summer, NAR symptoms in autumn/ winter. The prevalences of NAR phenotypes: idiopathic 39%, rhinitis medicamentosa 14%, occupational 8%, smokers' 6%, hormonal 4%, rhinorrhoea of the elderly 4%, gustatory 4% and drug-induced 1%. Nineteen percent could not be defined in neither of the phenotypes.

1 | INTRODUCTION

Non-allergic rhinitis (NAR) is a form of chronic rhinitis (CR) that is characterized by rhinorrhea, blocked nose, sneezing, and/or itchy nose without clinical signs of infection or allergy.^{1,2} NAR decreases the quality of life³ and has a substantial financial impact.⁴ We have recently suggested that CR (and, consequently, NAR) should be defined in epidemiological studies as the presence of at least one nasal symptom for more than 21 days per year.⁵ When this definition was applied to the general population, the prevalence of CR was 40% (65% NAR, 28% allergic rhinitis (AR), and in 7% allergy status was unknown).⁵

NAR is an umbrella term that covers a heterogeneous group of patients whose complaints have different phenotypes and endotypes and, therefore, require different treatment strategies. To choose the best treatment for each patient, it is necessary to understand the underlying endotype. While in clinical practice, determination of endotype in each patient is not technically feasible, and phenotypes may serve as a predictor¹ and, therefore, may help to tailor the treatment. Additionally, understanding the (prevalence of the) phenotypes of NAR may help clinicians to expect what kind of NAR patients they will most often encounter. The phenotypes of NAR defined/mentioned in the EAACI taskforce paper include rhinitis (rhinorrhea) of the elderly, *rhinitis medicamentosa*, smokers', occupational, hormonal, drug-induced, gustatory, and idiopathic rhinitis. Rhinitis medicamentosa is a subtype of drug-induced rhinitis, which we propose to distinguish as a separate phenotype due to a different route of the drug administration and the underlying mechanism.


The term "idiopathic rhinitis" is used when a patient does not fit into any of the aforementioned phenotypes and suffers from NHR. NHR is the stimulation of one or more nasal symptoms upon encounter of unspecific environmental stimuli, such as temperature or humidity changes, strong odors, or smoke¹ and is objectively confirmed by cold-dry air provocation.⁶ NHR is a common phenomenon in rhinitis and rhinosinusitis.⁷⁻⁹

The most important endotypes of NAR are inflammatory and neurogenic (Figure 1).¹

Some NAR phenotypes clearly belong to one endotype: idiopathic,¹⁰ rhinorrhea of the elderly, and gustatory rhinitis are of the neurogenic endotype.¹¹ Other phenotypes, for example, occupational rhinitis, may belong to both inflammatory and neurogenic, depending on the agent that has led to the development of the symptoms.¹ Mechanisms behind several phenotypes, for example, hormonal, are not completely understood.^{11,12}

Apart from the obvious subtypes of AR and NAR, some potential intermediate phenotypes, both with an inflammatory endotype,¹ exist: local allergic rhinitis (LAR) and non-allergic rhinitis with eosinophilia syndrome (NARES). LAR is characterized by local production of specific IgE during nasal exposure to aeroallergens¹³⁻¹⁶ in the absence of systemic atopy.¹⁷ Whether LAR should be considered a subtype of AR, NAR, or an independent entity is a subject of debate.¹⁸ On the one hand, LAR is similar to AR in terms of pulmonary comorbidities,¹⁷ sensitizations to aeroallergens tested by nasal provocation testing,¹³ and treatment strategies.^{19,20} On the other hand, LAR does not evolve into systematic atopy over time.¹⁷ Besides, LAR and AR may co-exist in a form of

FIGURE 1 Endotype and phenotype of NAR



so-called dual-allergic rhinitis.^{21,22} NARES is characterized by the presence of eosinophils in nasal mucosa in patients with NAR.²³ Seeing that local eosinophilia is present in both conditions²⁴ and the fact that studies describing NARES did not perform intranasal allergen provocations nor measurement of nasal IgE,^{25,26} LAR is possibly a subtype of NARES. On the other hand, some studies claim that these are two independent subtypes¹¹: in the study by Meng et al., NARES was characterized by high eosinophil cationic protein, whereas LAR was characterized by high histamine.²⁴ Another study demonstrated the absence of allergen-specific IgE in NARES,¹² which is characteristic of LAR. For this study, we consider LAR to be a form of AR because it cannot be discriminated with a questionnaire only from AR.

We found a prevalence of NAR of 28% in general population.⁵ To the best of our knowledge, the prevalence of the different phenotypes of NAR is currently unknown. Seeing that phenotypes may serve as an indicator of an underlying endotype, knowledge about their prevalence may help clinicians to expect the phenotypes they will most often encounter in daily practice and to choose tailored treatment strategies. The aim of this study was to determine the prevalence of NAR phenotypes in the general population.

2 | METHODS

We performed a cross-sectional, questionnaire-based study in a random sample of participants representing the general population of the Netherlands. Informed consent was obtained from all participants. The same subset of participants was used for an early published paper on the definition of CR.⁵ The questionnaire (Attachment 1) included questions intended to define each of the phenotypes.

2.1 | Definitions used

2.1.1 | Chronic rhinitis

CR was defined as the presence of nasal complaints about at least 21 days/year or a history of a positive allergy test and nasal medication use (outside of periods with common cold), irrespective of duration of the complaints.

2.1.2 | AR and NAR

When the participants fulfilled the CR criteria and answered affirmatively on the question whether they had allergic rhinitis or hay fever, they were considered as having (self-reported) AR. When the answer was negative, they were considered as having NAR, unless they had only eye symptoms ($N = 14$). Every patient who missed the question "Do you have allergic rhinitis" was excluded from this analysis.

2.1.3 | Nasal complaints

We asked participants which of the nasal complaints they had for at least one hour per day on most days of the week (hereinafter termed "regular nasal complaints"). Additionally, we asked them which of the aforementioned complaints they were experiencing at the time of filling in the questionnaire (hereinafter "current nasal complaints").

2.1.4 | Definitions of phenotypes

The phenotypes were defined according to EAACI position paper on NAR.¹ The algorithm used for phenotypes definitions is




FIGURE 2 Algorithm behind definitions of NAR phenotypes. Green arrows with a "+" sign represent "Yes," and red arrows with a "-" sign represent "No." NHR – nasal hyperreactivity

presented in Figure 2. Based on the algorithm, each subject was assigned exclusively to one phenotype. If a participant was not defined as having rhinorrhea of the elderly (ROE), occupational rhinitis (OR), rhinitis medicamentosa (RM), smokers' rhinitis (SR), hormonal rhinitis (HR), drug-induced rhinitis (DR), or gustatory rhinitis (GR) and had CR and NHR, he/she was classified in idiopathic rhinitis (IR) group. If NHR was absent, he/she was classified into the "other" group.

2.2 | Statistical analysis

Data are summarized as frequencies, means and standard deviations, medians, and interquartile ranges. To detect the differences between the groups, we used an unpaired sample *t*-test for normally distributed numerical variables (age), and an independent-sample Mann-Whitney U-test (MWU-test) for non-normally distributed numerical variables (VAS score) (SPSS).

3 | RESULTS

In December 2019, five thousand questionnaires were sent out to the residents of one municipality in the Netherlands. One thousand thirty-three participants filled in the questionnaire (response rate 27%).

3.1 | Common cold

In the whole study group, the median number of common colds was 2 (IQR 1–3). The median number of total days with common colds was 10 (IQR 4–20).

3.2 | AR and NAR

There were 363 participants with NAR, 159 with AR, and 790 controls. Twenty-two participants who did not answer the question “Do you have allergic rhinitis” were excluded from this analysis. Figure 3 shows the distribution of months with worst complaints for AR and NAR. Individuals with NAR had significantly more complaints in October–February, (chi-square, $p < .001$). Those with AR had significantly more complaints in April–August (chi-square, $p < .05$).

3.3 | NAR phenotypes

The prevalence of NAR phenotypes is presented in Figure 4. The most prevalent phenotype was IR (39% of NAR group). A comparison of baseline characteristics of NAR phenotypes is presented in Table 1. Nasal obstruction, rhinorrhea, and postnasal drip were the most common regular complaints (Figure 5).

3.4 | Rhinorrhea of the elderly

- In the general population, participants with rhinorrhea as the only regular complaint were significantly older compared to the rest (64 ± 15 vs 59 ± 17 years, $p = .028$). In NAR, the same trend was observed, though not statistically significant (61 ± 16 vs 58 ± 18).
- In NAR, participants with rhinorrhea as the most bothering complaint were older than the rest (62 ± 19 vs 56 ± 18 , T-test $p = .026$, MWU-test $p = .008$).
- Regardless of the age cut off used, the prevalence of RoE was around 10% for the corresponding age-group (Attachment 2). The remaining 90% of participants of the same age had other phenotypes of NAR (with the same prevalence of each phenotype as for the whole NAR group).

3.5 | Occupational rhinitis

The majority of working participants in the NAR group reported having nasal complaints at work (Attachment 2), but only a third of them reported an improvement during weekend/vacation.

From all the participants defined as having OR ($N = 30$), only one had nasal complaints related to work-specific triggers. In the majority of participants ($N = 11$), nasal complaints were attributable to hyperreactivity to non-specific triggers encountered at work (such as air conditioning, dust, and dry air). Seven participants thought that their complaints were caused by HDM or other allergies, and in eleven participants, the trigger was unknown.

3.6 | Rhinitis medicamentosa

- Sixty-one participants in the NAR group (17%) and 42 controls (5%) reported regular use of nasal decongestants (ND) (Attachment 2).




FIGURE 3 Percentages of individuals with worst nasal complaints per month of the year, for AR and NAR. Vertical bars represent 95% CI. The differences were statistically more significant in January, February, April, May, June, July, August, October, November, and December




FIGURE 4 The prevalence of NAR phenotypes (NAR group, $N = 363$)

Fifty-one participants with NAR were defined as having *rinitis medicamentosa* (RM), and ten participants classified into the OR group (Figure 2).

- Within participants who reported the regular use of ND, those with NAR had a significantly higher VAS compared to those from the control group (45 ± 26 vs 19 ± 24 , t -test $p < .001$; MWU test $p < .001$). Similarly, there were significantly more moderate/severe cases (ARIA) among ND users from the NAR group (84% vs 36%, chi-square $p < .001$).

3.7 | Smokers' rhinitis

3.7.1 | Current smokers

- Current smokers were significantly more likely to have NAR compared to non-smokers (chi-square $p = .034$, OR 1.7 (1.04–2.8); RR 1.6 (1.004–2.6)).
- Current smokers in the NAR group had significantly more pack-years of smoking compared to currently smoking controls (32 ± 29 vs 14 ± 14 ; t -test $p = .014$; MWU test $p = .04$).

3.7.2 | Former smokers

- Former smokers had the same probability of having NAR compared with those who never smoked.
- Pack/year of smoking for former smokers was not different between NAR and controls.

3.8 | Hormonal rhinitis

Sixteen women were classified as having HR. Two of them had nasal complaints during both pregnancy and menstruation, eight during pregnancy only, and six during menstruation only.

3.9 | Drug-induced rhinitis

- There were significantly more medication users in NAR compared to the controls (chi-square $p = .008$, OR 1.4 95% CI 1.1–1.8, RR 1.2 95% CI 1.1–1.5).
- The proportions of users of painkillers (including paracetamol and NSAIDs), β -blockers, antidepressants, sedatives, and "other" medications were not significantly different between NAR and controls. There was also no difference in medication intolerance.
- Thirty-six participants have developed their nasal symptoms after initiation of a medication. Of those, in six participants nasal complaints have started the same year as they have started using the medication. Of the whole group of participants who developed symptoms after initiation of a medication, five were using medication enlisted in EAACI position paper and were defined as having DR. The medications that they reported (could be more than one per subject) were NSAIDs ($N = 1$), β -blockers ($N = 3$), antidepressants ($N = 2$), and "other" ($N = 4$).
- The majority of NAR population used medications classified as "other": anticoagulants, calcium channel blockers, ACE inhibitors, statins, glucose-lowering medications, thiazide diuretics, PPI inhibitors, P2Y12 inhibitors, etc.

3.10 | Gustatory rhinitis

- Sixteen participants (4% of NAR) were defined as having GR.
- In the GR group, rhinorrhea was the most prevalent regular and current complaint ($N = 10$, 63%, for both). The most bothering complaints were rhinorrhea ($N = 7$, 44%) and nasal obstruction ($N = 6$, 38%).
- The majority ($N = 11$, 69%) of the GR group reported NHR.

3.11 | Idiopathic rhinitis

- One hundred and forty-one participants (39% of NAR) were classified as having IR.

3.12 | Other

- Sixty-eight participants (19% of NAR) could not be classified into any of the phenotypes.

TABLE 1 Characteristics of NAR phenotypes*

		Rhinorrhea of the elderly		Occupational	Medicamentosa	Smokers'	Hormonal	Gustatory	Idiopathic	Other	Controls
N (%) of the NAR group)		15 (4%)		30 (8%)	51 (14%)	21 (6%)	16 (4%)	16 (4%)	141 (39%)	68 (19%)	790
Demographic characteristics											
Age	Mean $\pm$ SD	78 $\pm$ 6.3		52 $\pm$ 13	60 $\pm$ 17	61 $\pm$ 15	45 $\pm$ 18	52 $\pm$ 19	57 $\pm$ 18	63 $\pm$ 17	60 $\pm$ 16
	Median (IQR)	77 (73; 82)		54 (43.5; 64)	65 (53.5; 73)	62 (55.5; 69.5)	47.5 (30; 57.25)	60 (35.25; 66.75)	59 (48.5; 70)	68 (54; 75)	63 (51; 72)
Gender: Female N (%)		5 (33%)		18 (60%)	26 (51%)	9 (43%)	16 (100%)	8 (50%)	62 (44%)	24 (35%)	427 (54%)
Current smoker		0		4 (13%)	5 (10%)	21 (100%)	-	-	-	-	38 (5%)
Former smoker		7 (47%)		15 (50%)	24 (47%)	-	3 (19%)	3 (19%)	76 (54%)	37 (54%)	337 (43%)
Self-reported asthma		0		5 (17%)	6 (12%)	0	0	0	7 (5%)	4 (6%)	26 (3%)
Other pulmonary complaints (cough, dyspnea, shortness of breath, and wheezing)		6 (40%)		13 (43%)	20 (39%)	12 (57%)	8 (50%)	6 (38%)	39 (28%)	15 (22%)	91 (11%)
Number of common colds per year	Mean $\pm$ SD	11 $\pm$ 27		6 $\pm$ 20	6 $\pm$ 9	3 $\pm$ 5	4 $\pm$ 4	5 $\pm$ 8	5 $\pm$ 9	2 $\pm$ 3	2 $\pm$ 4
	Median (IQR)	3 (1; 7)		3 (2; 4)	3 (2; 5)	2 (1; 3)	2.5 (1; 10)	2 (1; 5)	2 (1; 4)	2 (1; 2)	1 (1; 2)
Nasal medication use											
Regular use of nasal decongestants		0		10 (33%)	51	-	-	-	-	-	42 (5%)
Regular use of nasal medication		2 (13%)		8 (27%)	16 (31%)	5 (24%)	2 (13%)	0	24 (17%)	6 (9%)	28 (4%)
Systemic antihistamines		0		1 (3%)	0	1 (5%)	0	0	1 (1%)	0	6 (1%)
Topical steroids and/or antihistamines		1		4 (13%)	11 (22%)	0	0	0	12 (9%)	5 (7%)	9 (1%)
Saline nasal spray/rinse		0		2 (7%)	1 (2%)	1 (5%)	1 (6%)	0	8 (6%)	0	8 (1%)
Other/unknown		1		3 (10%)	6 (12%)	3 (14%)	1 (6%)	0	7 (5%)	1 (1%)	9 (1%)
Nasal complaints											
The number of regular nasal complaints	Mean $\pm$ SD	2.6 $\pm$ 1.5		2.4 $\pm$ 1.6	2.3 $\pm$ 1.6	2.3 $\pm$ 1.3	2.4 $\pm$ 1.3	2.2 $\pm$ 1.5	1.9 $\pm$ 1.6	1.5 $\pm$ 1.1	0.3 $\pm$ 0.8
	Median (IQR)	3 (1; 4)		2.0 (1.0; 4.0)	2 (1; 3)	2 (1; 3)	2 (1.25; 3.75)	2 (1; 3.75)	2 (1; 3)	1 (1; 2)	0 (0; 0)
The number of current nasal complaints	Mean $\pm$ SD	2.4 $\pm$ 1.2		2.4 $\pm$ 1.6	2.3 $\pm$ 1.5	1.6 $\pm$ 1.3	2 $\pm$ 0.8	2.1 $\pm$ 1.3	2 $\pm$ 1.4	1.8 $\pm$ 1.2	0.4 $\pm$ 1.0
	Median (IQR)	2 (1; 3)		1.6 (1; 4)	2 (1; 3)	1 (1; 2.5)	2 (1; 2.75)	2 (1; 3)	2 (1; 3)	1 (1; 2)	0 (0; 0)

(Continues)


TABLE 1 (Continued)

		Rhinorrhea of the elderly								
		Occupational	Medicamentosa	Smokers'	Hormonal	Gustatory	Idiopathic	Other	Controls	
Duration of nasal complaints per day	<1 h per day	8 (27%)	20 (39%)	9 (43%)	6 (38%)	6 (38%)	54 (38%)	29 (43%)	84 (11%)	
	≥1 h per day	22 (73%)	25 (4%)	12 (57%)	10 (63%)	9 (56%)	80 (57%)	31 (46%)	55 (7%)	
	No nasal complaints/Unknown	0	2 (4%)	0	0	1 (6%)	3 (2%)	8 (12%)	440 (56%)	
Duration of nasal complaints per year (days per year)	Mean ± SD	180 ± 129	177 ± 127.45	218 ± 121	268 ± 111	239 ± 117	224 ± 131	219 ± 133	17 ± 47	
	Median (IQR)	110 (63.75; 300)	150 (50; 300)	200 (100; 365)	325 (150; 365)	300 (112.5; 348.75)	200 (100; 365)	200 (100; 365)	8 (0; 14)	
Years with nasal complaints	Mean ± SD	15 ± 13	17 ± 17	15 ± 9	24 ± 18	12.5 ± 16	18 ± 17	16 ± 17	14 ± 19	
	Median (IQR)	11 (4.75; 20.25)	10 (4.5; 22.5)	17 (5.5; 20)	21 (11; 38)	5 (1; 15)	12.5 (5; 21.75)	10 (3; 20)	5 (1; 20)	
ARIA										
Severity	Mild	3 (10%)	8 (16%)	8 (38%)	5 (31%)	3 (19%)	35 (25%)	29 (43%)	333 (42%)	
	Moderate/severe	26 (87%)	42 (82%)	11 (52%)	9 (56%)	12 (75)	94 (67%)	36 (52%)	119 (15%)	
	Unknown	1 (3%)	1 (2%)	2 (10%)	2 (13%)	1 (6%)	12 (9%)	3 (4%)	319 (41%)	
Duration	Intermittent	15 (50%)	25 (49%)	9 (43%)	3 (19%)	7 (44%)	59 (42%)	32 (47%)	266 (34%)	
	Persistent	15 (50%)	24 (47%)	12 (57%)	13 (81%)	8 (50%)	79 (56%)	33 (49%)	22 (3%)	
	Unknown	0	2 (4%)	0	0	1 (6%)	3 (2%)	3 (4%)	502 (64%)	
VAS nasal complaints (0–100, mm)	Mean ± SD	44 ± 25	43 ± 27	34 ± 29	45 ± 28	45 ± 24	38 ± 26	30 ± 27	10 ± 17	
	Median (IQR)	53 (19.5; 54.5)	41.5 (19.75; 65)	29 (7.25; 58.75)	35.5 (21.5; 73.75)	49.5 (19.75; 54.75)	31.5 (17; 60.75)	22 (7; 48)	2 (0; 12)	

Abbreviations: DR, drug-induced rhinitis; GR, gustatory rhinitis; HR, hormonal rhinitis; NAR, non-allergic rhinitis; OR, occupational rhinitis; RM, rhinitis medicamentosa; RoE, rhinorrhea of the elderly; SR, smokers' rhinitis.

*Drug-induced rhinitis (N = 5) is not presented due to a small size.

FIGURE 5 The prevalence of regular nasal complaints per phenotype in NAR group ($N = 363$). Drug-induced rhinitis and occupational rhinitis are not presented due to small group sizes



- This group was mostly represented by participants with milder complaints in terms of the number of complaints, VAS score, and the proportion of mild cases (ARIA) (Table 1).

4 | DISCUSSION

This is the first paper that describes the prevalence of NAR phenotypes in the general population. Though some data on prevalence of individual phenotypes in particular populations are available (such as HR in pregnancy,²⁷ SR^{28,29} and OR in bakers³⁰), the distribution of each phenotype (as described by EAACI position paper¹) in the general population has not been previously reported. Phenotyping of NAR is essential for choosing the best treatment option. For example, IR is effectively treated with capsaicin,³¹ which is totally different from the treatment of RoE with ipratropium bromide³² or the treatment of RM by discontinuing the ND abuse.

For studies on prevalence of diseases, data from the general population are mandatory. In this study, we found a prevalence of CR of 40% what is comparable to other European studies.³³⁻³⁵ Ideally, some form of allergy testing should be used for the differentiation between AR and NAR in studies, but that is not always feasible. However, Saviouré et al. have shown that the question "Have you ever had allergic rhinitis?" or "Have you ever had hay fever?" has a positive predictive value of 0.71 (0.64–0.78) and a negative predictive value of 0.77 (0.71–0.84) in never asthmatics.³⁶ We expect that our study where we used not only the question about allergic rhinitis/hay fever but also whether the patient was tested for allergies will have at least comparable results. Our studies showed 70% of the subjects with CR to have NAR. This is higher than in a study by Bachert et al., where the prevalences of AR and NAR had a proportion of 3:1,³⁵ or the study of Bozek et al., where the proportion was 1:1,¹³ and is comparable with the study

by Jessen and Janzon, where the proportion was 1:4.³⁷ One of the reasons for this discrepancy could be that we only evaluated subjects with CR. It is possible that more subjects with NAR than AR have symptoms for more than 21 days per year. Furthermore, we do not expect that AR/NAR proportion would influence the results of NAR phenotype analysis. In this study, as far as we know, for the first time we show the difference between two distinct seasonality patterns between the AR and the NAR, with AR complaints being more prevalent during pollen season³⁸ and NAR during winter, when viral infections and factors that may correspond to NHR are more apparent (e.g., dry air and temperature differences between inside and outside). Possibly, a proportion of NAR patients is represented by unconfirmed mono-sensitized HDM allergy, though mono-sensitization to HDM is quite rare.³⁸ Therefore, only a minority of NAR group could have been wrongly assigned because of an unrecognized mono-sensitization to HDM.

To further phenotype NAR, we used definitions as proposed by the EAACI position paper.¹ Some subjects (7% of NAR group) can be phenotyped in different categories. For example, out of 31 currently smoking participants, 24 were classified as SR and seven as RM, though the latter seven participants most likely have a combination of SR and RM. For this study, we have chosen to exclusively include subjects in the most likely category and used an algorithm as described in this manuscript (Figure 2). Our definition, therefore, did not account for possible overlaps and the interactions between the provoking factors, what is more likely in reality. Hence, the reader should keep in mind that these data are an oversimplification of the truth. On the other hand, our approach to use NHR as a criterion for IR is a valid attempt to separate rhinitis patients from those with other diagnoses, which is reflected by a lower disease burden in the "other" group.

There are still many gaps in knowledge about NAR phenotypes. To facilitate further research in the field, robust definitions for each

phenotype should be developed (e.g., what duration and frequency of a medication use lead to the development of drug-induced rhinitis; after what age do we consider nasal discharge a RoE, etc.). Moreover, although there is increasing information about the relation between phenotypes and endotypes in NAR with the consequences for therapy, the picture is far from complete. In Figure 1, we propose underlying endotypes like neurogenic and inflammatory for the different phenotypes. We do realize, however, that significant overlap can be found (e.g., OR and DR can be inflammatory or neurogenic), of which it is unclear whether that is present in all subjects. Additionally, RM, apart from having a neurogenic component, also has a vascular one. In order to systematically fill in those gaps and to refine existing definitions and classification, and due to a shortage of effective NAR treatment options, we suggest that a new EAACI taskforce addresses these issues.

Van Rijswijk hypothesized that IR accounts for about 50% of NAR.³⁹ Our findings confirm this hypothesis: IR was the most prevalent phenotype (39% of NAR group). IR is a purely neurogenic phenotype, and the recognition of the disease with NHR as the most prominent symptom⁴⁰ has significant implications for the treatment.^{31,40-42}

Twenty percent of the NAR group could not be classified into any of the phenotypes. We were not able to identify a pattern that would explain the nature of nasal complaints in this group. If NHR was not used as a defining factor for IR, these patients would have been classified as having IR. Possibly, this group includes participants with anatomical reasons for nasal symptoms like a septal deviation and/or inferior turbinate hypertrophy and/or some (yet) unrecognized phenotypes.⁴³

The second most common phenotype was RM (17% of NAR). Strictly speaking, RM is a subtype of DR,¹ though we have analyzed them separately due to the fact that RM is caused by a very specific medication that acts locally, has a different pathophysiological mechanism,¹¹ and is associated with psychiatric conditions, such as anxiety⁴⁴ or opioid use disorders.⁴⁵ Mehuys et al. have also demonstrated an alarmingly high rates of ND (ab)use: about a half of individuals with persistent rhinitis purchasing over-the-counter medication for their nasal complaints were overusing ND, even though the majority was educated about the limit on duration of use.⁴⁶ A high prevalence of RM is a major concern, seeing that this is a preventable phenotype, and warrants attention in our daily practice because it is something often not spontaneously mentioned by our patients.

Interestingly, the pattern of complaints does not predict phenotype nor endotype (Figure 5) (except for ROE, which was defined by the presence of rhinorrhea). As such, OR, RM, and SR have a similar pattern with nasal obstruction and post-nasal drip being the most prominent complaints, but they all belong to different endotypes (inflammatory or neurogenic, neurogenic, and inflammatory, respectively). Likewise, HR and GR are similar in terms of rhinorrhea and nasal obstruction but, then again, belong to different endotypes (inflammatory and neurogenic, respectively). This could be explained by the presence of (sub)endotypes or gaps in knowledge regarding the pathophysiology of the endotypes.

Our study has a number of limitations that were described earlier and in our previous publication,⁵ such as inability to objectively exclude NARES, LAR, septal deviations, or CRS. Though due to the low prevalence of CRS (~4% of general population when confirmed objectively⁴⁷⁻⁴⁹) and NARES (2% in Dutch primary care patients with recurrent nasal symptoms²⁵), we expect that only a minority of the NAR group was represented by these conditions. Seeing that the severity of septal deviation does not correlate with subjective complaints⁵⁰ and that CR was defined by the presence of nasal symptoms, we do not expect that inability to exclude this anatomical feature would influence the results.

Another major limitation is the lack of a more thorough medical history, which may be necessary for accurate phenotyping. For example, we assigned all subjects that developed nasal symptoms after starting medication that can induce DR¹ into this phenotype. But subjects might have had other reasons for their symptoms not related to the use of medication, and we were not able to stop medication to prove the relation. Additionally, none of the participants reported the use of phosphodiesterase-5 inhibitors, probably because they do not consider it as a "medication," though it may cause DR.⁵¹

Moreover, the limited size of some of our groups made further subgroup analyses impossible. Finally, seasonality patterns described for AR and NAR should be validated for other climate zones.

It was previously hypothesized that up to 50%–70% of AR may be represented by mixed rhinitis.⁵² Unfortunately, the design of the questionnaire did not allow for the differentiation of this group.

5 | CONCLUSIONS

This is the first study to describe the prevalence of NAR phenotypes in general population. AR and NAR have distinct seasonality patterns with NAR being more prevalent in autumn/winter and AR in spring/summer. Idiopathic rhinitis is the most common phenotype of NAR, followed by rhinitis medicamentosa. The high prevalence of rhinitis medicamentosa is alarming. A new EAACI taskforce should systematically fill in gaps in knowledge about NAR phenotypes.

ACKNOWLEDGMENTS

The authors thank I. Bruins and Y. te Winkel for their assistance in digitalization of the questionnaires.

CONFLICT OF INTEREST

None.

ORCID

Klementina S. Avdeeva  <https://orcid.org/0000-0002-3910-4371>

Wyske J. Fokkens  <https://orcid.org/0000-0003-4852-229X>

REFERENCES

1. Hellings PW, Klimek L, Cingi C, et al. Non-allergic rhinitis: position paper of the European Academy of Allergy and Clinical Immunology. *Allergy*. 2017;72(11):1657-1665.

2. Bousquet J, Khaltaev N, Cruz AA, et al. Allergic Rhinitis and its Impact on Asthma (ARIA) 2008 update (in collaboration with the World Health Organization, GA(2)LEN and AllerGen). *Allergy*. 2008;63(Suppl 86):8-160.
3. Segboer CL, Terreehorst I, Gevorgyan A, Hellings PW, van Drunen CM, Fokkens WJ. Quality of life is significantly impaired in nonallergic rhinitis patients. *Allergy*. 2018;73(5):1094-1100.
4. Avdeeva KS, Reitsma S, Fokkens WJ. Direct and indirect costs of allergic and non-allergic rhinitis in the Netherlands. *Allergy*. 2020;75(11):2993-2996.
5. Avdeeva KS, Fokkens WJ, Reitsma S. Towards a new epidemiological definition of chronic rhinitis: prevalence of nasal complaints in the general population. *Rhinology*. 2021;59(3):258-266.
6. Huang Y, Lou H, Wang C, Zhang L. Cold dry air provocation is a reliable diagnostic tool in nonallergic rhinitis. *Rhinology*. 2019;57(3):225-230.
7. Backaert W, Steelant B, Jorissen M, et al. Self-reported nasal hyperreactivity is common in all chronic upper airway inflammatory phenotypes and not related to general well-being. *Allergy*. 2021;76(12):3806-3809.
8. Doulaptsi M, Steelant B, Prokopakis E, et al. Prevalence and impact of nasal hyperreactivity in chronic rhinosinusitis. *Allergy*. 2020;75(7):1768-1771.
9. Segboer CL, Holland CT, Reinartz SM, et al. Nasal hyper-reactivity is a common feature in both allergic and nonallergic rhinitis. *Allergy*. 2013;68(11):1427-1434.
10. van Rijswijk JB, Blom HM, KleinJan A, Mulder PG, Rijntjes E, Fokkens WJ. Inflammatory cells seem not to be involved in idiopathic rhinitis. *Rhinology*. 2003;41(1):25-30.
11. Papadopoulos NG, Bernstein JA, Demoly P, et al. Phenotypes and endotypes of rhinitis and their impact on management: a PRACTALL report. *Allergy*. 2015;70(5):474-494.
12. Becker S, Rasp J, Eder K, Berghaus A, Kramer MF, Gröger M. Non-allergic rhinitis with eosinophilia syndrome is not associated with local production of specific IgE in nasal mucosa. *Eur Arch Otorhinolaryngol*. 2016;273(6):1469-1475.
13. Bozek A, Scierski W, Ignasiak B, Jarzab J, Misiolek M. The prevalence and characteristics of local allergic rhinitis in Poland. *Rhinology*. 2019;57(3):213-218.
14. Eguiluz-Gracia I, Testera-Montes A, Rondon C. Medical algorithm: diagnosis and treatment of local allergic rhinitis. *Allergy*. 2021;76(9):2927-2930.
15. Meng Y, Wang Y, Lou H, et al. Specific immunoglobulin E in nasal secretions for the diagnosis of local allergic rhinitis. *Rhinology*. 2019;57(4):313-320.
16. Hamizan AW, Rimmer J, Husain S, et al. Local specific Immunoglobulin E among patients with nonallergic rhinitis: a systematic review. *Rhinology*. 2019;57(1):10-20.
17. Rondon C, Campo P, Eguiluz-Gracia I, et al. Local allergic rhinitis is an independent rhinitis phenotype: the results of a 10-year follow-up study. *Allergy*. 2018;73(2):470-478.
18. Reitsma S, Subramaniam S, Fokkens WWJ, Wang Y. Recent developments and highlights in rhinitis and allergen immunotherapy. *Allergy*. 2018;73(12):2306-2313.
19. Rondón C, Blanca-López N, Campo P, et al. Specific immunotherapy in local allergic rhinitis: a randomized, double-blind placebo-controlled trial with Phleum pratense subcutaneous allergen immunotherapy. *Allergy*. 2018;73(4):905-915.
20. Hellings PW, Scadding G, Bachert C, et al. EUFOREA treatment algorithm for allergic rhinitis. *Rhinology*. 2020;58(6):618-622.
21. Eguiluz-Gracia I, Fernandez-Santamaria R, Testera-Montes A, et al. Coexistence of nasal reactivity to allergens with and without IgE sensitization in patients with allergic rhinitis. *Allergy*. 2020;75(7):1689-1698.
22. Zhang Y, Lan F, Zhang L. Advances and highlights in allergic rhinitis. *Allergy*. 2021;76(11):3383-3389.
23. Ellis AK, Keith PK. Nonallergic rhinitis with eosinophilia syndrome. *Curr Allergy Asthma Rep*. 2006;6(3):215-220.
24. Meng Y, Lou H, Wang Y, et al. Endotypes of chronic rhinitis: a cluster analysis study. *Allergy*. 2019;74(4):720-730.
25. Crobach M, Hermans J, Kaptein A, Ridderikhoff J, Mulder J. Nasal smear eosinophilia for the diagnosis of allergic rhinitis and eosinophilic non-allergic rhinitis. *Scand J Prim Health Care*. 1996;14(2):116-121.
26. Purello-D'Ambrosio F, Isola S, Ricciardi L, Gangemi S, Barresi L, Bagnato GF. A controlled study on the effectiveness of loratadine in combination with flunisolide in the treatment of nonallergic rhinitis with eosinophilia (NARES). *Clin Exp Allergy*. 1999;29(8):1143-1147.
27. Baudoin T, Šimunjak T, Bacan N, Jelavić B, Kuna K, Košec A. Redefining pregnancy-induced rhinitis. *Am J Rhinol Allergy*. 2020;35(3):315-322.
28. Eriksson J, Ekerljung L, Sundblad BM, et al. Cigarette smoking is associated with high prevalence of chronic rhinitis and low prevalence of allergic rhinitis in men. *Allergy*. 2013;68(3):347-354.
29. Chung SJ, Kim BK, Oh JH, et al. Novel tobacco products including electronic cigarette and heated tobacco products increase risk of allergic rhinitis and asthma in adolescents: analysis of Korean youth survey. *Allergy*. 2020;75(7):1640-1648.
30. Storaas T, Steinsvåg SK, Florvaag E, Irgens A, Aasen TB. Occupational rhinitis: diagnostic criteria, relation to lower airway symptoms and IgE sensitization in bakery workers. *Acta Otolaryngol*. 2005;125(11):1211-1217.
31. Gevorgyan A, Segboer C, Gorissen R, van Drunen CM, Fokkens W. Capsaicin for non-allergic rhinitis. *Cochrane Database Syst Rev*. 2015;7:Cd010591.
32. Malmberg H, Grahne B, Holopainen E, Binder E. Ipratropium (Atrovent) in the treatment of vasomotor rhinitis of elderly patients. *Clin Otolaryngol Allied Sci*. 1983;8(4):273-276.
33. Eriksson J, Ekerljung L, Pullerits T, et al. Prevalence of chronic nasal symptoms in West Sweden: risk factors and relation to self-reported allergic rhinitis and lower respiratory symptoms. *Int Arch Allergy Immunol*. 2011;154(2):155-163.
34. Bauchau V, Durham SR. Prevalence and rate of diagnosis of allergic rhinitis in Europe. *Eur Respir J*. 2004;24(5):758-764.
35. Bachert C, van Cauwenberge P, Olbrecht J, van Schoor J. Prevalence, classification and perception of allergic and nonallergic rhinitis in Belgium. *Allergy*. 2006;61(6):693-698.
36. Savouré M, Bousquet J, Burte E, et al. Questionnaire as an alternative of skin prick tests to differentiate allergic from non-allergic rhinitis in epidemiological studies. *Allergy*. 2021;76(7):2291-2294.
37. Jessen M, Janzon L. Prevalence of non-allergic nasal complaints in an urban and a rural population in Sweden. *Allergy*. 1989;44(8):582-587.
38. Pfaar O, Karatzas K, Bastl K, et al. Pollen season is reflected on symptom load for grass and birch pollen-induced allergic rhinitis in different geographic areas-An EAACI Task Force Report. *Allergy*. 2020;75(5):1099-1106.
39. van Rijswijk JB, Blom HM, Fokkens WJ. Idiopathic rhinitis, the ongoing quest. *Allergy*. 2005;60(12):1471-1481.
40. Van Gerven L, Alpizar YA, Steelant B, et al. Enhanced chemosensory sensitivity in patients with idiopathic rhinitis and its reversal by nasal capsaicin treatment. *J Allergy Clin Immunol*. 2017;140(2):437-46.e2.
41. Van Gerven L, Alpizar YA, Wouters MM, et al. Capsaicin treatment reduces nasal hyperreactivity and transient receptor potential cation channel subfamily V, receptor 1 (TRPV1) overexpression in patients with idiopathic rhinitis. *J Allergy Clin Immunol*. 2014;133(5):1332-1399.e1-3.
42. Van Gerven L, Steelant B, Cools L, et al. Low-dose capsaicin (0.01 mM) nasal spray is equally effective as the current standard treatment for idiopathic rhinitis: a randomized, double-blind,

- placebo-controlled trial. *J Allergy Clin Immunol*. 2021;147(1):397-400.e4.
43. Clark DW, Del Signore AG, Raithatha R, Senior BA. Nasal airway obstruction: prevalence and anatomic contributors. *Ear Nose Throat J*. 2018;97(6):173-176.
44. De Corso E, Mastrapasqua RF, Tricarico L, et al. Predisposing factors of rhinitis medicamentosa: what can influence drug discontinuation? *Rhinology*. 2020;58(3):233-240.
45. Patel A, Levi JR, Brook CD. Should excess topical decongestant use raise a red flag? Rhinitis medicamentosa and opioid use disorder. *Ann Otol Rhinol Laryngol*. 2020;129(2):164-169.
46. Mehuys E, Gevaert P, Brusselle G, et al. Self-medication in persistent rhinitis: overuse of decongestants in half of the patients. *J Allergy Clin Immunol Pract*. 2014;2(3):313-319.
47. Hirsch AG, Nordberg C, Bandeen-Roche K, et al. Radiologic sinus inflammation and symptoms of chronic rhinosinusitis in a population-based sample. *Allergy*. 2020;75(4):911-920.
48. Dietz de Loos D, Louijssen ES, Wildeman MAM, et al. Prevalence of chronic rhinosinusitis in the general population based on sinus radiology and symptomatology. *J Allergy Clin Immunol*. 2019;143(3):1207-1214.
49. Ahn JC, Kim JW, Lee CH, Rhee CS. Prevalence and risk factors of chronic rhinosinusitis, allergic rhinitis, and nasal septal deviation: results of the Korean National Health and Nutrition Survey 2008-2012. *JAMA Otolaryngol Head Neck Surg*. 2016;142(2):162-167.
50. Verhoeven S, Schmelzer B. Type and severity of septal deviation are not related with the degree of subjective nasal obstruction. *Rhinology*. 2016;54(4):355-360.
51. Tadalafil. *LiverTox: Clinical and Research Information on Drug-Induced Liver Injury*. National Institute of Diabetes and Digestive and Kidney Diseases; 2012.
52. Bernstein JA. Allergic and mixed rhinitis: epidemiology and natural history. *Allergy Asthma Proc*. 2010;31(5):365-369.

SUPPORTING INFORMATION

Additional supporting information may be found in the online version of the article at the publisher's website.

How to cite this article: Avdeeva KS, Fokkens WJ, Segboer CL, Reitsma S. The prevalence of non-allergic rhinitis phenotypes in the general population: A cross-sectional study. *Allergy*. 2022;00:1-12. doi:[10.1111/all.15223](https://doi.org/10.1111/all.15223)

The questionnaire on nasal and pulmonary complaints

(Translated from Dutch)

1. What is your age? ____years
2. What is your gender? *Male/Female/I do not want to answer this question*
3. Do you have any of the following complaints present on the most days of the week for more than 1 hour per day:
 - a. Runny nose: *Yes/No*
 - b. Mucus in the throat: *Yes/No*
 - c. Blocked nose: *Yes/No*
 - d. Sneezing: *Yes/No*
 - e. Itchy nose or throat: *Yes/No*
 - f. Facial pain/pressure: *Yes/No*
 - g. Reduced smell/taste: *Yes/No*
 - h. Itchy/teary eyes: *Yes/No*
4. Which nasal complaints do you have at the moment? (Please tick all applicable answers):
 - a. Blocked nose
 - b. Runny nose
 - c. Mucus in the throat
 - d. Sneezing
 - e. Itchy nose or throat
 - f. Facial pain or pressure
 - g. Reduced smell/taste
 - h. Itchy or teary eyes
 - i. I do not have any nasal complaints
5. From which nasal complaints do you suffer the most? _____
6. How long do you have these complaints per day: *<1 hour per day / ≥1 hour per day / I do not have nasal complaints*
7. Altogether, on how many days per year do you have these complaints? ____ *days per year*
8. When did your complaints begin? *In* ____ *(year)*
9. We ask you to indicate whether the following statements are correct:
 - a. My nasal complaints (and eye complaints) have an influence on my sleep:
Yes/No/I do not know
 - b. My nasal complaints (and eye complaints) have an influence on my daily activities:
Yes/No/I do not know
 - c. My nasal complaints (and eye complaints) impair me at work or school:
Yes/No/I do not know
 - d. My nasal complaints (and eye complaints) are troublesome:
Yes/No/I do not know
10. On how many days per week do you suffer from your nasal complaints?
Less than 4 days per week/more than 4 days per week
11. If you have nasal complaints, is this a period of:
Less than 4 weeks in a row/4 or more weeks in a row

12. How much do you suffer today from your nasal complaints altogether? The line is 0-100 mm: 0 for absolutely no complaints and 100 is for the worst complaints possible. Please indicate with a vertical line:

Absolutely no complaints	Worst complaints possible	Worst complaints possible
-----------------------------	--	------------------------------

13. What do you think is the reason for your nasal complaints? _____
14. Are there any factors that worsen your nasal complaints? _____
15. How often did you have a cold last year? ___ *times*
16. On how many days in total have you had a cold last year? ___ *days*
17. Do you regularly use otrivin nasal spray or nasal drops (xylomethasoline)? *Yes/No*
18. Do you use any medications for your nose, such as nasal spray (other than otrivin) or tablets?
Yes/No
19. If yes, which medications do you use? _____
20. Do you have hay fever/allergic rhinitis? *Yes/No*
21. Have you ever had a test (skin-prick test or blood test) for inhalational allergens (think about tree pollen, grass pollen, cat, dog, house dust mite, etc.)? *Yes/No*
22. If yes, were you found to be allergic by this test? *Yes/No*
23. If yes, for which allergens? _____
24. Do you have asthma? *Yes/No*
25. If yes, since when? (year) _____
26. Do you have any other lung complaints, such as cough, dyspnea, shortness of breath, or wheezing? *Yes/No*
27. If yes, which lung complaints do you have? *Cough/Dyspnea/Shortness of breath/Wheezing*
28. Do you smoke? *Yes/No*
29. Have you ever smoked? *Yes/No*
30. If yes, how long have you been smoking? *Started in year* ___; *Stopped in year* ___
31. What do you smoke? *Cigarettes/cigars/pipe/other*
32. How many cigarettes on average per day have you been smoking? ___ *per week*
33. How many cigars on average per day have you been smoking? ___ *per week*
34. How many pipes on average per day have you been smoking? ___ *per week*
35. Do you use drugs? *Yes/No*
36. If yes, what kind of drugs do you use? _____
37. Do you ever sniff the drugs in your nose? *Yes/No*
38. If yes, on which side? *Left/Right*
39. Do you work? *Yes/No*
40. What kind of work do you do? _____
41. Do you have nasal complaints at work? *Yes/No*
42. Do your nasal complaints get worse at work? *Yes/No*
43. What do you think is the reason for these nasal complaints? _____
44. Do you have these complaints during vacations or at weekends? *Yes/No*
45. Have you ever had a period at work without these nasal complaints? *Yes/No*
46. When have you started your current job? ___ *month* ___ *year*

47. When did your nasal complaints start? __month__ year
48. Do you have nasal complaints when you are eating? *Yes/No*
49. If yes, with what type of food do you get these symptoms? _____
50. Do you have nasal complaints when you are drinking alcohol? *Yes/No*
51. If yes, with what type of alcohol drinks do you get these symptoms (think about wine, beer, strong drinks)? _____
52. How bad are your nasal complaints when you are exposed to cold dry air (such as in a room with air conditioner or in winter)? *None/A little/Quite a lot/A lot/Extremely a lot*
53. How bad are your pulmonary complaints (cough, dyspnea, shortness of breath or wheezing) when you are exposed to cold dry air (such as in a room with air conditioner or in winter)? *None/A little/Quite a lot/A lot/Extremely a lot*
54. Do you have an acute worsening of nasal complaints when exposed to...
- a. ...temperature changes? *Yes/No*
 - b. ...when smoking or when someone smokes next to you? *Yes/No*
 - c. ...when smelling strong odors such as perfume? *Yes/No*
 - d. ...during exercise or exertion? *Yes/No*
 - e. ...when having emotional stress? *Yes/No*
 - f. ...with changes in humidity? *Yes/No*
55. For women: do you have menstruations? *Yes/No*
56. If yes, do you have complaints such as blocked or runny nose during menstruation? *Yes/No*
57. Are you currently pregnant? *Yes/No*
58. Have you encountered nasal complaints during your pregnancy? *Yes/No*
59. Have you ever been pregnant? *Yes/No*
60. Have you had during the past pregnancy nasal complaints such as blocked or runny nose? *Yes/No*

Attachment 2. Characteristics of the subgroups, relevant for definitions of the phenotypes.

	NAR		Controls
N	364 (27% of the general population)		788
RHINORRHOEA OF THE ELDERLY			
Rhinorrhoea of the elderly defined by	N of participants with RoE	N of participants in the age group	% of the age group
≥ 70 years old	9	108	8%
≥ 65 years old	15	157	10%
≥ 60 years old	21	193	11%
≥ 55 years old	23	232	10%
≥ 50 years old	31	260	12%
OCCUPATIONAL RHINITIS			
N of currently working participants	196 (54%)		407 (52%)
Nasal complaints during work	107 (55% of currently working participants)		31 (8% of currently working participants)
Nasal complaints get worse during work	26 (13% of currently working participants)		18 (4% of currently working participants)
Participants who have nasal complaints at work/experience worsening of their nasal complaints at work			
Improvement of nasal complaints during weekend or vacation	30 (29% of participants who experience nasal complaints at work/worsening of nasal complaints at work)		
Self-reported reasons for work-related nasal complaints:			
Hyperreactivity	23 (21%)		
HDM or other allergies	12 (11%)		
Work-specific triggers	2 (2%)		
Other/unknown	71 (64%)		
RHINITIS MEDICAMENTOSA			
Regular use of nasal decongestants	61 (17%)		42 (5%)
		Regular nasal decongestant users in NAR	Regular nasal decongestant users in Control group
VAS (nasal complaints), mm	Mean ± SD	45 ± 26^	19 ± 24^
1-100	Median (IQR)	51 (24; 65)^	11 (1;26)^
ARIA classification			
	Mild	9 (15%)^	21 (50%)^
	Moderate/severe	51 (84%)^	15 (36%) ^
	Unknown	1 (1%)	6 (12%)
Frequency of nasal decongestant use			
Twice or more times per day	17 (28%)		
Daily	17 (28%)		
Every other day	4 (7%)		
Weekly	23 (38%)		

^a*P*<0.05

SMOKERS' RHINITIS			
Current smokers			
Current smokers, N		30 (8%)	38 (5%)
VAS, current	Mean ± SD	34±28	13±20
smoker, mm	Median (IQR)	25 (8; 59)	6 (0.0001; 14)
Pack-years of	Mean ± SD	32±29	14±14
smoking	Median (IQR)	29(3;57)	12 (2; 20)
Former smokers			
Former smokers, N		168 (46%)	335 (43%)
VAS, current	Mean ± SD	40±27	10±17
smoker, mm	Median (IQR)	36 (17; 61)	2 (0.0001; 13)
Pack-years of	Mean ± SD	13±16	12±16
smoking	Median (IQR)	7 (2; 17)	6 (1; 18)
HORMONAL RHINITIS			
Females, N (%)		178 (47%)	280 (36%)
Females who reported still having menstruation		44 (25% of women)	110 (26%)
Nasal complaints when having a menstruation		10 (23% of women who have menstruation)	5 (5%)
History of pregnancy		120 (67%)	299 (70%)
Nasal complaints during pregnancy		20 (17% of women with a history of pregnancy)	18 (6%)
DRUG-INDUCED RHINITIS			
Medication users (other than nasal, pulmonary or NSAIDs)		163 (45%)	272 (35%)
Beta-blockers		34 (21% of medication users)	46 (16% of medication users)
Sedatives		4 (3%)	2 (1%)
Antidepressants		8 (5%)	9 (3%)
Other medications		147 (90%)	246 (88%)
Painkillers users (NSAIDs and paracetamol)		253 (70%)	503 (64%)
NSAIDs		15 (6% of painkillers users)	27 (5%)
Paracetamol		173 (68%)	362 (72%)
NSAIDs and paracetamol		52 (21%)	79 (16%)
Aspirin as antiplatelet		13 (5%)	24 (5%)
Other/Unknown		4 (2%)	10 (2%)
Nasal or pulmonary complaints due to an ingestion (use) of a medication		6 (1.6%)	11 (1.4%)
Type of medication that caused adverse reaction			
INCS	1		1
Nasal decongestants	2		1
ACE inhibitor	1		1
Chemotherapy	1		0
NSAID	0		2
Prostaglandin F analogue	1		
GUSTATORY RHINITIS			
Nasal complaints when eating		61 (17%)	21 (3%)
Missing data		19 (5%)	113 (14%)

Nasal complaints are present when drinking alcohol	21 (6%)	11 (1%)
<i>Missing</i>	127 (35%)	363 (46%)
Types of food that cause nasal complaints		
Hot or spicy	26 (43% of participants with nasal complaints when eating)	13 (62%)
All kinds of food	13 (21%)	2 (10%)
Nasal complaints are present all the time, also when eating	5 (8%)	1 (5%)
Unknown	16	4 (19%)
IDIOPATHIC RHINITIS		
Any degree of nasal hyperreactivity	252 (69%)	283 (36%)
Degree of nasal hyperreactivity		
<i>Missing</i>	142 (39%)	420 (53%)
	% for participants who have filled in this question	
None	37 (17%)	178 (48%)
A little	92 (41%)	152 (41%)
Some	52 (23%)	26 (7%)
A lot	24 (11%)	8 (2%)
Extremely a lot	17 (8%)	4 (1%)
Degree of bronchial hyperreactivity		
<i>Missing</i>	54 (15%)	154 (20%)
	% for participants who have filled in this question	
None	155 (50%)	470 (74%)
A little	101 (28%)	132 (21%)
Some	36 (10%)	22 (4%)
A lot	10 (3%)	9 (1%)
Extremely a lot	8 (3%)	1 (0.2%)
Presence of nasal complaints with...		
...temperature changes	138 (38%)	102 (13%)
<i>missing</i>	58 (16%)	192 (24%)
...smoking (or when someone around smokes)	79 (22%)	65 (8%)
<i>missing</i>	70 (19%)	212 (27%)
...strong odors	56 (15%)	44 (6%)
<i>missing</i>	72 (20%)	207 (26%)
...exercise or exertion	72 (20%)	39 (5%)
<i>missing</i>	66 (18%)	210 (27%)
...emotional stress	45 (12%)	33 (4%)
<i>missing</i>	75 (21%)	210 (27%)
...humidity changes	121 (33%)	80 (10%)
<i>missing</i>	72 (20%)	206 (26%)