



ORIGINAL ARTICLE

WILEY

Asthma and Rhinitis

Effects of personal air pollution exposure on asthma symptoms, lung function and airway inflammation

L. Chambers¹ | J. Finch¹ | K. Edwards¹ | A. Jeanjean² | R. Leigh² | S. Gonem¹ ¹NIHR Leicester Biomedical Research Centre, Glenfield Hospital, Leicester, UK²Department of Physics and Astronomy, University of Leicester, Leicester, UK

Correspondence

Sherif Gonem, NIHR Biomedical Research Centre (Respiratory), Glenfield Hospital, Leicester, UK.

Email: sg330@le.ac.uk

Funding information

This paper presents independent research funded by the National Institute for Health Research (NIHR). The views expressed are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health. This work was funded by an Academy of Medical Sciences Starter Grant, supported by the Academy of Medical Sciences, Wellcome Trust, Medical Research Council, British Heart Foundation, Arthritis Research UK, Royal College of Physicians and Diabetes UK, and by the Midlands Asthma and Allergy Research Association. SG is funded by an NIHR Clinical Lectureship. LC was supported by a Wolfson Foundation Intercolated Degree Award. The authors thank the University of Leicester and EarthSense System Ltd for funding the development of the satellite dataset used in this study.

Summary

Background: There is evidence that air pollution increases the risk of asthma hospitalizations and healthcare utilization, but the effects on day-to-day asthma control are not fully understood.

Objective: We undertook a prospective single-centre panel study to test the hypothesis that personal air pollution exposure is associated with asthma symptoms, lung function and airway inflammation.

Methods: Thirty-two patients with a clinical diagnosis of asthma were provided with a personal air pollution monitor (Cairclip NO₂/O₃) which was kept on or around their person throughout the 12-week follow-up period. Ambient levels of NO₂ and particulate matter were modelled based upon satellite imaging data. Directly measured ozone, NO₂ and particulate matter levels were obtained from a monitoring station in central Leicester. Participants made daily electronic records of asthma symptoms, peak expiratory flow and exhaled nitric oxide. Spirometry and asthma symptom questionnaires were completed at fortnightly study visits. Data were analysed using linear mixed effects models and cross-correlation.

Results: Cairclip exposure data were of good quality with clear evidence of diurnal variability and a missing data rate of approximately 20%. We were unable to detect consistent relationships between personal air pollution exposure and clinical outcomes in the group as a whole. In an exploratory subgroup analysis, total oxidant exposure was associated with increased daytime symptoms in women but not men.

Conclusions and clinical relevance: We did not find compelling evidence that air pollution exposure impacts on day-to-day clinical control in an unselected asthma population, but further studies are required in larger populations with higher exposure levels. Women may be more susceptible than men to the effects of air pollution, an observation which requires confirmation in future studies.

KEYWORDS

asthma, environment and hygiene hypothesis, epidemiology

1 | INTRODUCTION

Ambient air pollution exposure contributes substantially to the global burden of disease, with particulate matter pollution being the fifth-ranking mortality risk factor world-wide in 2015.¹ The majority of

this excess mortality is attributable to cardiovascular and respiratory conditions such as ischaemic heart disease and chronic obstructive airways disease. Air pollution is mainly derived from the combustion of fossil fuels and comprises a number of distinct chemical constituents. These include gaseous pollutants such as nitrogen oxides

and ozone, and particulate matter which may be further subdivided into fine (aerodynamic diameter $< 2.5 \mu\text{m}$, $\text{PM}_{2.5}$) and coarse (aerodynamic diameter between 2.5 and $10 \mu\text{m}$, PM_{10}).

There is at present limited information available regarding the effects of air pollution exposure on individual patients in a real-life setting. This is primarily due to the difficulty of accurately evaluating personal exposures and correlating these with day-to-day variations in disease control. However, increasing miniaturization of air quality monitors over the past 5 years has now made personal exposure monitoring a realistic prospect.² Furthermore, satellite remote sensing with geospatial modelling of ground-level air pollution is emerging as an important source of exposure data in epidemiological and cohort studies.³

Evidence from epidemiological and experimental studies suggests that air pollution may contribute to exacerbations of asthma through oxidative injury to the airways, increased sensitization to aeroallergens, airway remodelling and airway hyper-responsiveness.⁴ We therefore hypothesized that personal air pollution exposure is associated with day-to-day variations in asthma symptoms, lung function and airway inflammation. To test this hypothesis, we undertook a prospective single-centre panel study, making use of portable sensors and satellite-based geospatial modelling.

2 | METHODS

2.1 | Participants

Thirty-two patients with asthma were recruited from general respiratory and specialist clinics at Glenfield Hospital, Leicester. Patients were seen at baseline in the stable state, with no changes having been made to their regular inhaled or oral asthma therapy within the preceding 6 weeks. All participants were never smokers or ex-smokers with less than 10 pack-years' smoking history. Asthma was diagnosed in a secondary or tertiary care setting according to British Thoracic Society guidelines.⁵ The study was approved by the National Research and Ethics Committee—East Midlands, Leicester, and all participants gave their written informed consent. Participants were recruited and studied on a rolling basis from May 2016 to April 2017.

2.2 | Exposure measurements

Participants were provided with a Cairclip NO_2/O_3 monitor (Cairpol, La Roche-Blanche, France), a cylindrical unit with a weight of 55 g, length of 62 mm and diameter of 32 mm. The unit uses an electrochemical sensor to sample total oxidants (the sum of NO_2 and O_3) in the ambient air once per minute, and stores readings with an accurate date and timestamp. Cairclip NO_2/O_3 has undergone detailed technical validation by the Joint Research Centre of the European Commission⁶ and the United States Environment Protection Agency.⁷ The Cairclip unit was kept within a fabric holder which could be easily attached to a belt or handbag. Participants were instructed to keep the Cairclip on their person when outdoors, and

in close proximity to them when indoors. The battery life of the unit was approximately 24 hours and it therefore required charging overnight using a universal serial bus (USB) cable. The same USB connection was used to extract data from the unit at study visits. The unit was set to store 15-minute average readings so as not to exceed its data storage capacity between the 2-weekly study visits.

Directly measured daily average levels of NO_2 , O_3 and $\text{PM}_{2.5}$ at the primary air quality monitoring station in central Leicester were obtained from the Department for the Environment, Food and Rural Affairs (DEFRA) (<https://uk-air.defra.gov.uk/data/>). Modelled average daily levels of NO_2 , PM_{10} and $\text{PM}_{2.5}$ were produced by the Earth Observation Science Group at the University of Leicester with a spatial resolution of 1 km^2 . This data set was derived from satellite data from the European ENSEMBLE model, which corresponds to the median of seven European air quality models provided by Copernicus Atmosphere.^{8,9} The ENSEMBLE model provides surface levels of daily mean values of NO_2 , $\text{PM}_{2.5}$ and PM_{10} at a spatial resolution of 0.1° (around 7 km). These were interpolated onto a 1 km grid scale based upon DEFRA national pollution climate mapping.¹⁰ Figure 1 shows annual mean NO_2 concentrations in 2016 for the City of Leicester calculated using this methodology. Exposures for each participant were calculated based upon the grid square in which their home address was located.

2.3 | Daily home monitoring

To provide daily time series of clinical outcomes, participants were provided with an electronic symptom diary and peak expiratory flow (PEF) meter (Micro Diary; Carefusion, Basingstoke, UK) and a portable exhaled nitric oxide (FeNO) monitor (NOBreath; Bedfont, Maidstone, UK). Participants used the Micro Diary unit twice per day in the morning and evening to record visual analogue scores for cough, breathlessness and wheeze over the preceding overnight or daytime period. These were recorded on an integer scale from zero (no symptoms) to ten (severe symptoms). Total night-time and daytime symptom scores (on an integer scale of zero to thirty) were calculated as the sum of the three visual analogue scores for the preceding overnight or daytime period. Participants also undertook three PEF measurements in the morning and evening using the same unit, and the best of the three readings was recorded on each occasion.

Two measurements of FeNO were made each morning using the NOBreath unit, at an exhalation rate of 50 mL/s. These were manually entered by patients into the Micro Diary as NOBreath does not have the facility for storing multiple test results. The average of the two daily readings was utilized in statistical analysis. We have previously shown that home FeNO monitoring using NOBreath produces reproducible results which correlate well with blood and sputum eosinophil counts.¹¹

2.4 | Study visit schedule

Participants attended a baseline visit at which clinical and demographic details were recorded and the home monitoring equipment




FIGURE 1 Annual mean NO₂ concentrations in 2016 for the City of Leicester. Raw satellite pixels from the ENSEMBLE model data (0.1 degrees resolution in latitude and longitude) were interpolated on a 1 km grid using spatial information from the DEFRA PCM database.¹⁰ Map data are from Google Maps

was introduced and explained. Participants were then followed up for a total of 12 weeks and attended study visits at 2-week intervals, at which time data were downloaded from the Cairclip and Micro Diary units. Participants also completed the Asthma Control Questionnaire¹² and Asthma Quality of Life Questionnaire,¹³ and underwent spirometry. Spirometry was performed according to American Thoracic Society/European Respiratory Society guidelines¹⁴ with predicted values calculated using Global Lung Initiative reference equations.¹⁵ Short- and long-acting bronchodilators were withheld for 4 and 12 hours, respectively, before study visits.

2.5 | Statistical analysis

Statistical analysis was performed using SPSS 24 (IBM, Armonk, New York, USA). Relationships between exposure and outcome variables were analysed using linear mixed effects models and cross-correlation, with the two techniques giving complementary information.

Linear mixed effects models were used to assess the relationship between linear trends in the exposure and outcome variables over the study period at a group level, with separate models constructed for each clinical outcome. Clinical outcomes were entered as dependent variables in the model, with exposures specified as fixed effects. An autoregressive covariance structure was specified to account for the observed autocorrelation in the clinical time series. For outcomes measured daily (symptom scores, PEF and FeNO), the corresponding exposure variable was the average total oxidant reading on the same day measured using the Cairclip unit. For outcomes measured at study visits, the average total oxidant over the previous

2 weeks was used as the corresponding exposure variable for spirometry and the Asthma Quality of Life Questionnaire, and over the previous week for the Asthma Control Questionnaire.

Cross-correlation was used to assess the relationship between day-to-day variations in exposure and outcome variables in individual participants over a range of lag times (−7 to +7 days). Cross-correlation results were assessed individually, and were combined across the group to give the mean and 95% confidence interval cross-correlation at each lag time. Exploratory subgroup analyses were performed, stratifying the group by sex and smoking history (ex-smokers vs never smokers).

2.6 | Statistical power calculation

The primary analysis was estimation of the effect of average total oxidant exposure on peak expiratory flow across the cohort, analysed using a linear mixed effects model. Calculation of statistical power was performed by Dr. Matthew Richardson (University of Leicester) using a simulation approach, based upon an internal pilot comprising the first six participants in the study. A linear mixed effects model was fitted to the pilot data; the dependent variable was morning PEF (PEF_{am}), and the independent variable was the daily average personal total oxidant concentration (AverageOX). A random intercept for each subject was also included in the model. Estimates for the fixed effect intercept, variance of the random intercept and variance of the error term were obtained from the pilot sample. Values for PEF_{am} were then generated according to the regression equation derived from the pilot data, but allowing the

regression coefficient for AverageOX to vary. The random intercept for each subject was assumed to follow a normal distribution, and a normally distributed error term was included in the regression equation. The *P*-value for the regression coefficient of AverageOX was obtained using 100 simulations, with sample size fixed at *n* = 30. Based on the simulation results, it was estimated that this sample size would be sufficient to detect an effect size of AverageOx (p.p.b.) on PEFam (L/min) of approximately 0.7 units with power = 80% at the 5% significance level.

3 | RESULTS

Thirty-one participants completed the study, with one participant withdrawing at an early stage due to technical problems with the Cairclip unit. Clinical and demographic characteristics of the participants are shown in Table 1. Participants were followed up for an average of 78 days, with a total of 2406 person-days of follow-up. The average missing data rate for daily monitoring were 15.5% for morning readings, 21.7% for evening readings and 20.6% for Cairclip total oxidant data. Missing Cairclip data were usually caused by complete discharge of the internal battery, due to the unit being left off charge overnight. This resulted in loss of the timestamp and necessitated return of the unit to the study centre to be reset. Figure 2 shows an example time series of personal total oxidant exposure in a single participant measured using Cairclip. This shows the typical diurnal pattern of exposure, with peak levels during daylight hours, and considerable day-to-day variability with regard to average daily exposure. Figure 3 shows the average total oxidant levels measured using Cairclip at different times of the day over 1911 person-days of observations. A peak is seen during daytime hours, with additional small peaks corresponding to the morning and evening rush hours on weekdays. Although exposure levels were slightly lower on weekends compared to weekdays, clinical outcomes did not differ significantly across the 7 days of the week, as shown in Table S1.

Table 2 shows the results of linear mixed effects models in which the fixed effects of average daily total oxidant exposure measured using Cairclip on clinical outcomes were estimated over the whole study group. In each case, no clinically important or statistically significant effects were observed. Cross-correlation analysis did not reveal any consistent associations between exposure and outcome variables in the study group as a whole, as shown in Figure 4 and Figures S1-S6. When the group was stratified by sex, we observed a positive correlation on the same day (lag 0) between total oxidant exposure using Cairclip and daytime symptoms in women, but not in men (Figure 5). Stratifying the group by smoking history did not reveal any significant associations.

4 | DISCUSSION

We report the first panel study of the effects of personal air pollution exposure on adults with asthma in which exposures and

TABLE 1 Clinical and demographic characteristics

Age (y)	55.2 (13.1)
Sex	Female <i>n</i> = 15 (48.4%) Male <i>n</i> = 16 (51.6%)
Body mass index (kg/m ²)	30.4 (6.7)
Positive atopic status ^a (<i>n</i> [%])	24 (77.4)
Duration of asthma (y)	32.2 (21.5)
Age of onset of asthma (y)	23.0 (21.4)
Number of asthma exacerbations in the previous year	2.4 (4.5)
Regular inhaled corticosteroid use (<i>n</i> [%])	29 (93.5)
Inhaled corticosteroid dose (beclometasone dipropionate equivalent [μg])	1000 (400-1600)
Regular oral prednisolone use (<i>n</i> [%])	7 (22.6)
Leukotriene receptor antagonist use (<i>n</i> [%])	7 (22.6)
Oral theophylline use (<i>n</i> [%])	3 (9.7)
Long-acting β ₂ agonist use (<i>n</i> [%])	21 (67.7)
Asthma Control Questionnaire score	1.41 (1.15)
Asthma Quality of Life Questionnaire score	5.32 (1.19)
Forced expiratory volume in one-second (% predicted)	78.6 (22.5)
Forced vital capacity (% predicted)	91.1 (17.3)
Ratio of forced expiratory volume in one-second to forced vital capacity (%)	67 (12)
Exhaled nitric oxide at a flow rate of 50 mL/s (p.p.b.)	28.1 (22.3)

Data are *n* (%), mean (standard deviation) or median (interquartile range) unless specified otherwise.

^aDefined as a positive skin test for any of a panel of specified aeroallergens (grass pollen, tree pollen [alder, silver birch, hazel], moulds [*Aspergillus fumigatus*, *Alternaria tenius*, *Cladosporium*, *Penicillium notatum*], cat fur, dog dander, and house dust mite [*Dermatophagoides pteronyssinus*]).




FIGURE 2 Personal total oxidant exposure measured using a Cairclip NO₂/O₃ unit in an individual with asthma

outcomes were assessed concurrently over an extended 12-week period. We assessed air pollution exposure using three separate methodologies, namely (i) a small portable sensor carried on the




FIGURE 3 Diurnal variation in total oxidant levels measured using Cairclip. Average total oxidant levels are shown during the 24-h daily cycle over 1911 person-days of observations. Results are presented for all days, weekdays (Monday to Friday), and weekends (Saturday and Sunday)

TABLE 2 Linear mixed models estimating fixed effects of total oxidant exposure on daily clinical outcomes

Outcome measure (units/range)	Estimate of fixed effects ^a	95% Confidence interval		P-value
		Lower bound	Upper bound	
Morning PEF (L/min)	0.038	-0.416	0.491	.870
Evening PEF (L/min)	-0.088	-0.576	0.400	.723
FeNO (p.p.b.)	0.012	-0.138	0.162	.875
Night-time symptoms (0-30)	0.0036	-0.0182	0.0253	.746
Daytime symptoms (0-30)	0.0022	-0.0246	0.0290	.872
ACQ (0-6)	0.0032	-0.0250	0.0314	.822
AQLQ (1-7)	-0.0246	-0.0613	0.0121	.187
Pre-bronchodilator FEV ₁ (% predicted)	-0.213	-0.766	0.340	.447
Post-bronchodilator FEV ₁ (% predicted)	-0.140	-0.586	0.306	.535
Pre-bronchodilator FVC (% predicted)	0.261	-0.316	0.839	.372
Post-bronchodilator FVC (% predicted)	0.083	-0.590	0.757	.807
Pre-bronchodilator FEV ₁ /FVC (% predicted)	-0.0017	-0.0054	0.0021	.385
Post-bronchodilator FEV ₁ /FVC (% predicted)	0.0008	-0.0023	0.0038	.618

ACQ, Asthma Control Questionnaire score; AQLQ, Asthma Quality of Life Questionnaire score; FeNO, exhaled nitric oxide measured at an expiratory flow rate of 50 mL/s; FEV₁, forced expiratory volume in one second; PEF, peak expiratory flow rate.

^aExposure variable was measured in parts per billion using Cairclip NO₂/O₃ unit.

person (Cairclip NO₂/O₃), (ii) modelled data based on satellite imaging and (iii) directly measured ambient levels at a central monitoring station. We measured a comprehensive panel of daily clinical outcomes, including asthma symptom scores, peak expiratory flow and exhaled nitric oxide. Most previous longitudinal studies of air pollution effects on asthma have relied solely on central monitoring stations to estimate the exposure of an entire community. A small number of studies have utilized personal air pollution sensors,¹⁶⁻¹⁹ but due to the bulky nature of the monitoring equipment, the follow-up duration has necessarily been limited. We have for the first time demonstrated the feasibility of long-term personal air pollution exposure monitoring.

We did not observe strong associations between exposures and clinical outcomes, although an exploratory subgroup analysis suggested the possibility of increased daytime symptoms in women exposed to air pollution. A number of possible explanations may be suggested to explain our findings. The participants in our study were recruited from Leicester and the surrounding area. To increase the generalizability of our results, we did not restrict participation to individuals residing in heavily polluted districts. Therefore, a number of study participants lived in rural areas and may not have had sufficient air pollution exposure to impact on their asthma control. Indeed, average peak daily exposure to total oxidant measured using Cairclip monitors was approximately 10 p.p.b. in our study cohort, which is well within United Kingdom and European air quality limits.²⁰ Future studies should therefore focus more specifically on high-risk populations, such as those living and working in inner-city areas. Moreover, our data collection and statistical methods relied on detecting associations between exposure and outcome variables produced by day-to-day variability and linear trends arising naturally. We did not impose an experimental change in exposure, as for instance was performed by McCreanor et al who measured changes in forced expiratory volume in one second caused by walking down a busy London street (Oxford Street) compared to a nearby park (Hyde Park).²¹ Therefore, the lack of observed associations could also have been due to low levels of day-to-day variability in exposures and/or clinical outcomes.

With 31 participants, we had sufficient statistical power to detect a 21 L/min change in peak expiratory flow in association with a 30 p.p.b. change in total oxidant exposure. It is possible that a true effect existed but that it was too small to detect with our sample size. Indeed, previous cohort studies have often reported smaller effect sizes than this.²² However, it has been shown that changes in peak expiratory flow of less than 20 L/min are unlikely to result in perceptibly increased symptoms.²³ Therefore, we considered that our sample size was sufficient to detect clinically important effects on lung function.

We endeavoured to reduce the missing data rate by seeing participants frequently at the study centre (every 2 weeks) and providing them with contact details of investigators who could quickly resolve technical issues with the home monitoring equipment. Nevertheless, approximately 15%-20% of daily exposure or outcome




FIGURE 4 Group-level cross-correlations between total oxidant levels measured using Cairclip and clinical outcomes. Group-level cross-correlations are shown between total oxidant levels measured using Cairclip and clinical outcomes. The group mean and 95% confidence interval of the cross-correlation are shown for each lag time (d). A lag of zero refers to effects occurring on the same day; positive lags refer to changes in outcome preceded by changes in exposure; negative lags refer to changes in exposure preceded by changes in outcome. PEF = peak expiratory flow; FeNO = fractional exhaled nitric oxide




FIGURE 5 Cross-correlation between total oxidant levels measured using Cairclip and daytime symptoms stratified by sex. Cross-correlations are shown between total oxidant levels measured using Cairclip and daytime symptoms, in female and male participants (Panels A and B, respectively). The group mean and 95% confidence interval of the cross-correlation are shown for each lag time (d). A lag of zero refers to effects occurring on the same day; positive lags refer to changes in outcome preceded by changes in exposure; negative lags refer to changes in exposure preceded by changes in outcome

data were missing, and this may have reduced the chance of observing positive associations.

In this study, we particularly investigated the effects of air pollution exposure on day-to-day variations in asthma symptoms, lung function and airway inflammation. It is possible that the major effects of air pollution on asthma relate to other aspects of the condition which our study was not designed to measure, such as exacerbations, hospital admissions and lung function decline. Indeed, a recent meta-analysis of 87 epidemiological studies conducted in multiple cities world-wide concluded that short-term exposure to a number of air pollutants results in significantly increased risks of asthma-related hospital admissions.²⁴ It has been reported in a

recent large population-based cohort study that cumulative air pollution exposure is a significant risk factor for progression from asthma to chronic obstructive pulmonary disease.²⁵ While significant at a population level, these effects may be too subtle to detect in small panel studies.

Susceptibility to the effects of air pollution varies between individuals and is most likely related to genetic factors.²⁶ Therefore, it is possible that clinically significant effects of air pollution may be confined to a small but important group of individuals with a combination of biological susceptibility and higher-than-average levels of exposure. Future research should seek to detect individuals at particular risk from air pollution and to develop strategies to mitigate this

risk. Our data suggest that women may be more susceptible to the effects of air pollution than men, an observation which is shared with a number of previous studies,²⁷ and which requires further investigation.

We chose to use total oxidant levels measured using the Cairclip NO₂/O₃ monitor as our primary exposure variable. This device was chosen due to a combination of portability, relatively low cost and good measurement properties (precision, linearity and limits of detection) under laboratory and field conditions.^{6,7,28} However, these validation studies were performed with the sensor in the static state, and it is known that movement of air pollution sensors can have a significant effect on the measurements.²⁹ A further consideration for mobile monitoring is the sensor response time, which although relatively fast at between 1.5 and 4 minutes for Cairclip^{6,7} may still not be sufficient to detect very transient increases in air pollution exposure. Therefore, sensors used in future personal exposure studies should first undergo validation under realistic field conditions.³⁰ Jaio et al found that Cairclip NO₂/O₃ accurately measured the sum of NO₂ and O₃ (total oxidants) when compared against reference sensors.³¹ However, when an estimate of NO₂ was derived by subtracting reference values of O₃ from Cairclip total oxidant measurements, these did not correlate with reference values of NO₂. This suggests that Cairclip NO₂/O₃ measurements are not entirely additive and cannot be used to reliably measure the individual NO₂ and O₃ components. This is a potential limitation of the study, as it is possible that measuring NO₂ or ozone alone would have resulted in greater correlation with clinical outcomes. However, Cairclip NO₂ was still in development when our study commenced and is less well validated than Cairclip NO₂/O₃.²⁸ There are now a number of sensors in development or commercially available, measuring either particulate matter³² or gaseous pollutants,^{33,34} which may be used in future personal exposure studies. It is likely that the measurement properties of these sensors will continue to improve as technology develops.

The current challenges and opportunities in the use of sensors to estimate environmental exposures have been recently summarized by Loh et al.² While small portable air pollution monitors have not yet been fully validated in the ambulatory setting, an alternative approach is to collect location and activity data using global positioning system-enabled smartphone applications, and to link these to ambient levels of air pollution measured using static ground-level monitors or satellite imaging. Furthermore, personal exposure to air pollution is dependent not only on ambient levels but also on the activity being undertaken and the rate of ventilation, which can also be estimated using wearable technology.

In conclusion, we were unable to find compelling evidence that air pollution exposure impacts on day-to-day clinical control in an unselected asthma population, but these results should be taken in the context of the relatively small study population and the low levels of exposure experienced. Future studies should focus on selected individuals who are highly exposed to air pollution due to their lifestyle or place of residence. Our results suggest the possibility of increased susceptibility to air pollution in women compared to

men, but this observation requires confirmation in further prospective studies. We have for the first time demonstrated the feasibility of long-term personal exposure monitoring. Sensor technology and geospatial data processing is developing rapidly and will in the near future enable us to further understand the relationship between health and the environment in a variety of long-term medical conditions.

ACKNOWLEDGEMENT

The authors gratefully acknowledge Dr. Matthew Richardson (University of Leicester) who performed the statistical power calculation for the study.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ORCID

S. Gonem  <http://orcid.org/0000-0002-6080-2246>

REFERENCES

1. Cohen AJ, Brauer M, Burnett R, et al. Estimates and 25-year trends of the global burden of disease attributable to ambient air pollution: an analysis of data from the Global Burden of Diseases Study 2015. *Lancet*. 2017;389:1907-1918.
2. Loh M, Sarigiannis D, Gotti A, et al. How sensors might help define the external exposome. *Int J Environ Res Public Health*. 2017;14:434.
3. Sorek-Hamer M, Just AC, Kloog I. Satellite remote sensing in epidemiological studies. *Curr Opin Pediatr*. 2016;28:228-234.
4. Guarnieri M, Balmes JR. Outdoor air pollution and asthma. *Lancet*. 2014;383:1581-1592.
5. British Thoracic Society/Scottish Intercollegiate Guidelines Network; 2016. British Guideline on the Management of Asthma. <https://www.brit-thoracic.org.uk/standards-of-care/guidelines/btsgn-british-guideline-on-the-management-of-asthma/>. Accessed December 1, 2017.
6. Joint Research Centre; 2013. Report of laboratory and in of micro-sensor for monitoring ambient air pollution: CairClipO3/NO2 of CAIRPOL. <http://publications.jrc.ec.europa.eu/repository/>. Accessed December 1, 2017.
7. Environmental Protection Agency; 2014. Washington, DC. Sensor Evaluation Report; Report EPA/600/R-14/143. <https://cfpub.epa.gov/si/>. Accessed December 1, 2017.
8. Marécal V, Peuch VH, Andersson C, et al. A regional air quality forecasting system over Europe: the MACC-II daily ensemble production. *Geosci Model Dev*. 2015;8:2777-2813.
9. Copernicus Atmosphere; 2016. Final Report MACC-III (Monitoring Atmospheric Composition and Climate 3). <http://atmosphere.copernicus.eu/reports>. Accessed December 1, 2017.
10. Department for the Environment, Food and Rural Affairs; 2015. 1 km DEFRA national pollution climate mapping for NO₂, PM₁₀ and PM_{2.5}. <https://uk-air.defra.gov.uk/data/pcm-data>. Accessed December 1, 2017.
11. Nanda CR, Singapuri A, Soares M, Monteiro W, Siddiqui S, Gonem S. Domiciliary exhaled nitric oxide and eosinophilic airway inflammation in adults with asthma. *Eur Respir J*. 2016;48:242-244.

12. Juniper EF, O'Byrne PM, Guyatt GH, Ferrie PJ, King DR. Development and validation of a questionnaire to measure asthma control. *Eur Respir J*. 1999;14:902-907.
13. Juniper EF, Buist AS, Cox FM, Ferrie PJ, King DR. Validation of a standardized version of the Asthma Quality of Life Questionnaire. *Chest*. 1999;115:1265-1270.
14. Miller MR, Hankinson J, Brusasco V, et al. Standardisation of spirometry. *Eur Respir J*. 2005;26:319-338.
15. Quanjer PH, Stanojevic S, Cole TJ, et al. Multi-ethnic reference values for spirometry for the 3-95 year age range: the global lung function 2012 equations. *Eur Respir J*. 2012;40:1324-1343.
16. Delfino RJ, Staimer N, Gillen D, et al. Personal and ambient air pollution is associated with increased exhaled nitric oxide in children with asthma. *Environ Health Perspect*. 2006;114:1736-1743.
17. Delfino RJ, Staimer N, Tjoa T, et al. Personal and ambient air pollution exposures and lung function decrements in children with asthma. *Environ Health Perspect*. 2008;116:550-558.
18. Spira-Cohen A, Chen LC, Kendall M, Lall R, Thurston GD. Personal exposures to traffic-related air pollution and acute respiratory health among Bronx schoolchildren with asthma. *Environ Health Perspect*. 2011;119:559-565.
19. Smargiassi A, Goldberg MS, Wheeler AJ, et al. Associations between personal exposure to air pollutants and lung function tests and cardiovascular indices among children with asthma living near an industrial complex and petroleum refineries. *Environ Res*. 2014;132:38-45.
20. Department for the Environment, Food and Rural Affairs; 2017. National air quality objectives. <https://uk-air.defra.gov.uk/air-pollution/uk-eu-limits>. Accessed December 1, 2017.
21. McCreanor J, Cullinan P, Nieuwenhuijsen MJ, et al. Respiratory effects of exposure to diesel traffic in persons with asthma. *N Engl J Med*. 2007;357:2348-2358.
22. United States Environmental Protection Agency; 2013. Integrated Science Assessment for Ozone and Related Photochemical Oxidants. <https://www.epa.gov/isa/integrated-science-assessment-isa-ozone-and-related-photochemical-oxidants>. Accessed December 1, 2017.
23. Santanello NC, Zhang J, Seidenberg B, Reiss TF, Barber BL. What are minimal important changes for asthma measures in a clinical trial? *Eur Respir J*. 1999;14:23-27.
24. Zheng XY, Ding H, Jiang LN, et al. Association between air pollutants and asthma emergency room visits and hospital admissions in time series studies: a systematic review and meta-analysis. *PLoS ONE*. 2015;10:e0138146.
25. To T, Zhu J, Larsen K, et al. Progression from asthma to chronic obstructive pulmonary disease. Is air pollution a risk factor? *Am J Respir Crit Care Med*. 2016;194:429-438.
26. Rava M, Smit LA, Nadif R. Gene-environment interactions in the study of asthma in the postgenomewide association studies era. *Curr Opin Allergy Clin Immunol*. 2015;15:70-78.
27. Clougherty JE. A growing role for gender analysis in air pollution epidemiology. *Environ Health Perspect*. 2010;118:167-176.
28. Duvall RM, Long RW, Beaver MR, Kronmiller KG, Wheeler ML, Szykman JJ. Performance evaluation and community application of low-cost sensors for ozone and nitrogen dioxide. *Sensors (Basel)*. 2016;16:pii: E1698.
29. Lerner U, Yacobi T, Levy I, Moltchanov SA, Cole-Hunter T, Fishbain B. The effect of ego-motion on environmental monitoring. *Sci Total Environ*. 2015;533:8-16.
30. Jerrett M, Donaire-Gonzalez D, Popoola O, et al. Validating novel air pollution sensors to improve exposure estimates for epidemiological analyses and citizen science. *Environ Res*. 2017;158:286-294.
31. Jiao W, Hagler G, Williams R, et al. Community Air Sensor Network (CAIRSENSE) project: evaluation of low-cost sensor performance in a suburban environment in the southeastern United States. *Atmos Meas Tech*. 2016;9:5281-5292.
32. Koehler KA, Peters TM. New methods for personal exposure monitoring for airborne particles. *Curr Environ Health Rep*. 2015;2:399-411.
33. McKercher GR, Salmond JA, Vanos JK. Characteristics and applications of small, portable gaseous air pollution monitors. *Environ Pollut*. 2017;223:102-110.
34. Peterson PJD, Aujla A, Grant KH, et al. Practical use of metal oxide semiconductor gas sensors for measuring nitrogen dioxide and ozone in urban environments. *Sensors (Basel)*. 2017;17:pii: E1653.

SUPPORTING INFORMATION

Additional Supporting Information may be found online in the supporting information tab for this article.

How to cite this article: Chambers L, Finch J, Edwards K, Jeanjean A, Leigh R, Gonem S. Effects of personal air pollution exposure on asthma symptoms, lung function and airway inflammation. *Clin Exp Allergy*. 2018;48:798-805. <https://doi.org/10.1111/cea.13130>

Figure S1: Group-level cross-correlations between directly measured ambient nitrogen dioxide and clinical outcomes




Figure S2: Group-level cross-correlations between directly measured ambient ozone and clinical outcomes




Figure S3: Group-level cross-correlations between directly measured ambient PM_{2.5} and clinical outcomes




Figure S4: Group-level cross-correlations between modelled ambient nitrogen dioxide and clinical outcomes




Figure S5: Group-level cross-correlations between modelled ambient PM₁₀ and clinical outcomes




Figure S6: Group-level cross-correlations between modelled ambient PM_{2.5} and clinical outcomes




Table S1: Mean values of clinical outcome measures by days of the week

	Mon	Tue	Wed	Thu	Fri	Sat	Sun
PEFam (L/min)	362	366	368	369	368	364	366
PEFpm (L/min)	376	376	374	378	377	370	369
PEFpm-am (L/min)	12	9	7	7	9	5	9
FeNO (ppb)	24	24	24	25	25	23	24
Night-time symptoms (0 – 30)	4.4	4.5	4.3	4.2	4.2	4.4	4.6
Daytime symptoms (0 – 30)	5.2	5.3	5.0	5.0	5.1	5.2	5.2

Mean values for each clinical outcome are shown for all observation days across the study cohort, split by day of the week.

PEFam/pm = morning / evening peak expiratory flow; PEFpm-am = evening peak expiratory flow minus morning peak expiratory flow; FeNO = exhaled nitric oxide.