

Macular-type cutaneous adverse reaction due to atezolizumab and pembrolizumab


Hitomi Sugino MD, Yu Sawada MD, PhD , Motonobu Nakamura MD, PhD

First published: 29 November 2021

<https://doi-org.bibliosan.clas.cineca.it/10.1002/cia2.12206>

Abstract

Anti-PD-1/PD-L1 targeted therapy is widely used for the treatment of various intractable malignancies and develops the therapeutic option in the frontier fields of oncology treatment. On the contrary, these immune checkpoint therapies sometimes develop undesirable adverse reaction as the results of activation of immune reaction to various organs. Herein, we report a case of cutaneous adverse reactions during PD-1/PD-L1 targeted treatment in both atezolizumab and pembrolizumab in a patient with lung cancer.



Dear Editor,

Anti-PD-1-/PD-L1-targeted therapy is widely used for the treatment of various intractable malignancies and develops the therapeutic option in the frontier fields of oncology treatment.^{1,2} On the contrary, these immune checkpoint therapies sometimes develop an

undesirable adverse reaction as a result of activation of immune reaction to various organs.^{3, 4} A cutaneous adverse reaction is a representative immune checkpoint treatment-mediated reaction.³ Herein, we report a case of cutaneous adverse reactions during PD-1/PD-L1-targeted treatment in both atezolizumab and pembrolizumab in a patient with lung cancer.

A 68-year-old man noticed erythematous plaques and papules on his trunk and extremities 4 days after first atezolizumab administration for the treatment of his right lung adenocarcinoma clinical stage IV, and his skin eruption gradually developed. On physical examination, erythematous plaques were localized in trunk and extremities without the involvement of the mucosal membrane (Figure 1A). A skin biopsy revealed liquefaction and spongiosis in the epidermis and lymphocyte infiltration in the dermis (Figure 1B). After the treatment with an antihistamine agent, topical application of betamethasone dipropionate, and the discontinuation of additional administration of atezolizumab, his skin eruption was gradually improved without recurrence of skin eruption. As an alternative treatment for his lung cancer, pembrolizumab was next administrated; however, he again recognized erythematous plaques on trunk and extremities after the first administration of pembrolizumab (Figure 1C). The second time cutaneous adverse reaction was more widely spread in his whole body. A skin biopsy again revealed that spongiosis and inflammatory infiltration in the interface of epidermis and dermis (Figure 1D). After the antihistamine agent, topical application of betamethasone dipropionate, and the discontinuation of additional pembrolizumab administration, his skin eruption was improved in several days.




FIGURE 1

[Open in figure viewer](#) | [PowerPoint](#)

Clinical manifestations and histological examinations. (A) The clinical manifestation of cutaneous adverse reaction

during atezolizumab treatment and (B) H&E histological examination (×10). (C) The clinical manifestation of cutaneous adverse reaction during pembrolizumab administration and (D) H&E histological examination (×10)

One of the unique characteristics of this case was to cause similar cutaneous adverse reactions by the different action point targeted treatment, PD-1 and PD-L1. Although the detailed molecular mechanism remains unclear in our case, we thought 3 possible mechanisms to cause cutaneous adverse reaction by both atezolizumab and pembrolizumab. The first reason is that atezolizumab and pembrolizumab have same additives in the drug, such as L-histidine, white soft sugar, and polysorbate. In particular, polysorbate includes in various drugs and influenza vaccine and is known to cause allergic reaction in some cases.^{5, 6} Therefore, we speculated that cutaneous allergic reaction to polysorbate might be enhanced by atezolizumab and pembrolizumab. The second reason is another drug contribution to cause adverse reaction. However, he ordinarily continued to intake acetaminophen without cutaneous adverse reaction. The third reason is atezolizumab and pembrolizumab might become a trigger to cause a cutaneous autoimmune-like reaction. Because skin eruption is relatively mild form and is improved by topical corticosteroid application and oral antihistamine agent, it seemed to be easy to recover his skin eruption even if his skin eruption is an autoimmunity-related skin eruption. Taken together, clinicians should keep in mind that adverse reaction might occur after switching into another anti-PD-1-/PD-L1-related inhibitor.

1 DECLARATION SECTION

Approval of the research protocol: No human participant was involved in this study.

Informed Consent: N/A.

Registry and the Registration No. of the study/trial: N/A.

Animal Studies: N/A.

CONFLICTS OF INTEREST

The authors declare no conflicts of interest.

REFERENCES



1 Janjigian YY, Shitara K, Moehler M, Garrido M, Salman P, Shen L, et al. First-line nivolumab plus chemotherapy versus chemotherapy alone for advanced gastric, gastro-oesophageal junction, and oesophageal adenocarcinoma (CheckMate 649): a randomised, open-label, phase 3 trial. *Lancet*. 2021; **398**: 27–40.

[Crossref](#) | [CAS](#) | [PubMed](#) | [Web of Science®](#) | [Google Scholar](#)

2 Pires da Silva I, Ahmed T, Reijers ILM, Weppler AM, Betof Warner A, Patrinely JR, et al. Ipilimumab alone or ipilimumab plus anti-PD-1 therapy in patients with metastatic melanoma resistant to anti-PD-(L)1 monotherapy: a multicentre, retrospective, cohort study. *Lancet Oncol.* 2021; **22**(6): 836– 47.

[Crossref](#) | [CAS](#) | [PubMed](#) | [Web of Science®](#) | [Google Scholar](#)

3 Brunot A, Grob JJ, Jeudy G, Grange F, Guillot B, Kramkimel N, et al. Association of anti-programmed cell death 1 antibody treatment with risk of recurrence of toxic effects after immune-related adverse events of ipilimumab in patients with metastatic melanoma. *JAMA Dermatol.* 2020; **156**(9): 982– 6.

[Crossref](#) | [PubMed](#) | [Web of Science®](#) | [Google Scholar](#)

4 Oda T, Sawada Y, Okada E, Yamaguchi T, Ohmori S, Haruyama S, et al. Hypopituitarism and hypothyroidism following atrioventricular block during nivolumab treatment. *J Dermatol.* 2017; **44**(6): e144– e5.

[Wiley Online Library](#) | [PubMed](#) | [Web of Science®](#) | [Google Scholar](#)

5 Palacios Castaño MI, Venturini Díaz M, Lobera Labairu T, González Mahave I, Del Pozo Gil MD, Blasco SA. Anaphylaxis due to the excipient polysorbate 80. *J Investig Allergol Clin Immunol.* 2016; **26**(6): 394– 6.

[Crossref](#) | [CAS](#) | [PubMed](#) | [Web of Science®](#) | [Google Scholar](#)

6 Shelley WB, Talanin N, Shelley ED. Polysorbate 80 hypersensitivity. *Lancet.* 1995; **345**(8960): 1312– 3.

[Crossref](#) | [CAS](#) | [PubMed](#) | [Web of Science®](#) | [Google Scholar](#)

[Download PDF](#)

About Wiley Online Library

Privacy Policy

Terms of Use

Cookies

Accessibility

Publishing Policies

Help & Support

Contact Us

Training and Support

DMCA & Reporting Piracy

Opportunities

Subscription Agents
Advertisers & Corporate Partners

Connect with Wiley

The Wiley Network
Wiley Press Room

Copyright © 1999-2021 John Wiley & Sons, Inc. All rights reserved