



Original article

Fish and seafood consumption during pregnancy and the risk of asthma and allergic rhinitis in childhood: a pooled analysis of 18 European and US birth cohorts

Nikos Stratakis,^{1,2} Theano Roumeliotaki,¹ Emily Oken,³ Ferran Ballester,^{4,5} Henrique Barros,^{6,7} Mikel Basterrechea,^{4,8} Sylvaine Cordier,⁹ Renate de Groot,^{2,10} Herman T. den Dekker,^{11,12} Liesbeth Duijts,^{11,12} Merete Eggesbø,¹³ Maria Pia Fantini,¹⁴ Francesco Forastiere,¹⁵ Ulrike Gehring,¹⁶ Marij Gielen,² Davide Gori,¹⁴ Eva Govarts,¹⁷ Hazel M. Inskip,¹⁸ Nina Iszatt,¹³ Maria Jansen,^{19,20} Cecily Kelleher,²¹ John Mehegan,²¹ Carolina Moltó-Puigmartí,¹⁹ Monique Mommers,¹⁹ Andreia Oliveira,^{6,7} Sjurður F. Olsen,^{22,23} Fabienne Pelé,⁹ Costanza Pizzi,²⁴ Daniela Porta,¹⁵ Lorenzo Richiardi,²⁴ Sheryl L. Rifas-Shiman,³ Sian M. Robinson,^{18,25} Greet Schoeters,^{17,26,27} Marin Strøm,^{22,28} Jordi Sunyer,^{4,29,30} Carel Thijs,¹⁹ Martine Vrijheid,^{4,29,30} Tanja G.M. Vrijkotte,³¹ Alet H. Wijga,³² Manolis Kogevinas,^{4,29,33} Maurice P. Zeegers^{2,19} and Leda Chatzi^{1,2*}

¹Department of Social Medicine, University of Crete, Heraklion, Greece, ²NUTRIM School of Nutrition and Translational Research in Metabolism, Maastricht University, Maastricht, The Netherlands, ³Obesity Prevention Program, Department of Population Medicine, Harvard Medical School and Harvard Pilgrim Health Care Institute, Boston, MA, USA, ⁴Spanish Consortium for Research in Epidemiology and Public Health (CIBERESP), Madrid, Spain, ⁵FISABIO–Universitat Jaume I–Universitat de València Joint Research Unit of Epidemiology and Environmental Health, Valencia, Spain, ⁶Department of Clinical Epidemiology, University of Porto Medical School, Porto, Portugal, ⁷EPIUnit – Institute of Public Health, University of Porto, Porto, Portugal, ⁸Public Health Division of Gipuzkoa, Basque Government, San Sebastián, Spain, ⁹Epidemiological Research in Environment, Reproduction and Child Development, University of Rennes I, Rennes, France, ¹⁰Welten Institute, Open University of the Netherlands, Heerlen, The Netherlands, ¹¹Generation R Study Group, Erasmus MC, University Medical Centre Rotterdam, Rotterdam, The Netherlands, ¹²Department of Pediatrics, Erasmus MC, University Medical Centre Rotterdam, Rotterdam, The Netherlands, ¹³Norwegian Institute of Public Health, Oslo, Norway, ¹⁴Department of Biomedical and Neuromotor Sciences (DIBINEM), University of Bologna, Bologna, Italy, ¹⁵Department of Epidemiology, Lazio Regional Health System, Rome, Italy, ¹⁶Institute for Risk Assessment Sciences, Utrecht University, Utrecht, The Netherlands, ¹⁷Environmental Risk and Health, Flemish Institute for Technological Research (VITO), Mol, Belgium, ¹⁸Medical Research Council (MRC) Lifecourse Epidemiology Unit, University of Southampton, Southampton, UK, ¹⁹CAPHRI School for Public Health and Primary Care, Maastricht University, Maastricht, The Netherlands, ²⁰Academic Collaborative Center for Public Health Limburg, Public

Health Services, Geleen, The Netherlands, ²¹Physiotherapy and Sports Science, University College Dublin, Dublin, Ireland, ²²Centre for Fetal Programming, Statens Serum Institut, Copenhagen, Denmark, ²³Department of Nutrition, Harvard T.H. Chan School of Public Health, Boston, MA, USA, ²⁴Cancer Epidemiology Unit, University of Turin and CPO-Piemonte, Turin, Italy, ²⁵National Institute for Health Research (NIHR), University of Southampton and University Hospital, Southampton, UK, ²⁶Department of Biomedical Sciences, University of Antwerp, Antwerp, Belgium, ²⁷Department of Environmental Medicine, University of Southern Denmark, Odense, Denmark, ²⁸Faculty of Natural and Health Sciences, University of the Faroe Islands, Tórshavn, Faroe Islands, ²⁹ISGlobal, Centre for Research in Environmental Epidemiology (CREAL), Barcelona, Spain, ³⁰Department of Preventive Medicine and Public Health, Pompeu Fabra University, Barcelona, Spain, ³¹Department of Public Health, University of Amsterdam, Amsterdam, The Netherlands, ³²Centre for Nutrition, Prevention and Health Services, National Institute for Public Health and the Environment, Bilthoven, The Netherlands and ³³IMIM Hospital del Mar Medicine Research Institute, Barcelona, Spain

*Corresponding author. Department of Social Medicine, Faculty of Medicine, University of Crete, PO Box 2208, Heraklion, 71003, Crete, Greece. E-mail: lchatzi@uoc.gr

Accepted 9 January 2017

Abstract

Background: It has been suggested that prenatal exposure to n-3 long-chain fatty acids protects against asthma and other allergy-related diseases later in childhood. The extent to which fish intake in pregnancy protects against child asthma and rhinitis symptoms remains unclear. We aimed to assess whether fish and seafood consumption in pregnancy is associated with childhood wheeze, asthma and allergic rhinitis.

Methods: We pooled individual data from 60 774 mother-child pairs participating in 18 European and US birth cohort studies. Information on wheeze, asthma and allergic rhinitis prevalence was collected using validated questionnaires. The time periods of interest were: infancy (0-2 years), preschool age (3-4 years), and school age (5-8 years). We used multivariable generalized models to assess associations of fish and seafood (other than fish) consumption during pregnancy with child respiratory outcomes in cohort-specific analyses, with subsequent random-effects meta-analyses.

Results: The median fish consumption during pregnancy ranged from 0.44 times/week in The Netherlands to 4.46 times/week in Spain. Maternal fish intake during pregnancy was not associated with offspring wheeze symptoms in any age group nor with the risk of child asthma [adjusted meta-analysis relative risk (RR) per 1-time/week = 1.01, 95% confidence interval 0.97-1.05] and allergic rhinitis at school age (RR = 1.01, 0.99-1.03). These results were consistently found in further analyses by type of fish and seafood consumption and in sensitivity analyses.

Conclusion: We found no evidence supporting a protective association of fish and seafood consumption during pregnancy with offspring symptoms of wheeze, asthma and allergic rhinitis from infancy to mid childhood.

Key words: Wheezing, asthma, allergic rhinitis, children, fish, seafood, pregnancy

Key Messages

- Findings from previous studies on the association of fish intake during pregnancy with offspring asthma and other allergy-related diseases are inconsistent.
- Heterogeneity between previous findings might be partly explained by small sample sizes and differences in diet assessment and health outcome definitions.
- We studied the association of fish and seafood consumption in pregnancy with child asthma and rhinitis outcomes in a combined analysis of 18 European and US birth cohorts with harmonized exposure and outcome variables according to a common protocol.
- Our results do not provide evidence of a beneficial association between fish and seafood consumption in pregnancy and development of asthma and allergic rhinitis symptoms in children up to 8 years of age.

Introduction

Asthma and allergic rhinitis are two of the most common chronic diseases in childhood, and their prevalence rates have increased over the past decades.¹ Emerging evidence suggests that a nutritional stressor or stimulus applied in intrauterine and early postnatal life can affect the development and maturation of respiratory and immune systems.² In this context, there has been a strong debate about the role of fish intake during pregnancy in the prevention of child asthma and other allergic diseases. Fish, and especially oily fish, is a rich source of n-3 long-chain fatty acids, which have been suggested to decrease the risk of allergic disease by exerting anti-inflammatory properties and modulating immune responses.³ However, it is also a major source of human exposure to harmful environmental contaminants, such as methylmercury, polychlorinated biphenyls and dioxins.⁴ Findings from birth cohort studies on the association of fish intake during pregnancy with offspring allergy-related disease symptoms are inconsistent, with reports of beneficial (i.e. lower incidence),^{5–8} null^{9–16} or even harmful effects.¹⁷ Likewise, prospective studies assessing n-3 fatty acid intake¹⁸ or biomarker levels in pregnancy^{19–25} and trials of n-3 long-chain fatty acid supplementation in pregnancy^{18,26,27} have yielded divergent results.

Most previous studies relied on relatively small sample sizes with a short duration of follow-up, and findings from one study were not replicated across populations with different fish consumption patterns. In the present analysis, we addressed these limitations by harmonizing and pooling for the first time individual data from a large network of cohorts in Europe and the USA, involving 60 774 pregnant women and their children with repeated follow-ups until the age of 8 years, to assess the strength and consistency of the association of fish and seafood consumption during pregnancy with the risk of childhood wheeze, asthma and allergic rhinitis.

Methods

Study population

European birth cohort studies participating in our previous analysis on fish intake in pregnancy and birth outcomes²⁸ were invited to participate if information on symptoms of asthma and rhinitis in childhood was available. From the 19 potentially eligible cohorts, 16 cohorts provided relevant data for the present analysis: ABCD (The Netherlands); DNBC (Denmark); FLEHS I (Belgium); GASPII (Italy); Generation R study (The Netherlands); Generation XXI study (Portugal); HUMIS (Norway); INMA (Spain); KOALA (The Netherlands); Lifeways Cross Generation (Ireland); LucKi (The Netherlands); NINFEA (Italy); PELAGIE (France); PIAMA (The Netherlands); RHEA (Greece); and SWS (UK). In addition, the Project Viva cohort from Massachusetts and the Bologna Birth Cohort agreed to take part in the current analysis. All participating cohorts covered deliveries from 1996 to 2011. In all cohorts, participants' parents or legal guardians provided written informed consent, and ethical approval was obtained from the local authorized institutional review boards. A data transfer agreement document was signed by each cohort study, and anonymized datasets were transferred to the University of Crete for analysis.

Overall, the study population included 60 774 mother-child pairs with information on fish intake during pregnancy, selected confounding variables and at least one of the health outcomes studied. Characteristics of the participating cohorts are shown in the eTable 1 and eFigure 1 (available as [Supplementary data](#) at *IJE* online).

Fish and seafood consumption during pregnancy

The exposure of interest was the frequency (times/week) of total fish, fatty fish, lean fish and seafood (other than fish) intake during pregnancy derived from cohort-specific food

frequency questionnaires or questionnaires specifically designed to assess fish intake during pregnancy (eTable 1, available as [Supplementary data](#)). Salmon, herring, mackerel, trout, sardines, Greenland halibut, anchovy, gurnard and tuna were classified as fatty fishes, whereas cod, pollock, plaice, flounder, garfish and similar species were classified as lean fishes. To align our analysis with the recent advice from the United States Food and Drug Administration and Environmental Protection Agency²⁹ on fish intake during pregnancy, and examine a potential dose-response relationship, we categorized total fish intake into the following categories; low: ≤ 1 time/week, moderate: > 1 but ≤ 3 times/week and high: > 3 times/week.

Symptoms of asthma and rhinitis in childhood

Participating cohorts provided information on wheeze, asthma and allergic rhinitis occurrence, mostly obtained by questionnaires adapted from the International Study on Asthma and Allergy in Childhood (ISAAC),³⁰ up to a maximum follow-up of 8 years. Cohort-specific information on outcome collection is shown in eTable 2 (available as [Supplementary data](#)). We defined wheeze in infancy as the presence of any episode of wheezing or whistling in the chest during the first 2 years of life, and wheeze in preschool and school-age children was defined as the presence of wheezing or whistling in the chest in the past 12 months reported at the ages of 3-4 years and 5-8 years, respectively, depending on the cohort-specific time points of assessment. Persistent wheeze at preschool and school age was defined as presence of wheeze both in infancy and the respective time period examined.³¹ We defined asthma at preschool and school age as satisfying at least two of the three following criteria for each time period: (i) ever-reported doctor diagnosis of asthma; (ii) presence of wheezing or whistling in the chest in the past 12 months; or (iii) asthma medication in the past 12 months at the ages of 3-4 years and 5-8 years, respectively.³² Allergic rhinitis at school age was defined as ever-reported diagnosis of allergic rhinitis or hay fever at the age of 5-8 years. If participants among the cohorts had repeatedly collected data within the same period, we used data collected at the oldest age.

Covariates

Important covariates or effect modifiers were defined as similarly as possible among the cohorts. Information on maternal pre-pregnancy body mass index (BMI; in kg/m²), maternal smoking during pregnancy (yes/no), gestational weight gain (in kg), maternal age at delivery (in years), birthweight (in grams), gestational age (in weeks) and child

sex (male or female) was collected by means of interviews or self-administered questionnaires, *ad hoc* measurements, birth records or medical registries. Information on parity (primiparous or multiparous), breastfeeding (yes/no), maternal education (cohort-specific definitions of low, medium or high), attendance at nursery school during infancy (yes/no), exposure to mould or dampness at home during infancy (yes/no) and history of parent asthma or hay fever (yes/no) was obtained through interviews or self-administered questionnaires.

Statistical analysis

We used multivariable generalized linear models for binary outcomes (modified Poisson³³) to estimate relative risks (RRs) and their 95% confidence intervals (CIs) for the associations of maternal fish and seafood consumption during pregnancy with offspring wheeze, asthma and allergic rhinitis in childhood. Selection of suitable covariates for model adjustment was based on previous knowledge and a directed acyclic graph (DAG) approach (eFigure 2, available as [Supplementary data](#) at *IJE* online). The set of variables identified by the DAG were maternal education, maternal smoking during pregnancy, parity, breastfeeding and parent asthma or hay fever (maternal allergy for PELAGIE). This list of variables plus maternal age and child sex, which are considered important determinants of child health outcomes, were the covariates included in multivariable models. For the cohort Generation XXI, the adjusted models did not include parent history of asthma or hay fever, owing to a large proportion of missing information on this variable. Further adjustment for maternal pre-pregnancy BMI, attendance at nursery school during infancy and exposure to mould or dampness at home during infancy did not materially change estimates for fish intake but reduced substantially the sample size; thus, we did not to include these factors in our final models.³⁴ Additionally, we did not include birthweight or gestational age in our main models, as we assumed that they are potential mediators in the pathway between fish intake during pregnancy and childhood respiratory health outcomes.^{28,35}

We analysed associations by undertaking cohort-specific analyses with subsequent random-effects meta-analysis. We assessed heterogeneity between cohort-specific effect estimates with the I^2 statistic and χ^2 test from Cochran's Q.³⁶ We also did pooled analysis including all participating cohorts by using mixed models with a random cohort intercept.

We performed several sensitivity analyses. First, we assessed the robustness of the results by repeating the meta-analyses while excluding one cohort at a time. Second, we examined the potential effect of the geographical location




Figure 1. Frequency of fish consumption during pregnancy (times/week) in participating cohorts. The line within the box marks the median; the boundaries of the box indicate the 25th and 75th percentiles; horizontal bars denote the variability outside the upper and lower quartiles (i.e., within 1.5 IQR of the lower and upper quartiles); and circles represent outliers.

of participating cohorts, based on the United Nations' classification (Southern Europe, Western Europe, Northern Europe and Northern America), by conducting meta-regression analyses. Third, we made further adjustment for gestational weight gain, used as a proxy for energy intake during pregnancy, for cohorts with available information. Fourth, we included birthweight and gestational age in the multivariable models to account for potential mediating effect of fetal growth. Finally, we assessed whether the effect estimates for wheeze in infancy and asthma in childhood varied by parent asthma or hay fever (yes vs no), and child sex (boy vs girl) by introducing interaction terms (one at a time), and conducting stratified analyses.³⁴ We performed analyses with STATA version 13 (StataCorp) and R version R3.1 (R Foundation).

Results

Participants' characteristics are shown in eTables 3 to 6 (available as [Supplementary data](#) at *IJE* online). Mothers had a median age of 30.4 years, were predominantly non-smokers, gave birth mainly to normal weight neonates and most of them breastfed their children. The ratio of boys to girls was 1.04 (eTable 3). The median fish consumption during pregnancy varied between cohorts and ranged from 0.44 times/week in the Generation R Study to 4.46 times/week in the INMA study (Figure 1; eTable 4). The highest median seafood intake was also reported in the INMA study (0.92 times/week) (eTable 4). We observed no differences in fish consumption levels between mothers with and without asthma or hay fever across cohorts (eTable 5).

In total, 17 518 (29.2%), 1949 (15.4%), and 3050 (13.1%) children had wheeze in infancy (0-2 years), at preschool (3-4 years) and school age (5-8 years), respectively (eTable 6); 921 (9.2%) and 2479 (10.5%) children had asthma at preschool and school age, respectively, and 1914 (5.4%) children had allergic rhinitis at school age (eTable 6).

Maternal fish intake during pregnancy was not associated with offspring wheeze symptoms in any age group (Figure 2 and Table 1). Also, no association was observed for persistent wheeze at preschool or school age (eTable 7, available as [Supplementary data](#) at *IJE* online). When we examined the risk of asthma at either preschool or school age, we observed no association with fish intake during pregnancy [adjusted meta-analysis RR per 1-time/week 1.02 (95% CI 0.97-1.07) and 1.01 (0.97-1.05), respectively; Figure 3 and Table 2]. Similarly, fish intake was not associated with the diagnosis of allergic rhinitis among school-age children (Table 2).

Further analysis by type of fish and seafood (other than fish) consumption did not modify the observed associations (Tables 1 and 2). Crude and adjusted estimates did not differ substantially (data not shown). Heterogeneity between cohort-specific risk estimates was low for most exposure-outcome associations (Tables 1 and 2; eTable 7), and the combined effect estimates from random-effect meta-analyses were broadly similar to those from the pooled analysis (eTables 8 and 9, available as [Supplementary data](#) at *IJE* online).

In sensitivity analyses, omitting one cohort at a time did not materially change the results (eTables 10 and 11, available as [Supplementary data](#) at *IJE* online). The risk estimates also remained similar when we took into account the geographical location of participating cohorts in meta-regression analyses, and when additional adjustment was made for gestational weight gain or birthweight and gestational age in multivariable models (data not shown). In stratified analyses, we found that fish intake in pregnancy was associated with a lower risk of asthma in preschool children with no history of parental asthma or hay fever [RR per 1 time/week 0.91 (95% CI 0.86-0.96)], but not in those of parents having either of these conditions [1.04 (0.98-1.10)]; *P* for interaction = 0.03; eTable 12, available as [Supplementary data](#) at *IJE* online). However, this interaction with parental history of asthma or hay fever was not observed for associations with wheeze or with asthma and allergic rhinitis at school age. Fish intake during pregnancy tended to be more protective against asthma in school-age boys [RR per 1 time/week 0.99 (0.99-1.00)] but not in girls [1.01 (0.99-1.03); *P* for interaction = 0.03], but associations were small in magnitude and not clinically relevant (eTable 12).




Figure 2. Adjusted associations of fish consumption during pregnancy (times/week) with offspring wheeze in infancy, preschool age, and school age. Wheeze in infancy was defined as presence of any episode of wheezing or whistling in the chest during the first 2 years of life. Wheeze at preschool age was defined as presence of wheezing or whistling in the chest in the past 12 months at the age of 3–4 years. Wheeze at school age was defined as presence of wheezing or whistling in the chest in the past 12 months at the age of 5–8 years. RRs (95% CIs) by cohort were obtained by using generalized linear models for binary outcomes (modified Poisson) adjusted for maternal age, maternal education, breastfeeding, smoking during pregnancy, parity, parent asthma or hay fever, and child sex. Combined estimates were obtained by using a random-effects meta-analysis. The names of the cohorts and number of participants per each cohort are shown on the left. Cohort-specific RRs (95% CIs), and their weights in meta-analysis are shown on the right. Squares represent the point estimate of each cohort, while the size of the square is proportional to the weight with which each cohort contributed to the overall RR; horizontal lines denote 95% CIs; and diamonds represent overall estimates. RR= relative risk.

Discussion

In this study of more than 60 000 mother child pairs from several locations in Europe and the USA with repeated follow-up data from birth up to 8 years, we found no protective association between fish and seafood consumption during pregnancy and risk of childhood wheeze, asthma

and allergic rhinitis. The present study is a major extension of previous collaborative work examining fish intake during pregnancy and child health.^{28,37}

Fetal life is a critical time period for airway and immune development, during which dietary exposures could have irreversible long-term influences on the development of

Table 1. Adjusted combined associations^a of fish and seafood (other than fish) consumption in pregnancy with childhood wheeze

Wheeze												
Infancy ^b												
	Preschool age ^c				School age ^d							
	Cohorts	Participants	RR (95% CI)	I ² (P heter) ^e	Cohorts	Participants	RR (95% CI)	I ² (P heter) ^e				
Fish consumption (times/week)	17	59986	1.00 (0.99-1.01)	11.2 (0.32)	11	12673	1.01 (0.99-1.04)	0.0 (0.81)	13	23317	1.01 (0.98-1.03)	8.1 (0.37)
Fish consumption categories ^f												
> 1 but ≤ 3 times/week	17	25144	0.98 (0.94-1.03)	17.7 (0.25)	11	3308	0.98 (0.88-1.10)	0.0 (0.65)	11	8318	1.01 (0.93-1.10)	0.0 (0.59)
> 3 times/week	14	13311	0.96 (0.89-1.03)	16.8 (0.27)	8	2039	0.98 (0.84-1.14)	0.0 (0.44)	9	3382	1.05 (0.95-1.17)	0.0 (0.69)
Type of fish consumption (times/week)												
Fatty fish	11	46492	1.00 (0.99-1.01)	0.0 (0.79)	8	7623	1.00 (0.96-1.04)	0.0 (0.78)	7	14156	0.99 (0.96-1.02)	0.0 (0.88)
Lean fish	11	46492	1.00 (0.99-1.01)	0.0 (0.76)	8	7623	1.04 (0.99-1.10)	25.9 (0.22)	7	14156	1.01 (0.98-1.03)	0.0 (0.44)
Seafood consumption (times/week)	14	49579	1.04 (0.99-1.10)	56.8 (0.01)	9	7737	1.03 (0.97-1.10)	0.0 (0.82)	10	15572	1.06 (0.94-1.20)	23.1 (0.23)

^aRelative risks (95% CIs) were estimated by random-effects meta-analysis by cohort. Generalized linear models for binary outcomes (modified Poisson) were adjusted for maternal age, maternal education, breastfeeding, smoking during pregnancy, parity, parent asthma or hay fever and child sex.

^bWheeze in infancy was defined as presence of any episode of wheezing or whistling in the chest during the first 2 years of life.

^cWheeze at preschool age (3-4 years) was defined as presence of wheezing or whistling in the chest during the first 12 months.

^dWheeze at school age (5-8 years) was defined as presence of wheezing or whistling in the chest in the past 12 months.

^eI² statistic and P-value for heterogeneity estimated by the χ^2 test from Cochran's Q.

^fReference category: ≤ 1 time/week.




Figure 3. Adjusted associations of fish consumption during pregnancy (times/week) with offspring asthma at preschool age and school age. Asthma at preschool age (3–4 years) and school age (5–8 years) was defined as satisfying at least two of the three following criteria for each time period: (1) ever-reported diagnosis of asthma, (2) presence of wheezing or whistling in the chest in the past 12 months, or (3) asthma medication in the past 12 months. RRs (95% CIs) by cohort were obtained by using generalized linear models for binary outcomes (modified Poisson) adjusted for maternal age, maternal education, breastfeeding, smoking during pregnancy, parity, parent asthma or hay fever, and child sex. Combined estimates were obtained by using random-effects meta-analysis. The names of the cohorts and number of participants per each cohort are shown on the left. Cohort-specific RRs (95% CIs), and their weights in meta-analysis are shown on the right. Squares represent the point estimate of each cohort, while the size of the square is proportional to the weight with which each cohort contributed to the overall RR; horizontal lines denote 95% CIs; and diamonds represent overall estimates. RR= relative risk.

asthma and other allergy-related diseases.³⁸ The n-3 long-chain fatty acids contained within fish can be transferred across the placenta and have been proposed to exert a protective effect through involvement in several immune regulatory pathways. Animal and *in vitro* studies have shown n-3 fatty acids to alter the expression of inflammatory genes by modifying transcription factor activation, inhibit T cell signalling through alteration of membrane lipid rafts and decrease the production of proinflammatory eicosanoids, cytokines and reactive oxygen species, as well as to give rise to a family of anti-inflammatory mediators termed resolvins (reviewed in Calder *et al.*³⁹). A beneficial effect of n-3 long-chain fatty supplementation in fetal or early life on childhood allergic disease symptoms has been shown in many, but not all, randomized trials.^{18,26,27,40} A relation between supplementation dose and effect becomes more

apparent in the very low end of the exposure spectrum of the n-3 long-chain fatty acids. Although fish and seafood is the major source of preformed n-3 long-chain fatty acids in the diet in most populations worldwide, maternal status is also determined by endogenous conversion from dietary alpha-linolenic acid from plant sources. A minority of the population has a genetically determined low endogenous conversion rate, and hence a higher dependence of preformed n-3 long-chain fatty acid intake from the diet.⁴¹ Therefore, our results do not exclude the possibility of a protective effect of fish intake against asthma and allergy in vulnerable subgroups.

Several birth cohort studies have examined the association of prenatal fish intake with asthma and rhinitis symptoms in childhood and have produced mixed results. High fish intake during pregnancy was associated with a

Asthma ^b							Allergic rhinitis					
Preschool age				School age			School age ^c					
Cohorts	Participants	RR (95% CI)	I ² (<i>P</i> heter) ^d	Cohorts	Participants	RR (95% CI)	I ² (<i>P</i> heter) ^d	Cohorts	Participants	RR (95% CI)	I ² (<i>P</i> heter) ^d	
Fish consumption (times/week)	10	10046	1.02 (0.97-1.07)	11.0 (0.34)	13	23532	1.01 (0.97-1.05)	27.1 (0.17)	10	35789	1.01 (0.99-1.03)	0.0 (0.44)
Fish consumption categories ^e												
> 1 but ≤ 3 times/week	8	2717	0.93 (0.78-1.10)	0.0 (0.52)	11	8472	1.02 (0.92-1.12)	0.0 (0.46)	10	14777	0.99 (0.84-1.16)	20.1 (0.26)
> 3 times/week	7	1643	1.00 (0.68-1.45)	37.1 (0.15)	9	3585	1.09 (0.97-1.23)	0.0 (0.61)	7	7827	0.99 (0.77-1.26)	25.7 (0.23)
Type of fish consumption (times/week)												
Fatty fish	8	6914	0.96 (0.88-1.06)	12.4 (0.33)	7	14661	1.00 (0.97-1.03)	0.0 (0.63)	6	26853	1.02 (0.98-1.05)	0.0 (0.83)
Lean fish	8	6914	1.06 (0.99-1.14)	0.0 (0.81)	7	14661	1.04 (0.96-1.12)	34.3 (0.17)	6	26853	0.99 (0.86-1.14)	47.7 (0.09)
Seafood consumption (times/week)	9	6976	1.03 (0.92-1.15)	0.0 (0.51)	10	16002	1.07 (0.91-1.25)	37.1 (0.11)	7	27101	1.04 (0.87-1.24)	39.2 (0.13)

^eReference category: ≤ 1 time/week.

lower risk of wheeze up to the age of 6.5 years in the Infancia y Medio Ambiente-Environment and Childhood Project-Menorca cohort,^{5,7} and the Danish National Birth Cohort study showed a protective effect of fish intake on ever asthma at the age of 7 years, by using hospital admission and prescription registry data.⁶ Other prospective studies conducted in The Netherlands,¹² Finland,^{11,16} Norway,¹⁴ Spain and Greece,¹³ France¹⁵ and Japan^{9,10} did not find an association of prenatal fish consumption with wheeze or asthma occurrence in childhood. Only few previous studies have assessed symptoms of childhood allergic rhinitis in association with fish intake in pregnancy; one UK study found that higher maternal intake of oily fish was protective against hay fever at age 5 years,⁸ whereas the Danish National Birth Cohort study⁶ and a Finnish study¹¹ found no association with allergic rhinitis up to the age of 7 years. Heterogeneity in previous findings may arise from differences in sample sizes, exposure assessment, health outcome definition and adjustment for confounding variables. In the present study, we had a large sample size and harmonized information on exposure variables, potential confounders and child asthma and rhinitis symptoms. We examined child outcomes at different time points from early infancy to mid childhood, and found little evidence for an association with fish intake during pregnancy.

Fish is generally considered an integral component of a healthy diet. However, it is a complex nutritional exposure. Fish is a good source of beneficial nutrients, including n-3 long-chain fatty acids, but is also a common route of exposure to persistent organic pollutants, such as polychlorinated biphenyls and dioxins.⁴ There is emerging evidence suggesting that prenatal exposure to these chemicals can exert adverse immunomodulatory effects and increase the risk of developing allergic disease symptoms in childhood.^{42,43} Hence, it is possible that pollutants bio-accumulating in fish counterbalance the effects of beneficial nutrients, moving the association between fish intake and child respiratory health towards the null. We did not have the possibility to collect information on amounts of nutrients and environmental chemicals contained in fish. Although we collected information on different fish types (fatty fish, lean fish, shellfish), we did not have enough data to distinguish between consumption of big and small species and fish origin, which would be relevant with respect to toxicant exposure. Further analyses using biomarker information on both nutrients and toxicants in fish are required to examine their interplay, as well as to disentangle potential opposite effect on child outcomes.

Genetic predisposition plays a major role in asthma.^{44,45} A recent systematic review concluded that family history of allergy should be considered as an important effect modifier when studying the association of

diet with the development of childhood asthma or allergies.³⁴ It has been proposed that atopic heredity is associated with abnormalities in fatty acid metabolism,⁴⁶ and this might affect body utilization of the fatty acids contained in fish. Previous studies examining a potential modifying effect of maternal history of asthma or atopy on prenatal fish intake have produced inconsistent results.^{47,48} In our study, we found that, among children of parents with no history of asthma or hay fever, prenatal fish intake was associated with lower asthma risk at preschool age, but not later in childhood or with wheeze or allergic rhinitis. Thus we treat this result with caution, as we cannot rule out the possibility that the observed association is due to chance. Further prospective studies incorporating genetic analysis are needed to evaluate whether genetic predisposition can modify the effect of fish intake during pregnancy on child allergy symptoms.

As in any observational study, there is the possibility of unmeasured residual confounding accounting for at least part of the observed findings. We had no data available to adjust for energy intake or dietary patterns during pregnancy; however, adjustment for gestational weight gain, as a good proxy for total energy intake in pregnancy,^{49,50} did not appreciably change the effect estimates. Moreover, although we examined breastfeeding as a potential confounder, other differences in child diet or lifestyle factors might have influenced the observed findings. However, the absence of important heterogeneity across cohorts and the similar effect estimates observed in the meta-analysis and the pooled analysis provide evidence in support of more robust and generalized conclusions.

Similar to most studies assessing the health effects of diet, we used self-reported dietary information, and we cannot rule out the possibility of exposure measurement error, especially in the absence of biomarkers of fish intake. However, most cohorts assessed fish intake using food frequency questionnaires that were developed and validated for use in pregnancy. Additionally, previous analyses in many participating cohorts have shown fish intake, as assessed by questionnaires, to positively correlate with exposure to n-3 long-chain fatty acids or environmental pollutants commonly found in fish.^{51–57} Any measurement error in the assessment of fish intake itself is likely to be non-differential with respect to the outcomes of interest, given that fish intake levels did not differ between mothers with and without asthma or hay fever across the cohorts, and dietary data were collected before child respiratory outcomes were known. Child asthma and allergic rhinitis symptoms were ascertained by symptom-based questionnaires, which could lead to outcome misclassification. However, most participating cohorts used questionnaires adapted from the ISAAC study.³⁰ To enhance asthma

outcome accuracy, we used a definition developed by a panel of experts within the Mechanisms of the Development of Allergy consortium³² that requires at least two conditions to be met. We assessed several asthma-related phenotypes, raising concern about multiple testing. However, an application of correction to take into account multiple comparisons would be inappropriate given that we studied outcomes that are highly correlated.⁵⁸

In conclusion, this large multicohort study found no evidence supporting a protective association between fish intake during pregnancy and offspring symptoms of wheeze, asthma and allergic rhinitis from infancy to mid childhood.

Supplementary Data

Supplementary data are available at *IJE* online.

Funding

This work was supported by the European Community's Seventh Framework Program [EU- FP7- HEALTH-2009-single-stage-241604]. Details of funding per cohort are available at *IJE* online.

Author Contributions

N.S., T.R. and L.C. had full access to all of the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis. Study concept and design: L.C. Acquisition of data: all authors. Statistical analysis: T.R. and N.S. Interpretation of data: all authors. Drafting of the manuscript: N.S. and L.C. Critical revision of the manuscript for important intellectual content: all authors. Study supervision: L.C.

Conflict of interest: Members of the Southampton Women's Survey team have received income for research (not related to this work) from Abbott Nutrition, Nestec and Danone Nutricia. L.D. received funding from the Lung Foundation Netherlands (no. 3.2.12.089; 2012). All other authors declare that they have no conflict of interest.

References

- Bousquet J, Khaltaev N (eds). *Global Surveillance, Prevention and Control of Chronic Respiratory Diseases: A Comprehensive Approach*. 2007. World Health Organization. <http://www.who.int/gard/publications/GARD Book 2007.pdf> (8 December 2016, date last accessed).
- Devereux G. The increase in the prevalence of asthma and allergy: food for thought. *Nat Rev Immunol* 2006;**6**:869-74.
- Calder PC, Kremmyda LS, Vlachava M, Noakes PS, Miles EA. Is there a role for fatty acids in early life programming of the immune system? *Proc Nutr Soc* 2010;**69**:373-80.
- Mozaffarian D, Rimm EB. Fish intake, contaminants, and human health: evaluating the risks and the benefits. *JAMA* 2006;**296**:1885-99.
- Romieu I, Torrent M, Garcia-Esteban R *et al*. Maternal fish intake during pregnancy and atopy and asthma in infancy. *Clin Exp Allergy* 2007;**37**:518-25.
- Maslova E, Strøm M, Oken E *et al*. Fish intake during pregnancy and the risk of child asthma and allergic rhinitis - longitudinal evidence from the Danish National Birth Cohort. *Br J Nutr* 2013;**110**:1313-25.
- Chatzi L, Torrent M, Romieu I, *et al*. Mediterranean diet in pregnancy is protective for wheeze and atopy in childhood. *Thorax* 2008;**63**:507-13.
- Willers SM, Devereux G, Craig LC *et al*. Maternal food consumption during pregnancy and asthma, respiratory and atopic symptoms in 5-year-old children. *Thorax* 2007;**62**:773-79.
- Miyake Y, Tanaka K, Okubo H, Sasaki S, Arakawa M. Maternal fat intake during pregnancy and wheeze and eczema in Japanese infants: the Kyushu Okinawa Maternal and Child Health Study. *Ann Epidemiol* 2013;**23**:674-80.
- Miyake Y, Sasaki S, Tanaka K, Ohfuji S, Hirota Y. Maternal fat consumption during pregnancy and risk of wheeze and eczema in Japanese infants aged 16-24 months: the Osaka Maternal and Child Health Study. *Thorax* 2009;**64**:815-21.
- Erkkola M, Nwaru BI, Kaila M *et al*. Risk of asthma and allergic outcomes in the offspring in relation to maternal food consumption during pregnancy: a Finnish birth cohort study. *Pediatr Allergy Immunol* 2012;**23**:186-94.
- Willers SM, Wijga AH, Brunekreef B *et al*. Maternal food consumption during pregnancy and the longitudinal development of childhood asthma. *Am J Respir Crit Care Med* 2008;**178**:12-31.
- Chatzi L, Garcia R, Roumeliotaki T *et al*. Mediterranean diet adherence during pregnancy and risk of wheeze and eczema in the first year of life: INMA (Spain) and RHEA (Greece) mother-child cohort studies. *Br J Nutr* 2013;**110**:2058-68.
- Oien T, Storro O, Johnsen R. Do early intake of fish and fish oil protect against eczema and doctor-diagnosed asthma at 2 years of age? A cohort study. *J Epidemiol Community Health* 2010;**64**:124-29.
- Pele F, Bajoux E, Gendron H *et al*. Maternal fish and shellfish consumption and wheeze, eczema and food allergy at age two: a prospective cohort study in Brittany, France. *Environ Health* 2013;**12**:102.
- Lumia M, Luukkainen P, Tapanainen H *et al*. Dietary fatty acid composition during pregnancy and the risk of asthma in the offspring. *Pediatr Allergy Immunol* 2011;**22**:827-35.
- Leermakers ET, Sonnenschein-van der Voort AM, Heppe DH *et al*. Maternal fish consumption during pregnancy and risks of wheezing and eczema in childhood: the Generation R Study. *Eur J Clin Nutr* 2013;**67**:353-59.
- Best KP, Gold M, Kennedy D, Martin J, Makrides M. Omega-3 long-chain PUFA intake during pregnancy and allergic disease outcomes in the offspring: a systematic review and meta-analysis of observational studies and randomized controlled trials. *Am J Clin Nutr* 2016;**103**:128-43.
- Rucci E, den Dekker HT, de Jongste JC *et al*. Maternal fatty acid levels during pregnancy, childhood lung function and atopic

- diseases. The Generation R Study. *Clin Exp Allergy* 2016;**46**:461-71.
20. Yu YM, Chan YH, Calder PC *et al.* Maternal PUFA status and offspring allergic diseases up to the age of 18 months. *Br J Nutr* 2015;**113**:975-83.
 21. Barman M, Johansson S, Hesselmar B, Wold AE, Sandberg AS, Sandin A. High levels of both n-3 and n-6 long-chain polyunsaturated fatty acids in cord serum phospholipids predict allergy development. *PLoS One* 2013;**8**:e67920.
 22. Notenboom ML, Mommers M, Jansen EH, Penders J, Thijs C. Maternal fatty acid status in pregnancy and childhood atopic manifestations: KOALA Birth Cohort Study. *Clin Exp Allergy* 2011;**41**:407-16.
 23. Pike KC, Calder PC, Inskip HM *et al.* Maternal plasma phosphatidylcholine fatty acids and atopy and wheeze in the offspring at age of 6 years. *Clin Dev Immunol* 2012;**2012**:474613.
 24. Newson RB, Shaheen SO, Henderson AJ, Emmett PM, Sherriff A, Calder PC. Umbilical cord and maternal blood red cell fatty acids and early childhood wheezing and eczema. *J Allergy Clin Immunol* 2004;**114**:531-37.
 25. Standl M, Demmelmair H, Koletzko B, Heinrich J. Cord blood LC-PUFA composition and allergic diseases during the first 10 yr. Results from the LISApplus study. *Pediatr Allergy Immunol* 2014;**25**:344-50.
 26. Best KP, Sullivan T, Palmer D *et al.* Prenatal fish oil supplementation and allergy: 6-year follow-up of a randomized controlled trial. *Pediatrics* 2016;**137**:e20154443.
 27. Hansen S, Strom M, Maslova E *et al.* Fish oil supplementation during pregnancy and allergic respiratory disease in the adult offspring. *J Allergy Clin Immunol* 2016;**139**:104-11.
 28. Leventakou V, Roumeliotaki T, Martinez D *et al.* Fish intake during pregnancy, fetal growth, and gestational length in 19 European birth cohort studies. *Am J Clin Nutr* 2014;**99**:506-16.
 29. *Fish: What Pregnant Women and Parents Should Know*. Draft updated advice by FDA and EPA. June 2014. <http://www.fda.gov/Food/FoodborneIllnessContaminants/Metals/ucm393070.htm> (8 December 2016, date last accessed).
 30. Asher MI, Keil U, Anderson HR *et al.* International Study of Asthma and Allergies in Childhood (ISAAC): rationale and methods. *Eur Respir J* 1995;**8**:483-91.
 31. Martinez FD, Wright AL, Taussig LM, Holberg CJ, Halonen M, Morgan WJ. Asthma and wheezing in the first six years of life. The Group Health Medical Associates. *N Engl J Med* 1995;**332**:133-38.
 32. Pinart M, Benet M, Annesi-Maesano I *et al.* Comorbidity of eczema, rhinitis, and asthma in IgE-sensitized and non-IgE-sensitized children in MeDALL: a population-based cohort study. *Lancet Respir Med* 2014;**2**:131-40.
 33. Zou G. A modified Poisson regression approach to prospective studies with binary data. *Am J Epidemiol* 2004;**159**:702-06.
 34. Nurmatov U, Nwaru BI, Devereux G, Sheikh A. Confounding and effect modification in studies of diet and childhood asthma and allergies. *Allergy* 2012;**67**:1041-59.
 35. den Dekker HT, Sonnenschein-van der Voort AM, de Jongste JC *et al.* Early growth characteristics and the risk of reduced lung function and asthma: A meta-analysis of 25,000 children. *J Allergy Clin Immunol* 2016;**137**:1026-35.
 36. Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med* 2002;**21**:1539-58.
 37. Stratakis N, Roumeliotaki T, Oken E *et al.* Fish intake in pregnancy and child growth: a pooled analysis of 15 European and US birth cohorts. *JAMA Pediatr* 2016;**170**:381-90.
 38. Devereux G. Early life events in asthma-diet. *Pediatr Pulmonol* 2007;**42**:663-73.
 39. Calder PC, Krauss-Etschmann S, de Jong EC *et al.* Early nutrition and immunity - progress and perspectives. *Br J Nutr* 2006;**96**:774-90.
 40. Gunaratne AW, Makrides M, Collins CT. Maternal prenatal and/or postnatal n-3 long chain polyunsaturated fatty acids (LCPUFA) supplementation for preventing allergies in early childhood. *Cochrane Database Syst Rev* 2015:CD010085.
 41. Minihane AM. Impact of genotype on EPA and DHA status and responsiveness to increased intakes. *Nutrients* 2016;**8**:123.
 42. Gascon M, Sunyer J, Casas M *et al.* Prenatal exposure to DDE and PCB 153 and respiratory health in early childhood: a meta-analysis. *Epidemiology* 2014;**25**:544-53.
 43. Hansen S, Strom M, Olsen SF *et al.* Maternal concentrations of persistent organochlorine pollutants and the risk of asthma in offspring: results from a prospective cohort with 20 years of follow-up. *Environ Health Perspect* 2014;**122**:93-99.
 44. Litonjua AA, Carey VJ, Burge HA, Weiss ST, Gold DR. Parental history and the risk for childhood asthma -Does mother confer more risk than father? *Am J Respir Crit Care Med* 1998;**158**:176-81.
 45. Bjerg A, Hedman L, Perzanowski MS, Platts-Mills T, Lundback B, Ronmark E. Family history of asthma and atopy: in-depth analyses of the impact on asthma and wheeze in 7- to 8-year-old children. *Pediatrics* 2007;**120**:741-48.
 46. Duchon K, Bjorksten B. Polyunsaturated n-3 fatty acids and the development of atopic disease. *Lipids* 2001;**36**:1033-42.
 47. Salam MT, Li YF, Langholz B, Gilliland FD. Maternal fish consumption during pregnancy and risk of early childhood asthma. *J Asthma* 2005;**42**:513-18.
 48. Calvani M, Alessandri C, Sopo SM *et al.* Consumption of fish, butter and margarine during pregnancy and development of allergic sensitizations in the offspring: role of maternal atopy. *Pediatr Allergy Immunol* 2006;**17**:94-102.
 49. Streuling I, Beyerlein A, Rosenfeld E, Schukat B, von Kries R. Weight gain and dietary intake during pregnancy in industrialized countries—a systematic review of observational studies. *J Perinat Med* 2011;**39**:123-29.
 50. Stuebe AM, Oken E, Gillman MW. Associations of diet and physical activity during pregnancy with risk for excessive gestational weight gain. *Am J Obstet Gynecol* 2009;**201**:58 e1-8.
 51. Papadopoulou E, Vafeiadi M, Agramunt S *et al.* Maternal diet, prenatal exposure to dioxins and other persistent organic pollutants and anogenital distance in children. *Sci Total Environ* 2013;**461-462**:222-29.
 52. van den Berg SW, Wijga AH, van Rossem L *et al.* Maternal fish consumption during pregnancy and BMI in children from birth up to age 14 years: the PIAMA cohort study. *Eur J Nutr* 2016;**55**:799-808.
 53. Bilau M, Matthys C, Baeyens W *et al.* Dietary exposure to dioxin-like compounds in three age groups: results from the Flemish environment and health study. *Chemosphere* 2008;**70**:584-92.

54. Halldorsson TI, Thorsdottir I, Meltzer HM, Nielsen F, Olsen SF. Linking exposure to polychlorinated biphenyls with fatty fish consumption and reduced fetal growth among Danish pregnant women: a cause for concern? *Am J Epidemiol* 2008;**168**:958-65.
55. Molto-Puigmarti C, Plat J, Mensink RP *et al.* FADS1 FADS2 gene variants modify the association between fish intake and the docosahexaenoic acid proportions in human milk. *Am J Clin Nutr* 2010;**91**:1368-76.
56. Ramon R, Murcia M, Aguinagalde X *et al.* Prenatal mercury exposure in a multicenter cohort study in Spain. *Environ Int* 2011;**37**:597-604.
57. Donahue SMA, Rifas-Shiman SL, Gold DR, Jouni ZE, Gillman MW, Oken E. Prenatal fatty acid status and child adiposity at age 3 y: results from a US pregnancy cohort. *Am J Clin Nutr* 2011;**93**:780-88.
58. Rothman KJ. No adjustments are needed for multiple comparisons. *Epidemiology* 1990;**1**:43-46.

Fish and seafood consumption during pregnancy and the risk of asthma and allergic rhinitis in childhood: a pooled analysis of 18 European and US birth cohorts

Table of contents

eTable 1: Description of participating cohorts.....	2
eTable 2: Data collection on respiratory outcomes in the participating cohorts.....	3
eTable 3: Description of maternal and child characteristics in participating cohorts....	5
eTable 4: Distribution of fish and seafood consumption in participating cohorts	6
eTable 5: Distribution of fish consumption levels among mothers with and without asthma or hayfever in participating cohorts	7
eTable 6: Distribution of respiratory outcomes in participating cohorts	8
eTable 7: Adjusted combined associations of fish and seafood (other than fish) consumption in pregnancy with childhood persistent wheeze.....	9
eTable 8: Adjusted combined associations (pooled analysis) of fish and seafood (other than fish) consumption in pregnancy with childhood wheeze.....	10
eTable 9: Adjusted combined associations (pooled analysis) of fish and seafood (other than fish) consumption in pregnancy with asthma and allergic rhinitis in childhood ..	11
eTable 10: Adjusted combined association of fish consumption during pregnancy (times/week) with childhood wheeze after omitting one cohort at a time	12
eTable 11: Adjusted combined association of fish consumption during pregnancy (times/week) with asthma and allergic rhinitis in childhood after omitting one cohort at a time.....	13
eTable 12: Adjusted combined associations (pooled analysis) of fish consumption in pregnancy (times/week) with wheeze in infancy and asthma in childhood by parent asthma or hay fever, and child sex.....	14
eFigure 1. Location of participating birth cohorts.. ..	15
eFigure 2: Directed acyclic graph for the association of fish consumption during pregnancy with child asthma or rhinitis symptoms	16

Supplementary material

eTable 1: Description of participating cohorts

Cohort	Recruitment	Available data on respiratory outcomes ^a	Available data on fish and seafood consumption during pregnancy	Method of dietary assessment	No. of participants included ^b
ABCD, Amsterdam, NL	2003-2004	5830	7719	Questionnaire	5029
CONER, Bologna, IT	2004-2005	402	402	FFQ	386
DNBC, nationwide, DK	1996-2002	55674	57921	FFQ	36158
FLEHS I, Flanders, BE	2002-2004	346	1056	FFQ	337
GASPII, Rome, IT	2003-2004	530	536	FFQ	530
Generation R, Rotterdam, NL	2001-2006	2451	2677	FFQ	2161
Generation XXI, Porto, PT ^c	2005-2006	257	276	FFQ	231
HUMIS, regional, NO	2003-2009	1530	1569	FFQ	357
INMA, Gipuzkoa, Sabadell, Valencia, SP	2003-2008	1844	1973	FFQ	1765
KOALA, regional, NL	2000-2003	2569	2697	FFQ	2508
Lifeways Cross Generation, Dublin, IR	2001-2003	187	662	FFQ	132
LucKi, regional, NL	2006-current	377	543	Questionnaire	229
NINFEA, nationwide, IT	2005-2016	2171	2214	Questionnaire	2105
PELAGIE, regional, FR	2002-2006	931	3228	Questionnaire	869
PIAMA, nationwide, NL	1996-1997	3335	3866	Questionnaire	3300
Project Viva, Massachusetts, US	1999-2002	1630	1777	FFQ	1329
RHEA, Heraklion, GR	2007-2008	907	970	FFQ	882
SWS, Southampton, UK	1998-2007	2510	2596	FFQ	2466
Overall		83624	92831		60774

ABCD: Amsterdam Born Children and their Development study; BE: Belgium; CONER: Bologna Birth Cohort; DK: Denmark; DNBC: Danish National Birth Cohort; FFQ: Food-frequency questionnaire; FLEHS I: Flemish Environment and Health Study I; FR: France; GASPII: Genetica e Ambiente: Studio Prospettico dell'Infanzia in Italia; Generation R: the Generation R Study; Generation XXI: the Generation XXI Birth Cohort; GR: Greece; HUMIS: Norwegian Human Milk Study; INMA: Infancia y Medio Ambiente-Environment and Childhood Project; IR: Ireland; IT: Italy; KOALA: Kind: Ouders en gezondheid: Aandacht voor Leefstijl en Aanleg Birth Cohort Study; Lifeways: Lifeways Cross Generation Cohort Study; LucKi: Luchtwegklachten bij Kinderen; NINFEA: Nascita e INFanzia: gli Effetti dell'Ambiente; NL: Netherlands; NO: Norway; PELAGIE: Endocrine disruptors: Longitudinal study on pregnancy abnormalities, infertility, and childhood; PIAMA: Prevention and Incidence of Asthma and Mite Allergy; Project Viva: the Project Viva study; PT: Portugal; RHEA: Mother Child Cohort in Crete; SP: Spain; SWS: Southampton Women's Survey; UK: United Kingdom; US, United States.

^a Participants with available information on at least one outcome of interest.

^b Participants with full information on exposure variables, selected confounding variables, and at least one outcome of interest.

^c The cohort Generation XXI involves a total of 8647 children, but only a sub-sample followed since the first trimester of pregnancy was included in this analysis.

Supplementary material

Table 2: Data collection on respiratory outcomes in the participating cohorts

Cohort	Wheeze	Asthma	Allergic rhinitis
ABCD, Netherlands	Information collected at 3-5 months: ever presence of wheezing Information collected at 7 years: presence of wheezing in the last 12 months	Information collected at 7 years: ever physician-diagnosed asthma	Information collected at 5 years: ever physician-diagnosed hay fever or allergic rhinitis
CONER, Italy	Information collected at 6 and 15 months: presence of wheezing or whistling in the chest at least once in the previous 6 months [from 6 to 15 months] Information collected at 3 years: presence of presence of wheezing or whistling in the chest at least once in the previous 12 months	Information collected at 3 years: - ever physician-diagnosed asthma - asthma medication in the last 12 months	Information collected at 8 years: ever physician-diagnosed hay fever or allergic rhinitis
DNBC, Denmark	Information collected at 6 months: presence of episodes of wheezing in the first 6 months Information collected at 1 and 7 years: presence of wheezing in the past 12 months	Information collected at 7 years: -ever physician-diagnosed asthma - asthma medication in the last 12 months	Information collected at 7 years: ever physician-diagnosed hay fever
FLEHS I, Belgium	Information collected at 6, 12, 18, 24, 30 and 36 months: presence of wheezing or whistling in the chest in the past 6 months Information collected at 4, 5, 6, 7, and 8 years: presence of wheezing or whistling in the chest in the past 12 months	Information collected at 4 years: medication for wheezing, coughing, rattling, shortness of breath or asthma in the last 12 months Information collected at 6 and 8 years: - ever physician-diagnosed asthma - asthma medication in the last 12 months	NA
GASPII, Italy	Information collected at 6 and 15 months: presence of wheezing or whistling in the chest at least once in the first 6 months of life [from 6 to 15 months] Information collected at 4, 7, and 8 years: presence of wheezing or whistling in the chest at least once since the previous follow-up	Information collected at 4, 7 and 8 years: - ever physician-diagnosed asthma - asthma medication in the last 12 months	Information collected at 4, 7 and 8 years: ever physician-diagnosed hay fever or allergic rhinitis
Generation R, Netherlands	Information collected at 1 year: presence of a whistling noise in the chest Information collected at 2, 4 and 6 years: presence of problems with a wheezing chest during the last year	Information collected at 6 years: ever physician-diagnosed asthma	Information collected at 6 years: physician-diagnosed respiratory allergy (pollen, house dust mite, and pets)
Generation XXI, Portugal	Information collected at 1 and 2 years: ever presence of wheezing Information collected at 4 and 7 years: presence of wheezing in the last 12 months	Information collected at 4 and 7 years: ever physician-diagnosed asthma	Information collected at 4 and 7 years: ever physician-diagnosed rhinitis
HUMIS, Norway	Information collected at 5 years: ever or current presence of obstructed/wheezing chest Information collected at 7 years: ever experience of wheezing/tightness in the chest after physical exercise in the last	Information collected at 5 and 7 years: ever physician-diagnosed asthma Information collected at 7 years: asthma medication in the last 12 months	Information collected at 5 and 7 years: ever physician-diagnosed pollen allergy

Supplementary material

	12 months		
INMA, Spain	Information collected at 1, 2 and 4 years: presence of whistling or wheezing from the chest, but not noisy breathing from the nose in the past 12 months	Information collected at 4 years: presence of asthma in the last 12 months	NA
KOALA, Netherlands	Information collected at 7 and 12 months: presence of wheezing at any time in the past [from 7 to 12 months] Information collected at 2, 4, 6, 8 years: presence of wheezing in the past 12 months	Information collected at 4, 6 and 8 years: - ever physician-diagnosed asthma - asthma medication in the last 12 months	NA
Lifeways Cross Generation, Ireland	Information collected at 1, and 2 years from Parent Held Child Study Records	NA	NA
LucKi, Netherlands	Information collected at 7 and 14 months: presence of wheezing or whistling in the chest in the previous 7 months	NA	NA
NINFEA, Italy	Information collected at 6 months: presence of episodes of wheezing or whistling in the chest in the first 6 months of life Information collected at 18 months: presence of episodes of wheezing or whistling in the chest from 6 to 18 months Information collected at 7 years: presence of episodes of wheezing in the past 12 months	Information collected at 7 years: - ever physician-diagnosed asthma - asthma medication in the last 12 months	Information collected at 7 years: ever physician-diagnosed allergic rhinitis
PELAGIE, France	Information collected at 2 years: presence of wheezing or whistling in the chest between birth to 24 months Information collected at 6 years: presence of wheezing or whistling in the chest in the past 12 months	Information collected at 6 years: ever physician-diagnosed asthma	NA
PIAMA, Netherlands	Information collected at 1, 2, 4, 6 and 8 years: presence of wheezing or whistling in the chest during the past 12 months	Information collected at 4, 6, and 8 years: - ever physician-diagnosed asthma - asthma medication in the last 12 months	Information collected at 6 and 8 years: ever diagnosis of hay fever
Project Viva, United States	Information collected at 1, 2, 3, 4, 5, 6 and 7 years: presence of wheezing or whistling in the chest in the past 12 months	Information collected at 3, 4, 5, 6 and 7 years: - physician diagnosis of asthma - asthma medication in the last 12 months	Information collected at 5, 6 and 7 years: physician diagnosis of hay fever, seasonal allergies or allergic rhinitis
RHEA, Greece	Information collected at 1 and 4 years: presence wheezing or whistling in the chest in the past 12 months	Information collected at 4 years: - ever physician-diagnosed asthma - asthma medication in the last 12 months	NA
SWS, United Kingdom	Information collected at 1, 2, 3, and 6 years: presence of any episodes of chestiness associated with wheezing or whistling in the chest in the last 12 months	Information collected at 3: ever physician-diagnosed asthma Information collected at 6 years: - ever physician-diagnosed asthma - asthma medication in the last 12 months	NA

Supplementary material

eTable 3: Description of maternal and child characteristics in participating cohorts

	Maternal age (years)	Smoking during pregnancy (yes)	Maternal education (low/medium)	Parity (multiparous)	Birth weight (grams)	Breastfeeding (yes)	Child sex (males)	Maternal asthma or hay fever (yes)	Paternal asthma or hay fever (yes)
	Median (IQR)	N (%)	N (%)	N (%)	Median (IQR)	N (%)	N (%)	N (%)	N (%)
ABCD, Netherlands	32.0 (29.0-35.0)	449 (8.9)	2646 (52.6)	2106 (41.9)	3480 (3150-3810)	4165 (82.8)	2523 (50.2)	728 (14.5)	593 (12.4)
CONER, Italy	33.0 (30.0-36.0)	47 (12.2)	237 (61.4)	166 (43.0)	3360 (3030-3618)	352 (91.2)	189 (49.0)	89 (23.1)	89 (23.2)
DNBC, Denmark	30.0 (27.0-33.0)	8442 (23.4)	11500 (31.8)	18767 (51.9)	3600 (3267-3950)	34001 (94.0)	18517 (51.2)	6415 (17.7)	2938 (8.2)
FLEHS I, Belgium	30.1 (27.7-32.5)	33 (9.8)	264 (78.3)	135 (40.0)	3400 (3115-3690)	243 (71.9)	175 (51.8)	107 (31.7)	116 (34.4)
GASPII, Italy	33.0 (30.0-36.0)	66 (12.5)	335 (63.2)	223 (42.1)	3350 (3020-3663)	463 (87.4)	272 (51.3)	165 (31.1)	171 (32.5)
Generation R, Netherlands	31.9 (29.6-34.3)	456 (21.1)	727 (33.6)	847 (39.2)	3540 (3200-3870)	1969 (91.1)	1057 (48.9)	670 (31.2)	560 (31.4)
Generation XXI, Portugal*	30.0 (25.0-34.0)	54 (23.4)	201 (87.0)	98 (42.4)	3250 (2955-3508)	215 (93.7)	112 (48.5)	13 (18.6)	7 (18.0)
HUMIS, Norway	30.0 (27.0-33.0)	24 (6.7)	96 (27.0)	213 (59.7)	3650 (3300-4050)	352 (98.6)	172 (48.2)	77 (21.6)	93 (26.1)
INMA, Spain	31.7 (29.2-34.5)	560 (31.7)	1160 (65.7)	778 (44.1)	3285 (2990-3550)	1561 (88.4)	913 (51.7)	309 (17.5)	283 (16.0)
KOALA, Netherlands	31.0 (29.0-34.0)	182 (7.3)	1289 (51.4)	1405 (56.0)	3510 (3200-3820)	2121 (84.6)	1274 (50.8)	928 (36.9)	899 (36.0)
Lifeways Cross Generation, Ireland	32.9 (30.4-36.2)	18 (13.6)	40 (30.3)	82 (62.1)	3605 (3233-3930)	95 (72.0)	64 (48.9)	15 (11.6)	12 (14.1)
LucKi, Netherlands	30.0 (28.0-33.0)	19 (8.3)	103 (45.0)	121 (52.6)	3375 (3065-3722)	157 (62.6)	110 (48.0)	88 (38.4)	74 (33.2)
NINFEA, Italy	33.0 (31.0-36.0)	170 (8.1)	838 (39.8)	494 (23.5)	3250 (2970-3540)	1933 (91.8)	1072 (50.9)	380 (18.1)	NA
PELAGIE, France ^a	30.6 (27.8-33.4)	191 (21.9)	500 (57.5)	513 (59.0)	3420 (3120-3725)	593 (68.2)	454 (52.2)	NA	NA
PIAMA, Netherlands	30.0 (28.0-33.0)	560 (17.0)	2099 (63.6)	1665 (50.5)	3510 (3200-3850)	2741 (83.1)	1700 (51.5)	787 (23.8)	815 (24.8)
Project Viva, United States	33.3 (30.6-36.5)	132 (9.9)	369 (27.8)	683 (51.4)	3509 (3175-3827)	1165 (87.7)	669 (50.3)	161 (12.1)	NA
RHEA, Greece	30.0 (26.0-33.0)	144 (16.3)	611 (69.3)	521 (59.1)	3210 (2950-3490)	751 (85.2)	448 (50.8)	215 (24.4)	163 (18.5)
SWS, United Kingdom	30.7 (27.8-33.4)	406 (16.5)	1773 (72.0)	1268 (51.4)	3468 (3170-3805)	2022 (82.0)	1286 (52.2)	556 (22.6)	459 (18.8)
Overall	30.4 (28.0-33.2)	11954 (19.7)	24788 (41.0)	30085 (49.5)	3540 (3200-3885)	54888 (90.3)	31006 (51.0)	11703 (19.6)	7272 (13.1)

NA, not available.

^a For the cohort PELAGIE, information on maternal allergy was provided for 869 participants, 94 (10.8%) of whom had allergy.

Supplementary material

eTable 4: Distribution of fish and seafood consumption in participating cohorts

	Total fish consumption (times/week)	Fatty fish consumption (times/week)	Lean fish consumption (times/week)	Seafood consumption (other than fish) (times/week)
	Median (IQR)	Median (IQR)	Median (IQR)	Median (IQR)
ABCD, Netherlands	1.00 (0.50-1.50)	NA	NA	NA
CONER, Italy	1.36 (0.92-2.00)	0.12 (0.00-0.34)	0.41 (0.12-0.75)	0.23 (0.1-0.48)
DNBC, Denmark	1.91 (1.10-3.56)	0.71 (0.29-1.46)	1.09 (0.56-1.89)	0.09 (0.00-0.25)
FLEHS I, Belgium	0.50 (0.25-1.00)	NA	NA	0.00 (0.00-0.25)
GASPII, Italy	2.23 (1.48-3.24)	1.00 (0.47-1.39)	0.50 (0.20-0.90)	0.38 (0.11-0.7)
Generation R, Netherlands	0.44 (0.15-0.74)	NA	NA	NA
Generation XXI, Portugal	4.00 (2.50-6.00)	1.00 (0.50-3.00)	2.00 (1.50-4.00)	0.50 (0.00-1.00)
HUMIS, Norway	1.77 (1.06-2.49)	0.37 (0.14-0.75)	1.23 (0.69-1.85)	0.17 (0.06-0.31)
INMA, Spain	4.46 (3.00-6.46)	0.92 (0.00-1.46)	3.46 (2.00-4.46)	0.92 (0.46-1.46)
KOALA, Netherlands	0.98 (0.49-1.96)	0.49 (0.00-0.49)	0.98 (0.49-0.98)	0.00 (0.00-0.00)
Lifeways Cross Generation, Ireland	1.00 (0.50-1.50)	0.50 (0.00-1.00)	0.50 (0.00-0.50)	0.00 (0.00-0.00)
LucKi, Netherlands	1.00 (0.50-1.50)	0.50 (0.00-1.00)	0.50 (0.00-1.00)	0.00 (0.00-0.50)
NINFEA, Italy	1.50 (0.50-1.50)	NA	NA	0.00 (0.00-0.50)
PELAGIE, France	0.67 (0.67-2.35)	NA	NA	0.35 (0.35-0.9)
PIAMA, Netherlands	0.50 (0.00-0.50)	NA	NA	NA
Project Viva, United States	0.98 (0.49-1.47)	0.49 (0.00-0.98)	0.49 (0.00-0.49)	0.49 (0.00-0.49)
RHEA, Greece	1.00 (0.46-1.37)	0.46 (0.00-1.00)	0.31 (0.08-0.69)	0.00 (0.00-0.23)
SWS, United Kingdom	1.77 (0.75-3.00)	0.50 (0.10-1.50)	1.00 (0.50-1.85)	0.00 (0.00-0.25)
Overall	1.50 (0.69-2.80)	0.64 (0.25-1.39)	1.00 (0.49-1.87)	0.08 (0.00-0.31)

NA, not available.

Supplementary material

eTable 5: Distribution of fish consumption levels among mothers with and without asthma or hayfever in participating cohorts

	Mothers without asthma or hayfever	Mothers with asthma or hayfever	P value for difference^a
	Total fish consumption in times/week	Total fish consumption in times/week	
	Median (IQR)	Median (IQR)	
ABCD, Netherlands	1.00 (0.50-1.50)	1.00 (0.50-1.62)	0.36
CONER, Italy	1.40 (0.93-2.00)	1.18 (0.66-1.88)	0.06
DNBC, Denmark	1.91 (1.11-3.57)	1.88 (1.07-3.54)	0.04
FLEHS I, Belgium	0.50 (0.25-1.00)	0.50 (0.25-1.00)	0.54
GASPII, Italy	2.19 (1.47-3.18)	2.29 (1.53-3.37)	0.72
Generation R, Netherlands	0.43 (0.15-0.77)	0.44 (0.15-0.80)	0.20
Generation XXI, Portugal	4.50 (3.00-6.00)	3.00 (2.00-6.50)	0.35
HUMIS, Norway	1.79 (1.02-2.51)	1.67 (1.14-2.25)	0.52
INMA, Spain	4.46 (3.00-6.46)	4.46 (3.00-6.89)	0.79
KOALA, Netherlands	0.98 (0.49-1.96)	1.47 (0.49-1.96)	0.17
Lifeways Cross Generation, Ireland	1.00 (0.50-1.50)	1.00 (0.50-1.00)	0.96
LucKi, Netherlands	1.00 (0.50-2.00)	1.00 (0.50-1.62)	0.82
NINFEA, Italy	1.50 (0.50-1.50)	1.50 (0.50-1.50)	0.38
PELAGIE, France	0.67 (0.67-2.35)	0.67 (0.67-2.35)	0.10
PIAMA, Netherlands	0.50 (0.00-0.50)	0.50 (0.00-1.00)	0.10
Project Viva, United States	0.98 (0.49-1.47)	0.98 (0.49-1.47)	0.28
RHEA, Greece	1.00 (0.46-1.34)	1.00 (0.46-1.46)	0.23
SWS, United Kingdom	1.75 (0.75-3.00)	1.85 (0.75-3.00)	0.69

^a P values for difference were estimated using the Mann-Whitney U test.

Supplementary material

eTable 6: Distribution of respiratory outcomes in participating cohorts

	Wheeze (yes) ^a			Persistent wheeze (yes) ^b		Asthma (yes) ^c		Allergic rhinitis at school age (yes) ^d
	Infancy	Preschool age	School age	Preschool age	School age	Preschool age	School age	
	N (%)	N (%)	N (%)	N (%)	N (%)	N (%)	N (%)	N (%)
ABCD, Netherlands	536 (10.7)	NA	186 (6.4)	NA	33 (1.3)	NA	179 (6.2)	54 (1.5)
CONER, Italy	129 (33.4)	42 (12.1)	NA	23 (9.7)	NA	42 (12.1)	NA	8 (5.1)
DNBC, Denmark	10955 (30.3)	NA	1771 (21.0)	NA	893 (20.9)	NA	1376 (15.2)	1185 (4.8)
FLEHS I, Belgium	44 (36.7)	14 (12.3)	18 (6.1)	9 (11.5)	3 (5.5)	8 (12.9)	21 (7.1)	NA
GASPII, Italy	211 (39.8)	84 (18.5)	36 (8.5)	51 (17.5)	23 (8.9)	79 (17.4)	30 (7.1)	36 (8.6)
Generation R, Netherlands	713 (34.3)	187 (10.1)	140 (8.0)	116 (9.4)	82 (7.2)	NA	55 (3.6)	144 (7.7)
Generation XXI, Portugal	62 (40.0)	42 (22.6)	25 (18.1)	18 (20.7)	14 (20.3)	7 (3.5)	9 (6.7)	12 (8.8)
HUMIS, Norway	NA	NA	17 (4.8)	NA	NA	NA	23 (6.7)	29 (8.1)
INMA, Spain	766 (43.4)	266 (18.5)	NA	186 (20.6)	NA	35 (2.9)	NA	NA
KOALA, Netherlands	757 (30.3)	124 (10.9)	139 (6.7)	74 (8.8)	82 (5.6)	101 (9.0)	114 (5.6)	NA
Lifeways Cross Generation, Ireland	27 (20.5)	NA	NA	NA	NA	NA	NA	NA
LucKi, Netherlands	43 (18.9)	NA	NA	NA	NA	NA	NA	NA
NINFEA, Italy	473 (22.5)	NA	14 (5.6)	NA	6 (3.2)	NA	6 (2.4)	15 (6.1)
PELAGIE, France	146 (16.9)	NA	78 (9.0)	NA	32 (4.6)	NA	53 (6.7)	NA
PIAMA, Netherlands	955 (29.0)	360 (11.7)	205 (6.6)	235 (10.2)	114 (5.1)	271 (8.8)	229 (7.4)	160 (5.2)
Project Viva, United States	388 (29.8)	258 (21.0)	194 (16.8)	155 (17.2)	108 (13.1)	202 (16.0)	220 (18.8)	271 (23.0)
RHEA, Greece	226 (25.9)	26 (4.5)	NA	10 (2.5)	NA	48 (8.1)	NA	NA
SWS, United Kingdom	1087 (44.0)	546 (24.2)	227 (14.2)	351 (24.8)	135 (14.2)	128 (7.4)	164 (11.0)	NA
Overall	17518 (29.2)	1949 (15.4)	3050 (13.1)	1228 (14.1)	1525 (10.4)	921 (9.2)	2479 (10.5)	1914 (5.4)

NA, not available.

^a Wheeze in infancy was defined as presence of any episode of wheezing or whistling in the chest during the first 2 years of life, while wheeze at preschool age (3-4 years) and school age (5-8 years) were defined as presence of wheezing or whistling in the chest in the past 12 months.

^b Persistent wheeze at preschool (3-4 years) and school age (5-8 years) was defined as presence of any episode of wheezing or whistling in the chest during the first 2 years of life, and wheezing or whistling in the chest in the past 12 months at the respective time period examined.

^c Asthma at preschool age (3-4 years) and school age (5-8 years) was defined as satisfying at least two of the three following criteria for each time period: (1) ever-reported diagnosis of asthma, (2) presence of wheezing or whistling in the chest in the past 12 months, or (3) asthma medication in the past 12 months.

^d Allergic rhinitis at school age was defined as ever-reported diagnosis of allergic rhinitis or hay fever at the age of 5-8 years.

eTable 7: Adjusted combined associations^a of fish and seafood (other than fish) consumption in pregnancy with childhood persistent wheeze

	Persistent wheeze							
	Pre-school age ^b				School age ^c			
	Cohorts	Participants	RR (95% CI)	I ² (p heter) ^d	Cohorts	Participants	RR (95% CI)	I ² (p heter) ^d
Fish consumption (times/week)	11	8708	1.02 (0.98-1.05)	0.0 (0.90)	11	14595	1.01 (0.99-1.03)	0.0 (0.88)
Fish consumption categories ^e								
>1 but ≤3 times/week	10	2221	1.00 (0.87-1.14)	0.0 (0.70)	9	4892	1.00 (0.88-1.15)	6.9 (0.39)
>3 times/week	8	1290	0.97 (0.79-1.20)	4.0 (0.40)	6	1755	1.08 (0.94-1.24)	0.0 (0.96)
Type of fish consumption (times/week)								
Fatty fish	8	5088	0.99 (0.94-1.05)	0.0 (0.97)	6	7849	0.99 (0.95-1.03)	0.0 (0.98)
Lean fish	8	5088	1.04 (0.98-1.10)	13.1 (0.33)	6	7849	1.03 (0.99-1.05)	0.0 (0.83)
Seafood consumption (times/week)	9	5166	1.04 (0.97-1.12)	0.0 (0.65)	8	8738	1.12 (0.98-1.28)	13.1 (0.33)

^a Relative risks (95% CIs) were estimated by random-effects meta-analysis by cohort. Generalized linear models for binary outcomes

(modified Poisson) were adjusted for maternal age, maternal education, breastfeeding, smoking during pregnancy, parity, parent asthma or hay fever, and child sex.

^b Persistent wheeze at preschool age was defined as presence of any episode of wheezing or whistling in the chest during the first 2 years of life, and wheezing or whistling in the chest in the past 12 months at the age of 3-4 years.

^c Persistent wheeze at school age was defined as presence of any episode of wheezing or whistling in the chest during the first 2 years of life, and wheezing or whistling in the chest in the past 12 months at the age of 5-8 years.

^d I² statistic and p value for heterogeneity estimated by the χ^2 test from Cochran's Q.

^e Reference category: ≤1 time/week.

eTable 8: Adjusted combined associations (pooled analysis)^a of fish and seafood (other than fish) consumption in pregnancy with childhood wheeze

	Wheeze								
	Infancy ^b			Preschool age ^c			School age ^d		
	Cohorts	Participants	RR (95% CI)	Cohorts	Participants	RR (95% CI)	Cohorts	Participants	RR (95% CI)
Fish consumption (times/week)	17	59889	1.00 (0.99-1.00)	11	12572	1.01 (1.00-1.03)	13	23230	1.00 (0.99-1.01)
Fish consumption categories ^e									
>1 but ≤3 times/week	17	25110	0.99 (0.97-1.01)	11	3271	0.98 (0.87-1.12)	13	8346	1.02 (0.96-1.09)
>3 times/week	17	13266	0.99 (0.97-1.01)	11	1987	0.97 (0.87-1.08)	13	3433	1.04 (1.01-1.08)
Type of fish consumption (times/week)									
Fatty fish	11	46395	1.00 (0.99-1.00)	8	7522	1.00 (0.96-1.03)	7	14069	0.99 (0.98-1.00)
Lean fish	11	46395	1.01 (1.00-1.01)	8	7522	1.03 (1.01-1.05)	7	14069	1.01 (1.00-1.02)
Seafood consumption (times/week)	14	49482	1.00 (0.97-1.03)	9	7636	1.02 (0.97-1.06)	10	15485	1.06 (0.96-1.17)

^a Pooled relative risks (95% CIs) were estimated by using multilevel mixed-effects generalized linear models for binary outcomes (modified Poisson) with a random cohort intercept adjusted for maternal age, maternal education, breastfeeding, smoking during pregnancy, parity, parent asthma or hay fever, and child sex.

^b Wheeze in infancy was defined as presence of any episode of wheezing or whistling in the chest during the first 2 years of life.

^c Wheeze at preschool age was defined as presence of wheezing or whistling in the chest in the past 12 months at the age of 3-4 years.

^d Wheeze at school age was defined as presence of wheezing or whistling in the chest in the past 12 months at the age of 5-8 years.

^e Reference category: ≤1 time/week.

eTable 9: Adjusted combined associations (pooled analysis)^a of fish and seafood (other than fish) consumption in pregnancy with asthma and allergic rhinitis in childhood

	Asthma ^b						Allergic rhinitis at school age ^c		
	Preschool age			School age					
	Cohorts	Participants	RR (95% CI)	Cohorts	Participants	RR (95% CI)	Cohorts	Participants	RR (95% CI)
Fish consumption (times/week)	10	9940	0.99 (0.94-1.04)	13	23442	1.00 (0.99-1.01)	10	35703	1.01 (1.00-1.02)
Fish consumption categories ^d									
>1 but ≤3 times/week	10	2748	0.90 (0.74-1.11)	14	8499	1.02 (0.91-1.14)	10	14745	1.03 (0.98-1.08)
>3 times/week	10	1709	0.88 (0.63-1.24)	14	3629	1.05 (0.97-1.15)	10	7801	1.02 (0.96-1.09)
Type of fish consumption (times/week)									
Fatty fish	8	6808	0.94 (0.84-1.05)	7	14571	0.99 (0.98-1.01)	6	26767	1.02 (1.01-1.02)
Lean fish	8	6808	1.04 (1.00-1.07)	7	14571	1.00 (0.98-1.03)	6	26767	1.01 (1.00-1.01)
Seafood consumption (times/week)	9	6870	0.94 (0.79-1.13)	10	15912	1.03 (0.93-1.13)	7	27015	1.00 (0.87-1.15)

^a Pooled relative risks (95% CIs) were estimated by using multilevel mixed-effects generalized linear models for binary outcomes (modified Poisson) with a random cohort intercept adjusted for maternal age, maternal education, breastfeeding, smoking during pregnancy, parity, parent asthma or hay fever, and child sex.

^b Asthma at preschool age (3-4 years) and school age (5-8 years) was defined as satisfying at least two of the three following criteria for each time period: (1) ever-reported diagnosis of asthma, (2) presence of wheezing or whistling in the chest in the past 12 months, or (3) asthma medication in the past 12 months.

^c Allergic rhinitis at school age (5-8 years) was defined as ever-reported diagnosis of allergic rhinitis or hay fever.

^d Reference category: ≤1 time/week.

eTable 10: Adjusted combined association^a of fish consumption during pregnancy (times/week) with childhood wheeze after omitting one cohort at a time

	Wheeze		
	Infancy ^b	Preschool age ^c	School age ^d
	RR (95% CI)	RR (95% CI)	RR (95% CI)
Overall	1.00 (0.99-1.01)	1.01 (0.99-1.04)	1.01 (0.98-1.03)
Cohort omitted			
ABCD, NL	1.00 (0.99-1.01)	NA	1.00 (0.97-1.04)
CONER, IT	1.00 (0.99-1.01)	1.01 (0.99-1.04)	NA
DNBC, DK	1.00 (0.99-1.02)	NA	1.01 (0.97-1.06)
FLEHS I, BE	1.00 (1.00-1.01)	1.01 (0.99-1.04)	1.00 (0.99-1.02)
GASPII, IT	1.00 (0.99-1.01)	1.01 (0.99-1.04)	1.01 (0.98-1.04)
Generation R Study, NL	1.00 (0.99-1.01)	1.01 (0.99-1.04)	1.00 (0.98-1.03)
Generation XXI, PT	1.00 (0.99-1.01)	1.01 (0.99-1.04)	1.01 (0.98-1.04)
HUMIS, NO	NA	NA	1.01 (0.98-1.03)
INMA, ES	1.00 (0.99-1.02)	1.01 (0.97-1.04)	NA
KOALA, NL	1.00 (0.99-1.01)	1.01 (0.99-1.04)	1.00 (0.99-1.02)
Lifeways Cross Generation, IR	1.00 (0.99-1.01)	NA	NA
LucKi, NL	1.00 (0.99-1.01)	NA	NA
NINFEA, IT	1.00 (0.99-1.01)	NA	1.00 (0.98-1.03)
PELAGIE, FR	1.00 (0.99-1.01)	NA	1.00 (0.98-1.02)
PIAMA, NL	1.00 (1.00-1.01)	1.01 (0.98-1.03)	1.00 (0.98-1.03)
Project Viva, US	1.00 (0.99-1.01)	1.01 (0.99-1.04)	1.01 (0.97-1.04)
RHEA, GR	1.00 (0.99-1.01)	1.01 (0.98-1.03)	NA
SWS, UK	1.01 (0.99-1.02)	1.01 (0.98-1.05)	1.00 (0.97-1.04)

^a All effect estimates correspond to an increase of 1 time/week in fish consumption in pregnancy.

Relative risks (95% CIs) were estimated by random-effects meta-analysis by cohort. Generalized linear models for binary outcomes (modified Poisson) were adjusted for maternal age, maternal education, breastfeeding, smoking during pregnancy, parity, parent asthma or hay fever, and child sex.

^b Wheeze in infancy was defined as presence of any episode of wheezing or whistling in the chest during the first 2 years of life.

^c Wheeze at preschool age was defined as presence of wheezing or whistling in the chest in the past 12 months at the age of 3-4 years.

^d Wheeze at school age was defined as presence of wheezing or whistling in the chest in the past 12 months at the age of 5-8 years.

eTable 11: Adjusted combined association^a of fish consumption during pregnancy (times/week) with asthma and allergic rhinitis in childhood after omitting one cohort at a time

	Asthma ^b		Allergic rhinitis at school age ^c
	Preschool age RR (95% CI)	School age RR (95% CI)	RR (95% CI)
Overall	1.02 (0.97-1.07)	1.01 (0.97-1.05)	1.01 (0.99-1.03)
Cohort omitted			
ABCD, NL	NA	1.01 (0.96-1.06)	1.02 (0.98-1.06)
CONER, IT	1.02 (0.96-1.08)	NA	1.01 (0.97-1.05)
DNBC, DK	NA	1.01 (0.95-1.07)	1.02 (0.95-1.09)
FLEHS I, BE	1.02 (0.97-1.07)	1.00 (0.98-1.02)	NA
GASPII, IT	1.03 (0.97-1.08)	1.01 (0.96-1.05)	1.01 (0.99-1.03)
Generation R Study, NL	NA	1.01 (0.96-1.05)	1.01 (0.99-1.03)
Generation XXI, PT	1.01 (0.96-1.06)	1.01 (0.96-1.05)	1.01 (0.97-1.05)
HUMIS, NO	NA	1.01 (0.96-1.05)	1.01 (0.98-1.05)
INMA, ES	1.03 (0.96-1.09)	NA	NA
KOALA, NL	1.02 (0.96-1.08)	1.01 (0.98-1.05)	NA
Lifeways Cross Generation, IR	NA	NA	NA
NINFEA, IT	NA	1.01 (0.96-1.05)	1.01 (0.99-1.03)
PELAGIE, FR	NA	1.01 (0.97-1.05)	NA
PIAMA, NL	1.01 (0.96-1.07)	1.01 (0.96-1.05)	1.01 (0.98-1.04)
Project Viva, US	1.03 (0.99-1.09)	1.00 (0.96-1.05)	1.01 (0.96-1.06)
RHEA, GR	1.02 (0.96-1.08)	NA	NA
SWS, UK	1.01 (0.95-1.07)	0.99 (0.96-1.03)	NA

^a All effect estimates correspond to an increase of 1 time/week in fish consumption in pregnancy. Relative risks (95% CIs) were estimated by random-effects meta-analysis by cohort. Generalized linear models for binary outcomes (modified Poisson) were adjusted for maternal age, maternal education, breastfeeding, smoking during pregnancy, parity, parent asthma or hay fever, and child sex.

^b Asthma at preschool age (3-4 years) and school age (5-8 years) was defined as satisfying at least two of the three following criteria for each time period: (1) ever-reported diagnosis of asthma, (2) presence of wheezing or whistling in the chest in the past 12 months, or (3) asthma medication in the past 12 months.

^c Allergic rhinitis at school age (5-8 years) was defined as ever-reported diagnosis of allergic rhinitis or hay fever.


eTable 12: Adjusted combined associations (pooled analysis)^a of fish consumption in pregnancy (times/week) with wheeze in infancy and asthma in childhood by parent asthma or hay fever, and child sex

	Wheeze in infancy ^b		Asthma ^c			
	N	RR (95% CI)	Preschool age		School age	
			N	RR (95% CI)	N	RR (95% CI)
Parent asthma or hay fever						
No	43069	1.00 (0.99-1.00)	6114	0.91 (0.86-0.96)	15783	0.99 (0.98-1.01)
Yes	16820	1.00 (1.00-1.01)	3826	1.04 (0.98-1.10)	7659	1.01 (1.00-1.02)
P-interaction ^d		0.67		0.03		0.11
Child sex						
Male	30585	1.00 (1.00-1.01)	5084	0.97 (0.89-1.06)	12352	0.99 (0.99-1.00)
Female	29304	1.00 (0.99-1.00)	4856	1.00 (0.95-1.06)	11090	1.01 (0.99-1.03)
P-interaction ^d		0.84		0.71		0.03


^a All effect estimates correspond to an increase of 1 time/week in fish consumption in pregnancy. Pooled relative risks (95% CIs) were estimated by using multilevel mixed-effects generalized linear models for binary outcomes (modified Poisson) with a random cohort intercept adjusted for maternal age, maternal education, breastfeeding, smoking during pregnancy, parity, parent asthma or hay fever (except for the models run separately by parent asthma or hay fever), and child sex (except for the models run separately by child sex).

^b Wheeze in infancy was defined as presence of any episode of wheezing or whistling in the chest during the first 2 years of life.

^c Asthma at preschool age (3-4 years) and school age (5-8 years) was defined as satisfying at least two of the three following criteria for each time period: (1) ever-reported diagnosis of asthma, (2) presence of wheezing or whistling in the chest in the past 12 months, or (3) asthma medication in the past 12 months. ^d P-interaction was estimated by Wald test.



eFigure 1. Location of participating birth cohorts. The size of the square is indicative of the cohort size. ABCD: Amsterdam Born Children and their Development study; CONER: Bologna Birth Cohort; DNBC: Danish National Birth Cohort; FLEHS I: Flemish Environment and Health Study I; GASPII: Genetica e Ambiente: Studio Prospettico dell'Infanzia in Italia; Generation R: the Generation R Study; Generation XXI: the Generation XXI Birth Cohort; HUMIS: Norwegian Human Milk Study; INMA: Infancia y Medio Ambiente-Environment and Childhood Project; IR: Ireland; KOALA: Kind: Ouders en gezondheid: Aandacht voor Leefstijl en Aanleg Birth Cohort Study; Lifeways: Lifeways Cross Generation Cohort Study; LucKi: Luchtwegklachten bij Kinderen; NINFEA: Nascita e INFanzia: gli Effetti dell'Ambiente; PELAGIE: Endocrine disruptors: Longitudinal study on pregnancy abnormalities, infertility, and childhood; PIAMA: Prevention and Incidence of Asthma and Mite Allergy; Project Viva: the Project Viva study; RHEA: Mother Child Cohort in Crete; SWS: Southampton Women's Survey.



eFigure 2: Directed acyclic graph for the association of fish consumption during pregnancy with child asthma or rhinitis symptoms