



Neutrophil recruitment by allergens contribute to allergic sensitization and allergic inflammation

Koa Hosoki^a, Toshiko Itazawa^a, Istvan Boldogh^b, and Sanjiv Sur^a

Purpose of review

To discuss the presence and role of neutrophils in asthma and allergic diseases, and outline the importance of pollen and cat dander-induced innate neutrophil recruitment in induction of allergic sensitization and allergic inflammation.

Recent findings

Uncontrolled asthma is associated with elevated numbers of neutrophils, and levels of neutrophil-attracting chemokine IL-8 and IL-17 in bronchoalveolar lavage fluids. These parameters negatively correlate with lung function. Pollen allergens and cat dander recruit neutrophils to the airways in a toll-like receptor 4, myeloid differentiation protein-2, and chemokine (C-X-C motif) receptor (CXCR) 2-dependent manner. Repeated recruitment of activated neutrophils by these allergens facilitates allergic sensitization and airway inflammation. Inhibition of neutrophil recruitment with CXCR2 inhibitor, disruption of toll-like receptor 4, or small interfering RNA against myeloid differentiation protein-2 also inhibits allergic inflammation. The molecular mechanisms by which innately recruited neutrophils contribute to shifting the airway inflammatory response induced by allergens from neutrophilic to an eosinophilic-allergic is an area of active research.

Summary

Recent studies have revealed that neutrophil recruitment is important in the development of allergic sensitization and inflammation. Inhibition of neutrophils recruitment may be a strategy to control allergic inflammation.

Keywords

cat dander, myeloid differentiation protein-2, neutrophils, pollens, toll-like receptor 4

INTRODUCTION

Asthma is a common inflammatory disease of the airways that affects 10% of the population in North America [1]. In patients with asthma, only a small 5–10% subset have uncontrolled asthma [2]; yet this subset accounts for a remarkable 30–50% of morbidity from this disease [3]. The type of inflammatory cells in the airways in uncontrolled asthma is different from controlled asthma, with controlled asthma having more eosinophils and Th2 cytokines than uncontrolled asthma [4–11]. The percentage of neutrophils in the sputum in uncontrolled asthma is higher compared with controlled asthma [6,7]. Study participants with acute severe asthma have a higher percentage of neutrophils in their tracheal aspirate compared with control patients undergoing nonpulmonary surgical procedures [8]. Similarly, patients intubated for status asthmaticus have a higher percentage of neutrophils in their bronchoalveolar lavage (BAL) fluid compared with that from patients with stable controlled asthma [9]. Sudden-

onset fatal asthma is characterized by more neutrophils in the airways than eosinophils in peribronchial lung tissues, which is different from slow-onset fatal asthma which demonstrates a preponderance of eosinophils in the airways [5]. Neutrophils have long been viewed as terminally differentiated cells that clear extracellular pathogens like bacterium. However, a growing body of literature indicates that neutrophils have numerous additional effects that regulate innate and adaptive immune responses [12,13]. Exposure to allergens can rapidly induce

^aDepartment of Internal Medicine, Division of Allergy and Immunology and
^bDepartment of Microbiology and Immunology, University of Texas Medical Branch, Galveston, Texas, USA

Correspondence to Sanjiv Sur, Department of Internal Medicine, Division of Allergy and Immunology, University of Texas Medical Branch, 301 University Boulevard, Galveston, TX 77555-0144, USA. Tel: +1 409 772 3410; fax: +1 409 772 5841; e-mail: sasur@UTMB.EDU

Curr Opin Allergy Clin Immunol 2016, 16:45–50

DOI:10.1097/ACI.0000000000000231

KEY POINTS

- Uncontrolled asthma is associated with increased numbers of neutrophils in the airways and elevated levels of neutrophil-attracting chemokine IL-8 and IL-17. These parameters negatively correlate with lung function.
- Neutrophil recruitment is crucial for development of skin allergic inflammation.
- Pollen allergenic extracts and cat dander innately recruit neutrophils to the airway in a TLR4, MD2, and CXCR2-dependent manner.
- Repeated recruitment of activated neutrophils following exposure to pollen allergenic extracts or cat dander facilitates allergic sensitization and airway inflammation.
- The molecular mechanisms by which repeated recruitment neutrophils shift the airway inflammation to an allergic phenotype is an active area of research.
- Inhibition of CXCR2, Leukotriene B₄, MD2, and TLR4 may serve as strategies to inhibit induction of allergic sensitization and inflammation.

an innate inflammatory response consisting of recruitment of neutrophils to the airways [14–17]. However, the molecular mechanisms by which allergens induce rapid recruitment of neutrophils are not well understood. In this review, we summarize recent data describing the molecular mechanisms that control allergen-induced neutrophil recruitment, and the contribution of these cells to allergic sensitization and allergic airway inflammation.

Uncontrolled asthma is associated with elevated level of IL-8 and IL-17

IL-8, a chemokine that attracts neutrophils, is associated with uncontrolled asthma. The concentration of IL-8 in sputum of study participants with uncontrolled asthma is higher than controlled asthma [6]. The levels of IL-8 in tracheal aspirates from the study participants intubated for acute uncontrolled asthma are higher compared with patients undergoing surgical procedures unrelated to the lung [8]. The patients intubated for status asthmaticus have higher levels of IL-8 in BAL fluid compared with controlled asthma [9]. We evaluated a panel of 48 cytokines and chemokines in BAL fluid from healthy control study participants and study participants with controlled and uncontrolled asthma, and reported that neutrophils and IL-8 are the only inflammatory components in BAL fluids that

distinguish controlled asthma from uncontrolled asthma, and both correlate inversely with forced expiratory volume (FEV)₁ [18[¶]].

IL-17 is also another chemokine that attracts neutrophils. Like IL-8, it is associated with uncontrolled asthma. The levels of IL-17 are elevated in the study participants with uncontrolled asthma compared with study participants with controlled asthma or normal controls [19]. Study participants with asthma have greater percentages of IL-17A positive compared with healthy control study participants [20]. Supernatants from patients with steroid resistant asthma have more IL-17A compared with patients with steroid sensitive asthma [20]. Asthma exacerbations in atopic severe asthma increase IL-17F and IL-17A expression in lamina propria in the airway and stimulate neutrophil recruitment to the airways [21[¶]]. The number of bronchial IL-17F positive cells correlates with the number of bronchial neutrophils and FEV₁ (%predicted) [21[¶]]. The levels of IL-17 in BAL fluid correlate negatively with FEV₁ in asthmatic study participants [22^{¶¶}]. Together these studies suggest a role for IL-8 and IL-17 in the pathogenesis of uncontrolled asthma.

Neutrophil recruitment stimulates development of allergic cutaneous inflammation

Neutrophil recruitment to the skin can also stimulate allergic skin inflammation [23]. Neutrophil recruitment to the skin is stimulated by tape stripped in wild-type (WT) mice. This tape-stripped neutrophil recruitment is reduced in leukotriene B₄ (Ltb₄) r1^{-/-} mice. Ltb₄r1^{-/-} mice mount reduced ovalbumin-induced allergic skin inflammation [23]. Transfer of neutrophils from WT mice to Ltb₄r1^{-/-} recipient mice exacerbates allergic skin inflammation, indicating that expression of leukotriene B₄ receptor 1 on neutrophils is important for acceleration of allergic skin inflammation [23]. Intradermal administration of neutrophils from donor WT mice to Ltb₄r1^{-/-} recipient mice enhances allergic skin inflammation [23]. Taken together, recruitment of neutrophil is crucial to develop allergic skin inflammation through their Leukotriene B₄ (LTB₄) production and their expression of leukotriene B₄ receptor 1. Another group reported a requirement of neutrophils for sensitization and elicitation of contact hypersensitivity. Treating the skin with 2,4,6-trinitrochlorobenzene stimulates infiltration of neutrophils [24^{¶¶}]. Neutrophil depletion by anti-lymphocyte antigen 6 complex, locus G antibody during the phase of sensitization and mice with a myeloid-specific conditional

deletion of the antiapoptotic myeloid cell leukemia 1 (Mcl-1) protein (Mcl-1^{ΔMyelo} mice) inhibits the 2,4,6-trinitrochlorobenzene-induced inflammatory skin response [24^{***}]. Together, these studies suggest that neutrophils are important in facilitating allergic cutaneous inflammation.

Pollen allergenic extracts stimulate CXCL chemokines and recruit neutrophils to the airways

Several in-vitro and ex-vivo studies [14,25–27] have shown that diverse allergenic extracts induce secretion of IL-8 from airway cells. Thus, microarray analysis revealed that timothy grass pollen extract induces elevation of IL-8 expression in NCI-H292 cells, a mucoepidermoid carcinoma cell line derived from human lungs [25]. Human airway epithelial cells secrete IL-8 following incubation with timothy grass pollen allergen [26], and this IL-8 secretion in bronchial epithelial cells occurred from not only study participants with uncontrolled asthma, but healthy volunteers [27]. Allergenic pollens have intrinsic NADPH oxidases, and these NADPH oxidases raised oxidized glutathione and 4-hydroxynonenal in BAL fluids of challenged mice. These oxidases stimulate tyrosine phosphorylation of p38 MAPK and secretion of IL-8 [14]. Intranasal challenge with ragweed pollen NADPH oxidase induces neutrophil recruitment to the airway in naïve mice and mice lacking T cells, B cells or mast cells [14]. Similarly, in human study participants with asthma or seasonal allergic rhinitis, allergen challenge increases the level of IL-8 [28–30] and neutrophil recruitment [15–17] in the airways.

Toll-like receptor 4, myeloid differentiation protein-2, and chemokine (C-X-C motif) receptor 2 mediate pollen and cat dander-induced innate neutrophil recruitment

We have recently reported that ragweed pollen extract can bind and stimulate CXCL-8 secretion from cells that express both toll-like receptor 4 (TLR4) and Myeloid differentiation protein-2 (MD2), but not cells that only express TLR4 [31^{***}]. Suppression of *Md2* in human bronchial epithelial cells by small interfering RNA (siRNA) inhibits ragweed pollen extract, Amb a 1, the main IgE-inducing allergenic protein in ragweed pollen extract, Bermuda, rye, firebush, pigweed, and cat dander extract-induced CXCL8 secretion [31^{***}]. Challenge of mice with ragweed pollen extract induces TLR4, chemokine (C-X-C motif) receptor (CXCR) 2, and nuclear factor-kappa B (NF-κB)-dependent recruitment of reactive oxygen species (ROS)-generating neutrophils to the airways [32^{***}].

Suppression of *Md2* mRNA by siRNA administration inhibits ragweed pollen extract and cat dander extract-induced neutrophil recruitment in naïve mice [31^{***}]. One of the major allergenic components of house dust mite, Der p2, has structural homology with MD2 that facilitates allergic inflammation by enhancing TLR4 inflammatory cascade [33,34]. Because there is no similarity of structure between Amb a1 and Der p2, its ability to stimulate CXCL8 suggests a distinct mechanism from the structural mimicry of MD2 used by Der p2 [33]. These data indicate that TLR4 and MD2 together regulate pollen extract and cat dander-induced NF-κB activation, CXCL 1/2 secretion, and recruitment of neutrophils to the airway, likely utilizing a mechanism that is distinct from molecular mimicry.

Toll-like receptor 4, myeloid differentiation protein-2, and CXCR2 mediate pollen and cat dander-induced allergic sensitization and allergic airway inflammation

We have reported that MD2, TLR4, and CXCR2 mediated neutrophil recruitment is crucial for induction of allergic sensitization and allergic inflammation [31^{***},32^{***}]. Thus, blocking ragweed pollen extract-induced recruitment of activated neutrophils by deletion of TLR4 inhibits induction of allergic airway inflammation [32^{***}]. Similarly, intranasal administration of CXCR2 inhibitor during repeated intranasal administration of ragweed pollen extract to the airway reduces allergic sensitization and inflammation [32^{***}]. Furthermore, forcing recruitment of neutrophils by administration of superoxide generator in *Tlr4* Knock out (KO) mice reconstitutes ragweed pollen extract-induced allergic inflammation [32^{***}]. Direct intranasal administration of neutrophils from WT donor mice together with ragweed pollen extract into *Tlr4* KO recipient mice reconstitutes ragweed pollen extract-induced allergic sensitization and airway inflammation [32^{***}]. In mice treated with siRNA to *Md2*, repeated challenge with ragweed pollen extract induces minimal allergic sensitization and airway inflammation [31^{***}], indicating MD2 plays an important role in induction of allergic sensitization to ragweed pollen extract. As diverse allergenic extracts require both MD2 and TLR4 to induce innate inflammation [31^{***},32^{***}], competition of LPS and allergens to bind MD2/TLR4 might explain why exposure to endotoxin reduces the risk of developing allergies [35,36]. Together these results strongly suggest that ragweed pollen extract and cat dander extract-induced innate recruitment of neutrophils stimulates allergic sensitization and allergic airway inflammation.




FIGURE 1. Schematic diagram showing allergen-induced innate neutrophil recruitment is critical for induction of allergic sensitization and airway inflammation. Recognition of pollens and cat dander by myeloid differentiation protein-2 in the context of toll-like receptor 4 activates NF- κ B mediated secretion of CXCL1, CXCL2, and CXCL8 from the airway epithelium. These chemokines recruit reactive oxygen species-generating neutrophils to the epithelium via CXCR2. The neutrophil-activated airway epithelium secretes TSLP and IL-33 upon subsequent contact with the same allergens, thereby facilitating allergic sensitization and allergic airway inflammation.

Molecular mechanisms by which neutrophils shifts to an allergic phenotype

The molecular mechanisms by which neutrophils stimulate allergic inflammation is an active area of research. Recently, several studies have shown that neutrophils can regulate the induction of allergic inflammation through their innate responses. Neutrophils are a major source of oxidative stress by their respiratory chain. Because mice deficient in gp91phox showed less Th2 cytokine secretion and allergic inflammation [37], recruitment of ROS-generating neutrophils after allergen challenge may have an important role in subsequent allergic inflammation. Several studies have shown that chronic oxidative stress can worsen allergic asthma [38–41], alter dendritic cell function, and modify Th1/Th2 balance [42,43]. By contrast, administration of antioxidants such as ascorbic acid, N-acetyl-cysteine, tocopherol, or lactoferrin with antigen suppresses antigen-induced allergic airway inflammation [44–46]. As neutrophils recruited by intranasal allergen challenge generate sustained ROS in the airways, it is likely that this property of neutrophil promotes allergic inflammation through ROS generation. This is suggested by studies that mice deficient in gp91phox, the dominant superoxide generating enzyme in neutrophils, have

decreased ROS, and mount an attenuated allergic inflammatory response to allergen challenge [37]. LTB₄ generation from recruited neutrophil to the skin stimulates accumulation of effector CD4⁺ T cells to ovalbumin-challenged skin, thereby facilitating allergic skin inflammation [23]. Neutrophils may have contributed to allergic inflammation by increasing microvascular permeability [47], inducing proinflammatory cytokines [48], matrix metalloproteinase 9 [49], and mucin 5AC [50].

Small molecule inhibitors of neutrophil recruitment inhibit allergic inflammation

Neutrophil recruitment from blood to a sterile extravascular site has been shown to be dependent specifically on LTB₄ [51]. In addition, recruited neutrophil-derived LTB₄ is crucial for induction of subsequent allergic skin inflammation [23]. Pharmacologic blockade of LTB₄ synthesis by besatin suppresses allergic skin inflammation elicited by cutaneous antigen challenge in sensitized mice [23]. Because the level of LTB₄ is elevated in BAL fluid from the patients with asthma [52], suppression of LTB₄ synthesis or using an LTB₄ inhibitor could be effective in inhibiting allergic airway inflammation. However, as intranasal ragweed

pollen extract challenge does not induce a significant increase of LTB₄ in BAL fluid in allergic model mice [32[■]], further studies about the role of LTB₄ in induction of allergic airway inflammation are required.

CXCR2 is the receptor for CXCL1 and CXCL2. Prior murine and human studies suggests an importance of CXCR2 signaling in induction of asthma and allergic inflammation. Thus, treatment of mice with anti-CXCR2 mAb inhibits late-phase airway obstruction, airway hyperresponsiveness, eosinophilic inflammation, and goblet cell hyperplasia [53[■]]. Furthermore, study participants with uncontrolled asthma treated with the CXCR2 inhibitor 2-hydroxy-N,N-dimethyl-3-{2-[[[(R)-1-(5-methylfuran-2-yl)-propyl]amino]-3,4-dioxo-cyclobut-1-enylamino]-benzamide have been reported to have fewer mild exacerbations [54]. We recently reported that this receptor plays a critical role in recruitment of pollen-induced neutrophils to the airways and pollen-induced allergic sensitization and allergic airway inflammation [55]. SB225002 is a small molecule inhibitor of ligand binding to CXCR2 and inhibits CXCR2-mediated neutrophilic recruitment [55]. Administration of SB225002 before intranasal ragweed pollen extract challenge inhibits superoxide generating neutrophil recruitment to the airway [32[■]]. Adminstrating SB225002 prior to each repeated ragweed pollen extract challenge suppresses ragweed pollen extract-induced allergic inflammation, accumulation of mucin in epithelial cells, production of ragweed pollen extract-specific IgE in serum, and secretion of IL-5, IL-13, thymic stromal lymphopoietin (TSLP), and IL-33 in BAL fluid [32[■]]. Future studies will have to validate the results of these studies in humans.

CONCLUSION

An increasing body of literature indicates that neutrophils and neutrophil-attracting chemokines like IL-8 and IL-17 are increased in uncontrolled asthma. Recent studies have shown that neutrophils are closely associated with not only severity, but initiation of allergic inflammation and allergic sensitization. The innate immune responses stimulated by exposure of the airways to pollen and cat dander allergenic extracts recruits neutrophils to the airways that in turn stimulate allergic sensitization and inflammation (Fig. 1). The mechanisms by which recruited neutrophils contribute to the induction of allergic sensitization and inflammation is not yet understood, but could be because of its ability to produce ROS, leukotrienes, cytokines like IL-33 or TSLP, or other factors that stimulate allergic inflammation. Further, basic and

translational research is required to characterize the role and contribution of neutrophils in induction of allergic sensitization and allergic airway inflammation. Future studies should evaluate targets like TLR4/MD2, CXCR2, or LTB₄ to inhibit recruitment of activated neutrophils as a novel strategy to prevent development of allergic sensitization and allergic airway inflammation.

Acknowledgements

None.

Financial support and sponsorship

This work was supported by National Institute of Allergy and Infectious Diseases P01 AI062885-06, National Institute of Environmental Health Sciences RO1 ES18948, National Heart, Lung and Blood Institute Proteomic Center, N01HV00245, National Institute of Environmental Health Sciences T32 ES007254 and Leon Bromberg Professorship at UTMB.

Conflicts of interest

There are no conflicts of interest.

REFERENCES AND RECOMMENDED READING

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Braman SS. The global burden of asthma. *Chest* 2006; 130:4S–12S.
2. Godard P, Chanez P, Siraudin L, *et al.* Costs of asthma are correlated with severity: a 1-yr prospective study. *Eur Respir J* 2002; 19:61–67.
3. Anagnostou K, Harrison B, Iles R, Nasser S. Risk factors for childhood asthma deaths from the UK Eastern Region Confidential Enquiry. *Prim Care Respir J* 2012; 21:71–77.
4. Wenzel SE, Schwartz LB, Langmack EL, *et al.* Evidence that severe asthma can be divided pathologically into two inflammatory subtypes with distinct physiologic and clinical characteristics. *Am J Respir Crit Care Med* 1999; 160:1001–1008.
5. Sur S, Crotty TB, Kephart GM, *et al.* Sudden-onset fatal asthma. A distinct entity with few eosinophils and relatively more neutrophils in the airway submucosa? *Am Rev Respir Dis* 1993; 148:713–719.
6. Jatakanon A, Uasuf C, Maziak W, *et al.* Neutrophilic inflammation in severe persistent asthma. *Am J Respir Crit Care Med* 1999; 160:1532–1539.
7. Moore WC, Hastie AT, Li X, Li H, Busse WW, Jarjour NN, Wenzel SE, Peters SP, Meyers DA, Bleecker ER. Sputum neutrophil counts are associated with more severe asthma phenotypes using cluster analysis. *J Allergy Clin Immunol* 2014; 133:1557–1563; e1555.
8. Ordonez CL, Shaughnessy TE, Matthay MA, Fahy JV. Increased neutrophil numbers and IL-8 levels in airway secretions in acute severe asthma: clinical and biologic significance. *Am J Respir Crit Care Med* 2000; 161:1185–1190.
9. Lamblin C, Gosset P, Tillie-Leblond I, *et al.* Bronchial neutrophilia in patients with noninfectious status asthmaticus. *Am J Respir Crit Care Med* 1998; 157:394–402.
10. Sur S, Gleich GJ, Swanson MC, *et al.* Eosinophilic inflammation is associated with elevation of interleukin-5 in the airways of patients with spontaneous symptomatic asthma. *J Allergy Clin Immunol* 1995; 96:661–668.
11. Sur S, Kita H, Gleich GJ, *et al.* Eosinophil recruitment is associated with IL-5, but not with RANTES, twenty-four hours after allergen challenge. *J Allergy Clin Immunol* 1996; 97:1272–1278.
12. Mantovani A, Cassatella MA, Costantini C, Jaillon S. Neutrophils in the activation and regulation of innate and adaptive immunity. *Nat Rev Immunol* 2011; 11:519–531.
13. Mocsai A. Diverse novel functions of neutrophils in immunity, inflammation, and beyond. *J Exp Med* 2013; 210:1283–1299.

14. Boldogh I, Bacsí A, Choudhury BK, *et al.* ROS generated by pollen NADPH oxidase provide a signal that augments antigen-induced allergic airway inflammation. *J Clin Invest* 2005; 115:2169–2179.
15. Lommatzsch M, Julius P, Kuepper M, *et al.* The course of allergen-induced leukocyte infiltration in human and experimental asthma. *J Allergy Clin Immunol* 2006; 118:91–97.
16. Hoskins A, Reiss S, Wu P, *et al.* Asthmatic airway neutrophilia after allergen challenge is associated with the glutathione S-transferase M1 genotype. *Am J Respir Crit Care Med* 2013; 187:34–41.
17. Metzger WJ, Richerson HB, Worden K, *et al.* Bronchoalveolar lavage of allergic asthmatic patients following allergen bronchoprovocation. *Chest* 1986; 89:477–483.
18. Hosoki K, Ying S, Corrigan C, *et al.* Analysis of a panel of 48 cytokines in BAL fluids specifically identifies IL-8 levels as the only cytokine that distinguishes controlled asthma from uncontrolled asthma, and correlates inversely with FEV1. *PLoS ONE* 2015; 10:e0126035.
- The publication identified IL-8 as the only significantly elevated cytokine among 48 measured and elevated neutrophil numbers in BAL fluid in uncontrolled asthma compared with controlled asthma and both were inversely correlated with FEV1.
19. Chakir J, Shannon J, Molet S, *et al.* Airway remodeling-associated mediators in moderate to severe asthma: effect of steroids on TGF- β , IL-11, IL-17, and type I and type III collagen expression. *J Allergy Clin Immunol* 2003; 111:1293–1298.
20. Nanzer AM, Chambers ES, Ryanna K, *et al.* Enhanced production of IL-17A in patients with severe asthma is inhibited by 1 α ,25-dihydroxyvitamin D3 in a glucocorticoid-independent fashion. *J Allergy Clin Immunol* 2013; 132:297–304.
21. Sorbello V, Ciprandi G, Di Stefano A, *et al.* Nasal IL-17F is related to bronchial IL-17F/neutrophilia and exacerbations in stable atopic severe asthma. *Allergy* 2015; 70:236–240.
- The manuscript demonstrated that IL-17F and IL-17A expression in nasal/bronchial lamina propria and bronchial neutrophils was increased in uncontrolled atopic asthma, indicating that bronchial IL-17-driven neutrophilia is important in uncontrolled asthma.
22. Irvin C, Zafar I, Good J, *et al.* Increased frequency of dual-positive TH2/TH17 cells in bronchoalveolar lavage fluid characterizes a population of patients with severe asthma. *J Allergy Clin Immunol* 2014; 134:1175–1186; e1177.
- The manuscript identified that dual-positive TH2/TH17 cells in bronchoalveolar lavage fluid from asthmatic patients reflects the resistance to glucocorticoids and asthma severity.
23. Oyoshi MK, He R, Li Y, *et al.* Leukotriene B4-driven neutrophil recruitment to the skin is essential for allergic skin inflammation. *Immunity* 2012; 37:747–758.
24. Weber FC, Nemeth T, Csepregi JZ, *et al.* Neutrophils are required for both the sensitization and elicitation phase of contact hypersensitivity. *J Exp Med* 2015; 212:15–22.
- The manuscript demonstrates a critical role of neutrophils in the elicitation phase of contact hypersensitivity by depletion or transfer the neutrophils.
25. Roschmann KI, Luiten S, Jonker MJ, *et al.* Timothy grass pollen extract-induced gene expression and signalling pathways in airway epithelial cells. *Clin Exp Allergy* 2011; 41:830–841.
26. Roschmann K, Farhat K, König P, *et al.* Timothy grass pollen major allergen Phl p 1 activates respiratory epithelial cells by a nonprotease mechanism. *Clin Exp Allergy* 2009; 39:1358–1369.
27. Blume C, Swindle EJ, Dennison P, *et al.* Barrier responses of human bronchial epithelial cells to grass pollen exposure. *Eur Respir J* 2013; 42:87–97.
28. Virchow JC Jr, Walker C, Hafner D, *et al.* T cells and cytokines in bronchoalveolar lavage fluid after segmental allergen provocation in atopic asthma. *Am J Respir Crit Care Med* 1995; 151:960–968.
29. Teran LM, Carroll MP, Frew AJ, *et al.* Leukocyte recruitment after local endobronchial allergen challenge in asthma. Relationship to procedure and to airway interleukin-8 release. *Am J Respir Crit Care Med* 1996; 154:469–476.
30. KleinJan A, Dijkstra MD, Boks SS, *et al.* Increase in IL-8, IL-10, IL-13, and RANTES mRNA levels (in situ hybridization) in the nasal mucosa after nasal allergen provocation. *J Allergy Clin Immunol* 1999; 103:441–450.
31. Hosoki K, Boldogh I, Leopoldo A-A, *et al.* MD2 facilitates pollen and cat dander-induced innate and allergic airway inflammation. *J Allergy Clin Immunol*.
- The article demonstrated a critical rate-limiting role of MD2 in induction of innate immune responses, allergic sensitization, and allergic airway inflammation by diverse allergenic extract from cat dander and pollens, such as grasses, weeds, and trees.
32. Hosoki K, Aguilera-Aguirre L, Brasier AR, *et al.* Pollen-induced innate recruitment of neutrophils facilitates induction of allergic sensitization and airway inflammation. *Am J Respir Cell Mol Biol* 2015. (in press).
- The study demonstrates that pollen-induced repeated recruitment of neutrophils is critical for induction of allergic sensitization to pollens and stimulation of pollen-induced allergic airway inflammation.
33. Trompette A, Divanovic S, Visintin A, *et al.* Allergenicity resulting from functional mimicry of a Toll-like receptor complex protein. *Nature* 2009; 457:585–588.
34. Hammad H, Chieppa M, Perros F, *et al.* House dust mite allergen induces asthma via Toll-like receptor 4 triggering of airway structural cells. *Nat Med* 2009; 15:410–416.
35. Braun-Fahrlander C, Riedler J, Herz U, *et al.* Environmental exposure to endotoxin and its relation to asthma in school-age children. *N Engl J Med* 2002; 347:869–877.
36. Eisenbarth SC, Piggott DA, Huleatt JW, *et al.* Lipopolysaccharide-enhanced, toll-like receptor 4-dependent T helper cell type 2 responses to inhaled antigen. *J Exp Med* 2002; 196:1645–1651.
37. Sevin CM, Newcomb DC, Toki S, *et al.* Deficiency of gp91phox inhibits allergic airway inflammation. *Am J Respir Cell Mol Biol* 2013; 49:396–402.
38. McDonnell WF, Abbey DE, Nishino N, Lebowitz MD. Long-term ambient ozone concentration and the incidence of asthma in nonsmoking adults: the AHSMOG study. *Environ Res* 1999; 80:110–121.
39. Kim BJ, Kwon JW, Seo JH, *et al.* Association of ozone exposure with asthma, allergic rhinitis, and allergic sensitization. *Ann Allergy Asthma Immunol* 2011; 107:214–219.
40. McConnell R, Berhane K, Gilliland F, *et al.* Asthma in exercising children exposed to ozone: a cohort study. *Lancet* 2002; 359:386–391.
41. Yamazaki S, Shima M, Nakadate T, *et al.* Association between traffic-related air pollution and development of asthma in school children: Cohort study in Japan. *J Expo Sci Environ Epidemiol* 2014; 24:372–379.
42. Peterson JD, Herzenberg LA, Vasquez K, Waltenbaugh C. Glutathione levels in antigen-presenting cells modulate Th1 versus Th2 response patterns. *Proc Natl Acad Sci U S A* 1998; 95:3071–3076.
43. Csillag A, Boldogh I, Pazmandi K, *et al.* Pollen-induced oxidative stress influences both innate and adaptive immune responses via altering dendritic cell functions. *J Immunol* 2010; 184:2377–2385.
44. Dharajani N, Choudhury BK, Bacsí A, *et al.* Inhibiting pollen reduced nicotinamide adenine dinucleotide phosphate oxidase-induced signal by intrapulmonary administration of antioxidants blocks allergic airway inflammation. *J Allergy Clin Immunol* 2007; 119:646–653.
45. Kruzel ML, Bacsí A, Choudhury B, *et al.* Lactoferrin decreases pollen antigen-induced allergic airway inflammation in a murine model of asthma. *Immunology* 2006; 119:159–166.
46. Chodaczek G, Saavedra-Molina A, Bacsí A, *et al.* Iron-mediated dismutation of superoxide anion augments antigen-induced allergic inflammation: effect of lactoferrin. *Postepy Hig Med Dosw (Online)* 2007; 61:268–276.
47. Chignard M, Balloy V. Neutrophil recruitment and increased permeability during acute lung injury induced by lipopolysaccharide. *Am J Physiol Lung Cell Mol Physiol* 2000; 279:L1083–1090.
48. Sibille Y, Marchandise FX. Pulmonary immune cells in health and disease: polymorphonuclear neutrophils. *Eur Respir J* 1993; 6:1529–1543.
49. Park SJ, Wiekowski MT, Lira SA, Mehrad B. Neutrophils regulate airway responses in a model of fungal allergic airways disease. *J Immunol* 2006; 176:2538–2545.
50. Voynow JA, Young LR, Wang Y, *et al.* Neutrophil elastase increases MUC5AC mRNA and protein expression in respiratory epithelial cells. *Am J Physiol* 1999; 276:L835–L843.
51. Lammernann T, Afonso PV, Angermann BR, *et al.* Neutrophil swarms require LTB4 and integrins at sites of cell death in vivo. *Nature* 2013; 498:371–375.
52. Wenzel SE, Trudeau JB, Kaminsky DA, *et al.* Effect of 5-lipoxygenase inhibition on bronchoconstriction and airway inflammation in nocturnal asthma. *Am J Respir Crit Care Med* 1995; 152:897–905.
53. Mizutani N, Nabe T, Yoshino S. IL-17A promotes the exacerbation of IL-33-induced airway hyperresponsiveness by enhancing neutrophilic inflammation via CXCR2 signaling in mice. *J Immunol* 2014; 192:1372–1384.
- The manuscript demonstrates that IL-33 cooperates with IL-17A to induce neutrophilic inflammation via CXCR2 signaling, and exacerbates airway hyperresponsiveness.
54. Nair P, Gaga M, Zervas E, *et al.* Safety and efficacy of a CXCR2 antagonist in patients with severe asthma and sputum neutrophils: a randomized, placebo-controlled clinical trial. *Clin Exp Allergy* 2012; 42:1097–1103.
55. Braber S, Overbeek SA, Koelink PJ, *et al.* CXCR2 antagonists block the N-AcP-GP-induced neutrophil influx in the airways of mice, but not the production of the chemokine CXCL1. *Eur J Pharmacol* 2011; 668:443–449.