

Allergic contact dermatitis from dithranol

VISHAL CHADHA AND SHRUTAKIRTHI D. SHENOI

Department of Skin & STD, Kasturba Medical College & Hospital, Manipal 576 119, India

Key words: allergic contact dermatitis; dithranol; anthralin; medicaments; psoriasis. © Munksgaard, 1999.

Dithranol (anthralin) is a synthetic derivative of a substance originally extracted from the araroba tree and is frequently used in the treatment of psoriasis (and alopecia areata). It is a known irritant and stains the skin.

Case Report

A 60-year-old man, with psoriasis vulgaris for the last 12 years, presented with an exacerbation. He was on methotrexate 2.5 mg, 3× weekly and to date had received 1210 mg of the drug. Bath water PUVA was commenced. For a few resistant plaques on the thighs, dithranol 0.01% cream was prescribed. 2 days later, erythema with vesiculation and exudation developed at the

sites treated. There was a past history of dithranol intolerance. Patch testing with various strengths of dithranol was done (Table 1). Patch testing with the same concentrations in 5 healthy volunteers was negative.

Discussion

Few cases of allergic contact dermatitis from dithranol have been reported (1–3). There is disagreement over its irritant threshold. Burden et al. (3) proposed 0.001% as the appropriate patch test concentration, while De Groot et al. (1) stated that 0.05% pet. gave sufficiently reliable results.

References

1. De Groot A C, Nater J P. Contact allergy to dithranol. *Contact Dermatitis* 1981; 7: 5–8.
2. Di Landro A, Valsecchi R, Cainelli T. Contact allergy to dithranol. *Contact Dermatitis* 1992; 26: 49–50.
3. Burden A D, Stapleton M, Beck M H. Dithranol allergy: fact or fiction? *Contact Dermatitis* 1992; 27: 291–293.

Table 1. Results of patch testing with dithranol

	0.01% pet.	0.005% pet.	0.0025% pet.
D2	++	+	+
D3	++	+	+

Active sensitization to sodium metabisulfite in hydrocortisone cream

SIAMAK HESHMATI AND HOWARD I. MAIBACH

Department of Dermatology, School of Medicine, University of California, San Francisco, Box 0989, Surge 110, San Francisco, CA 94143, USA

Key words: active sensitization; sodium metabisulfite; topical corticosteroids; medicaments; preservatives; patch testing technique; allergic contact dermatitis. © Munksgaard, 1999.

In performing assays for cumulative irritancy of topical corticosteroid formulations, we have actively sensitized 2 out of a panel of 50 to sodium metabisulfite. The formulations were 1% hydrocortisone (for over-the-counter distribution) in which metabisulfite was included at an unstated concentration. An edematous patch test response led to component testing: 2 subjects had a ++ response to sodium metabisulfite 1% pet., utilizing Finn Chambers (Epitest, Helsinki), the upper back and readings at D2 and D4.

Numerous adverse reactions have been attributed to sulfiting agents, including GI symptoms, urticaria, pruritus, angioedema, headache, asthma, and non-specific cutaneous lesions. Metabisulfites, along with other sulfiting agents, are used in the food and beverage industries as antioxidants and preservatives. They are also used in topical pharmaceuticals. The bisulfite of commerce consists chiefly of sodium metabisulfite, and for

all practical purposes, possesses the same properties as the true bisulfite (9). Intolerance reactions, of both immediate and delayed type, to sulfiting agents are increasingly being reported. Results of patch testing with various formulations containing sodium metabisulfite are given in Table 1.

Giorgini et al. (1) recently reported induction of contact dermatitis to sodium metabisulfite and Melcloderm cream. Doooms-Goosens et al. (2) found a strong positive reaction to sodium metabisulfite in 2 patients sensitive to ketoconazole cream, and concluded that there probably existed a reaction with features of both the immediate and delayed-type allergy to sodium metabisulfite. Vissers-Croughs et al. (8) patch tested a man with recurrent facial seborrhoeic dermatitis, who had developed an itchy papulovesicular dermatitis on treatment with ketoconazole cream. Patch tests with the control cream base, as well as sodium sulfite at the concentrations given in