

Do Urban Minority Parents and Children Agree on Asthma Symptoms with Exercise, Worries, and Confidence in Disease Management?

Florinda Islamovic BA , Ellen Johnson Silver PhD ,
Marina Reznik MD, MS

PII: S1876-2859(19)30133-0
DOI: <https://doi.org/10.1016/j.acap.2019.05.007>
Reference: ACAP 1350



To appear in: *Academic Pediatrics*

Received date: 3 January 2019
Accepted date: 11 May 2019

Please cite this article as: Florinda Islamovic BA , Ellen Johnson Silver PhD , Marina Reznik MD, MS , Do Urban Minority Parents and Children Agree on Asthma Symptoms with Exercise, Worries, and Confidence in Disease Management?, *Academic Pediatrics* (2019), doi: <https://doi.org/10.1016/j.acap.2019.05.007>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

TITLE PAGE

Title: Do Urban Minority Parents and Children Agree on Asthma Symptoms with Exercise, Worries, and Confidence in Disease Management?

Florinda Islamovic, BA¹

Ellen Johnson Silver, PhD¹

Marina Reznik, MD, MS¹

¹Department of Pediatrics, Division of Academic General Pediatrics, Children's Hospital at Montefiore, Albert Einstein College of Medicine

Florinda Islamovic is currently a medical student at St. Georges University School of Medicine

Word Count:

Abstract: 247

Main Text: 2,804

Key Words: caregivers, urban, pediatric, agreement, asthma

Funding: 5K23HD065742 (PI Reznik); P60 MD000514 for Spanish translation services of informed consent materials.

Running header: Parent and Child Agreement on Asthma Symptoms with Exercise

Conflicts of interest: The authors report no conflicts of interest. The authors alone are responsible for the content and writing of this paper.

Corresponding Author: Marina Reznik, MD, MS; Associate Professor of Pediatrics, Albert Einstein College of Medicine, Children's Hospital at Montefiore, 3411 Wayne Ave, 8th Floor, Bronx, NY 10467; Phone: 718-741-2494; Fax: 718-654-6692; Email: mreznik@montefiore.org

What's New:

Our results revealed parent-child disagreement regarding exercise-induced asthma symptoms, asthma-related worries, and confidence in disease management suggesting that including children's reports may help providers identify those needing enhanced asthma management and open conversations to address concerns.

ABSTRACT

Objective: Providers commonly rely on parent report of child's asthma symptoms. However, children as young as 7 years can report on their own health. Our study compared parent and child reports of asthma symptoms with exercise, worry about developing an asthma attack, and confidence in disease management.

Methods: Children aged 7-10 years with asthma and their parents were recruited from four Bronx, NY schools. Parents and children completed interview surveys with parallel questions regarding perceived asthma symptoms, asthma-related worries and confidence in disease management. We used McNemar's tests for paired data to compare parent and child responses.

Results: 105 parent-child pairs' data were analyzed. Mean child age was 8.5 yrs (SD .99); 53% were male, 82% were Hispanic. Children were more likely than their parents to report ever having an exercise-induced asthma attack (85% vs. 48%, $p<.001$) and that they "worry a lot" about developing an asthma attack during exercise (63% vs. 45%, $p=.01$). Children also felt more confident to use an inhaler correctly (76% children were "very sure" vs. 60% parents, $p=.009$) and to report having an inhaler available in case of an attack (38% children were "very sure" vs. 20% parents, $p=.003$).

Conclusions: In this school-based sample of urban children with asthma, we found disagreement in parent and child reports of symptoms with exercise, worry about developing an attack, and

confidence in medication use. These findings suggest that including child reports during history taking might help providers identify children in need of enhanced asthma management.

INTRODUCTION

Asthma is the most common pediatric chronic respiratory disease that affects approximately 7.0 million children in the United States.¹ The prevalence of asthma is higher in minority children living in inner cities, such as the Bronx, New York (NY), where 20-25% of schoolchildren have asthma.^{2,3} Many urban children with asthma experience poor symptom control and asthma management remains a challenge.

Children as young as 7 years of age are able to report on their health status.⁴ However, physicians often rely only on parent report to determine if the child experiences symptoms with exercise, especially for children younger than 12 years of age.⁵⁻⁷ Previous studies analyzing parent and child agreement on asthma diagnosis, asthma symptoms, and quality of life show significant differences in agreement.⁸⁻¹⁰ Whereas there is a high level of agreement between parents and children on the diagnosis of asthma, there are varying levels of agreement regarding reported asthma symptoms.^{9,11,12} Although parent report is important, parents can only provide an indirect account of children's experiences, which may lead to an underestimation of asthma symptoms and activity limitations. This in turn may lead to sub-optimal treatment plans, poor disease management, and decreased exercise.

Exercise is important for children with asthma and has been associated with decreased symptom severity, improved quality of life, and reduced school absenteeism.¹³⁻¹⁷ Prior studies have revealed that children are more likely than parents to report wheezing during or after exercise,¹⁸ but parents are less likely to report the impact of exercise on the child's

asthma.¹⁰ However, there is no literature comparing parent-child report on worry about developing an asthma attack with exercise or about using inhalers in front of others.

Asthma management often relies on many factors, including medication availability, proper medication use and self-efficacy. There are multiple barriers to asthma management, especially in Hispanic and other urban minority populations.¹⁹ Furthermore, asthma management is often a shared responsibility between parents and children. Previous studies examined parent report of medication adherence and have shown decreased adherence to medication use, especially in urban populations.¹⁹ However, there is little literature comparing parent and child report on medication use. A study comparing children 10 to 12 years of age, and their parents, showed a discrepancy on medication use.¹⁰ In prior studies examining the medication administration and asthma management self-efficacy of parents and adolescents, the parents reported a higher level of confidence and self-efficacy on medication administration and lower confidence with more complex tasks involving judgment and decision making.²⁰⁻²² However, there are no studies that have explored parent-child agreement on child confidence and self-efficacy in regards to either having asthma medication available during an attack or its proper use.

By obtaining both parent and child report, physicians may obtain a more complete picture of the frequency of asthma symptoms and medication use as well as the impact asthma has on children and their families. There is scant literature comparing urban minority parent and child reports about asthma symptoms during exercise, worry about having an asthma attack and medication use, or self-efficacy in inhaler use. Thus, the objective of our study was to compare these types of parent and child reports in a school-based sample of urban minority children with asthma and their parents.

METHODS

Participants and Settings

This study used cross-sectional baseline data from 7- to 10-year-old children who were enrolled in a school-based asthma intervention trial and from their parents. Prior to study enrollment, we identified children with asthma by conducting an asthma screening of all students attending 2nd through 4th grades in four Bronx, New York (NY) elementary schools.³ To determine asthma intervention trial eligibility, research assistants completed a telephone screening with parents or primary caregivers who indicated that their child had asthma on the screening survey and described the study. Research assistants then met with parent and child at school or home to obtain informed consent/assent and complete baseline survey. Families were eligible for participation if they had children 7-10 years of age with: (1) parent report of asthma diagnosis made by a physician; (2) prescription for asthma medication in the past 12 months; and (3) English or Spanish speaking caregiver. We will use the term parent and caregiver interchangeably in this report. Children with other chronic pulmonary diseases (e.g., cystic fibrosis, bronchopulmonary dysplasia) were excluded.

The study was approved by both the university's and the New York City Department of Education (NYC DOE) Institutional Review Boards. Written consent from caregivers and assent from children were obtained prior to study participation.

Measures

Child and Parent Surveys

All child and parent surveys were administered verbally, one-on-one by trained research assistants. Child surveys were conducted at the school and parent surveys (in English or Spanish)

were administered either at school or home. The child and parent surveys included parallel questions regarding perceived asthma symptoms with exercise and the child's competence in addressing them.

Survey questions were developed based on the results of a qualitative study that explored barriers to exercise in children with asthma in a school setting,²³ expert opinion and literature review. Cognitive interviews were conducted with three children to test child understanding of the questions and answers. Questions were then revised based on the results of cognitive interviews. Each child survey proceeded by first asking the child to define "asthma attack," "exercise," "asthma medicine pump," what it means "to worry," and "to be sure of something." Definitions of these terms were then read to the child. Asthma attack was defined as asthma symptoms (cough, wheezing, shortness of breath) experienced by children.

Survey Questions

Survey questions were adapted from several validated scales.²⁴⁻²⁶ Survey questions asked about the child ever having an asthma attack with exercise (yes/no); child's worries about developing an asthma attack during exercise and not having an asthma inhaler during an attack, as well as about asthma-related stigma (worry about peers seeing them use an inhaler and laughing or saying hurtful things) (7 questions); and child's self-efficacy or confidence in using an inhaler, inhaler availability in case of having an asthma attack during exercise and child's knowledge of which medication to use during an attack (7 questions).

Response categories for questions about the child's worries included "worry a lot," "worry somewhere in between," "worry a little" or "not at all". We dichotomized these responses into "high" ("worry a lot") and "low" ("worry somewhere in between," "worry a little" or "not at all"). Response categories for self-efficacy questions included "very sure," "a little bit sure" or

“not sure at all”. We dichotomized these responses into “very sure” or “not sure” (“a little bit sure”/“not sure at all”).

Demographic Data

The parents provided information about the children’s demographic characteristics, including age, gender, ethnicity (Hispanic, non-Hispanic) and race (African American, White, Unspecified/Other), duration of asthma diagnosis, medications prescribed by the child’s primary care physician, and insurance status (Medicaid/Not Medicaid) as well as several parental variables, including age, relationship to the child, ethnicity/race, whether the family was a single parent household, history of asthma and education (recorded as “less than high school,” “graduated high school,” “1-3 years of college,” and “4-year college degree or more”). Medication availability at school was assessed via NYC Department of Education Medication Administration Forms (MAF) which gives a permission to a school nurse to administer asthma medications to a student in school.²⁷

Statistical Analysis

The level of agreement between children and parents on each pair of items was assessed by the kappa coefficient, which is a statistic that indicates whether agreement is beyond the level of chance.²⁸ We used weighted kappa when the items being compared had more than two categories; weighted kappa accounts for relative agreement of categories on an ordinal scale by allocating less weight to the categories farther apart.^{28,29} Kappa values less than .20 are considered to indicate slight or poor agreement; .21 to .40 fair agreement; .41 to .60, moderate agreement; .61 to .80 substantial agreement; and more than .80, almost perfect agreement²⁹. We also used McNemar’s test for paired data to determine whether the child and caregiver responses differed significantly. Findings with p-values $\leq .05$ were considered as statistically significant.

Analyses were conducted using SPSS (IBM SPSS Statistics for Windows, Version 25.0.

Armonk, NY: IBM Corp).

RESULTS

Demographic Characteristics

To identify potential participants, we screened 1,270 students through caregiver-completed surveys sent home with students.²⁷ Of the 1,270 surveys, 769 (61%) were returned. Overall, 192 children (25%) had physician-diagnosed asthma and 158 met criteria and were eligible for enrollment. We enrolled 108 children and their parents (enrollment rate of 68%). Two children and their parents withdrew from the study prior to the baseline data collection. One child was discharged from the school prior to the baseline survey completion.

Survey responses were evaluated for 105 child and parent pairs for whom data were available. Table 1 provides the demographic characteristics of the participants. The mean age of children was 8.5 years (range, 7-11 years) with the average duration of asthma diagnosis of 6.4 years. Slightly over half were male (53.3%) and 81.9% were of Hispanic background. Over 90% of parents were mothers and 41.9% indicated having asthma themselves.

Child and Parent Agreement on Asthma Attack with Exercise

Children were significantly more likely than parents (84.6% vs. 48.1%) to report that the child ever had an asthma attack with exercise ($p < 0.001$). The kappa statistic ($\kappa = 0.03$) indicated “poor” agreement between children and parents on this item.

Child and Parent Agreement on Asthma-Related Worries

As shown in Table 2, there were significant differences between parents and children on four of seven questions that assessed perception of being worried about asthma. Children were more likely than their parents to report that they “worry a lot” about developing an asthma attack during exercise, about other children seeing them take medication, and about other children laughing or saying hurtful things. However, parents were more likely than children to say they “worry a lot” that the child would not stop wheezing when he/she exercises. Weighted kappa statistics indicated “poor” parent-child agreement for all questions pertaining to perception of risk (kappa coefficients ranged from -0.03 to 0.14).

Child and Parent Agreement on Child’s Asthma Self-Efficacy

Table 3 provides results on parent-child agreement on child’s self-efficacy in managing and preventing an asthma attack, including medication availability during an attack and confidence in using an inhaler. Children were more likely than their parents to say they were “very sure” that they would have an inhaler available in case of an asthma attack and know how to use it correctly. Compared to their parents, children also were more likely to feel “very sure” that they could slow down during exercise to prevent an attack and would know what to do during an attack when parents were not present with them. Parent-child responses did not differ on perceptions of the child’s ability to know which medication to use in case of an attack. This pattern did not change in comparisons of parent-child responses when children were prescribed both controller and rescue inhalers versus rescue inhaler only. Low kappa coefficients indicated poor agreement between parent and child responses for all 7 asthma self-efficacy questions (kappa coefficient ranged between -0.02 and 0.18).

DISCUSSION

Our study findings suggest disagreement between children with asthma and their parents on the topics of symptoms with exercise, feelings of worry, and confidence in managing and preventing asthma exacerbation. Children were more likely to report symptoms with exercise and greater worry about developing an asthma attack when they exercise, but parents were more worried that their children would not stop wheezing when they exercise. Children also expressed greater concern about stigmatization (feelings of worry that peers would see them use an inhaler and laugh or say hurtful things), but more confidence in their ability to manage and prevent an asthma attack.

In concordance with our findings, the literature suggests that the parents under-report their child's asthma symptoms with exercise.^{10,12,18,30} Children and teenagers have been found to report more symptoms with exercise, such as coughing, wheezing and breathing problems, than their parents report.^{10,12,18,30} Parent reports of child symptoms may vary because most school-aged children are away from their parents for many hours during the day. School-aged children spend approximately 6.5 hours in school each day away from their parents (US DOE) and 28% of NYC children spending additional time in afterschool programs.^{31,32} As part of curriculum, they participate in physical activities during gym class or recess which may induce an asthma exacerbation.³³ While parent report is important, our findings and that of prior research suggest that obtaining both parent and child report may provide a more complete picture of how disease impacts children and their families.³⁴

Due to amount of time spent in school, parents may not be aware of their child's symptoms, medication use while in school or child's relationships with peers. Our results indicate that children are more likely than their parents to report asthma symptoms with exercise and; thus, it would be logical that children feel more worried that they will develop symptoms

and experience fear of stigmatization when participating in sports alongside their healthy peers. This coincides with our findings that children report greater worry about stigmatization and are more likely to report these feelings than parents. Other studies have also shown that elementary school children and adolescents have reported feelings of worry and embarrassment with taking their medication in the presence of other children.^{35,36} The stigma and worry associated with asthma symptoms during exercise may lead to avoidance of physical activity. Based on NHLBI guidelines, it is important to promote exercise in children with asthma.³⁷ Physicians play an important role in promoting physical activity in this population by ensuring proper medication prescription (e.g. pre-exercise rescue inhaler), evaluating medication technique and adherence, as well as, providing education to parents and children. Obtaining report from children may shine light on this issue and allow parents and physicians the ability to address stigma thereby promoting exercise. However, there is a lack of studies comparing parent and child reports of worry about having an asthma attack and using asthma medications.

Although our study showed that children were more worried about issues related to their asthma, they were more confident about their abilities to manage symptoms, slow down during exercise to prevent exacerbation and to correctly use the inhaler than their parents felt they would be. Studies also have shown that parents and children separately feel confident in asthma inhaler technique as well as in managing asthma attacks.^{20,22} Also, separate studies show that there is a positive correlation between self-efficacy and asthma control in parents³⁸ and minority teenagers.³⁹ Previous studies of mainly White participants from non-US countries that examined self-efficacy in medication administration technique in school-age children^{21,22,40} and parents of school-aged children^{20,41} indicated poor inhaler technique in this population. A study of mainly Black and Hispanic teens and caregivers found that teens were less likely to identify correct

medication but there was no difference between teens and caregivers to describe correct indications for medication use.⁴² Based on these studies, while children may feel confident in their ability to identify and use their inhaler, it has been shown that their confidence may be misguided. This highlights the need to evaluate child confidence, beliefs and correlate it to self-efficacy of the child. It has been recommended that physicians increase communication directly with children about medications.⁴³ However, there is a lack of literature comparing inner-city, minority parent and child report on confidence in asthma management and medication administration.³⁸

Our study had some limitations. It was conducted in urban public elementary schools in the Bronx, NY, which is predominately Hispanic, thus, the conclusions may not be generalizable to non-minority children in other schools or in different communities. It is possible that some children may have attributed shortness of breath due to not being physically fit as asthma-related symptoms. This misconception may lead to an increase of affirmative responses regarding asthma symptom experience from children. Additionally, questions reflecting confidence or worry regarding asthma inhaler use or availability did not specify use of a spacer along with the inhaler. Finally, it is unclear if the children's or parents' perceptions are more accurate or whether the children's higher levels of confidence that we observed are warranted.

CONCLUSIONS:

Children were more likely than their parents to report that they had exercise-induced asthma symptoms and concerns about exacerbation during exercise. Parents felt less confident about their children using an inhaler correctly and about having an inhaler with them in case of an asthma attack. These findings suggest that including children's reports regarding symptoms with exercise and worries about embarrassment and assessing their inhaler technique during visits

might help providers to identify those in need of enhanced asthma management, address parent and child concerns, and open conversations about stigma. The differences in parent and child responses indicate that communication about asthma, particularly in the context of exercise, among the child, parent, and physician is warranted.

REFERENCES

1. Moorman JE AL, Baily CM, et al. National Surveillance of Asthma: United States 2001-20102012.
2. Webber MP, Carpinello KE, Oruwariye T, Lo Y, Burton WB, Appel DK. Burden of asthma in inner-city elementary schoolchildren: do school-based health centers make a difference? *Archives of pediatrics & adolescent medicine* 2003;157:125-9.
3. Reznik M, Bauman LJ, Okelo SO, Halterman JS. Asthma identification and medication administration forms in New York City schools. *Annals of allergy, asthma & immunology : official publication of the American College of Allergy, Asthma, & Immunology* 2015;114:67-8.e1.
4. Olson LM, Radecki L, Frintner MP, Weiss KB, Korfmacher J, Siegel RM. At what age can children report dependably on their asthma health status? *Pediatrics* 2007;119:e93-102.
5. Bates K, Elbers E, Meeuwesen L, Bensing J. Doctor-parent-child relationships: a 'pas de trois'. *Patient education and counseling* 2002;48:5-14.
6. Pantell RH, Stewart TJ, Dias JK, Wells P, Ross AW. Physician communication with children and parents. *Pediatrics* 1982;70:396-402.
7. Yawn BP, Brennum SK, Allen-Ramey FC, Cabana MD, Markson LE. Assessment of asthma severity and asthma control in children. *Pediatrics* 2006;118:322-9.
8. Varni JW, Thissen D, Stucky BD, et al. Item-level informant discrepancies between children and their parents on the PROMIS((R)) pediatric scales. *Quality of life research : an international journal of quality of life aspects of treatment, care and rehabilitation* 2015;24:1921-37.
9. Magzamen S, Mortimer KM, Davis A, Tager IB. School-based asthma surveillance: a comparison of student and parental report. *Pediatric allergy and immunology : official publication of the European Society of Pediatric Allergy and Immunology* 2005;16:669-78.
10. Davis KJ, Disantostefano R, Peden DB. Is Johnny wheezing? Parent-child agreement in the Childhood Asthma in America survey. *Pediatric allergy and immunology : official publication of the European Society of Pediatric Allergy and Immunology* 2011;22:31-5.
11. Wittich AR, Li Y, Gerald LB. Comparison of parent and student responses to asthma surveys: students grades 1-4 and their parents from an urban public school setting. *J Sch Health* 2006;76:236-40.
12. Yawn BP, Wollan P, Kurland M, Bertram S. Comparison of parent and student responses to asthma surveys: students grades 3-12 and their parents from a suburban private school setting. *J Sch Health* 2006;76:241-5.

13. Joschtel B, Gomersall SR, Tweedy S, Petsky H, Chang AB, Trost SG. Effects of exercise training on physical and psychosocial health in children with chronic respiratory disease: a systematic review and meta-analysis. *BMJ open sport & exercise medicine* 2018;4:e000409.
14. Fanelli A, Cabral AL, Neder JA, Martins MA, Carvalho CR. Exercise training on disease control and quality of life in asthmatic children. *Medicine and science in sports and exercise* 2007;39:1474-80.
15. Welsh L, Roberts RG, Kemp JG. Fitness and physical activity in children with asthma. *Sports medicine (Auckland, NZ)* 2004;34:861-70.
16. Welsh L, Kemp JG, Roberts RG. Effects of physical conditioning on children and adolescents with asthma. *Sports medicine (Auckland, NZ)* 2005;35:127-41.
17. Basaran S, Guler-Uysal F, Ergen N, Seydaoglu G, Bingol-Karakoc G, Ufuk Altintas D. Effects of physical exercise on quality of life, exercise capacity and pulmonary function in children with asthma. *Journal of rehabilitation medicine* 2006;38:130-5.
18. Hedman L, Lindgren B, Perzanowski M, Ronmark E. Agreement between parental and self-completed questionnaires about asthma in teenagers. *Pediatric allergy and immunology : official publication of the European Society of Pediatric Allergy and Immunology* 2005;16:176-81.
19. Rosser FJ, Forno E, Cooper PJ, Celedon JC. Asthma in Hispanics. An 8-year update. *Am J Respir Crit Care Med* 2014;189:1316-27.
20. Brown N, Gallagher R, Fowler C, Wales S. Asthma management self-efficacy in parents of primary school-age children. *Journal of child health care : for professionals working with children in the hospital and community* 2014;18:133-44.
21. Gillette C, Rockich-Winston N, Kuhn JA, Flesher S, Shepherd M. Inhaler Technique in Children With Asthma: A Systematic Review. *Academic pediatrics* 2016;16:605-15.
22. Alexander DS, Geryk L, Arrindell C, et al. Are children with asthma overconfident that they are using their inhalers correctly? *The Journal of asthma : official journal of the Association for the Care of Asthma* 2016;53:107-12.
23. Walker TJ, Reznik M. In-school asthma management and physical activity: children's perspectives. *J Asthma* 2014;51:1-6.
24. Mancuso CA, Sayles W, Allegrante JP. Development and testing of the Asthma Self-Management Questionnaire. *Annals of allergy, asthma & immunology : official publication of the American College of Allergy, Asthma, & Immunology* 2009;102:294-302.
25. Annesi JJ. Relations of age with changes in self-efficacy and physical self-concept in preadolescents participating in a physical activity intervention during afterschool care. *Percept Mot Skills* 2007;105:221-6.
26. Schutz RW, Smoll FL, Carre FA, Mosher RE. Inventories and Norms for Children's Attitudes Toward Physical Activity. *Res Q Exerc Sport*. 1985;56(3):256-265.138.
27. Reznik M, Bauman LJ, Okelo SO, Halterman JS. Asthma identification and medication administration forms in New York City schools. *Annals of allergy, asthma & immunology : official publication of the American College of Allergy, Asthma, & Immunology* 2015;114:67-8 e1.
28. Viera AJ, Garrett JM. Understanding interobserver agreement: the kappa statistic. *Family medicine* 2005;37:360-3.
29. Sim J, Wright CC. The kappa statistic in reliability studies: use, interpretation, and sample size requirements. *Physical therapy* 2005;85:257-68.

30. Lara M, Duan N, Sherbourne C, et al. Differences between child and parent reports of symptoms among Latino children with asthma. *Pediatrics* 1998;102:E68.
31. New York City After 3PM. at http://www.afterschoolalliance.org/documents/AA3PM_New_York_City_10202010.pdf.)
32. U.S Department of Education NCfES, Schools, and Staffing Survey (SASS). Public School Data File 2007-2008.
33. Storms WW. Review of exercise-induced asthma. *Med Sci Sports Exerc* 2003;35:1464-70.
34. Petsios K, Priftis KN, Tsoumakas C, et al. Level of parent-asthmatic child agreement on health-related quality of life. *J Asthma* 2011;48:286-97.
35. Walker TJ, Reznik M. In-school asthma management and physical activity: children's perspectives. *The Journal of asthma : official journal of the Association for the Care of Asthma* 2014;51:808-13.
36. Cohen R, Franco K, Motlow F, Reznik M, Ozuah PO. Perceptions and attitudes of adolescents with asthma. *The Journal of asthma : official journal of the Association for the Care of Asthma* 2003;40:207-11.
37. Health NIO. Expert Panel Report 3: Guidelines for the Diagnosis and Management of Asthma Summary Report 2007: National Heart, Lung and Blood Institute; 2007.
38. Gomes ALA, Lima KF, Mendes E, et al. Association of self-efficacy of parents/caregivers with childhood asthma control parameters. *Rev Esc Enferm USP* 2017;51:e03282.
39. Rhee H, Wicks MN, Dolgoff JS, Love TM, Harrington D. Cognitive factors predict medication adherence and asthma control in urban adolescents with asthma. *Patient Preference Adherence* 2018;12:929-37.
40. Burkhart PV, Rayens MK, Bowman RK. An evaluation of children's metered-dose inhaler technique for asthma medications. *The Nursing clinics of North America* 2005;40:167-82.
41. Aziz NA, Norzila MZ, Hamid MZ, Noorlaili MT. Skills amongst parents of children with asthma: a pilot interventional study in primary care setting. *The Medical journal of Malaysia* 2006;61:534-9.
42. Frey SM, Jones MR, Goldstein N, Riekert K, Fagnano M, Halterman JS. Knowledge of Inhaled Therapy and Responsibility for Asthma Management Among Young Teens With Uncontrolled Persistent Asthma. *Academic pediatrics* 2018;18:317-23.
43. De Maria C, Lussier MT, Bajcar J. What do children know about medications? A review of the literature to guide clinical practice. *Can Fam Physician* 2011;57:291-5.
44. National Heart, Lung, and Blood Institute. National Asthma Education and Prevention Program Expert Panel Report 3: Guidelines for the Diagnosis and Management of Asthma. U.S. Department of Health and Human Services. Full Report 2007, NIH publication 08-4051. Available at: <http://www.nhlbi.nih.gov/guidelines/asthma/asthgdln.htm> Accessed January 29, 2019.

Table 1. Child and Caregiver Demographic Characteristics (n=105 dyads)*

Characteristics	Number of subjects (%)
CHILD	
Child age, years (mean $\pm$ SD)	8.5 $\pm$.99
Sex	
Male	56 (53.3)
Ethnicity	
Hispanic	86 (81.9)
Non-Hispanic	19 (18.1)
Race	
African American	30 (28.6)
White	13 (12.4)
Unspecified/Other	62 (59.0)
Duration of asthma diagnosis, years (mean $\pm$ SD)	6.4 $\pm$ 2.4
Insurance	
Medicaid	90 (85.7)
Reported medication prescription**	
Rescue inhaler only	41 (39.0)
Controller + Rescue inhalers	60 (57.1)
Medication availability in school based on MAF forms***	24 (22.4)
Severity-Control Classification****	
Intermittent or Well Controlled	25 (23.8)
Mild- Moderate Persistent or Not Well Controlled	52 (49.5)
Severe Persistent or Poorly Controlled	28 (26.7)
CAREGIVER	
Caregiver age, years (mean $\pm$ SD)	36.2 $\pm$ 6.9
Relationship	
Mother	99 (93.9)
Ethnicity	
Hispanic	80 (76.2)
Non-Hispanic	25 (23.8)
Race	
African American	29 (27.6)
White	13 (12.4)
Unspecified/Other	63 (60.0)
Single parent household	59 (56.2)
History of Asthma Diagnosis	44 (41.9)

Caregiver education	
Less than high school	34 (32.4)
Graduated high school or GED	26 (24.8)
Some College (1-3 yrs)	36 (34.3)
College (4 yrs or more)	9 (8.6)

MAF, Medication Administration Form

* Of the 105 participants, 36 caregivers required surveys to be administered in Spanish

** Caregivers of 4 participants did not report rescue inhaler prescription

*** Data available on 107 students

*** As per NHLBI asthma diagnosis and management guidelines,⁴⁴ asthma severity was identified for children who were on rescue medication only and asthma control level was assessed for children who were prescribed preventive medications.

Table 2. Asthma-Related Worry

Question topics	Child question	Child response: "A Lot"	Caregiver question	Caregiver response: "A Lot"	McNemar P value
"Worry" about developing asthma attack during exercise	How worried are you that you will get asthma attack when you exercise?	62.5%	How worried are you that your child will get an asthma attack when [he/she] exercises?	45.2%	0.011
"Worry" about inhaler availability	How worried are you that you will need your asthma medicine pump but not have it with you?	44.8%	How worried are you that your child will need [his/her] asthma medication pump but not have it?	55.2%	0.161
"Worry" other children will see medication administration	How worried are you that other children will see you using your asthma medicine pump?	21.0%	How worried are you that other children will see your child using [his/her] asthma medication pump?	9.5%	0.029
"Worry" other children will laugh/say hurtful things	How worried are you that other children would laugh at you or say hurtful things?	64.4%	How worried are you that other children would laugh at [him/her] or say hurtful things to [him/her]?	41.3%	0.001

“Worry” about coughing during exercise	How worried are you that you will not stop coughing when you exercise?	44.2%	How worried are you that your child will not stop coughing when [he/she] exercises?	55.8%	0.134
“Worry” about wheezing during exercise	How worried are you that you will not stop wheezing or have noisy breathing in your chest when you exercise?	47.1%	How worried are you that your child will not stop wheezing or have noisy breathing in [his/her] chest when [he/she] exercises?	63.5%	0.033
“Worry” about risk of asthma attack when running up stairs	How worried are you that you will get asthma attack when you run upstairs?	46.7%	How worried are you that your child will get an asthma attack when [he/she] runs upstairs?	50.5%	0.659

Table 3: Asthma Self-Efficacy

Question topics	Child question	Child response: Very Sure	Caregiver question	Caregiver response: Very Sure	McNemar P value
Inhaler availability	Let's imagine you do exercise and you start having an asthma attack. How sure are you that you will have your asthma medicine pump with you?	38.1%	Let's imagine your child does exercise and [he/she] starts having an asthma attack. How sure you are that [he/she] will have [his/her] asthma medication pump with [him/her]?	20.0%	0.003
Inhaler use confidence	How sure are you that you can use your asthma medicine pump correctly?	76.2%	How sure are you that your child can use [his/her] asthma medication pump correctly?	60.0%	0.009
Medication knowledge	Let's imagine you are having an asthma attack. How sure are you that you will know which	41.3%	If your child is having an asthma attack. How sure are you that [he/she] will know	49.0%	0.291

	medicine to use?		which medication to use?		
Child asthma attack management when alone	Imagine you are having an asthma attack and your parents are not with you. How sure are you that you would know what to do?	48.5%	If your child has an asthma attack and you are not with [him/her], how sure are you that your child will know what to do?	30.1%	0.008
Child asthma attack prevention	Imagine you are having an asthma attack. How sure are you that you can keep your asthma from getting worse?	26.5%	If your child is having an asthma attack, how sure are you that your child can keep [his/her] asthma from getting worse?	17.6%	0.175
Child asthma attack management during exercise	Imagine you do exercise and you start having an asthma attack. How sure are you that you would know what to do?	48.5%	Let's imagine your child does exercise and [he/she] starts having an asthma attack. How sure are you that [he/she] will know what to do?	40.8%	0.302
Child asthma attack prevention during exercise	How sure are you that you can slow down when you exercise to prevent an asthma attack?	56.2%	How sure are you that your child will slow down when [he/she] exercises to prevent an asthma attack?	40.0%	0.030