




Available online at
ScienceDirect
www.sciencedirect.com

Elsevier Masson France
EM|consulte
www.em-consulte.com/en



Original article

Symptoms of allergic rhinitis in Parakou, Benin: Prevalence, severity and associated factors

M.-C. Flatin^{a,*}, S. Ade^a, S.-H.-R. Hounkpatin^a, B. Ametonou^a, U.-B. Vodouhe^b,
W. Adjibabi^b

^a Faculté de médecine, université de Parakou, BP 123, Parakou, Benin

^b Faculté des sciences de la santé, Université d'Abomey Calavi, Benin

ARTICLE INFO

Keywords:

Allergic rhinitis
Severity
Associated factors
Parakou

ABSTRACT

Aims: Allergic rhinitis (AR) is an extremely common disease. This study aimed to determine the prevalence and severity of AR symptoms and potential associated factors, in Parakou, a city in Benin.

Materials and methods: A cross-sectional study was conducted between April and July 2016, involving inhabitants of the city aged between 6 and 65 years. Cluster sampling was carried out. AR symptoms were defined as the simultaneous presence of rhinorrhea, nasal obstruction and sneezing without evidence of respiratory infection. Data were analyzed on EpiData Analysis v2.2.2.183 and then on R.3.2.1.

Results: Three hundred and ninety-five inhabitants were questioned, with a male/female ratio of 1.32, and median age of 19 years. One hundred and forty-one (35.7%) had AR symptoms, 47 (33.3%) had family history of atopy, and 87 (61.7%) had allergic conjunctivitis. The main triggers were house dust (139; 98.5%) and air pollution (111; 78.7%). On multivariate analysis, associated factors were age < 18 years ($P < 0.05$), exposure to house dust ($P < 0.001$), and personal history of urticaria ($P < 0.001$) or allergic conjunctivitis ($P < 0.01$). Twenty-eight subjects (19.9%) had persistent symptoms. Forty-three (30.5%) had moderate/severe symptoms, associated with age ≥ 18 years ($P < 0.01$), family history of atopy ($P < 0.01$), personal history of atopic eczema/dermatitis ($P < 0.01$), and presence of a dog or cat at home ($P = 0.01$).

Conclusion: The prevalence of AR symptoms in Parakou was high. A non-negligible proportion of symptoms were severe. This disease merits special attention in the city.

© 2017 Published by Elsevier Masson SAS.

1. Introduction

Allergic rhinitis (AR) is defined by the World Allergy Organization as the group of nasal symptoms resulting from an IgE-mediated immune reaction to allergen exposure. The main symptoms are rhinorrhea, nasal pruritus, nasal obstruction and sneezing fits, reversible spontaneously or under treatment [1]. It is the most frequent allergic manifestation, and is a public health issue. According to the World Health Organization, AR affects 10–30% of the general adult population and 40% of children [2]. More worryingly, prevalence is growing in most countries, as witnessed by the progression between phases I and III of the International Study of Asthma and Allergies in Childhood (ISAAC Study), in a mere 7 years on average [3].

AR can be a severe disease in that it induces psychological disorders such as fatigability, irritability, depression, sleep disorder and

sexual impairment, progressively isolating the patient socially and diminishing quality of life. The other non-negligible impact is economic, with both direct and indirect costs [2,4,5]. AR is a burden not only to the patient, but to the family and to society as a whole [4,5].

AR blights every part of the world, including sub-Saharan Africa, already ravaged by transmissible diseases [6]. Previous studies in Africa reported very varying prevalences: e.g., 11.4% in Yaoundé (Cameroon) and 29.6% in Ilorin (Nigeria) [7,8].

In Benin, in West Africa, there are no official data on AR. The aim of the present study was to assess AR symptoms in the city of Parakou. Objectives comprised:

- to determine the prevalence of AR symptoms in the population of Parakou;
- to identify associated factors;
- to classify patients according to symptom duration and severity;
- and to identify factors associated with severity.

* Corresponding author.
E-mail address: flatmar03@yahoo.fr (M.-C. Flatin).

2. Materials and method

2.1. Study type and duration

A cross-sectional study was conducted from April to July 2016.

2.2. Study context [9,10]

The study was performed in Parakou, the third largest city in Benin, in the north-central region, with an area of 441 km², and a population of 255,478. Parakou is a crossroads between Niger, Burkina-Faso and Mali inland, northern Nigeria to the east and northern Togo to the west. The climate is tropical, humid (south Soudan climate), with variations according to latitude, altitude and season. The humid southwest oceanic monsoon lasts from April to November, and the continental Harmattan, a dry wind, comes down from the northern Sahel between November and February, bringing orange ochre dust with it.

In health centers and hospitals, allergy treatment is basically clinical. Prick tests are not routinely used.

2.3. Study population

The study population was the inhabitants of Parakou.

2.3.1. Inclusion criteria

Inclusion criteria comprised:

- age 6 to 65 years
- and consent to participate (by parent or guardian for 6–18-year olds).

2.3.2. Sampling

2.3.2.1. Sample size. Sample size was calculated on Schwartz's formula $n = \frac{kZ_{\alpha}^2 p(1-p)}{i^2}$ (Schwartz D., *Méthodes statistiques à l'usage des médecins et biologistes*, ed. Flammarion Médecins Sciences, 1969), where n = sample size, p = 13.9% (prevalence reported in a similar study performed in D.R. Congo [11]), Z_{α} = 1.96, cluster effect k = 2, and precision i = 5%. The minimum sample size was thus 370, and a size of 395 was adopted to allow for 10% non-response.

2.3.2.2. Sampling technique. Thirty of the city's 41 administrative neighborhoods or villages were selected by clustering. Within each, the direction was chosen by an investigator standing in the center and throwing a pen in the air; in each direction, every second house was visited and in each house, every second subject was interviewed.

2.4. Data collection

Structured individual interviews were conducted to collect data on: sociodemographic characteristics; lifestyle; personal and/or familial history of bronchial asthma, allergic conjunctivitis, eczema or atopic dermatitis; otorhinolaryngologic symptoms; triggering factors according to the patient; main period of onset in the year; duration; and impact on everyday and occupational life. Subjects showing AR symptoms were then examined. Data were collected on a survey form filled out by the principal investigator.

2.5. Diagnostic criteria

The criteria were those used in the phase-I ISAAC study, with AR symptomatology defined as: simultaneous recurrent liquid rhinorrhea, nasal obstruction, sneezing fits and nasal pruritus, in the absence of respiratory infection [12].

Table 1

Characteristics of patients presenting symptoms of allergic rhinitis in Parakou, Benin, April–July 2016.

Characteristics	n
Sociodemographic data	
Male/female sex-ratio	1.2
Median age (IR) (years)	18 (12–24.5)
Median symptom duration (IR) (years)	4 (2–10)
History	
Familial atopy, n (%)	47 (35.1)
Personal history of allergic conjunctivitis, n (%)	87 (61.7)
Personal history of urticaria, n (%)	55 (39)
Personal history of bronchial asthma, n (%)	13 (9.2)
Personal history of eczema/atopic dermatitis, n (%)	13 (9.2)
Personal history of Quincke's edema, n (%)	1 (0.7)
Habits and lifestyle	
Active smoking, n (%)	2 (1.4)
Cat or dog at home, n (%)	45 (31.9)
Wood stove, n (%)	29 (20.6)
Triggers according to patient	
House dust, n (%)	139 (98.6)
Air pollution, n (%)	111 (78.7)
Grass pollen, n (%)	84 (59.6)
Mold, n (%)	72 (51.1)
Humidity, n (%)	31 (22)
Change of season, n (%)	23 (16.3)
Other ^a	11 (7.8)
ENT results	
Pale nasal mucosa, n (%)	137 (97.2)
Bilateral turbinate edema, n (%)	99 (70.2)
Pale soft palate, n (%)	23 (16.3)

^a Other triggers were: foods (peanuts), dog hair, cat hair, cereal dust, perfume.

Classification according to symptom duration and severity followed the Allergic Rhinitis and its Impact on Asthma (ARIA) guidelines:

- symptoms are “persistent” if occurring >4 days per week for >4 consecutive weeks, or otherwise “intermittent”;
- symptoms associated with abnormal sleep and/or impairment of daily activities, sport, leisure and/or impaired work and school and/or troublesome symptoms are “moderate to severe”, and those associated with normal sleep, no impairment of daily activities, sport, leisure, no impairment of work and school and no troublesome symptoms are “mild” [2,4].

2.6. Statistical analysis

Data were entered on EpiData Entry Client software v2.0.7.22 (EpiData Association, Odense, Denmark), and first analyzed on EpiData Analysis v2.2.2.183 (EpiData Association, Odense, Denmark). Frequencies and percentages were calculated for categorical variables, and medians for continuous variables with asymmetric distribution. Binary logistic regression was implemented on R.3.2.1 software to identify factors associated with AR symptoms and with symptom severity. The significance threshold was set at <5%.

3. Results

The male/female sex ratio in the 395 subjects was 1.32; median age was 19 years (interquartile range: 13–26 years; range: 6–64 years).

3.1. AR symptoms and associated factors

Based on the above criteria, 141 subjects (35.7%) showed AR symptoms: 38% of the 163 subjects aged under 18 and 34.1% of the 232 aged ≥ 18. Table 1 shows patient characteristics. Forty-seven (33.3%) had familial history of atopy and 87 (61.7%) personal history of allergic conjunctivitis. The most frequently claimed triggering

Table 2
Factors associated with allergic rhinitis symptoms in Parakou, Benin, April–July 2016.

Factors	Raw OR	Adj. OR	Adj. OR 95% CI	P-value
Age < 18 years	1.19	2.03	1.03–4.1	< 0.05
House dust exposure	199	239.6	48.1–1963	< 0.001
PH urticaria	6.42	3.40	1.53–8.15	< 0.001
PH allergic conjunctivitis	16.18	3.95	1.98–7.71	< 0.01

PH: personal history of; Adj. OR: adjusted odds ratio; Adj. OR 95%CI: 95% confidence interval of adjusted odds ratio.

factors were house dust (139; 98.5%) and air pollution (111; 78.7%). In all cases, symptoms worsened between October and February (Harmattan season). Examination mainly found pale nasal mucosa (97.2%) and bilateral turbinate edema (70.2%). Table 2 shows factors for AR symptoms on binary logistic regression: age < 18 years ($P < 0.05$), house dust exposure ($P < 0.001$), and personal history of urticaria ($P < 0.001$) or allergic conjunctivitis ($P < 0.01$).

3.2. Classification of AR symptoms and factors associated with severity

Twenty-eight of the 141 subjects with AR symptoms (19.9%) had persistent AR, and 43 (30.5%) had moderate to severe symptoms. Factors for severity are shown in Table 3: age > 18 years ($P < 0.01$), familial history of atopy ($P < 0.01$), personal history of eczema or atopic dermatitis ($P < 0.01$), and cat or dog at home ($P = 0.01$).

4. Discussion

This study, conducted in a small African city, found that 35.7% of the population showed AR symptoms. Prevalence in adults in the European Community Respiratory Health Survey varied widely, from 4% to 32%, between countries [13]. In the ISAAC study, which focused more on children, prevalence ranged from 0.8% to 14.9% in 6–7-year olds and from 1.4% to 39.7% in 13–14-year olds [12]. Prevalence increases with age, following the progression of atopy [14]. Prevalences as high as the present finding were reported by other African and non-African authors [8,15,16]. The lower prevalence of 10.4% reported by Pefura-Yone et al. in Cameroon may have been due to their use of sensitization screening by skin tests for airborne allergens [7]. At all events, these frequencies show that AR rates are high and require special attention.

House dust emerged as the main trigger in Parakou, and is known to be one of the most important factors for onset or aggravation of allergy symptoms in genetically predisposed subjects [2,4,17]. Parakou has a hot and often humid climate, providing good growth conditions for house mites, which are difficult and costly to eradicate [18]. Eradication requires a series of more or less complex environmental measures: controlled humidity, ventilation, removing carpeting and using anti-mite covers and miticides.

Air pollution was another trigger, and has been worsening in recent years with the increasing number of motor vehicles. More

Table 3
Factors associated with allergic rhinitis symptoms severity in Parakou, Benin, April–July 2016.

Factors	Raw OR	Adj. OR	Adj. OR 95% CI	P-value
Age ≥ 18 years	3.19	3.21	1.28–8.08	< 0.01
Familial atopy	2.66	3.67	1.56–8.63	< 0.01
PH eczema/atopic dermatitis	2.54	9.59	1.74–52.95	< 0.01
Cat or dog at home	2.43	3.16	1.30–7.72	0.01

PH: personal history of; Adj. OR: adjusted odds ratio; Adj. OR 95%CI: 95% confidence interval of adjusted odds ratio.

than three-quarters of patients were troubled by air pollution, in agreement with other reports [19,20].

Factors found to be associated with AR symptoms were: age < 18 years, house dust exposure, and personal history of urticaria or allergic conjunctivitis. Again, this is in agreement with other reports [4,17].

Symptoms were persistent in 19.9% of cases and moderate to severe in 30.5%, and notably impaired quality of life. Factors for severity comprised age ≥ 18 years, familial or personal history of atopy, and having a cat or dog at home. It is unsurprising that AR persisting from later childhood should be more severe in adulthood. While it is unfeasible to combat personal or familial atopy, removing or isolating domestic pets can be suggested, especially when symptoms are impairing quality of life and sensitization can be demonstrated.

The strong points of the present study were its prospective data collection and rigorous methodology in selecting subjects, allowing extrapolation to the population of Parakou as a whole. Also, the diagnostic criteria were those used in large-scale multi-center prevalence surveys such as the phase-I ISAAC study. However, the absence of prick tests for airborne allergens to refine diagnosis was a limitation, and mainly due to the lack of allergens in Benin; in the light of the present results, local production of allergens would be worth thinking about, especially as the allergen to be used in a given patient should take account of the specificities of circulating antigens, to avoid false positives.

5. Conclusion

AR symptoms were found in a third of the population of Parakou. They were notably triggered by mites, and aggravated by air pollution, which is worsening in the city. They were persistent in 1 in 5 patients and severe in one-third. Treatment strategies for this public health problem are needed, including local production of allergen extracts and measures to reduce allergen exposure, notably for house mites.

Disclosure of interest

The authors declare that they have no competing interest.

References

- [1] World Allergy Organization. Rhinitis: synopsis. Definition of allergic rhinitis [Available from <http://www.worldallergy.org/professional/allergic-iseases.center/rhinitis/rhinitissynopsis.php>; accessed March 2017].
- [2] Pawankar R, Canonica GW, Holgate ST, Lockey RF, Blaiss M. The WAO, white book on allergy (Update 2013). [Available from <http://www.worldallergy.org>; accessed March 2017].
- [3] Asher MI, Montefort S, Björkstén B, et al. Worldwide time trends in the prevalence of symptoms of asthma, allergic rhinoconjunctivitis, and eczema in childhood: ISAAC Phases One and Three repeat multicountry cross-sectional surveys. *Lancet* 2006;368:733–43.
- [4] Brozek JL, Bousquet J, Baena-Cagnani CE. Allergic rhinitis and its impact on asthma (ARIA) guidelines: 2010 revision. *J Allergy Clin Immunol* 2010;126:466–76.
- [5] Storms W, Meltzer EO, Nathan RA, Selner JC. The economic impact of allergic rhinitis. *J Allergy Clin Immunol* 1997;99:S820–4.
- [6] Mbugi EV, Chilongola JO. Allergic disorders in Africa and Africans: is it primarily a priority? *WAO J* 2010;3:175–81.
- [7] Pefura-Yone EW, Kengne AP, Balkissou AD, Boulelys-Nana JR, Efe-de-Melingui NR, Ndjoutecheu-Moualeu PI. Prevalence of asthma and allergic rhinitis among adults in Yaounde, Cameroon. *PLoS One* 2015;10(4):e0123099. <http://dx.doi.org/10.1371/journal.pone.0123099>.
- [8] Desalu OO, Salami AK, Iseh KR, Oluboyo PO. Prevalence of self-reported allergic rhinitis and its relationship with asthma among adult Nigerians. *J Invest Allergol Clin Immunol* 2009;19(6):474–80.
- [9] Annuaire Mairie. Ville de Parakou [Disponible sur : <http://www.annuaire-mairie.fr/ville-parakou.html>; accessible mars 2017].
- [10] Institut national de statistique et d'analyse économique (INSAE). Statistiques. Statistiques démographiques. Population. RGPH4 : Que retenir des effectifs de

- population en 2013 ? [Disponible <http://www.insae-bj.org/population.html>; accessible mars 2017].
- [11] Nyembue TD, Jorissen M, Hellings PW, et al. Prevalence and determinants of allergic disease in a Congolese population. *Int Forum Allergy Rhinol* 2012;2(4):285–93.
- [12] Strachan D, Sibbald B, Weiland S, et al. Worldwide variations in prevalence of symptoms of allergic rhinoconjunctivitis in children: the International Study of Asthma and Allergies in Childhood (ISAAC). *Pediatr Allergy Immunol* 1997;8:161–76.
- [13] Bousquet PJ, Leynaert B, Neukirch F, et al. Geographical distribution of atopic rhinitis in the European Community Respiratory Health Survey I. *Allergy* 2008;63(10):1301–9.
- [14] Bantz SK, Zhu Z, Zheng T. The atopic march: progression from atopic dermatitis to allergic rhinitis and asthma. *J Clin Cell Immunol* 2014;5(2):1–16, <http://dx.doi.org/10.4172/2155-9899.1000202>.
- [15] El Kettani S, Lotfi B, Aichane A. Prévalence de la rhinite allergique en milieu rural à Settat (Maroc). *East Mediterr Health* 2009;15(1):167–77.
- [16] Zhang Y, Zhang L. Prevalence of allergic rhinitis in China. *Allergy Asthma Immunol Res* 2014;6(2):105–13.
- [17] Ng CL, Wang DY. Latest developments in allergic rhinitis in allergy for clinicians and researchers. *Allergy* 2015;70:1521–30.
- [18] Martinot JB, Mercenier C. Mesures d'éviction des acariens [Disponible http://www.cmsenamur.be/files/files/mesures_d_eviction.pdf; accessible mars 2017].
- [19] Zhang F, Wang W, Lv J, Krafft T, Xu J. Time-series studies on air pollution and daily outpatient visits for allergic rhinitis in Beijing, China. *Sci Total Environ* 2011;409:2486–92.
- [20] Hwang BF, Jaakkola JJ, Lee YL, Lin YC, Guo Y-LL. Relation between air pollution and allergic rhinitis in Taiwanese school children. *Respir Res* 2006;7:23, <http://dx.doi.org/10.1186/1465-9921-7-23>.