



Behavioral interventions for asthma self-management in South Asian populations: a systematic review

Charlotte Lee, Emma Alexander, Rachel Lee, Nneka Okorochoa, Logan Manikam & Monica Lakhanpaul

To cite this article: Charlotte Lee, Emma Alexander, Rachel Lee, Nneka Okorochoa, Logan Manikam & Monica Lakhanpaul (2019): Behavioral interventions for asthma self-management in South Asian populations: a systematic review, Journal of Asthma, DOI: [10.1080/02770903.2019.1658209](https://doi.org/10.1080/02770903.2019.1658209)

To link to this article: <https://doi.org/10.1080/02770903.2019.1658209>

 View supplementary material 

 Published online: 18 Sep 2019.

 Submit your article to this journal 

 Article views: 20

 View related articles 

 View Crossmark data 

REVIEW



Behavioral interventions for asthma self-management in South Asian populations: a systematic review

Charlotte Lee, MSc^{a,b†} , Emma Alexander, MBBS^c , Rachel Lee, MBBS^{d*} , Nneka Okorocha, MSc^d, Logan Manikam, PhD^{e,f} , and Monica Lakhanpaul, MBBS, DM^{a,b} 

^aUCL Great Ormond Street Institute of Child Health, University College London, London, United Kingdom; ^bWhittington Health NHS Trust, London, United Kingdom; ^cSchool of Medical Education, King's College London GKT, London, United Kingdom; ^dInstitute of Medical and Biomedical Education, St George's, University of London, Cranmer Terrace, London, United Kingdom; ^eUCL Institute of Epidemiology and Health Care, University College London, London, United Kingdom; ^fAceso Global Health Consultants Limited, London, United Kingdom

ABSTRACT

Objective: Asthma outcomes are significantly worse for minority groups, including South Asians (SAs), in high-income settings. Despite this, comparatively few existing studies have focused on SAs when studying the effectiveness of behavioral interventions on asthma self-management, and no prior study has synthesized these findings. We review the effectiveness of behavioral interventions on asthma management in adults and children of SA origin across low- (LICs), middle- (MICs), and high-income countries (HICs).

Methods: Data sources included EMBASE, MEDLINE, Cochrane Library and Trial registries: WHO, ICTRP and ClinicalTrials.gov. Eligibility criteria: randomized controlled trials (RCTs), quasi-RCTs and non-RCTs (controlled before-after [CBA] studies), published in English, with no publication year or country restrictions in adults and children of South Asian origin. Exclusion criteria: those focusing solely on pharmacological interventions. Search terms were "asthma" and "South Asian".

Results: We included 33 studies, 27 from MICs and 6 from HICs (education [$n = 10$], self-management plans [$n = 6$], yoga/breathing exercises [$n = 10$]) organizational interventions [$n = 1$], diet therapy [$n = 1$] and combined interventions [$n = 5$]). Outcome measures included: blood biochemistry, lung function, healthcare utilization and quality of life. A meta-analysis was not performed due to significant study heterogeneity.

Conclusion: Behavioral interventions for asthma management in SAs are effective. Educational interventions that aim to optimize asthma knowledge, control, and inhaler technique, and yoga/breathing exercises are most effective for improved long-term outcomes in adults and children across LICs and MICs. Further research is needed to evaluate the effectiveness of all behavioral interventions for SAs in HICs to better inform current guidance by policy makers and health care providers.

ARTICLE HISTORY

Received 1 April 2019
Revised 5 August 2019
Accepted 17 August 2019

KEYWORDS

Asthma; Asian; South Asian; behavioral; effectiveness

Introduction

The global prevalence of asthma is 300 million and rising (1). As a significant source of morbidity and mortality, asthma represents a substantial burden for adults, children and health services. For example, in 2013 the total cost of asthma for the United States (U.S) was estimated to be \$81.9 billion, which included medical costs, absenteeism and mortality (2).

Asthma morbidity (3,4), under-recognized and uncontrolled symptoms, and hospital admissions for acute asthma (5,6) disproportionately affect South

Asian minority groups in high-income settings, compared with majority populations. For example, in a meta-analysis of ethnic variations in asthma outcomes in the United Kingdom (UK), South Asians had an odds ratio of 2.9 (95% CI 2.4–3.4) relative to white people for admission to hospital for asthma (3). Known exacerbating factors include inequitable healthcare access and alternative cultural understandings (7), and may contribute towards growing health inequalities.

Culturally appropriate behavioral interventions offer potential to address these barriers for optimal asthma

CONTACT Charlotte Lee  charlotte.lee.12@ucl.ac.uk  UCL Great Ormond Street Institute of Child Health at University College London, 30 Guilford Street, London WC1N 1EH, United Kingdom.

[†]Present address: Nuffield Department of Primary Care Health Sciences, University of Oxford, Oxford, Radcliffe Primary Care Building, Radcliffe Observatory Quarter, Woodstock Road, Oxford, United Kingdom.

*Current affiliation: Nottingham University Hospitals NHS Trust, Nottingham, United Kingdom.

 Supplemental data for this article is available online at <https://doi.org/10.1080/02770903.2019.1658209>.

© 2019 Taylor & Francis Group, LLC

self-management. Reviews assessing behavioral change techniques, irrespective of ethnicity, have been previously published (8). Yet there is limited evidence assessing the effectiveness of behavioral interventions for South Asians in high-income countries (HICs), or how successful behavioral interventions offered to South Asians in low (LICs) and middle-income countries (MICs) may be efficiently utilized in resource-stretched HIC settings. This is despite the inequities in outcomes for this group.

To address this knowledge gap, we undertook a systematic review to assess the effectiveness of behavioral interventions for South Asian adults and children across LICs, MICs and HICs. Our review may assist in the adoption or adaptation of culturally-sensitive, behavioral interventions for South Asian minority diasporas. Increased understanding of interventions that can encourage behavior change and improve agency may lead to reduced individual and societal inequities for this group.

Methods

The protocol for the systematic review is registered in the PROSPERO database and can be accessed with CRD Registration No: CRD42016033718.

Search question

- Patient or population: South Asian adults and children with asthma and/or caregivers of children with asthma in LICs, MICs or HICs
- Intervention: behavioral (i.e. non-pharmacological) interventions
- Comparator: routine standards of care
- Outcomes: all measures of asthma control

Eligibility criteria

We defined South Asians (hereafter SAs) as individuals who are native of the southern part of Asia (including India, Pakistan, Bangladesh, and other territories). LICs, MICs, and HICs were classified according to the World Bank assignment of the world's economies, which is based on gross national income (GNI) per capita (2019) (9). Studies were included if they met the following criteria:

- Participants: SA adults and children of all ages in LICs, MICs or HICs
- Language: studies published in English or with translation available
- Intervention: any behavioral intervention aimed at managing asthma

- Design: randomized controlled trials (RCTs), quasi-RCTs and non-RCTs (controlled before-after (CBA) studies)

Studies that solely focused on pharmacological interventions for asthma management were excluded, unless they assessed a behavioral element e.g. improving compliance towards pharmacological treatment using an educational program. We did not perform a meta-analysis of results owing to study heterogeneity in intervention delivery, statistical reporting and outcome. Outcome measures were not pre-specified. Instead, we identified and described interventions that emerged from individual full papers only.

Information sources

The following databases were searched by RL and NO supported by LM: MEDLINE, EMBASE and Cochrane Library. Trial registries WHO, ICTRP and ClinicalTrials.gov were also searched for completed and ongoing studies. Reference lists of all relevant articles were also hand-searched by RL and NO for possible additional publications, returning one eligible article. No year or country limits were set. Searches were conducted from February to May 2016 and updated by CL and EA between April and May 2018.

Search strategy

The search strategy comprised of the two overall terms “asthma” and “South Asian” (including terms specifying all major subgroups). “Southeast Asian” was also included to ensure a comprehensive search that may capture Sri Lanka, which is sometimes considered as part of Southeast Asia, as well as any maritime islands of Southeast Asia which are part of South Asian countries. The pilot search included the additional specific terms “Sri Lankan” and “Myanmar”. These terms did not yield additional results and hence were not included in the final search strategy. For example, the search strings used for MEDLINE were:

Term 1: Asthma

Asthma OR asthma* OR abnormal respiratory sound OR respiratory sound* AND

Term 2: South Asians

India OR india OR Pakistan OR pakistan* OR Bangladesh OR bangladesh* OR punjab* OR gujurat OR South Asia OR south asia* OR south* asia* OR Southeast Asia OR southeast asia* OR south-east asia*




Figure 1. Study selection flow diagram according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines.

OR indian subcontinent OR indian sub-continent OR indian subcontinent

Study selection and data extraction

The study selection process is denoted in Figure 1, which provides a summary of all articles shortlisted and excluded at each stage of the review. In total, 5896

articles were identified in the initial search. RL and NO screened titles and abstracts, excluding 22 duplicates and a further 5494 articles due to ineligibility in line with the inclusion and exclusion criteria. This left 378 potentially eligible articles, which underwent full-text review. Conflicts were resolved via discussion with LM. One additional article was included from hand searches of reference lists. After full-text review, 348 articles were

excluded leaving 31 articles to be quality assessed by RL and NO. The most common reason for exclusion at this stage was that the articles were not evaluating interventions. After quality assessment, one study was excluded due to high risk of bias (10) leaving 30 studies. In 2018, 428 title and abstracts were screened, leaving 17 potentially eligible articles shortlisted full-text review, and three additional articles to include for data extraction and quality assessment. Overall, 33 articles were deemed suitable for inclusion in the review.

Data was extracted in the initial review by RL using a piloted modified worksheet that included: study type; country; inclusion and exclusion criteria; participant age and ethnicity; numbers recruited, lost to follow up and analyzed; intervention design and delivery method; control; sampling; randomization; blinding; method of data collection; study duration or follow up; outcomes measured; factors contributing to intervention engagement and acceptability; theoretical models; data collection, statistical methods and results. NO checked each extraction. In the 2018 update, data was extracted using the modified worksheet and quality assessed by CL and checked by EA.

Risk of bias assessment

Risk of bias in both RCTs and non-randomized studies was measured using the relevant 'Risk of Bias' tools developed by Cochrane (11). Six components of RCT studies were assessed as either low, high or unclear risk of bias: (1) random sequence generation, (2) concealment of allocation, (3) blinding, (4) incomplete outcome data, (5) selective outcome reporting and (6) additional sources of bias. Components (1) and (2) accounted for sequence bias, (3) encompassed performance and detection bias, (4) attrition bias, and finally (5) covered reporting bias.

For non-randomized studies, five components were assessed as either low, moderate, severe, or critical: (1) bias in measurement of interventions, (2) bias due to departures from intended interventions, (3) bias due to missing data, (4) bias in measurement of outcomes, (5) bias in selection of the reported result. Overall bias in both RCTs and non-randomized studies was then determined based on the average weightings of each component. RCTs with an overall high risk of bias and non-randomized studies with a critical risk of bias was excluded from our analyses (11,12).

Results synthesis

In view of the heterogeneity across methods, participants, interventions and outcomes, we narratively

synthesized results. We use guidance developed from the University of York Center for Reviews and Dissemination (CRD) and the Economic and Social Research Council (ESRC).

Results

Study characteristics

Of the 33 studies included, 27 were based in MICs ($n = 25$ – India (13–37), $n = 1$ – Pakistan (38), $n = 1$ – Malaysia (39)), and 6 in HICs ($n = 2$ – Canada (40,41), $n = 3$ – UK (42–44), $n = 1$ – Singapore (45). [Supplementary File S1](#) denotes a summary of all studies included.

For RCTs, overall risk of bias was judged as low for eight studies (15,17,22,30,39,41,44), moderate for one (16) and unclear for 17 (14,18–21,24,27,28,32–38,40,42). For non-randomized studies, overall risk of bias was judged as moderate for two studies (25,29), and severe for five studies (13,23,26,31,45).

Interventions were grouped into the following: educational or counseling interventions ($n = 10$), self-management plans ($n = 6$), yoga therapy or breathing exercises ($n = 10$), organizational interventions ($n = 1$), diet therapy ($n = 1$), combined educational intervention and self-management plan ($n = 1$); combined self-management plan and written asthma action plan ($n = 1$), combined self-management plan and cognitive behavioral therapy (CBT; $n = 1$), combined nature care, diet and yoga therapy ($n = 2$). Outcome measures identified included pharmacological and blood biochemistry, lung function, healthcare utilization (i.e. hospital visits, work or school days missed) and qualitative measures (i.e. quality of life; QoL). [Supplementary File S2](#) denotes a results summary of all studies included.

Results of studies assessing individual interventions

Educational/counseling interventions

Significant improvements on inhaler technique performance scores. Two MIC-based studies in India used written instructions (23) or educational booklets (26), although both were judged at severe risk of bias. One further India-based study utilized personalized health education (25), judged at moderate risk of bias. All three studies reported significant improvements on inhaler technique performance scores. Mishra et al. (28) offered an education program and reported improvements across frequency of attacks and hospitalization. These authors note that booklets written in culturally

appropriate languages (i.e. Hindi) may aid participant's acceptability of educational interventions and that educational interventions in combination with inhaler technique demonstration may improve patient understanding.

Two large HIC-based RCTs, both conducted in Canada (97 and 87 total participants), made use of either a physician-led knowledge- or a patient-generated community video (40,41) – unclear and low risk of bias, respectively. Pourseslami et al. (41) reported a significant percentage change in correct inhaler use in groups who watched a physician-led video (52% change 3 months' post intervention, $p \leq 0.001$). However, this did not occur for groups who watched the patient-generated video or both, compared with controls who received standard care. Two other larger HIC-based RCTs conducted in the UK (42,43) (342 participants and 105 eligible general practices, respectively) found no significant improvements in asthma symptoms, unscheduled care or long-term improvements in QoL. The risk of bias in these two studies was judged as low.

Self-management plans

Three large RCTs in India (18–20), all at unclear risk of bias, found self-management training and self-care manuals as effective in asthma management across Indian adults. Behera et al. (18) provided self-care manuals and reported improved symptom scores, adopted self-care measures during acute attacks, and patient knowledge of asthma and its precipitating factors at all three follow-ups. Behera et al. (19) later reported self-care manuals which included ways to avoid asthma triggers were associated with an increased number of patients avoiding triggers. In this study, the percentage of patients who started avoiding cigarette smoke increased from 90.7% at first visit, 94% (2 weeks follow-up), to 100% (6 months and 1 year follow-up; $\chi^2=16.53$, $p \leq 0.001$). The corresponding figures for the control group were 95.6%, 96.6% and 98.4% ($\chi^2=2.45$, $p \geq 0.05$). This trend was similar for other triggers listed including dust, changes in weather, exercise and emotions. Another study conducted in Singapore, a HIC, reported no change in self-management (45), although this study was judged at severe risk of bias. No change in asthma exacerbations or QoL score were reported in children with asthma and their caregivers in the MIC Malaysia. Of note, the outcomes in most of these studies (18–20,39,45) were measured via self-reports.

Yoga therapy/breathing exercises

Three MIC-based RCTs (24,27,35), all at unclear risk, assessed the effectiveness of yoga training in Indian adults. These authors reported significant improvement in patients' forced expiratory volume (FEV1), forced vital capacity (FVC), FEV1/FVC ratio, peak expiratory flow rate (PEFR), chest expansion, 40 mmHg endurance test and respiratory rate. Similar results were reported in other MIC-based RCTs that assessed the effectiveness of yoga training in Indian children (16,17) – moderate and low risk respectively. Agnihotri et al. (15) found that hemoglobin levels increased by 7.52% in the case group after 6 months, compared to a 3.97% increase in the control group who received standard medical treatment. Long-term improvements in FVC, FEV1 and QoL have been reported in RCTs across India (34,37) – both unclear risk. Sodhi et al. (36) also report improvements in the number and severity of attacks – although this study was judged at severe risk of bias. Finally, Singh et al. (33) promoted breathing exercises and reported improvements in PEFR. Results suggest yoga and breathing therapy may be effective in improving asthma related symptoms practiced as an adjuvant therapy with standard medical treatment (14,33). However, the long-term benefits and cost-effectiveness remain to be fully explored.

Organizational/dietary interventions/cognitive behavioral therapy

One study (38) conducted in Pakistan involved 428 adults receiving an organization-wide intervention that included improved availability of context-sensitive guidelines and case management materials. Results from this study are yet to be released. Only one small study judged at severe risk (13) conducted an elimination diet based intervention. Significant improvements in IgE titers in 17 children were observed. However, there was no improvement for 25% of children observed and a negative difference in one child compared to the control group. One study (21) that used CBT found significant improvement in PEFR, asthma symptoms and QoL, compared with standard pharmacological treatment.

Results of studies assessing mixed interventions

Self-management plan and written asthma action plan or educational intervention and medication counseling

Mixed interventions involved a combination of educational and/or self-management plans with any other

behavioral approach e.g. diet or yoga/breathing exercises. One UK-based RCT (44) combined an educational intervention with individually tailored written self-management plans. This was delivered to 593 participants aged 11–59 who were of White European (WE) or Indian Sub-continent (ISC) ethnic groups in the UK. Significant improvements in hospital admissions, out of hour's home visits by GPs or deputizing services, number of GP consultations, steroids and antibiotics prescription and QoL across all participants were reported. Another RCT (32) based in India and judged at unclear risk combined education with a care diary and medication counseling and found improved outcomes in PEFR, asthma control scores and QoL. Although combined interventions may have their place in improving QoL, there appears limited long-term or cost benefit to adults at risk of asthma morbidity such as SAs.

Diet therapy, yoga therapy and nature care treatment

Two MIC-based CBAs assessed the effects of diet therapy, nature care treatment and yoga therapy on a small sample of Indian adults (29,31). Although one of these studies (31) was judged at severe risk of bias, both demonstrated a significant improvement in lung parameters such as FVC, FEV1, PEF and MVV. No significant changes in hemoglobin, white blood cell count, and red blood cell count measurements were reported (29).

Discussion

To our knowledge, this is the first systematic review to assess the effectiveness of behavioral interventions in SA adults and children across LICs, MICs and HICs. The majority of studies were conducted in the MIC India and used educational interventions, self-management plans and yoga/breathing exercises. These interventions generally improved asthma symptoms, knowledge, and QoL. In addition, educational interventions were most effective for long-term improved patient outcomes. At present, there appear no individual or cost benefit in diet elimination, CBT or combined interventions. Only six studies across three HICs were relevant to our review. There also exist no high-quality studies across the U.S. that evaluate the effectiveness of behavioral interventions for asthma management in SAs. This is of particular concern given that Indians (3.8 million; 14.1%) are one of the largest ethnic groups across the U.S and rising (46). Nonetheless, our results support the U.S

Department of Health and Human Services guidelines that encourage policy makers and health care providers to support provision of educational asthma self-management programs.

Implications of key findings

Behavioral management interventions, in some cases, may have other benefits for health. Educational interventions and self-care plans allow patients to avoid adverse side effects from typical pharmacological treatment including tremors and tachycardia induced by salbutamol inhalers (47) and oral candidiasis and dysphonia from steroid inhalers (48,49). As such, behavioral interventions may be used either in refractory cases of asthma or as an adjunct to current treatment.

The improvements observed may be attributed to the co-development of plans by patients and healthcare professionals, which involve personalization to the patient's needs as well as cultural beliefs. Some studies (40,41) suggested simple, culturally and linguistically appropriate media may make interventions more widely accepted by and accessible to participants. For instance, dietary advice is not advised by the U.S Department of Health and Human Sciences guidelines for asthma management (50). Yet, beliefs and practices concerning food and asthma are prevalent amongst SA communities (6). Personalized interventions that tackle these unhelpful beliefs and practices as well as stigma may be offered to minority groups to better improve their asthma self-management (35). It is concerning that for many of the reviewed studies, the quality appraisal process indicated an unclear risk or severe risk of bias. This indicates that SAs with asthma are not only under-researched but are being left out of high-quality studies with strong reporting standards.

Structural health reforms offer new opportunities to incorporate new and re-design existing services and to reduce health inequalities. Interventions aimed at improving knowledge alone are unlikely to lead to practice change. Rather, interventions that combine behavioral change techniques and involve active participant co-development will be important for long-term asthma management. Adopting patient-focused approaches may be necessary for upstream empowerment and improvements in asthma management for minority groups at high-risk of asthma morbidity such as SAs.

Study limitations

Our study has a number of limitations. Our data sources were confined to MEDLINE, EMBASE,

Cochrane Library and trial registries only. We excluded papers not published in English, although valuable studies may have been published in other languages. We excluded behavioral interventions targeted at other ethnic minority groups, which may have allowed direct comparison of non-pharmacological interventions across ethnic groups. Only six out of 33 studies included SA participants in HICs with the rest focusing on SA participants across MICs, and no studies conducted in LICs. This limits the generalizability of our findings particularly towards SAs living in LICs and partially towards SAs living in HICs. It is possible that individuals between LICs and MICs, and even between lower- and upper-middle income countries, experience different issues around asthma recognition and health service access. Furthermore, we focused on a select number of countries to refine the scope of our review, which may have limited the number of potentially eligible articles.

A further limitation is that many of the studies reported statistically significant results, but publication bias is likely to be present. Most studies did not use standardized measures or explain which aspects of their interventions made them more accepted by participants. There were some mixed interventions delivered to either adults or children, making it difficult to distinguish if any of the interventions worked separately for each of the age groups. Future longitudinal studies investigating specific features of successful behavioral and that use validated measures are now paramount.

Acknowledgements

The authors would like to thank Mr. Virad Kisan (Medical Student, University College London) for his assistance in the development of this systematic review.

Declarations

Ethics approval and consent to participate: not applicable.

Consent to publish

Not applicable.

Availability of data and materials

All data generated and/or analyzed during this study are included in this published article (and its [supplementary information](#) files).

Disclosure statement

The authors declare that they have no conflicts of interests.

Funding

This research received no specific grant from any funding agency, commercial or not-for-profit sectors. ML and CL were (in part) supported by the National Institute for Health Research (NIHR) Collaboration for Leadership in Applied Health Research and Care (CLAHRC) North Thames at Bart's Health National Health Service (NHS) Trust. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health and Social Care.

Author contributions

LM and ML conceived and participated in the design of the study. LM coordinated and RL and NO undertook the review. CL and EA coordinated and performed updated searches. All authors performed the data interpretation. CL and EA wrote the final manuscript. All authors read and approved the final manuscript. LM and ML are the study guarantors.

ORCID

Charlotte Lee  <http://orcid.org/0000-0001-8252-9538>
 Emma Alexander  <http://orcid.org/0000-0001-7063-6846>
 Rachel Lee  <http://orcid.org/0000-0003-4612-2786>
 Logan Manikam  <http://orcid.org/0000-0001-5288-3325>
 Monica Lakhanpaul  <http://orcid.org/0000-0002-9855-2043>

References

1. The Global Asthma Report (2014). Auckland, New Zealand: Global Asthma Network. Available from: http://www.globalasthmanetwork.org/publications/Global_Asthma_Report_2014.pdf [last accessed 30 August 2019].
2. Nurmagambetov T, Kuwahara R, Garbe P. The economic burden of asthma in the United States, 2008–2013. *Ann Am Thorac Soc* 2018;15:348–356. doi:10.1513/AnnalsATS.201703-259OC.
3. Netuveli G, Hurwitz B, Levy M, Fletcher M, Barnes G, Durham SR, Sheikh A. Ethnic variations in UK asthma frequency, morbidity, and health-service use: a systematic review and meta-analysis. *Lancet* 2005;365:312–317. doi:10.1016/S0140-6736(05)17785-X.
4. DePalo V, Mayo PH, Friedman P, Rosen MJ. Demographic influences on asthma hospital admission rates in New York City. *Chest* 1994;106:447–451. doi:10.1378/chest.106.2.447.
5. Kuehni CE, Strippoli M-PF, Low N, Brooke AM, Silverman M. Wheeze and asthma prevalence and related health-service use in white and South Asian pre-schoolchildren in the United Kingdom. *Clin Exp Allergy* 2007;37:1738–1746. doi:10.1111/j.1365-2222.2007.02784.x.

6. Lakhanpaul L, Bird D, Manikam L, Culley L, Perkins G, Hudson N, Wilson J, Johnson M. A systematic review of explanatory factors of barriers and facilitators to improving asthma management in South Asian children. *BMC Public Health* 2014;14:403. doi:10.1186/1471-2458-14-403.
7. Smeeton NC, Rona RJ, Gregory J, White P, Morgan M. Parental attitudes towards the management of asthma in ethnic minorities. *Arch Dis Child* 2007;92:1082–1087. doi:10.1136/adc.2006.112037.
8. Denford S, Taylor RS, Campbell JL, Greaves CJ. Effective behavior change techniques in asthma self-care interventions: systematic review and meta-regression. *Health Psychol* 2013. [PubMed: 23815765]. doi:10.1037/a0033080.
9. World Bank. World Development Indicators. The World Bank Group. Available from: <https://www.worldbank.org> [last accessed 13th July 2019].
10. Saxena T, Saxena M. The effect of various breathing exercises (pranayama) in patients with bronchial asthma of mild to moderate severity. *Int J Yoga* 2009; 2:22–25. doi:10.4103/0973-6131.53838.
11. Higgins JPT, Green S (editors). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 [updated March 2011]. The Cochrane Collaboration 2011. Available from: <http://www.handbook.cochrane.org> [last accessed 20 July 2016].
12. Sterne JAC, Higgins JPT, Reeves BC, On Behalf of the Development Group for ROBINS-I. ROBINS-I: A tool for assessing risk of bias in non-randomized studies of interventions, Version 7 March 2016. Available from <http://www.riskofbias.info> [last accessed 20 July 2016].
13. Agarkhedkar SR, Bapat HB, Bapat BN. Avoidance of food allergens in childhood asthma. *Indian Pediatr* 2005;42:362–366.
14. Aggarwal T, Khatri A, Siddiqui SS, Hasan SN, Singh D, Kulshreshtha M, Agarwal S. Pranayama has additive beneficial effects along with medication in bronchial asthma patients. *J Phys Pharm Adv* 2013;3: 292–297.
15. Agnihotri S, Kant S, Kumar S, Mishra RK, Mishra SD. Impact of yoga on biochemical profile of asthmatics: a randomized controlled study. *Int J Yoga* 2014;7:17–21. doi:10.4103/0973-6131.123473.
16. Agnihotri S, Kant S, Kumar S, Mishra RK. The assessment of effects of yoga on pulmonary functions in asthmatic patients: a randomized controlled study. *J Med Soc* 2016;30:98–102. doi:10.4103/0972-4958.182909.
17. Agrawal SK, Singh M, Mathew JL, Malhi P. Efficacy of an individualized written home-management plan in the control of moderate persistent asthma: a randomized, controlled trial. *Acta Paediatr* 2007;94: 1742–1746. doi:10.1080/08035250510039973.
18. Behera D, Kaur S, Gupta D, Verma SK. Evaluation of self-care manual in bronchial asthma. *Indian J Chest Dis Allied Sci* 2006;48:43–48.
19. Behera D, Kaur S, Gupta D, Verma SK. Avoidance of allergens by the patients with bronchial asthma. *J Assoc Phys India* 2008;56:325–328.
20. Ghosh CS, Ravindran P, Joshi M, Stearns SC. Reductions in hospital use from self management training for chronic asthmatics. *Soc Sci Med* 1998;46: 1087–1093. doi:10.1016/s0277-9536(97)10047-8.
21. Grover N, Kumaraiah V, Prasadrao PSDV, D'souza G. Cognitive behavioral intervention in bronchial asthma. *J Asso Phys India* 2002;50:896–900.
22. Grover C, Goel N, Armour C, Van Asperen PP, Gaur SN, Moles RJ, Saini B. Medication education program for Indian children with asthma: a feasibility study. *Niger J Clin Pract* 2016;19:76–84. doi:10.4103/1119-3077.173716.
23. Jolly GP, Mohan A, Guleria R, Poulouse R, George J. Evaluation of metered dose inhaler use technique and response to educational training. *Indian J Chest Dis Allied Sci* 2015;57:17–20.
24. Mandanmohan JL, Udupa K, Bhavanani AB. Effect of yoga training on handgrip, respiratory pressures and pulmonary function. *Indian J Physiol Pharmacol* 2003;47:387–392.
25. Maazuddin M, Hazari MAH, Hasan A, Javeedullah M, Nasir Mohiuddin M, Misba A, Baig M. Training in the use of inhalation devices and respiratory improvement in asthma and COPD patients: a before–after experimental study. *Pharmanest* 2014;5: 2314–2320.
26. Kumar A, Singh NV, Kaur S, Singh A. Skill development regarding use of metered dose inhaler (MDI) amongst school teachers of Northern India. *Indian J Pediatr* 2013;80:903–906. doi:10.1007/s12098-013-1075-y.
27. Lathadevi GV, Uma Maheswari T. Evaluation of pulmonary functions in asthmatics after six weeks of ujjayi pranayama and shavasana training. *Biomedicine* 2012;32:52–56.
28. Mishra Narayana KVR, Rao S, Padhi K. Asthma education for better compliance in disease management. *Indian J Allergy Asthma Immunol* 2005;19:25–28.
29. Rao YC, Kadam A, Jagannathan A, Babina N, Rao R, Nagendra HR. Efficacy of naturopathy and yoga in bronchial asthma. *Indian J Physiol Pharmacol* 2014; 58:233–239.
30. Saji MS, Ja Jiju AK, Simdaran S. Study on the impact of patient counseling on the quality of life and pulmonary function of asthmatic patient. *Int J Pharm Pharm Sci* 2012;4:300–304.
31. Sathyaprabha TN, Murthy H, Murthy BT. Efficacy of naturopathy and yoga in bronchial asthma – a self-controlled matched scientific study. *Indian J Physiol Pharmacol* 2001;45:80–86.
32. Shanmugam S, Varughese J, Nair MAS, Balasubramanian R, Velu S, Bhojan C, Devarajan V, Sabzghabae AM. Pharmaceutical care for asthma patients: a developing country's experience. *J Res Pharm Pract* 2012;1:66–71. doi:10.4103/2279-042X.108373.
33. Singh V, Wisniewski A, Britton J, Tattersfield A. Effect of yoga breathing exercises (pranayama) on airway reactivity in subjects with asthma. *Lancet* 1990; 335:1381–1383. doi:10.1016/0140-6736(90)91254-8.
34. Singh S, Soni R, Singh KP, Tandon OP. Effect of yoga practices on pulmonary function tests including

- transfer factor of lung for carbon monoxide (TLCO) in asthma patients. *Indian J Physiol Pharmacol* 2012; 56:63–68.
35. Sodhi C, Singh S, Dandona PK. A study of the effect of yoga training on pulmonary functions in patients with bronchial asthma. *Natl J Physiol Pharm Pharmacol* 2009;59:169–174. doi:[10.5455/njppp.2018.8.1033009112018](https://doi.org/10.5455/njppp.2018.8.1033009112018).
 36. Sodhi C, Singh S, Bery A. Assessment of the quality of life in patients with bronchial asthma, before and after yoga: a randomized trial. *Iran J Allergy Asthma Immunol* 2014;13:55–60.
 37. Vempati R, Bijlani RL, Deepak KK. The efficacy of a comprehensive lifestyle modification programme based on yoga in the management of bronchial asthma: a randomized controlled trial. *BMC Pulm Med* 2009;9:37. doi:[10.1186/1471-2466-9-37](https://doi.org/10.1186/1471-2466-9-37).
 38. Khan MA, Ahmed M, Anil S, Walley J. Strengthening the delivery of asthma and chronic obstructive pulmonary disease care at primary health-care facilities: study design of a cluster randomized controlled trial in Pakistan. *Glob Health Action* 2015;16:28225. doi:[10.3402/gha.v8.28225](https://doi.org/10.3402/gha.v8.28225).
 39. Wong SS, Nathan AM, de Bruyne J, Zaki R, Mohd Tahir SZ. Does a written asthma action plan reduce unscheduled doctor visits in children? *Indian J Pediatr* 2013;80:590–595. doi:[10.1007/s12098-012-0839-0](https://doi.org/10.1007/s12098-012-0839-0).
 40. Poureslami I, Nimmon L, Doyle-Waters M, Rootman I, Schulzer M, Kuramoto L, FitzGerald JM. Effectiveness of educational interventions on asthma self-management in Punjabi and Chinese asthma patients: a randomized controlled trial. *J Asthma* 2012;49:542–551. doi:[10.3109/02770903.2012.682125](https://doi.org/10.3109/02770903.2012.682125).
 41. Poureslami I, Shum J, Nimmon L, FitzGerald JM. Culturally specific evaluation of inhaler techniques in asthma. *Respir Care* 2016;61:1588–1596. doi:[10.4187/respcare.04853](https://doi.org/10.4187/respcare.04853).
 42. Griffiths C, Foster G, Barnes N, Eldridge N, Tate H, Behum S, Wiggins M, Dawson C, Livingston AE, Chambers M, et al. Specialist nurse intervention to reduce unscheduled asthma care in a deprived multi-ethnic area: the east London randomized controlled trial for high risk asthma (ELECTRA). *Br Med J* 2004; 328:144. doi:[10.1136/bmj.37950.784444.EE](https://doi.org/10.1136/bmj.37950.784444.EE).
 43. Griffiths C, Bremner S, Islam K, Sohanpal R, Vidal DL, Dawson C, Foster G, Ramsay J, Feder G, Taylor S, et al. Effect of an education programme for South Asians with asthma and their clinicians: a cluster randomized controlled trial (OEDIPUS). *PLoS One* 2016;11:e0158783. doi:[10.1371/journal.pone.0158783](https://doi.org/10.1371/journal.pone.0158783).
 44. Moudgil H, Marshall T, Honeybourne T. Asthma education and quality of life in the community: a randomized controlled study to evaluate the impact on white European and Indian subcontinent ethnic groups from socioeconomically deprived areas in Birmingham, UK. *Thorax* 2000;55:177–183. doi:[10.1136/thorax.55.3.177](https://doi.org/10.1136/thorax.55.3.177).
 45. Tan NC, Chen Z, Soo WF, Ngoh AS, Tai BC. Effects of a written asthma action plan on caregivers' management of children with asthma: a cross-sectional questionnaire survey. *Prim Care Respir J* 2013;22: 188–194. doi:[10.4104/pcrj.2013.00040](https://doi.org/10.4104/pcrj.2013.00040).
 46. U.S. Census Bureau. (2017). Washington, D.C.: Government Printing Office. U.S. Census Bureau. (2017). Population Estimates Programme. Retrieved from <http://www.census.gov>.
 47. Mosnaim GS, Pappalardo AA, Resnick SE, Codispoti CD, Bandi S, Nackers L, Malik RN, Vijayaraghavan V, Lynch EB, Powell LH. Behavioral interventions to improve asthma outcomes for adolescents: a systematic review. *J Allergy Clin Immunol Pract* 2016;4: 130–141. doi:[10.1016/j.jaip.2015.09.011](https://doi.org/10.1016/j.jaip.2015.09.011).
 48. Sears MB. Adverse effects of beta-agonists. *J Allergy Clin Immunol* 2002;110:S322–S328. doi:[10.1067/mai.2002.129966](https://doi.org/10.1067/mai.2002.129966).
 49. Roland NJ, Bhalla RK, Earis J. The local side effects of inhaled corticosteroids: current understanding and review of the literature. *Chest* 2004;126:213–219. doi:[10.1378/chest.126.1.213](https://doi.org/10.1378/chest.126.1.213).
 50. National Heart, Lung and Blood Institute – National Asthma Education and Prevention Programme Expert Panel Report 3 – Guidelines for the Diagnosis and Management of Asthma. U.S Department of Health and Human Sciences. Available from: <https://www.nhlbi.nih.gov/health-topics/guidelines-for-diagnosis-management-of-asthma> [last accessed 1 April 2019].

SUPPLEMENTARY MATERIAL

Asthma behavioral interventions in south Asians

SUPPLEMENTARY FILE S1

TABLE 1. SUMMARY OF INCLUDED STUDIES

Study (country)	Study Design	Patient Population and Age	Intervention Description	Delivery Method	Outcomes Assessed	Impact of Intervention on Outcomes
Middle Income Countries (MICs; including lower and upper middle)						
Agarkhedkar <i>et al.</i> 2005 [13] (India)	CBA	Moderate perennial asthma 3-15 years	Specific elimination diet based on results of in-vitro allergy tests for a selected food panel	NS	1. IgE titres	1. + (for 17 patients), ↔ (for 6 patients), - (for 1 patient)
Aggarwal <i>et al.</i> 2013 [14] (India)	RCT	Patients with bronchial asthma 20-45 years	Yoga (pranayama), breathing exercises and normal medical treatment, compared to the control group who did not receive breathing training	Trained expert	1. FEV 2. FVC 3. FEV1/FVC 4. PEFR	1. + 2. + 3. + 4. +
Agnihotri <i>et al.</i> 2014 [15] (India)	RCT	Non-smokers with mild to moderate persistent asthma 12-60 years	Yoga training for 30 minutes every morning (including standard medical treatment), compared to the control group who received standard medical treatment only	Yoga trainer	1. PEFR 2. Asthma control scores 3. Hb 4. TLC 5. Polymorphs 6. Lymphocytes 7. Eosinophils 8. Monocytes 9. Superoxide dismutase 10. QoL	1. + 2. + 3. + 4. + 5. + 6. + 7. + 8. + 9. + 10. +
Agnihotri <i>et al.</i> 2016 [16] (India)	RCT	Non-smokers with mild to moderate persistent bronchial	Yoga intervention along with standard medical treatment, compared to the control group who received standard medical treatment	NS	1. FEV 2. FVC 3. FEV1/FVC 4. PEFR	1. + 2. + 3. + 4. +

Asthma behavioral interventions in south Asians

		asthma 12-60 years				
Agrawal et al. 2005 [17] (India)	RCT	Patients with persistent asthma 5-12 years	Individualised written home- management plan (including standard asthma therapy), compared to the control group who received standard asthma therapy only	Trained physician and social scientist	1. The number of acute asthma events requiring emergency hospital visits 2. The number of school days missed 3. The number of night time awakenings 4. Symptom score	1. + 2. + 3. + 4. +
Behera et al. 2006 [18] (India)	RCT	Asthma patients 18-60 years	A self-care manual on bronchial asthma, compared to no specific instructions	NS	1. Symptom score 2. Patient knowledge 3. Self-care measures 4. Influencing factors 5. Indices of asthma morbidity	1. + 2. + 3. + 4. + 5. +
Behera et al. 2008 [19] (India)	RCT	Patients with bronchial asthma Mean age of 36.72 years in the intervention group, and 34.33 years in the control group	Self-care manual stating details of asthma triggers and ways to avoid triggers, compared to the control group who received no access to self-care manuals	Investigators	1. Number of patients whose symptoms were aggravated by listed factors 2. Number of patients avoiding triggers	1. + 2. +
Ghosh et al. 1998 [20] (India)	RCT	Patients with chronic asthma and at least one hospital visit in the last year 10-45 years	Self-management training, compared to the control group who did not receive training sessions	Social scientist and physician	1. PEFR 2. Loss of productive days 3. Hospital days 4. % hospitalized 5. ER visits 6. Intervention costs 7. Direct costs 8. Indirect costs 9. Average costs	1. + 2. + 3. + 4. + 5. + 6. + 7. + 8. + 9. +
Grover et al. 2002 [21]	RCT	Patients with asthma and at least two years	CBT, compared with standard pharmacological treatment	NS	1. PEFR 2. Symptom diary 3. QoL	1. + 2. + 3. +

Asthma behavioral interventions in south Asians

(India)		of illness 18-45 years				
Grover <i>et al.</i> 2016 [22] (India)	RCT	Patients with asthma and at least two hospital visits in the last 12 months 7-12 years	PowerPoint slides and a child workbook, compared to the control group who received the usual standard of care (a standard information pack given before usual doctor's appointment)	Asthma educator	1. Asthma knowledge of caregiver and child 2. Asthma control 3. Drug adherence 4. Inhaler technique 5. Action plan ownership 6. PACQL (QoL)	1. + 2. + 3. + 4. + 5. + 6. +
Jolly <i>et al.</i> 2015 [23] (India)	Randomised parallel group study	Patients with respiratory disorders ≥18 years	Written instructions regarding correct inhaler technique of MDI, compared with a practical demonstration	Instructors and trained investigators	1. Median score for inhaler technique	1. +
Mandanmohan <i>et al.</i> 2003 [24] (India)	RCT	School children with no yoga training experience and no respiratory or cardiac diseases 12-15 years	Yoga training, compared to the control group who did not receive any yoga training	NS	1. FEV 2. FEV1 3. PEFR 4. Maximum expiratory pressure 5. Maximum inspiratory pressure 6. Hand grip strength 7. Hand grip endurance	1. + 2. + 3. + 4. + 5. + 6. + 7. +
Maazuddin <i>et al.</i> 2014 [25] (India)	CBA	Patients with asthma or COPD and prescribed bronchodilators and/or corticosteroids Age NS	Personalised health education and training regarding inhaler use	Pharmacists	1. Symptom score 2. Activity score 3. Impact score 4. Total score	1. + 2. + 3. + 4. +
Kumar <i>et al.</i> 2013 [26] (India)	CBA	Teachers 24-57 years	Educational booklet on asthma management around developing MDI skills	NS	1. Performance score	1. +

Asthma behavioral interventions in south Asians

Lathadevi <i>et al.</i> 2012 [27] (India)	RCT	Patients with asthma 18-60 years	Yoga training, compared to control group who did not receive any yoga training	NS	1. FEV1 2. FVC 3. FEV1/FVC 4. PEFR 5. Chest expansion 6. 40mmHg endurance test 7. Respiratory rate	1. + 2. + 3. + 4. + 5. + 6. + 7. +
Mishra <i>et al.</i> 2005 [28] (India)	RCT	Patients with mild and persistent asthma ≥14 years	Asthma awareness education programme, compared to those who did not receive the education programme	NS	1. Frequency of asthma attacks prior to intervention 2. Hospitalization 3. PEFR	1. + 2. + 3. +
Rao <i>et al.</i> 2014 [29] (India)	CBA	Patients with bronchial asthma 18-70 years	Nature cure therapy, diet therapy and yoga therapy	NS	1. FEV1 2. FVC 3. FEV1/FVC 4. PEFR 5. MVV	1. + 2. + 3. + 4. + 5. +
Saji <i>et al.</i> 2012 [30] (India)	RCT	Patients with moderate to severe asthma 18-65 years	Counselling points such as procedures to be followed during inhalation therapy, self-management of acute exacerbations, time drugs should be taken and dosage, compared to the non-counselled control group	Counsellor	1. FEV1 2. PEFR 3. FVC 4. QoL	1. + 2. + 3. + 4. +
Sathyaprabha <i>et al.</i> 2001 [31] (India)	CBA	Bronchial asthma patients Men mean age 40 years; women mean age 35 years	Diet therapy, nature care treatment and yoga therapy	NS	1. Vital capacity, FEV1, Maximum voluntary ventilation (MVV), PEFR 2. Hb, WBC, RBC 3. ESR, absolute eosinophil count	1. + 2. ↔ 3. +

Asthma behavioral interventions in south Asians

Shanmugam <i>et al.</i> 2012 [32] (India)	RCT	With asthma diagnosis Age NS	An asthma care diary, asthma education and medication counselling	Clinical pharmacist	1. PEFR, asthma control scores 2. QoL	1. + 2. +
Singh <i>et al.</i> 1990 [33] (India)	RCT	Those diagnosed with asthma, manifesting nocturnal wheeze 11-58 years	Pranayama breathing exercises using a Pink City lung exerciser by itself or with additional hot humid air, compared to the control period where a matched placebo device was used	NS	1. PEFR 2. Frequency of nocturnal wheeze	1. + (more so in hot air group) 2. +
Singh <i>et al.</i> 2012 [34] (India)	RCT	Patients with mild to moderate, and stable asthma 18-60 years	Yoga breathing training (pranayama)	Yoga trainer	1. FVC 2. FEV1 3. PEFR 4. MVV 5. SVC 6. TLCO 7. QoL	1. + 2. + 3. + 4. + 5. + 6. + 7. +
Sodhi <i>et al.</i> 2009 [35] (India)	RCT	Non-smokers with mild to moderate asthma. 17-50 years	Yoga training sessions, compared to the control group who did not undertake regular yoga training sessions	Trained yoga instructor	1. FVC 2. FEV1 3. PEFR 4. Forced mid expiratory flow (FEF) in 0.25-0.75 seconds (FEF25-75) 5. FEV1/FVC ratio	1. + 2. + 3. + 4. + 5. +
Sodhi <i>et al.</i> 2014 [36] (India)	RCT	Non-smokers with well controlled asthma 17-50 years	Yoga training sessions, compared to the control group who did not receive regular yoga training sessions	Trained yoga instructor	1. Number of attacks 2. Severity of attacks 3. Dosage of medication 4. QoL	1. + 2. + 3. + 4. +

Asthma behavioral interventions in south Asians

Vempati <i>et al.</i> 2009 [37] (India)	RCT	Patients with mild to moderate asthma ≥18 years	A yoga-based lifestyle modification intervention, compared to wait-listed control group	NS	1. FVC 2. FEV1 3. FEV1/FVC 4. PEFR 5. Average forced expiratory flow rate during the expulsion of 25–75% of FVC (FEF25–75%) 6. Eosinophilic cationic protein 7. Exercise-induced bronchoconstriction 8. Urinary concentration of prostaglandin D2 9. QoL	1. + 2. + 3. + 4. + 5. + 6. ↔ 7. + 8. ↔ 9. +
Khan <i>et al.</i> 2015 [38] (Pakistan)	RCT	Patients with newly diagnosed with asthma and COPD ≥18 years	Improved availability of context-sensitive guidelines and materials for case management including patient education tools for awareness and smoking cessation, compared to control group who received routine care for asthma and COPD case management	NS	1. FEV1/FVC ratio 2. Beta-2 agonist use 3. Waking up at night 4. Restriction of daytime activity 5. Emergency asthma treatment 6. Missed work or school days	Results not available
Wong <i>et al.</i> 2013 [39] (Malaysia)	RCT	Children with asthma 6-17 years	Written asthma action plan, compared to the control group who received verbal counselling on the management of asthma exacerbations only	Investigator	1. Acute asthma exacerbations 2. Patients with controlled asthma 3. QoL	1. ↔ 2. + 3. ↔
High Income Countries (HICs)						
Poureslami <i>et al.</i> 2012 [40] (Canada)	RCT	Patients with asthma and regular medication use 21-87 years	Education via physician-led videos or a patient-generated community video or both, compared to the control group who received an educational pamphlet	NS	1. Asthma knowledge 2. Correct inhaler use 3. Understanding of physician's instructions	1. + 2. + 3. +

Asthma behavioral interventions in south Asians

Poureslami <i>et al.</i> 2016 [41] (Canada)	RCT	Patients with asthma and daily medication use 21-87 years	Culturally specific educational videos that were either physician-led or community videos, compared to the control group who received and educational pamphlets	Study facilitators	1. Skills in appropriate inhaler technique	1. +
Griffiths <i>et al.</i> 2004 [42] (UK)	RCT	Patients with acute asthma 4-60 years	A self-management plan and a written action plan, compared to the control group who received a visit promoting standard asthma guidelines	Specialist nurses	1. % receiving unscheduled care 2. Rates of unscheduled care attendance 3. Self-management behavior 4. QoL	1. + 2. ↔ 3. ↔ 4. ↔
Griffiths <i>et al.</i> 2016 [43] (UK)	RCT	Children with asthma and their GPs ≥3 years	Two theoretically-based educational interventions that were culturally adapted for South Asians	Specialist nurse	1. Time to first unscheduled contact 2. Proportion of participants without unscheduled appointment 3. Time to first asthma review in primary care 4. Prescription 5. QoL	1. ↔ 2. ↔ 3. + 4. ↔ 5. +
Moudgil <i>et al.</i> 2000 [44] (UK)	RCT	Asthma patients (both White European and Indian Subcontinent patients) 11-59 years	Asthma education programme, individually tailored written self-management plan and peak flow meters, compared to the control group who only attended the beginning and end of the programme, along with their usual care	Educator	1. Hospital admissions 2. GP or deputizing services home 3. GP consults 4. Steroids prescription 5. Antibiotics prescription 6. QoL	1-5. + (Significant in White European group) 6. +
Tan <i>et al.</i> 2013 [45] (Singapore)	CBA	Caregivers of children with asthma 30-50 years	Written asthma action plan for caregivers	Student nurses	1. Symptom understanding and recognition 2. Use of asthma medications 3. Consultation and self-management	1. + 2. + 3. ↔

Asthma behavioral interventions in south Asians

KEY

- + : Intervention had a positive impact
- : Intervention had a negative impact
- ↔ : Intervention did not produce significant change

LIST OF ABBREVIATIONS

CBT: Cognitive behavioural therapy

COPD: Chronic obstructive pulmonary disease

ESR: Erythrocyte sedimentation rate

FEV: Forced expiratory volume

FEV1: Forced expiratory volume in one second

FVC: Forced vital capacity

FEV1/FVC: Forced expiratory volume in one second/forced vital capacity

Hb: Haemoglobin

MDI: Metered-dose inhaler

MVV: Maximum voluntary ventilation

NS: Not specified

PACQL: Paediatric asthma caregiver quality of life

PEFR: Peak expiratory flow rate

QoL: Quality of life

RBC: Red blood cell

RCT: Randomised controlled trial

SVC: Slow vital capacity

TLC: Total lung capacity

TLCO: Transfer factor for the lung for carbon monoxide

WBC: White blood cell

Asthma behavioral interventions in south Asians

SUPPLEMENTARY FILE S2

TABLE 2. RESULTS SUMMARY OF INCLUDED STUDIES

Intervention classification of outcome measure	Intervention outcome measure	Significant improvement	No Change
Physiological symptoms	IgE titres	1 study - [13]	1 study - [13]
	FEV or FEV1	10 studies - [14, 16, 24, 27, 29, 30, 31, 34, 35, 37]	
	FVC	8 studies - [14, 16, 27, 29, 30, 34, 35, 37]	
	FEV1/FVC	4 studies - [14, 16, 27, 29, 37]	
	PEFR	16 studies [14, 15, 16, 20, 21, 24, 27, 28, 29, 30, 31, 32, 33, 34, 35, 37]	
	MVV	3 studies - [29, 31, 34]	
	Chest expansion, 40mm Hg endurance test, and respiratory rate	1 study [27]	
	Maximum expiratory pressure, and maximum inspiratory pressure	1 study - [24]	
	Hand grip strength, and hand grip endurance	1 study - [24]	
	Hb	1 study - [15]	1 study - [31]
Unscheduled impact	Asthma events / attacks	3 studies - [17, 28, 36]	
	Hospitalization / emergency visits and/treatment / unscheduled care	6 studies - [17, 20, 28, 38, 42, 44]	2 studies - [42, 43]
	Number of school days missed / productive days lost / restriction of day time activity	1 study - [17, 20]	
	The number of night time awakenings / nocturnal wheeze	1 study - [17, 33]	
Self-reported	Symptom score	5 studies - [17, 18, 19, 21, 25]	
	Patient knowledge / understanding / recognition	4 studies - [18, 22, 40, 44]	
	Asthma control / self-care / self-management / ownership	4 studies - [15, 18, 22, 32]	1 study - [42]

Asthma behavioral interventions in south Asians

	QoL	10 studies - [15, 21, 22, 30, 32, 34, 36, 37, 43, 44]	2 studies - [39, 42]
	Avoidance of triggers	1 study - [19]	
Existing treatment	Drug adherence	1 study - [22]	
	Prescriptions	1 study - [44]	2 studies - [43, 44]
	Inhaler technique / intervention performance score	1 studies - [22, 23, 40, 41]	
Intervention Specific	Cost	1 study - [20]	
	Performance score	1 study - [26]	
	Understanding instructions	1 study - [40]	