
Review article

Supported by a grant from AstraZeneca Pharmaceuticals

Update on nonallergic rhinitis

Russell A. Settipane, MD* and Philip Lieberman, MD†

Background: Although nonallergic rhinitis is a well recognized entity, its incidence and therapy have not been definitively studied. Recent epidemiologic studies and treatment trials have furthered our knowledge regarding the frequency of occurrence of this disorder and effective treatment modalities.

Objective: To review and put into perspective recent advances in our knowledge regarding the incidence and significance as well as therapy of chronic nonallergic rhinitis. In addition, based upon these data, to propose a classification of this disorder.

Data Sources: The MEDLINE database and the results of a national survey of allergists (National Rhinitis Task Force) conducted in 15 allergy practices involving 975 patients.

Conclusions: Nonallergic rhinitis is a common disease that probably affects as many as 17 million Americans. Of equal importance is that, based on available data, approximately 22 million people suffer with a combination of nonallergic rhinitis and allergic diseases (mixed rhinitis). Both nonallergic and mixed rhinitis occur more frequently in adults than in children, may be more common in female patients than in male patients, and are more likely to be perennial than seasonal. Agents demonstrating efficacy (based on controlled trials or having approval by the FDA) for the therapy of nonallergic rhinitis are azelastine and topical nasal steroids.

Ann Allergy Asthma Immunol 2001;86:494–508.

INTRODUCTION

Rhinitis is defined as inflammation of the membranes lining the nose, characterized by nasal symptoms, including itching, rhinorrhea, and/or nasal congestion. Chronic nasal symptoms without allergic causation are a broad classification of nasal diseases known as nonallergic rhinitis. As many as half of the patients presenting with nasal symptoms may have this disorder. Nasal symptoms characteristic of nonallergic rhinitis are often indistinguishable from those that occur in allergic rhinitis, and, therefore, negative testing

for IgE-mediated sensitivity to relevant aeroallergens is necessary to confirm this diagnosis. However, it is important to note that positive skin tests to irrelevant aeroallergens may occur in nonallergic rhinitis.

Nonallergic rhinitis may be subclassified on the basis of various characteristics. These include frequency of occurrence (Table 1),¹ immunologic and cytologic features (Table 2),² and etiologic and systemic disease association (Table 3). The latter subclassification is the subject of this review. The epidemiology and clinical presentation of nonallergic rhinitis will be presented, followed by a descriptive summary of each of the major nonallergic rhinitis syndromes.

EPIDEMIOLOGY OF NONALLERGIC RHINITIS

Rhinitis is a significant cause of morbidity worldwide; however, it is often

viewed as trivial by practitioners. Rhinitis symptoms significantly affect patients' quality of life because of systemic symptoms and include, among others, fatigue, headache, and cognitive impairment. Allergic rhinitis affects 20 to 40 million people in the United States each year. Ten percent to 30% of allergic rhinitis sufferers are adults and 40% are children.³ In contrast, the frequency of nonallergic rhinitis and the different syndromes that comprise this disorder are poorly defined and no true data have existed regarding prevalence. However, the Joint Task Force on Practice Parameters in Allergy, Asthma, and Immunology points out that whereas allergic rhinitis is very common, approximately 50% of patients presenting with rhinitis do not have allergic rhinitis.³ Several studies have helped to establish the frequency of occurrence of allergic versus nonallergic rhinitis as seen in the allergist's office (Table 4). Mullarkey et al⁴ found that 52% of 142 rhinitis patients seen in an allergy clinic could be classified as having nonallergic rhinitis. Enberg⁵ evaluated 152 consecutive adults with nasal symptoms and found a 30% frequency of perennial nonallergic rhinitis. This frequency would increase to 46% if these subjects had been more precisely classified with positive skin tests, without clinical correlation, as having nonallergic rhinitis. The European Community Respiratory Health Survey (ECRHS)⁶ reported a 25% frequency of nonallergic rhinitis among 1,412 subjects selected for having a history that suggested allergic rhinitis. Togias⁷ found that 17% of 362 rhinitis patients at an academic allergy clinic

* Brown University School of Medicine, Providence, RI.

† Division of Allergy and Immunology and Department of Pediatrics, University of Tennessee School of Medicine, Memphis, TN.

Received for publication June 2, 2000.

Accepted for publication in revised form February 15, 2001.

Table 1. Classification of Nonallergic Rhinitis Based on Frequency of Occurrence

Common	Infrequent
Vasomotor (perennial nonallergic rhinitis)	Hypothyroidism
Chronic sinusitis	Atrophic
Structural (septum, turbinates, valve)	Systemic immunologic disorders
NARES*/BENARS†	Cerebral spinal fluid rhinorrhea
Drug induced	Other structural disorders
Nasal polyps	Foreign body
Physical/chemical/irritant	Ciliary dyskinesia
Aspirin sensitivity	Nasal mastocytosis

* NARES-nonallergic rhinitis with eosinophilia syndrome.

† BENARS - blood eosinophilia nonallergic rhinitis syndrome.

had nonallergic rhinitis. These studies, however, may be skewed toward the allergic rhinitis because these patients are seen most often in the allergy practice.

Rough data from large population surveys indicate the prevalence of chronic sinusitis to be 13.5%⁸ and chronic rhinitis to be 20.4%.⁹ Presumably, a substantial portion of subjects surveyed would be more precisely labeled as having nonallergic rhinitis if skin testing and sinus imaging had been performed.

The Joint Task Force notes that data regarding the true prevalence of rhinitis are difficult to interpret. Physician-diagnosed rhinitis is the basis for most of the population surveys and may lead to a lower reporting of rhinitis whereas

epidemiologic studies lean toward seasonal allergic rhinitis or hay fever, because identification of seasonal symptom exacerbation is easier to identify.³

Settipane and Klein¹⁰ looked at the frequency of occurrence of nonallergic rhinitis subtypes in 78 patients and determined that nonallergic rhinitis with eosinophilia syndrome (NARES) was found in 33%, blood eosinophilia nonallergic rhinitis syndrome (BENARS) in 4%, VMR in 61%, sinusitis in 16%, elevated IgE in 12%, and hypothyroidism in 2%. In his series of nonallergic rhinitis patients, Enberg⁵ found that 10% had NARES.

Three additional observations of note have been reported regarding the clinical presentation of nonallergic rhinitis. These include associations with

age, sex, and seasonal occurrence/variation. Togias⁷ has observed that 70% of patients diagnosed with nonallergic nasal disease developed their condition in adult life (age >20 years), whereas approximately 70% of patients diagnosed with allergic rhinitis developed their condition in childhood (age <20 years). However, a causal relationship between aging and nonallergic rhinitis has not been confirmed by other investigators.¹¹

Sex may be a risk factor for the development of nonallergic rhinitis. In the series of Settipane and Klein,¹¹ 58% of the nonallergic rhinitis patients were female. Enberg⁵ found 74% of nonallergic rhinitis patients to be female. However, these two studies lacked a control group. The National Rhinitis Classification Task Force (NRCTF) study found that 71% of patients with nonallergic rhinitis were female compared with 55% in the allergic rhinitis group.¹²

Sibbald and Rink¹¹ found that patients with nonallergic rhinitis were more likely to experience perennial, rather than seasonal, symptoms. They reported the following frequencies of negative skin testing: 50% in patients with perennial rhinitis, 32% in patients with combined perennial/seasonal rhinitis, and 22% in patients with purely seasonal allergic rhinitis.

Table 2. Classification of Nonallergic Rhinitis Based on Immunologic and Nasal Cytologic Features

Perennial Nonallergic Rhinitis (Inflammatory)	Non-inflammatory, Nonallergic Rhinitis	Structurally-related Rhinitis
Eosinophilic nasal disease (NARES*, BENARS**)	Rhinitis medicamentosa	Septal deviation, turbinate deformation, nasal valve dysfunction
Basophilic/metachromatic nasal disease	Reflex-induced rhinitis (bright light or other physical modalities)	Neoplastic and non-neoplastic tumors
Infectious	Vasomotor rhinitis	Miscellaneous (choanal atresia/stenosis, trauma, foreign body, malformation, cleft palate, adenoid hypertrophy)
	Irritant rhinitis	
	Cold air rhinitis	
	Gustatory rhinitis	
Nasal polyps	Rhinitis sicca	
Atrophic rhinitis	Metabolic (restronin-related or hyperthyroid)	
Immunologic nasal disease (non-IgE mediated or secondary to systemic immunologic disorders)		

* NARES - nonallergic rhinitis with eosinophilia syndrome.

** BENARS - blood eosinophilia nonallergic rhinitis syndrome.

Table 3. Classification of Chronic Nonallergic Rhinitis Based on Etiology or Systemic Disease Association

Syndromes of Unknown Etiology
Vasomotor rhinitis
Nonallergic rhinitis with eosinophils (NARES*, BENARS†)
Basophilic/metachromatic nasal disease
Syndromes of Suggested Etiology
Chronic Sinusitis
Immunodeficiencies
Ostiomeatal obstruction
Metabolic Conditions
Estrogen related (oral contraceptive/hormone replacement therapy, pregnancy)
Hypothyroidism
Acromegaly
Vasculitides/Autoimmune & Granulomatous Diseases
Sjögren's syndrome
Systemic lupus erythematosus
Relapsing polychondritis
Churg-Strauss syndrome
Sarcoidosis
Wegener granulomatosis
Drug-Induced
Topical decongestants
Systemic medications
Rhinitis with Nasal Polyps
Aspirin intolerance
Chronic sinusitis
Churg-Strauss syndrome
Young's syndrome (sinopulmonary disease, azoospermia, nasal polyps)
Cystic Fibrosis
Kartagener syndrome (bronchiectasis, chronic sinusitis, nasal polyps)
Structurally Related Rhinitis
Septal deviation
Turbinate deformation
Nasal valve dysfunction
Obstructive adenoid hyperplasia
Trauma (e.g., cerebrospinal fluid rhinorrhea)
Congenital
Neoplastic
Atrophic Rhinitis
Resulting from surgery
Ozena
Physical/Chemical/Irritant-Induced
Dry air
Gustatory
Bright light
Air pollution
Occupational
Occupational Rhinitis
Annoyance
Irritant
Immunologic
Corrosive

* NARES-nonallergic rhinitis with eosinophilia syndrome.

† BENARS - blood eosinophilia nonallergic rhinitis syndrome.

Although these epidemiologic data are interesting, consideration must be given to the fact that most of these

studies classify patients as having only allergic or nonallergic disease while failing to accurately capture the signif-

icant number of patients with co-existing allergic/nonallergic rhinitis or "mixed rhinitis."

The NRCTF recently conducted a retrospective analysis of 975 patients in approximately 15 allergy practices with the goal of determining the prevalence of "pure" allergic, "pure" non-allergic, and mixed rhinitis in the allergy practice (Table 5).¹² Forty-three percent of patients were classified as having pure allergic rhinitis, 23% pure nonallergic rhinitis, and 34% mixed rhinitis. Of all patients presenting with chronic rhinitis, 57% were classified as having nonallergic rhinitis, either pure or mixed. (Table 6) Clearly, this relatively high incidence of nonallergic rhinitis and the similar symptomatic presentation of allergic rhinitis dictates that a definitive diagnosis be made before treatment is initiated.

CLINICAL PRESENTATION, DIAGNOSIS, PATHOPHYSIOLOGY, AND TREATMENT OF INDIVIDUAL NONALLERGIC RHINITIS SYNDROMES

Syndromes of Unknown Etiology

Vasomotor rhinitis (VMR). Vasomotor rhinitis is unrelated to allergy, infection, structural lesions, systemic disease, or drug abuse.³ It is a diagnosis of exclusion (eg, negative allergen testing) and is likely to result from many different etiologies. For the purposes of this review, the term VMR will be used to designate idiopathic, perennial nonallergic rhinitis associated with negative allergy skin tests to relevant allergens, normal serum IgE levels, and a lack of identifiable inflammation on nasal cytology.

The symptoms of VMR range from obstructive/congestive to secretory/rhinorrhea; sneezing and nasal pruritus are less common. In a survey of 678 rhinitis patients,¹³ nasal blockage was the predominant symptom in patients with VMR, whereas allergic rhinitis patients were more likely to suffer from eye irritation, some sneezing, and rhinorrhea. Concomitant asthma was more common in the allergic rhinitis

Table 4. Frequency of Occurrence: Allergic vs Nonallergic Rhinitis

Investigator (year)	N	Rhinitis Type		
		Allergic	Mixed	Nonallergic
Mullarkey '80	142	48%	Not studied	52%
Enberg '89	152 (128)*	54%	16%† Undetermined	30%
Togias '90	362	83%	Not studied	17%
ECRHS '99‡	1,412	75%	Not studied	25%

* Diagnosis determined in only 128.

† "Mixed" counted by allergic for total analysis except Enberg's "undetermined group."

‡ European Community Respiratory Health Survey.

group. Togias⁷ reported nonallergic rhinitis to be associated with fewer sneezes and conjunctival symptoms but, with regard to rhinorrhea or congestion, could not differentiate this disorder from perennial allergic rhinitis.

In terms of pathophysiology, little is known about VMR except that nonspecific nasal hyperreactivity occurs on exposure to nonimmunologic stimuli such as changes in temperature or relative humidity, alcohol ingestion, strong odors, and other airborne irritants. Hyperreactivity of the nasal mucosa to methacholine,¹⁴ capsaicin,¹⁵ and histamine¹⁶ has been demonstrated in the laboratory, but its cause remains unexplained. Recent studies show an increased number of mast cells but not goblet cells in patients with perennial nonallergic rhinitis. No significant difference has been found between the number of mast cells in allergic and nonallergic patients.^{17,18}

Options in the treatment of VMR include either nonspecific, "broad-based" therapy aimed at multiple symptoms or alternatively, therapy tailored to treat specific symptoms. Nonspecific treatment may be more advantageous because symptoms of VMR are often variable, alternating from obstructive/congestive to secretory/rhi-

norrhea. Examples of broad-based treatment include topical steroids and the topical antihistamine, azelastine.

Budesonide aerosol preparation, beclomethasone aqueous preparation, and fluticasone aqueous preparation are topical corticosteroids that have an FDA indication for the treatment of nonallergic rhinitis. The benefit derived from topical nasal steroids impact overall rhinitis symptoms rather than being limited to any one particular symptom. Because of their anti-inflammatory nature, it has been proposed that nasal steroids may have greater benefit when nasal mucosal inflammation is present. The pharmacologic effects of fluticasone nasal spray in nonallergic rhinitis have been studied. Unilaterally administered fluticasone nasal spray has been shown to have a beneficial effect on nasal inflammatory cells and cytokine profile in both allergic and nonallergic rhinitis patients.¹⁹ The number of CD3, major basic protein, and tryptase positive cells was significantly less in allergic and nonallergic treated patients as compared with controls, and IL-4 and IL-5 mRNA were downregulated on the treated side. Based on this information, a 2- to 4-week trial of a topical

Table 6. National Rhinitis Classification Task Force Survey of Nonallergic Rhinitis 1999 (N = 975)

Type of Rhinitis		
Pure Allergic	Mixed	Pure Nonallergic
43%	34%	23%
	57%	

nasal steroid is a therapeutic consideration in VMR.²⁰⁻²²

Azelastine nasal spray was recently approved by the FDA for the treatment of the symptoms of VMR such as rhinorrhea, nasal congestion, and postnasal drip. Azelastine has been shown, in clinical trials, to be effective across the total symptom complex, ie, rhinorrhea, sneezing, and nasal discharge, without discrimination as to the type of nasal symptoms that respond.²³ Two multicenter, placebo-controlled trials, using identical protocols for the use of azelastine nasal spray in VMR patients, have demonstrated significant improvement in postnasal drip, rhinorrhea, sneezing, and congestion (Table 7). These studies had a fairly high response rate to azelastine; between 82% and 85% of more than 200 subjects with VMR were shown to respond favorably to azelastine nasal spray treatment. The response rate in the placebo group was 73%. It is not uncommon to see such a high placebo response when placebo consists of a nasal saline spray, as was the case in this study. In fact, nasal saline can be considered an "active placebo" because it is often efficacious in nonallergic rhinitis. Clinical and preclinical studies suggest that the mechanism of azelastine effects may involve anti-inflammatory actions.²⁴⁻²⁸ These include azelastine's ability to reduce the effects of neurokinins such as substance P and vasointestinal peptide,²⁰ prevent histamine release in vitro and in vivo,²¹ diminish eosinophil activation and the expression of adhesion molecules,²² suppress the synthesis of inflammatory cytokines and nitric oxide via an inhibitory effect on nuclear factor- κ B (NF- κ B),²⁵ and reduce vascular permeabil-

Table 5. National Rhinitis Classification Task Force Survey of Nonallergic Rhinitis 1999 (N = 975)

	Pure Allergic	Pure Nonallergic	Mixed
% of samples	43	23	34
Female (%)	53	71	62
Mean age (years)	40	43	39
Caucasian (%)	95	98	96

ity.²⁴ These studies suggest that azelastine has significant effects on transcription of inflammatory factors, thus azelastine seems to have unique combination of anti-inflammatory effects in addition to those that have been traditionally cited, such as inhibition of histamine release, calcium transport, and the influx of inflammatory cells into the site of an allergic reaction. It is perhaps through a combination of all these activities, especially those involving decreased production of cytokines, that azelastine exerts its activity in VMR.

Although there are no direct comparison studies between azelastine and topical nasal steroids in perennial non-allergic (vasomotor) rhinitis, there is information which suggests that azelastine nasal spray has a more rapid onset of action than intranasal corticosteroids. This information consists of clinical studies of these individual therapies in allergic rhinitis as well as a review of their FDA-approved product information. In dose-ranging trials, azelastine nasal administration resulted in a decrease in symptoms which reached statistical significance from saline placebo within 3 hours after initial dosing. In comparison, a decrease in allergic rhinitis symptoms after fluticasone or budesonide may be seen in 12 to 24 hours and may even occur within 7 hours of the latter. Of further note, azelastine nasal spray is the only therapy shown in nonallergic rhinitis to improve all individual symptoms (congestion, rhinorrhea, postnasal drip, and sneezing) that are usually combined in a total symptom score for VMR. This may be because, in part, of the antihistaminic and anti-inflammatory activities of azelastine.

As mentioned previously, an alternate treatment plan for VMR is to target a specific symptom. For example, patients whose symptoms are primarily obstruction/congestion may be treated with decongestants as first-line agents. These can be administered topically on a short-term basis, but for longer term use, they should be administered systemically because of the frequent adverse effect of rebound nasal congestion.

Patients with predominantly secretory/rhinorrhea symptoms may be best treated with the anticholinergic agent ipratropium bromide in a nasal spray.²⁹ This is effective because of its ability to inhibit the parasympathetic (cholinergic) nervous system, which innervates the serous and seromucous glands of the nasal mucosa. Intranasal ipratropium bromide is well tolerated because there is little systemic absorption; no systemic anticholinergic drug-related adverse effects have been associated with its use.^{30,31} Topical side effects include infrequent episodes of nasal dryness and minor epistaxis.

Another treatment which may be useful for hypersecretion symptoms because of their anticholinergic, drying effect on nasal mucosa is first-generation antihistamines. However, there are no clinical data demonstrating the efficacy of this therapy in VMR. Further, first-generation oral antihistamines may be limited by frequently associated sedation and tachyphylaxis. The cost of treatment as well as the clinical implications suggest that oral antihistamines may not be the ideal approach to the treatment of rhinitis without a definitive diagnosis of allergic disease. Oral antihistamines would not be expected to be effective in pure nonallergic rhinitis and only partially or intermittently effective in mixed rhinitis.

Another therapy for VMR is topical saline in the form of a spray or irrigation device. Nasal saline may result in reduction of postnasal drip, sneezing, and congestion.³¹ In some cases, using the above therapies in various combinations may give added benefit. Experimental treatments for VMR include topical use of capsaicin intranasally.¹⁵ This is a pungent substance derived from pepper that acts to desensitize sensory neural fibers in the nose, reducing nasal hyperreactivity. Beneficial effects are delayed in onset with 63% and 69% reduction of nasal blockage and nasal discharge respectively, after 1 month of therapy. Topical application of 15% to 20% silver nitrate might also be effective.^{32,33} Surgical approaches to therapy include endoscopic vidian nerve section and/or electrocoagulation of the anterior ethmoidal nerve.^{34,35} In both surgical approaches, the parasympathetic supply to the nasal mucosa is divided, resulting in reduced nasal secretion. Although there may be recurrence of symptoms secondary to re-innervation, recent studies report long-term benefits.³⁶ Although sphenopalatine ganglion block has also been reported to relieve symptoms of VMR, the number of blocks required for complete relief range from 2 to 4.³⁷ In cases where congestion is the predominant symptom,

Table 7. Relative Contribution of Individual Symptoms to Overall Response

Ratio of Azelastine Nasal Spray: Placebo for Vasomotor Rhinitis Symptoms									
Primary Efficacy Variable† P = 0.002					OVERALL		1.8:1*		
					Post Nasal Drip		1.7:1*	Study 374‡	
					Rhinorrhea		1.6:1*		
					Sneezes		>2.0:1*		
					Congestion		2.0:1*		
Primary Efficacy Variable† P = 0.005					OVERALL		1.9:1*	Study 375‡	
					Post Nasal Drip		1.9:1*		
					Rhinorrhea		1.9:1*		
					Sneezes		1.6:1*		
					Congestion		1.5:1*		
2.0	1.5	1.0	0.5	0	0.5	1.0	1.5	2.0	
Placebo superior treatment					Azelastine nasal spray superior treatment				

* Clinically significant.
† Primary efficacy variable is the improvement in total vasomotor rhinitis symptom score from baseline versus placebo.
‡ Studies 374 and 375 shared an identical protocol.
From Banov C (23); with permission.

tom, turbinectomy is another treatment option. A variety of surgical procedures can be performed for the treatment of hypertrophic inferior turbinates. There is some concern about the extent to which the nasal mucosa return to a normal functioning state after radical turbinectomy. However, laser turbinectomy has been reported to result in preservation of normal nasal cytology and saccharin time.³⁸

Nonallergic rhinitis with eosinophilia syndrome (NARES). First described in 1981, Jacobs and colleagues³⁹ presented a series of 52 patients with perennial symptoms of sneezing paroxysms, profuse watery rhinorrhea, and nasal pruritus. Nasal smears showed marked eosinophilia, but allergic disease could not be identified by skin testing or by RAST. This condition may represent approximately 15% to 33% of adults with nonallergic rhinitis.^{4,10} In these patients, NARES usually occurs as an isolated disorder. However, it may be associated with non-IgE-mediated asthma, aspirin intolerance, and nasal polyps. BENARS seems to be a subtype of NARES.¹⁰ Defined as having characteristics similar to NARES, BENARS is also associated with elevated numbers of blood eosinophils (average 957/mm³ in three patients). In a study of 78 consecutive patients with nonallergic rhinitis (Table 8), NARES was found in 33% and BENARS in 4% (Table 9). Rare nonallergic disorders associated with nasal eosinophilia include phaeohyphomycosis of the maxilloethmoid sinus⁴⁰ and Churg-Strauss syndrome.⁴¹

NARES patients tend to have more intense nasal symptoms than patients with either VMR or allergic rhinitis.⁴² Additionally, the presence of anosmia is common. The nasal cytology of NARES is characterized by eosinophilic infiltration, which may range from mild to massive, and is frequently associated with basophilic/metachromatic cell infiltration.² The pathophysiology of NARES syndrome is not understood. Eosinophilia may contribute to nasal mucosal dysfunction. This may occur as a result of the release of toxic substances, such as major basic protein and eosinophil cationic protein contained in eosinophil granules. These toxic proteins may damage nasal ciliated epithelium and prolong mucociliary clearance.⁴³⁻⁴⁸ In 1992, Davidson et al⁴⁹ reported on 56 nonallergic rhinitis patients and found a significant correlation between nasal eosinophilia and prolonged nasal ciliary clearance time (as measured by saccharin clearance). Delayed mucociliary clearance may result in an increased propensity toward infection. Recurrent infections may be a factor predisposing to the development of nasal polyps. Because nasal polyps are frequently associated with nasal eosinophils, there is some concern that nasal eosinophilia may be a precursor for nasal polyps or aspirin intolerance.⁴² In cases where aspirin intolerance exists, the aspirin reaction is not the etiology of the eosinophilic rhinosinusitis but mainly a marker of a severe form of NARES that is often associated with asthma, sinusitis, and nasal polyps. The presence of nasal

eosinophilia in patients with nonallergic rhinitis is generally regarded as a good prognostic indicator for response to treatment with topical steroid therapy. Where eosinophilic infiltration is massive, which is often the case in aspirin sensitivity syndrome, intermittent to frequent use of oral glucocorticoid may be required to control symptomatology.

Basophilic/metachromatic nasal disease. Basophilic/metachromatic cell nasal disease, or nasal mastocytosis, is a subcategory of nonallergic rhinitis that, such as NARES, is a histologic diagnosis.^{50,51} Mast cell infiltration (frequently >2,000/mm³) without nasal eosinophilia is the hallmark of this syndrome. Nasal symptoms are more likely to be secretion/rhinorrhea and congestion/blockage without significant sneezing/pruritus. Physical examination reveals an especially pale nasal mucosa. Unlike patients with NARES, patients with this condition are not predisposed to develop aspirin sensitivity, nasal polyps, asthma, or sinusitis. The etiology of this condition is unknown. Because it is an inflammatory condition, treatment should include a trial of topical anti-inflammatories; some patients benefit from intranasal cromolyn or intranasal corticosteroids, whereas others may require oral corticosteroids.

Syndromes of Suggested Etiology
Chronic sinusitis. Chronic rhinitis symptoms may be associated with or result from chronic infection and/or inflammation of the paranasal sinuses. According to a 1975 health interview survey, chronic infectious rhinosinusitis was experienced by 4% of the population.⁵² In 1985 the U.S. National Health Interview Survey ranked chronic sinusitis first among the most common chronic diseases, with a prevalence of 13.5%.⁸

Chronic sinusitis in adults has been recently defined⁵³ as persistent symptoms and signs of sinusitis lasting >8 weeks, or four episodes per year of acute sinusitis, each lasting at least 10 days in association with persistent changes documented by computed tomography (CT). CT of the sinuses

Table 8. Characteristics of Nonallergic Rhinitis Cases

Characteristics	Number	(%)
Women	45	(58)
Men	33	(42)
Average age (years)	42	
Range (years)	16 to 85	
Average duration of rhinitis (years)	8	
Range* (months to years)	3* to 30	
Elevated IgE	9	(12)
Negative allergy skin tests	78	(100)

*Six patients had symptoms for less than 6 months and 14 patients had symptoms for less than 1 year.

From Settiple GA, Klein DE, et al with permission.

Table 9. Diagnosis of 78 Patients with Nonallergic Rhinitis

Diagnosis	Patients (N)	Patients considered (N)	% of patients
VMR	44	72	61
NARES	25	75	33
Sinusitis (radiographs)*	11	68	16
Possible allergy†	9	76	12
BENARS	3	76	4
Hypothyroidism	1	68	2

BENARS - blood eosinophilia nonallergic rhinitis syndrome.

NARES - nonallergic rhinitis with eosinophils syndrome.

* Overlapping of diagnosis present.

† Elevated serum IgE levels.

Adapted from Settiple GA and Klein DE with permission.

should be performed 4 weeks after medical therapy without intervening acute infection. In children, the respective figures are 12 weeks of symptoms or more than six episodes of sinusitis per year. The symptoms and signs of chronic sinusitis are nasal congestion, discharge, headache, facial pain or pressure, and olfactory disturbance, with fever and halitosis as minor symptoms, and cough and irritability as possible symptoms in children.

The etiology of chronic sinusitis often involves an infectious origin, but the extent to which chronic infection is important in the perpetuation of the disease is unclear. Further, it is not clear to what extent anaerobic bacteria are present in chronic sinusitis as a result of dysfunctional sinus clearance or whether, in some cases, anaerobes may be a primary cause of the condition.⁵⁴ It is important to note that bacteria found in cultures of nasal secretions are not representative of those found in the infected sinus cavity. The frequent failure of prolonged and intensive antimicrobial therapy suggests that bacterial infection is not the cause of the persistence of chronic sinusitis. It is more likely that chronic sinusitis represents chronic mucosal inflammation, which is multifactorial in etiology.

The inflammatory nature of chronic sinusitis has been well documented. In the sinus fluid of chronic sinusitis patients undergoing surgery, inflammatory cells are mainly the neutrophils normally observed in acute sinusitis,

but a low percentage of eosinophils, mast cells, and basophils may also be observed.^{55,56} High concentrations of histamine, leukotrienes C₄, D₄, and E₄, and prostaglandin D₂ have been found, suggesting mast-cell/basophil activation in chronically inflamed sinuses.

The diagnosis of chronic sinusitis in a nonallergic population necessitates a workup for an underlying etiology (Table 10). With proper diagnosis and treatment of the underlying disorder, chronic sinusitis is optimally controlled. One example is osteomeatal complex obstruction, which when identified and treated, first medically and, if that is unsuccessful, then by endoscopic surgery, may reverse the infectious process, at least temporarily. Another example is antibody deficiency syndromes that usually respond well to intravenous replacement therapy.

There are, however, many patients with chronic sinusitis in whom no underlying disorder can be identified. An underlying inflammatory process is often present and can contribute to inadequate sinus ventilation and drainage. These patients, who often respond poorly to therapy, present a therapeutic dilemma. Scarce data exist on the efficacy of treatment for chronic sinusitis. Intuitively, maintenance of sinus osteal patency is important for proper drainage of retained secretions. Treatment should include a long-term trial of a topical steroid nasal spray in an attempt to reduce tissue inflammation and edema in the region of the os-

teomeatal complex obstruction. Additional therapeutic modalities that may be beneficial include guaifenesin, decongestants, steam inhalation, and saline spray or irrigation, and for acute flares, intermittent 3-week courses of antibiotics sometimes combined with systemic glucocorticoids. However, treatment will not be successful until a more complete understanding of the pathomechanisms of inflammation in chronic sinusitis is achieved.

Metabolic conditions. Pregnancy, oral contraceptives, other estrogens, hypothyroidism, and acromegaly can contribute to rhinitis symptoms. In pregnancy rhinitis, it often turns out that rhinitis symptoms, usually of less severity, have been present since before conception, and often these symptoms can be attributed to causes unrelated to the pregnancy.⁵⁷ The extent to which pregnancy is a causal factor is unknown; it may be more of an aggravating factor. During pregnancy, physiologic changes may contribute to increased nasal congestion/obstruction. Among these changes are a massive expansion of circulating blood volume, which contributes to an increase in nasal vascular pooling, and possible progesterone-induced, vascular smooth muscle relaxation. Further, pregnancy-associated hormones may have a direct effect on the nasal mucosa, resulting in increased nasal secretion/rhinorrhea secondary to increased nasal mucous gland hyperactivity.⁵⁸ Additionally, episodes of bacterial rhinosinusitis have been noted to be more common during pregnancy.⁵⁹ A complete discussion of the treatment of rhinitis during pregnancy is beyond the scope of this review.⁶⁰ However, the major prin-

Table 10. Conditions That May Underlie Chronic Sinusitis

Anatomical
Rhinitis
Nasal polyps
Ciliary dyskinesia: Kartagener syndrome
Cystic fibrosis
Granulomatous disease
Neoplasm
Immunodeficiency

ciple underlying the treatment of rhinitis in pregnancy is caution with medication use. First-line treatment should include the safest therapies such as steam inhalation, nasal saline sprays, and an avoidance of irritants. Topical medical therapy is preferred to systemic. The FDA classification system for the safety of medications in pregnancy provides additional guidance.

Vasculitides/autoimmune and granulomatous diseases. Systemic autoimmune/vasculitis diseases such as Churg-Strauss vasculitis,⁴¹ systemic lupus erythematosus,⁶¹ relapsing polychondritis,⁶² and Sjögren syndrome⁶³ can also result in rhinitis. Additionally, granulomatous diseases such as sarcoidosis⁶⁴ and, more commonly, Wegener granulomatosis⁶⁵ can produce nasal manifestations in a large proportion of affected patients.

Rhinitis sicca is a related syndrome that commonly occurs in Sjögren syndrome. It may also occur as a normal consequence of aging. Treatment with

a liberal use of nasal saline sprays and moisturizing nasal gels is appropriate.

Drug-induced rhinitis. Drugs listed in Tables 11 and 12 can induce rhinitis in allergic and nonallergic individuals. The term "rhinitis medicamentosa" is most commonly used to describe the rebound nasal congestion that occurs with overuse of topical decongestants/vasoconstrictor nasal preparations (oxymetazoline, phenylephrine) as well as from abuse of cocaine. Usually there is an underlying form of rhinitis that has led to this form of self-treatment. Classically, the nasal mucosa is erythematous, congested, and granular, with areas of punctate bleeding because of tissue friability. Recent research supports the theory that the rebound swelling is attributable to interstitial edema rather than to vasodilation.⁶⁶ Every patient presenting with the complaint of nasal congestion should be carefully questioned about the extent of use of decongestant nose sprays and cocaine, especially if the

nasal mucosa is erythematous on examination. Treatment of rhinitis medicamentosa requires withdrawal from the topical decongestant as well as treatment of the underlying rhinitis disorder. This is often best accomplished with the use of topical or systemic glucocorticoid, providing that there is some component of inflammatory rhinitis present. In some instances, patients can be initiated on a topical steroid spray bilaterally with instructions to discontinue the decongestant spray in one nostril and, 1 week later, in the remaining nostril. Another often effective treatment is a 1-week tapering course of oral glucocorticoid, with discontinuation of the decongestant spray on days 2 or 3. Thereafter, the patient should be followed for recurrence, possibly maintained on a topical nasal steroid, and worked up for an underlying etiology of chronic rhinitis.

Nasal polyps. Nasal polyps, which occur in approximately 1% of the general population, are not any more frequent in patients with allergic rhinitis.⁶⁷ Nasal polyps are frequently associated with aspirin intolerance and aspirin tetrad syndrome (aspirin intolerance, nasal polyps, sinusitis, and asthma), intrinsic asthma, chronic sinusitis, Young syndrome, cystic fibrosis, Kartagener syndrome, Churg-Strauss syndrome, and allergic fungal sinusitis (Table 13). Children 16 years or younger with nasal polyps should be evaluated for cystic fibrosis. Nasal polyps are often bilateral, multiple, freely movable, and pale gray, and arise from the middle meatus of the nose. Polyps in the paranasal sinuses can lead to bone erosion of the sinus walls (Woakes syndrome).⁶⁸ Histologically, nasal polyps classically have a pseudostratified ciliated columnar epithelium, thickening of the epithelial basement membrane, a high stromal eosinophil count, mucin with neutral pH, few glands, and essentially no nerve endings.⁶⁹ Cells usually consist of a mixture of lymphocytes, plasma cells, and eosinophils. The majority of nasal polyps are associated with an eosinophilic inflammatory mechanism. However, in polyps from cystic fibro-

Table 11. Antihypertensive Drugs Causing Rhinitis

Trade Name	Generic	Symptoms
Aldomet*	Methyldopate	Nasal stuffiness
Apresoline†	Hydralazine	Nasal congestion, dyspnea
Cardura‡	Doxazosin	Nasal congestion
Catapres§	Clonidine	Dryness of nasal mucosa
Corgard¶	Nadolol	Nasal stuffiness, bronchospasm, cough
Hytrin**	Terazosin	Nasal congestion
Ismelin†	Guanethidine	Dyspnea, asthma in susceptible individuals, nasal congestion
Minipress‡	Prazosin	Nasal congestion, dyspnea
Moduretic*	Amiloride and hydrochlorothiazide	1% nasal congestion, 1% dryness
Normodyne††	Labetalol	Nasal stuffiness, wheezing in 1
Trandate‡‡	Labetalol	Nasal stuffiness, dyspnea, bronchospasm
Wyntensin§§	Guanabenz	Nasal stuffiness, dyspnea
—	Reserpine	Nasal congestion, dyspnea

* Merck and Company, West Point, PA.

† Novartis Pharmaceuticals, East Hanover, NJ.

‡ Pfizer, Inc, New York, NY.

§ Boehringer Ingelheim Pharmaceuticals, Ridgefield, CT.

¶ Bristol-Myers Squibb, Princeton, NJ.

** Abbott Laboratories, Abbott Park, IL.

†† Schering Corporation, Kenilworth, NJ.

‡‡ Glaxo-Wellcome, Research Triangle Park, NC.

§§ Wyeth-Ayerst Laboratories, Philadelphia, PA.

Adapted from Settiple GA and Klein SA with permission.

Table 12. Antihypertensive Drugs with Diuretics Causing Rhinitis

Trade Name	Generic	Symptoms
Aldoclor*	Combination of methyl dopa and chlorothiazide	Nasal stuffiness
Aldoril*	Methyl dopahydrochlorothiazide, MSD	Nasal stuffiness
Apresazide†	Hydralazine and hydrochlorothiazide	Conjunctivitis, eosinophilia, dyspnea, nasal congestion
Combipres‡	Clonidine and chlorthalidone	Dryness of nasal mucosa
Diupres*	Chlorothiazide and reserpine	Nasal congestion, dyspnea
Diutensin-R§	Methylchlorothiazide and reserpine	Frequent nasal congestion, dyspnea
Enduronyl tablets¶	Methylchlorothiazide and deserpidine	Asthma in asthmatic patients, nasal congestion, dyspnea
Hydropres*	Reserpine-hydrochlorothiazide	Nasal congestion, dyspnea
Rauzide tablets**	Rauwolfia serpentina with bendroflumethiazide	Nasal congestion, dyspnea
Ser-Ap-Es†	Reserpine, hydralazine hydrochlorothiazide	Dyspnea, nasal congestion
Minizide‡‡	Prazosin and polythiazide	Nasal congestion, dyspnea

* Merck and Company, West Point, PA.

† Novartis Pharmaceuticals, East Hanover, NJ.

‡ Boehringer Ingelheim Pharmaceuticals, Ridgefield, CT.

§ Wallace Laboratories, Cranbury, NJ.

¶ Abbott Laboratories, Abbott Park, IL.

** Bristol-Myers Squibb, Princeton, NJ.

‡‡ Pfizer, Inc. New York, NY.

Adapted from Settiple GA and Klein SA with permission.

sis patients, neutrophils predominate with insignificant eosinophils.⁷⁰ Chemical mediators found in nasal polyps are as follows: histamine, serotonin, leukotrienes (LTB₄, LTC₄, LTD₄, LTE₄), IL-5, norepinephrine, kinins, TAME-esterase, and possibly prostaglandin D₂.^{71,72} There is a higher concentration of histamine in nasal polyps than in normal nasal mucosa, and norepinephrine is present in greater concentration at the base of nasal polyps than in normal nasal mucosa.⁷³ The concentrations of IgA, IgE, and in some cases, IgG and IgM, are greater in polyp fluid than in serum.⁷⁴ At the present time, the pathogenesis of polyp formation is unknown. IgE-mediated disease is not the cause of nasal polyps. In fact, nasal polyps may be associated with a loss of susceptibility to potential additional effects of inhaled allergens.⁷⁵ Upper respiratory infec-

tions may contribute to the exacerbation of nasal polyps. Despite medical and surgical management, nasal polyps are frequently recurrent. Topical corticosteroids alone can often control symptoms.⁷⁶ If topical therapy is not effective, it is often worthwhile to try systemic corticosteroids before surgical polypectomy. A commonly used regimen starts with prednisone at 60 mg and tapers by 5 mg daily over a 12-day course. Beclomethasone has been approved by the FDA for the prevention of nasal polyp recurrence after surgical removal. Polypectomy does not increase the risk of developing asthma or making asthma worse.⁷⁷

Structurally related rhinitis. Common anatomical causes of nasal obstruction include deviated septum, enlarged turbinates, dysfunctional nasal valve, and in children, adenoid hypertrophy. Anatomical abnormalities ac-

count for approximately 5% to 10% of chronic nasal disorders. Nasal septal deviation is fairly common but usually mild and well tolerated. In cases where deviation is moderate to severe, particularly when there is concomitant nasal mucosal edema secondary to underlying rhinitis, symptoms of blockage will result and may be complicated by sinusitis, snoring, sleep apnea, and fatigue. Enlarged turbinates and nasal valve disease are also aggravated by concomitant rhinitis. Therefore whenever there is any question about the extent of anatomical cause, and before the physician embarks on a surgical procedure to reduce the bulk of nasal turbinates, straighten the septum, or correct a dysfunctional nasal valve, workup and treatment of associated rhinitis should be pursued.

Common structural causes of rhinitis symptoms were mentioned previously, but rare causes of obstruction/congestion deserve mention. The differential diagnosis includes tumors and neoplasms such as chordoma, chemodectoma, neurofibroma, angiofibroma, inverting papilloma, squamous cell carcinoma, sarcoma, and encephalocoeles or meningocele.

Structural defects may also result in rhinorrhea. A rare cause of rhinorrhea is cerebral spinal fluid (CSF) leak, occurring most commonly as a result of trauma, but also either spontaneously or as a complication of surgery.⁷⁸ Although traumatic CSF leaks usually present within 48 hours of trauma, delayed onset can be seen. However, 95% of cases present within 3 months. Because CSF contains sugar and nasal mucous does not, testing for the presence of glucose is helpful in making

Table 13. Etiology of Nasal Polyps

Aspirin intolerance
Aspirin tetrad syndrome
Intrinsic asthma
Chronic sinusitis
Young syndrome
Cystic fibrosis
Kartagener syndrome
Churg-Strauss syndrome
Allergic fungal sinusitis

the diagnosis. Quantitative glucose measurements accurately identify the fluid as CSF if it contains more than 30 mg/100 mL glucose. A more specific test for CSF rhinorrhea is β_2 -transferrin, which has come into common use in recent years.⁷⁹ Additionally, thin coronal CT, radionuclide cisternography, and magnetic resonance cisternography can help to localize the site of CSF leakage.⁸⁰

Atrophic rhinitis. In industrialized countries, atrophic rhinitis is usually only seen as a rare complication of radical nasal tissue removal by surgery aimed to relieve obstruction.^{81,82} In undeveloped countries, symptoms of epistaxis, severe crusting, and stuffiness are associated with a foul or fetid odor and *Klebsiella* colonization. Presumably, greater use of antibiotics has reduced the occurrence of this syndrome in developed countries. Atrophic rhinitis is resistant to treatment. The standard modality is antibiotic treatment of bacterial overgrowth and aggressive nasal saline irrigation. One suggested form of saline therapy is the use of a dental water-pick attached to a nasal irrigator nozzle. An alternative therapy is the implantation of bone chips, which reportedly reduces nasal stuffiness.

Physical/chemical/irritant-induced rhinitis. Physical triggers of rhinitis symptoms include cold and dry air, ingestion of spicy food, and exposure to bright light. These common triggers are most often markers of increased nasal hyperreactivity because of an underlying nasal disorder. However, patients without an underlying disorder, particularly those who work outdoors in the cold, occasionally present complaining chiefly of excessive nasal symptomatology in response to the above stimuli. Some insight into the pathophysiology of this disorder has been gained by reproducing the conditions in the laboratory. Studies with cold, dry air have demonstrated the involvement of both mast cell degranulation and an increased neuronal reflex mechanism.^{83,84} The underlying defect seems to be an inability of the nasal mucosa to humidify inhaled air at extreme atmospheric conditions. This

results in increased osmolality of nasal secretions and nasal epithelial desiccation and detachment, leading to mast cell degranulation and activation of irritant sensory nerves. Studies with spicy food ingestion have suggested

the pathophysiology involves a purely neurogenic reflex mechanism, with the efferent pathway being parasympathetic nerve fibers.⁸⁵ This condition is called gustatory rhinitis. Why some patients may have a more responsive

Table 14. Occupational Rhinitis: Causative Agents in the Workplace

Agent	Occupational environment
Annoyance rhinitis	
Perfumes	Beauty salons, department stores
Cooking odors	Restaurants, food manufacturers
Exhaust fumes	Garages, public transportation industry
Detergents	Supermarkets
Cleaning agents	Supermarkets
Room deodorizers	
Flower fragrances	Florist shops, gardening industry
Clothing fragrances	Clothing stores
Cosmetic odors (hair spray)	Cosmetic factories, stores, beauty salons
Tobacco smoke on clothing	
Irritation rhinitis	
Tobacco smoke	
Pollution	
Oxides of nitrogen	Chemical industry
Paint fumes	
Garden sprays	
Toluene, xylene	Petroleum, chemical, paint industries
Capsaicin	Food industry
Formaldehyde	Chemical industry
Sulfur dioxide	
Ozone	
Immunologic rhinitis	
Animal proteins	Animal laboratories, pet stores
Wheat	Food-processing plants
Green tea	
Pyrethrum	Insecticide, gardening industries
Cotton fibers	Cotton mills
Reactive dyes	
Toluene diisocyanate	Auto body spray painting
<i>Bacillus subtilis</i>	Detergent factories
Trypsin	
Papain	Meat-processing plants
Latex	Health care industry
Platinum salts	
Colophony	Metal, electronics industries
Acid anhydrides	Adhesive industry
Plicatic acid	Cedar sawmills
Corrosive rhinitis	
Ammonia	Tanneries; dye, bronzing industries
Chloride	Pulp, paper industry; swimming pool maintenance
Hydrochloric acid	Bleach, battery manufacturing
Vinyl chloride	
Organophosphide compounds	Pesticide industry
Acrylamide	Resins, plastics manufacturing
Cyanide, nitriles	Steel, plastics, herbicide industries

Organic sulfur-containing compounds.

Adapted from Bardana and Chan-Yeung and Lam with permission.

reflex mechanism is unknown. Effective treatment for both these conditions is prophylaxis with topical ipratropium bromide before cold air exposure or eating spicy food.

An increasing array of indoor and outdoor air pollutants is recognized as affecting the nose. Among these are dust, ozone, sulfur dioxide, formaldehyde, volatile organic compounds, wood smoke, and environmental tobacco smoke.⁸⁶ The relationship between frequency of exposure and the development of acute or chronic rhinitis syndromes is unknown. Nor has associated rhinitis been well characterized. Most of these triggers are nasal irritants causing dryness and reduced nasal airflow, rhinorrhea, and sneezing.⁸⁶ Chronic cigarette smoke exposure results in decreased cilia beat frequency, ineffective movement, and frank deciliation in experimental models.⁸⁷ Ciliostasis has been demonstrated in wood workers exposed to dust.⁸⁸ Neutrophils' influx has been shown to occur with ozone exposure and exposure to volatile organic compounds.^{86,89} Chronic exposure to formaldehyde vapors may result in nasal histologic abnormalities.⁹⁰ Prolonged occupational exposure to nickel, leather or wood dust, formaldehyde, and chlorophenol has been associated with hypertrophic rhinitis, metaplasia, and frank carcinoma.²

Occupational rhinitis. Occupational rhinitis is defined as the episodic, work-related occurrence of sneezing, rhinorrhea, and nasal obstruction.⁹¹ Occupational rhinitis can be classified by one of the following reaction types: annoyance, irritation, immunologic, or corrosive (Table 14).⁹² Nonallergic rhinitis in the workplace needs to be distinguished from allergic rhinitis secondary to exposure to workplace aeroallergens and nonoccupational, allergic rhinitis aggravated by workplace exposure. The greater the number of symptomatic workers, the more likely it is that the offending agent is nonimmunogenic.⁹³ Symptoms of annoyance reactions typically result from exposure to strong odors in patients with heightened olfactory awareness.

Symptoms of irritant reactions result from exposure to one or more known respiratory irritants at levels exceeding threshold limits. Symptoms of corrosive reactions result from exposure to high concentrations of irritating and soluble chemical gases. This results in significant nasal inflammation, often manifesting as mucosal burns, frank ulceration, and, at times, associated inflammation of the skin, mouth, and eyes. Environmental control is the mainstay of therapy, achieved by either removing the etiologic agent, improving ventilation, wearing protective masks, or changing the work site. Recommended medical therapy includes nasal saline lavage to remove accumulated particulates.

SUMMARY

Pure nonallergic rhinitis is a common disease that affects as many as 17 million Americans. Patients with this condition have negative allergy skin tests to relevant aeroallergens and normal serum IgE. Symptoms are similar to those of allergic rhinitis but are usually less likely to be associated with sneezing episodes and conjunctival complaints. Frequently, however, nonallergic rhinitis occurs in tandem with allergic disease and presents as mixed rhinitis. Approximately 22 million people suffer with a combination of allergic and nonallergic disease. Nonallergic rhinitis, pure and mixed, occurs more frequently in adults than in children, may be more common in women, and is more likely to be perennial than seasonal. Common causes of nonallergic rhinitis are VMR, chronic sinusitis, nasal septal deviation, NARES/BENARES, nasal polyps, and aspirin sensitivity; nonallergic rhinitis may also be drug-induced, estrogen-related, or physical/chemical/irritant-induced. Infrequent causes include hypothyroidism, atrophic mucosa, systemic immunologic disorders, CSF rhinorrhea, ciliary dyskinesia, and nasal mastocytosis. Frequency ranges from 2% for hypothyroidism to 61% for VMR. At least, in some cases, the pathophysiology of VMR seems to involve proinflammatory chemical medi-

ators. Given the often indistinguishable symptoms and high incidence of pure nonallergic and mixed rhinitis, treatment should ideally be based on a differential diagnosis that includes skin testing for allergens. However, rhinitis is often treated empirically and oral antihistamines, most often chosen for empirical treatment, have not been shown to be effective for nonallergic rhinitis and would only be partially or intermittently effective for mixed rhinitis. If empirical treatment is chosen, treatment options should be limited to those which have been demonstrated to be effective in allergic and nonallergic disease. For long-term control of symptoms, topical medications are preferred to systemic and treatment should address all symptoms whether allergic or nonallergic in etiology. Azelastine nasal spray and intranasal corticosteroids should be considered as first-line therapy if empirical treatment is chosen. Additionally, it is sometimes worthwhile to use nasal lavage (saline washing). Experimental treatment includes capsaicin and silver nitrate application. Experimental surgical treatment includes vidian nerve section, electrocoagulation of the anterior ethmoidal nerve, sphenopalatine ganglion block, and turbinectomy. As knowledge grows of the pathophysiology of nonallergic rhinitis subtypes, physicians will be better able to treat this common disorder.

REFERENCES

1. Settupane RA, Settupane GA. Nonallergic rhinitis. In: Kaliner MA, ed. *Current Review of Allergic Diseases*. Philadelphia: Current Medicine, 1999.
2. Zeiger RS. Differential diagnosis and classification of rhinosinusitis. In: Schatz M, Zeiger RS, Settupane GA, eds. *Nasal Manifestations of Systemic Diseases*. Providence, RI: OceanSide Publications, 1991.
3. Dykewicz MS, Fineman S, Skoner DP. Diagnosis and management of rhinitis: complete guidelines of the Joint Task Force on Practice Parameters in Allergy, Asthma, and Immunology. American Academy of Allergy, Asthma, and Immunology. *Ann Allergy Asthma Immunol* 1998;81:478-518.
4. Mullarkey MF, Hill JS, Webb DR. Al-

6. lergic and nonallergic rhinitis: their characterization with attention to the meaning of nasal eosinophilia. *J Allergy Clin Immunol* 1980;65:122–126.
5. Enberg RN. Perennial nonallergic rhinitis: a retrospective review. *Ann Allergy* 1989;63:513–516.
6. Leynaert B, Bousquet J, Neukirch C, et al. Perennial rhinitis: an independent risk factor for asthma in nonatopic subjects: results from the European Community Respiratory Health Survey. *J Allergy Clin Immunol* 1999;104:301–304.
7. Togias A. Age relationships and clinical features of nonallergic rhinitis. *JACI* 1990;85:182.
8. Collins JG. Prevalence of selected chronic conditions, United States, 1983–85; National Center for Health Statistics Advance Data 1988;155:1–16.
9. Turkeltaub PC, Gergen PJ. The prevalence of allergic and nonallergic respiratory symptoms in the U.S. population: data from the second national health and nutrition examination survey 1976–80 (NHANES II). *J Allergy Clin Immunol* 1988;81:305.
10. Settipane GA, Klein DE. Nonallergic rhinitis: demography of eosinophils in nasal smear, blood total eosinophil counts and IgE levels. *N Engl J Med* 1985;6:363–366.
11. Sibbald B, Rink E. Epidemiology of seasonal and perennial rhinitis: clinical presentation and medical history. *Thorax* 1991;46:895–901.
12. Data presented at the National Allergy Advisory Council Meeting (NAAC)-The broad spectrum of rhinitis: etiology, diagnosis, and advances in treatment. St. Thomas, U.S. Virgin Islands, October 16, 1999.
13. Lindberg S, Malm L. Comparison of allergic rhinitis and vasomotor rhinitis patients on the basis of a computer questionnaire. *Allergy* 1993;48:602–607.
14. Borum P. Nasal methacholine challenge. *J Allergy Clin Immunol* 1979;63:253–257.
15. Stjärne P, Lundblad L, Änggård A, et al. Local capsaicin treatment of the nasal mucosa reduces symptoms in patients with nonallergic nasal hyperactivity. *Am J Rhinology* 1991;5:145–151.
16. Togias A, Proud D, Kagey-Sobotka A, et al. Cold dry air (CDA) and histamine (HIST) induce more potent responses in perennial rhinitis compared to normal individuals. *J Allergy Clin Immunol* 1991;87:148.
17. Berger G, Marom Z, Ophir D. Goblet cell density of the inferior turbinates in patients with perennial allergic and nonallergic rhinitis. *Am J Rhinol* 1997;11:233–236.
18. Berger G, Goldberg A, Ophir D. The inferior turbinate mast cell population of patients with perennial allergic and nonallergic rhinitis. *Am J Rhinol* 1997;11:63–66.
19. Kondo H, Nachtigal D, Frenkiel S, et al. Effect of steroids on nasal inflammatory cells and cytokine profile. *Laryngoscope* 1999;109:91–97.
20. Malm L, Wihl JA. Intra-nasal beclomethasone dipropionate in vasomotor rhinitis. *Acta Allergol* 1976;31:245–253.
21. Wight RG, Jones AS, Beckingham E, et al. A double blind comparison of intranasal budesonide 400 micrograms and 800 micrograms in perennial rhinitis. *Clin Otolaryngol* 1992;17:354–358.
22. Small P, Black M, Frenkiel S. Effects of treatment with beclomethasone dipropionate in subpopulations of perennial rhinitis patients. *J Allergy Clin Immunol* 1982;70:178–182.
23. Banov C, Laforce C, Lieberman P. Double-blind trial of Astelin nasal spray in the treatment of vasomotor rhinitis. *Ann Allergy Asthma Immunol* 2000;84:138.
24. Shinoda M, Watanabe N, Suko T, et al. Effects of substance P (SP) and vasoactive intestinal peptide (VIP) in nasal secretions. *Am J Rhinol* 1997;11:237–241.
25. Takao A, Shimoda T, Matsuse H, et al. Inhibitory effects of azelastine hydrochloride in alcohol-induced asthma. *Ann Allergy Asthma Immunol* 1999;82:390–394.
26. Ciprandi G, Pronzato C, Passalacqua G, et al. Topical azelastine reduces eosinophil activation and intercellular adhesion molecule-1 expression on nasal epithelial cells: an antiallergic activity. *J Allergy Clin Immunol* 1996;98:1088–1096.
27. Yoneda K, Yamamoto T, Ueta E, and Osaki T. Suppression by azelastine hydrochloride of NF- κ B activation involved in generation of cytokines and nitric oxide. *Jpn J Pharmacol* 1997;73:145–153.
28. Tamaoki J, Yamawaki I, Tagaya E, et al. Effect of azelastine on platelet-activating factor-induced microvascular leakage in rat airways. *Am J Physiol* 1999;276:L351–L357.
29. Grossman J, Banov C, Boggs P, et al. Use of ipratropium bromide nasal spray in chronic treatment of nonallergic perennial rhinitis, alone and in combination with other perennial rhinitis medications. *J Allergy Clin Immunol* 1995;95:1123–1127.
30. Wood CC, Fireman P, Grossman J, et al. Product characteristics and pharmacokinetics of intranasal ipratropium bromide. *J Allergy Clin Immunol* 1995;95:1111–1116.
31. Bronsky EA, Druce H, Findlay SR, Hampel FC. A clinical trial of ipratropium bromide nasal spray in patients with perennial nonallergic rhinitis. *J Allergy Clin Immunol* 1995;95:1117–1122.
32. Bhargava KB, Shirali GN, Abhyankar US, Gadre KC. Treatment of allergic and vasomotor rhinitis by the local application of different concentrations of silver nitrate. *J Laryngol Otol* 1992;106:699–701.
33. al-Samarrae SM. Treatment of ‘vasomotor rhinitis’ by the local application of silver nitrate. *J Laryngol Otol* 1991;105:285–287.
34. el-Guindy A. Endoscopic transseptal vidian neurectomy. *Arch Otolaryngol Head Neck Surg* 1994;120:1347–1351.
35. Dong Z. Anterior ethmoidal electrocoagulation in the treatment of vasomotor rhinitis. *Zhonghua Er Bi Yan Hou Ke Za Zhi* 1991;26:358–359, 383.
36. Fernandes CM. Bilateral transnasal vidian neurectomy in the management of chronic rhinitis. *J Laryngol Otol* 1994;108:569–573.
37. Prasanna A, Murthy PS. Vasomotor rhinitis and sphenopalatine ganglion block. *J Pain Symptom Manage* 1997;13:332–338.
38. Mladina R, Risavi R, Subaric M. CO₂ laser anterior turbinectomy in the treatment of non-allergic vasomotor rhinopathy. A prospective study upon 78 patients. *Rhinology* 1991;29:267–271.
39. Jacobs RL, Freedman PM, Boswell RN. Nonallergic rhinitis with nasal eosinophilia (NARES syndrome). Clinical and immunologic presentation. *J Allergy Clin Immunol* 1981;67:

- 253–262.
40. Sobol SM, Love RG, Stutman HR, Pysher TJ. Phaeohyphomycosis of the maxilloethmoid sinus caused by *Drechslera spicifera*: a new fungal pathogen. *Laryngoscope* 1984;94:620–627.
41. Olsen KD, Neel HB, DeRemee RA, Weiland LH. Nasal manifestations of allergic granulomatosis and angiitis (Churg-Strauss syndrome). *Otolaryngol Head Neck Surg* 1980;88:85–89.
42. Moneret-Vautrin DA, Hsieh V, Wayoff M, et al. Nonallergic rhinitis with eosinophilia syndrome a precursor of the triad: nasal polyposis, intrinsic asthma, and intolerance to aspirin. *Ann Allergy* 1990;64:513–518.
43. Bousquet J, Chanez P, Lacoste JY, et al. Eosinophilic inflammation in asthma. *N Engl J Med* 1990;323:1033–1039.
44. Hastie AT, Loegering DA, Gleich GJ, Kueppers F. The effect of purified human eosinophil major basic protein on mammalian ciliary activity. *Am Rev Respir Dis* 1987;135:848–853.
45. Flavahan NA, Slifman NR, Gleich GJ, Vanhoutte PM. Human eosinophil major basic protein causes hyperreactivity of respiratory smooth muscle. *Am Rev Respir Dis* 1988;138:685–688.
46. Venge P, Dahl R, Fredens K, Peterson CG. Epithelial injury by human eosinophils. *Am Rev Respir Dis* 1988;138:S54–S57.
47. Spector SL, English G, Jones L. Clinical and nasal biopsy response to treatment of perennial rhinitis. *J Allergy Clin Immunol* 1980;66:129–137.
48. Ayars GH, Altman LC, McManus MM, et al. Injurious effect of the eosinophil peroxidase-hydrogen peroxide-halide system and major basic protein on human nasal epithelium in vitro. *Am Rev Respir Dis* 1989;140:125–131.
49. Davidson AE, Miller SD, Settipane RJ, et al. Delayed nasal mucociliary clearance in patients with nonallergic rhinitis and nasal eosinophilia. *Allergy Proc* 1992;13:81–84.
50. Connell JT. Nasal mastocytosis. *J Allergy* 1969;43:182.
51. McKenna EL. Nasal mastocytosis. *Laryngoscope* 1974;84:112–115.
52. Asthma and the Other Allergic Diseases. Washington DC. N.I.A.I.D. Task Force Report;1979:23–31. US Dept. Of Health, Education and Welfare, National Institutes of Health Publication 79–387.
53. Lund VJ, Kennedy DW. Quantification for staging sinusitis. *Ann Otol Rhinol Laryngol* 1995;104(Suppl):17–21.
54. Gwaltney JM Jr. Microbiology of sinusitis. In: Druce HM, ed. *Sinusitis*, New York: Marcel Dekker, 1994:50.
55. Stierna P, Carlsoo B. Histopathological observations in chronic maxillary sinusitis. *Acta Otolaryngol* 1990;110:450–458.
56. Georgitis JW, Matthews BL, Stone B. Chronic sinusitis: characterization of cellular influx and inflammatory mediators in sinus lavage fluid. *Int Arch Allergy Immunol* 1995;106:416–421.
57. Incaudo GA, Schatz M. Rhinosinusitis associated with endocrine conditions: hypothyroidism and pregnancy. In: Schatz M, Zeiger RS, Settipane GA, eds. *Nasal Manifestations of Systemic Diseases*, Providence, OceanSide Publications, 1991:54.
58. Topozada H, Michaels L, Topozada M, et al. The human respiratory nasal mucosa in pregnancy. An electron microscopic and histochemical study. *J Laryngol Otol* 1982;96:613–626.
59. Sorri M, Bortikanen-Sorri AL, Kanja J. Rhinitis during pregnancy. *Rhinology* 1980;18:83–86.
60. Schatz M, Hoffman CP, Zeiger RS, et al. The course and management of allergic diseases during pregnancy. In: Middleton E, Reed CE, Ellis EF, et al, eds. *Allergy Principles and Practice*, 4th ed. Vol II. St. Louis, MO: Mosby, 1997.
61. Reiter D, Myers AR. Asymptomatic septal perforations in systemic lupus erythematosus. *Ann Otol Rhinol Laryngol* 1980;89:78.
62. Hughes RA, Berry CL, Seifert M, Lesosof MH. Relapsing polychondritis. Three cases with a clinico-pathological study and literature review. *Q J Med* 1972;41:363–380.
63. Henkin RI, Talal N, Larson AL, Mattern CF. Abnormalities of taste and smell in Sjögren's syndrome. *Ann Intern Med* 1972;76:375–383.
64. Wilson R, Lund V, Sweatman M, et al. Upper respiratory tract involvement in sarcoidosis and its management. *Eur Respir J* 1988;1:269–272.
65. Fauci AS, Haynes BF, Katz P, Wolff SM. Wegener's granulomatosis: prospective clinical and therapeutic experience with 85 patients for 21 years. *Ann Intern Med* 1983;98:76–85.
66. Graf P, Hallen H, Juto JE. The pathophysiology and treatment of rhinitis medicamentosa. *Clin Otolaryngol* 1995;20:224–229.
67. Connell J. Nasal hypersensitivity in comprehensive immunology. In: Gupta S, Good RA, eds. *Cellular, Molecular and Clinical Aspects of Allergic Disorders*. Vol. 6. New York, NY: Plenum, 1979:397.
68. Wentges RT. Edward Woakes: the history of an eponym. *J Laryngol Otol* 1972;86:501–512.
69. Oppenheimer EH, Rosenstein BJ. Differential pathology of nasal polyps in cystic fibrosis and atopy. *Lab Invest* 1979;40:445–449.
70. Rowe-Jones JM, Shembekar M, Trendell-Smith N, et al. Polypoidal rhinosinusitis in cystic fibrosis: a clinical and histopathological study. *Clin Otolaryngol* 1997;22:167–171.
71. Salari H, Borgeat P, Steffenrud S, et al. Immunological and nonimmunological release of leukotrienes and histamine from human nasal polyps. *Clin Exp Immunol* 1986;63:711–717.
72. Bachert C, Wagenmann M, Hauser U, et al. IL-5 synthesis is up regulated in human nasal polyp tissue. *J Allergy Clin Immunol* 1997;99:837–842.
73. Bumsted RM, El-Ackad T, Smith JM, Brody MJ. Histamine, norepinephrine and serotonin content of nasal polyps. *Laryngoscope* 1979;89:832–843.
74. Chandra RK, Abrol BM. Immunopathology of nasal polypi. *J Laryngol Otol* 1974;88:1019–1024.
75. Keith PK, Conway M, Evans S, et al. Nasal polyps: effects of seasonal allergen exposure. *J Allergy Clin Immunol* 1994;93:567–574.
76. Lildholdt T, Rundcrantz H, Bende M, Larsen K. Glucocorticoid treatment for nasal polyps. The use of topical budesonide powder, intramuscular betamethasone, and surgical treatment. *Arch Otolaryngol Head Neck Surg* 1997;123:595–600.
77. Downing ET, Braman S, Settipane GA. Bronchial reactivity in patients with nasal polyps before and after polypectomy. *J Allergy Clin Immunol* 1982;69:102.
78. Zlab MK, Moore GF, Daly DT, Yonkers AJ. Cerebrospinal fluid rhinorrhea: a review of the literature. *Ear Nose Throat J* 1992;71:314–317.
79. Skedros DG, Cass SP, Hirsch BE,

- Kelly RH. Beta-2 transferrin assay in clinical management of cerebral spinal fluid and perilymphatic fluid leaks. *J Otolaryngol* 1993;22:341–344.
80. Sillers MJ, Morgan CE, el-Gammal T. Magnetic resonance cisternography and thin coronal computerized tomography in the evaluation of cerebrospinal fluid rhinorrhea. *Am J Rhinol* 1997;11:387–392.
81. Han-Sen C. The ozena problem. Clinical analysis of atrophic rhinitis in 100 cases. *Acta Otolaryngol* 1982;93:461–464.
82. Holopainen E. Nasal mucous membrane in atrophic rhinitis with reference to symptomfree nasal mucosa. Histology, histochemistry, and exfoliative cytology. *Acta Otolaryngol* 1967;227(Suppl):1–55.
83. Proud D, Bailey GS, Naclerio RM, et al. Tryptase and histamine as markers to evaluate mast cell activation during the responses to nasal challenge with allergen, cold, dry air, and hyperosmolar solutions. *J Allergy Clin Immunol* 1992;89:1098–1110.
84. Togias AG, Naclerio RM, Proud D, et al. Nasal challenge with cold, dry air results in the production of inflammatory mediators: possible mast cells involvement. *J Clin Invest* 1985;76:1375–1381.
85. Raphael GD, Hauptschein-Raphael M, Kaliner MA. Gustatory rhinitis. *Am J Rhinol* 1989;3:145.
86. Bascom R. Air Pollution: In Mygind N, Naclerio RM, eds. *Allergic and Nonallergic Rhinitis*. Copenhagen: Munksgaard, 1993:33.
87. Coggins CR, Fouillet XL, Lam R, Morgan KT. Cigarette smoke induced pathology of the rat respiratory tract: a comparison of the effects of the particulate and vapour phases. *Toxicology* 1980;16:83–101.
88. Black A, Evans JC, Hadfield EH, et al. Impairment of nasal mucociliary clearance in woodworkers in the furniture industry. *Br J Ind Med* 1974;31:10–17.
89. Graham D, Henderson F, House D. Neutrophil influx measured in nasal lavages of humans exposed to ozone. *Arch Environ Health* 1988;43:228–233.
90. Olsen JH, Jensen SP, Hink M, et al. Occupational formaldehyde exposure and increased nasal cancer risk in man. *Int J Cancer* 1984;34:639–644.
91. Slavin RG. Occupational rhinitis. *Immunol Allergy Clin North Am* 1992;12:769.
92. Bardana EJ. Occupational asthma and related respiratory disorders. *Dis Mon* 1995;41:143–199.
93. Bernstein DI. Clinical assessment and management of occupational asthma. In: Bernstein IL, Chan-Yeung M, Malo JL, eds. *Asthma in the Workplace*. New York, NY: Marcel Dekker, 1993:103–123.

Requests for reprints should be addressed to:
Russell A. Settipane, MD
95 Pittman Street
Providence, RI 02906
E-mail: setti5@aol.com

CME Examination

No XXX-XXX

1–5, Settipane RA and Lieberman P. 2001;86:494–508.

CME Test Questions

- True statements regarding chronic nonallergic rhinitis include which one of the following?
 - The occurrence of asthma in patients with nonallergic rhinitis is greater than the occurrence of asthma in allergic asthma.
 - Conjunctivitis occurs more frequently in nonallergic rhinitis patients than in those with allergic rhinitis.
 - Sneezing occurs more frequently in patients with nonallergic rhinitis than in those with allergic rhinitis.
 - Biopsy studies of patients with nonallergic rhinitis reveal large numbers of neutrophils invading vessel walls.
- Topical medications demonstrating efficacy in controlled studies and/or having FDA approval for the therapy of nonallergic rhinitis include:
 - Fluticasone.
 - Beclomethasone (aqueous).
 - Budesonide (aerosol).
 - Azelastine.
 - All of the above.
- Rhinorrhea occurring after the ingestion of food (gustatory rhinitis) is attributable to:
 - Mast cell degranulation.
 - Basophil degranulation.
 - Activation of the sympathetic nerve supply to the nose.
 - Activation of the parasympathetic nerve supply to the nose.
 - Activation of the nonadrenergic noncholinergic (NANC) nerve supply to the nose.
- Which of the following forms of rhinitis is characterized by an inflammatory infiltrate on biopsy?
 - Allergic rhinitis.
 - Rhinitis medicamentosa.
 - Gustatory rhinitis.
 - Rhinitis in pregnancy.
 - All of the above.

-
- | | | |
|--|---|--|
| 5. True statement(s) regarding the epidemiology of mixed rhinitis (a combination of nonallergic and allergic rhinitis occurring in the same patient) include(s): | A. It is more common than allergic rhinitis alone.
B. It is more common than the combined incidence of pure allergic rhinitis and pure nonallergic rhinitis. | C. It is more common in children than adults.
D. It is usually characterized by seasonal symptoms only.
E. All of the above. |
|--|---|--|
-

Answers to CME examination—Annals of Allergy, Asthma, and Immunology, May 2001 (Identification No XXX-XXX) Settipane RA and Lieberman P: Update on non-allergic rhinitis. *Ann Allergy Asthma Immunol* 2001;86: XXX-XXX.

1. e
2. e
3. d
4. a
5. a