

Received Date : 17-May-2015

Revised Date : 22-Jul-2015

Accepted Date : 17-Aug-2015

Article type : Original Article: Epidemiology and Genetics

Editor : Antonella Muraro

**Maternal nutrition during pregnancy and risk of asthma, wheeze and atopic diseases
during childhood: a systematic review and meta-analysis**

Andrea A. Beckhaus¹, Luis Garcia-Marcos², Erick Forno³, Rosa M. Pacheco-Gonzalez², Juan C. Celedón³, Jose A. Castro-Rodriguez.¹

¹School of Medicine, Pontificia Universidad Católica de Chile, Santiago, Chile; ²Respiratory and Allergy Units. Arrixaca University Children's Hospital, University of Murcia; and IMIB Bio-research Institute, Murcia, Spain; ³University of Pittsburgh, Pittsburgh, Pennsylvania, USA.

Corresponding author:

Jose A. Castro Rodriguez, MD, PhD

Departments of Pediatrics and Public Health, School of Medicine,

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/all.12729

This article is protected by copyright. All rights reserved.

Pontificia Universidad Católica de Chile, Lira 44, 1er Piso, casilla 114-D, Santiago, Chile.

Telephone: (56) 2 354 8189, FAX: (56) 2 354 8122, E-mail: jacastro17@hotmail.com

Funding Sources: This work was supported by the grant Fondecyt (Conicyt-Chile) #1141195.

Conflicts of interest: The authors declare that there are no conflicts of interest.

Running head: Maternal nutrition pregnancy and risk of wheeze

Key words: Nutrition, diet, pregnancy, asthma, wheeze, atopy, children.

ABSTRACT

BACKGROUND: Epidemiologic studies suggest a relationship between maternal nutrition during pregnancy and the occurrence of asthma and atopic conditions during childhood. However, individual study results are conflicting. The objective of this meta-analysis is to critically examine the current evidence for an association between nutrition (dietary patterns, food groups, vitamins or oligo-elements) ingestion during pregnancy and asthma, wheeze or atopic conditions in childhood. **METHODS:** The inclusion criteria were: (1) systematic recording of diet during the gestational period and (2) documentation of asthma, wheezing, eczema or other atopic disease in the offspring. The primary outcomes were prevalence of asthma or wheeze among the offspring during childhood; and secondary outcomes were prevalence of eczema, allergic rhinitis or other atopic conditions. **RESULTS:** We found 120 titles, abstracts and citations, and 32 studies (29 cohorts) were included in this analysis. Data on vitamins, oligo-elements, food groups and dietary patterns during pregnancy were collected. A meta-analysis revealed that higher maternal intake of vitamin D (odds ratio [OR]= 0.58, 95% confidence interval [CI]= 0.38 to 0.88), vitamin E (OR= 0.6, 95% CI=0.46 to 0.78) and zinc

(OR=0.62, 95% CI=0.40 to 0.97) were each associated with lower odds of wheeze during childhood. However, none of these or other nutrients was consistently associated with asthma *per-se* or other atopic conditions. **CONCLUSIONS:** Current evidence suggests a protective effect of maternal intake of each of three vitamins or nutrients (vitamin D, vitamin E and zinc) against childhood wheeze but is inconclusive for an effect on asthma or other atopic conditions.

INTRODUCTION

Asthma and allergies are a major public health burden worldwide. Over the last few decades, the prevalence of childhood asthma has noticeably increased¹. In the United States (U.S.), such prevalence increased from 7.3% in 2001 to 8.4% in 2010 (and from 3.5% in the 1970s). In 2010, an estimated 25.7 million people had current or active asthma in the U.S., comprising 18.7 million adults and 7 million children². This “asthma epidemic” is likely related to changes in environment or lifestyle, such as those in diet and nutrition³.

Changes in dietary patterns preceded the increased occurrence of allergic diseases in the United Kingdom and other industrialized countries⁴. In particular, consumption of a “westernized” diet with low intake of antioxidants (and thus increased oxidative stress) may explain this temporal trend⁵.

In utero exposures may markedly increase the risk of asthma or allergies in childhood⁶, and thus identifying such exposures is relevant to primary prevention of these common conditions. Findings from experimental and epidemiologic studies suggest that one such exposure -maternal diet or maternal nutrient status during pregnancy- affects the incidence of asthma and/or other allergic diseases (food allergy, eczema and allergic rhinitis) in childhood^{7,8}, perhaps through

effects on the development of the fetal airways or immune responses^{5,9}. However, the available evidence is inconclusive and of variable quality, depending on the dietary pattern or nutrient of interest and considering current evidence, no recommendations about increasing or avoiding the consumption of these nutrients to prevent or ameliorate allergic disease in their offspring can be made⁸.

The objective of this meta-analysis is to systematically evaluate prior published studies in addition to recently published evidence on whether dietary patterns, food groups and vitamins or oligo-elements ingestion during pregnancy are associated with increased risk of asthma, wheeze or atopic conditions during childhood.

METHODS

Search and Selection Criteria

We searched electronic databases (March and November 2014) Medline, CENTRAL (Cochrane Collaboration clinical trials register), LILACS, and CINAHL. The search process was conducted using the keywords: (asthma OR wheeze* OR dermatitis OR eczema OR rhinitis) AND ((maternal diet OR pregnancy diet OR (pregnancy AND diet)) AND (offspring OR infant OR child*). We also searched in other non-bibliographic data sources, such as web searching, references of included publications, and pharmaceutical industry web sites. If the information was incomplete, we attempted to contact the authors. The inclusion criteria were: (1) systematic recording of diet during the gestational period (daily register or food frequency questionnaire); (2) assessment of asthma, wheezing, eczema or other atopic disease in the offspring, and (3) no language restriction. The exclusion criteria were: (1) vitamin or oligo-element supplementation

to the pregnant mother or her offspring, (2) measurement of nutrients in blood (e.g. serum or plasma) samples with no recording of diet during pregnancy, (3) reviews or letters or abstracts, in order to have full information and results of all included studies. The primary outcomes of the study were asthma and wheeze during childhood. Secondary outcomes included eczema, allergic rhinitis or other atopic condition during childhood.

Data Abstraction and Assessment of Risk of Bias

This systematic review was performed according to the Cochrane methodology and the recommendations for reporting proposed by the Meta-analysis of Observational Studies in Epidemiology group¹⁰ and to the Preferred Reporting Items for Systematic Reviews and Meta-analyses guidelines (PRISMA)¹¹. Titles, abstracts and citations were independently analyzed by two independent investigators (JACR and LGM). From the full texts, the reviewers independently assessed all studies for inclusion, based on the criteria for population intervention, study design, and outcomes. After obtaining full reports about potentially relevant trials, they assessed eligibility. Disagreements were resolved through a mediator (EF). The risk of bias from including certain studies was assessed according to the Newcastle–Ottawa Scale¹².

Data Analysis

We calculated pooled odds ratios (ORs), relative risk ratios (RRs) and hazard ratios (HRs) with 95% confidence interval (CI) for categorical outcomes. Heterogeneity was assessed using the I^2 test and chi-square. For all of the outcomes measured, a fixed-effects meta-analysis was used when low heterogeneity was present ($I^2 < 40\%$), and a random-effects meta-analysis was performed to address the variation across the included studies when high heterogeneity was

detected ($I^2 \geq 40\%$). Data from cohort and retrospective studies were pooled into separate meta-analyses. For each food group or oligo-element, high consumption was compared with low consumption. Some studies reported the same associations with several outcomes at different time of follow up. When this happened, we used only one given estimate of effect (e.g. OR) to perform the meta-analysis, to avoid considering the same data twice. The meta-analysis was performed using Review Manager 5.1.6 software (The Nordic Cochrane Centre, The Cochrane Collaboration, 2011, Copenhagen, Denmark).

RESULTS

A total of 122 studies were initially identified from databases and other sources: 32 of these studies met our inclusion criteria and were included in the review, as listed in **Figure 1**. All of the 32 studies included in the analysis were published between 2002 and 2014: 29 were prospective cohort studies and 3 were retrospective cohort studies. A detailed description of these 32 studies is shown in **Table 1**, and a description of our quality assessment is presented in **Table 2**. Our main results are summarized in **Table 3**.

Vitamin A

Seven studies reported on maternal intake of vitamin A or its constituent carotenoids and asthma or atopic outcomes^{13,14,15,16,17,18,19}. Two studies^{17,19} found no significant association between maternal vitamin A intake during pregnancy and childhood asthma. Alpha-carotene consumption was not associated with wheezing^{15,16} or eczema^{16,17}, and β carotene was not associated with wheezing^{13,14,15,16,18} (pooled OR=0.97, 95% CI=0.68-1.37, I^2 0%), eczema^{16,17,18} (pooled OR=0.76, 95% CI=0.46-1.26, I^2 72%), food allergy¹⁸ or skin test reactivity (STR) to allergens¹⁸.

Vitamin B complex

Four cohort studies reported on maternal intake of folic acid (vitamin B9) during pregnancy and asthma or atopic outcomes^{15,17,20,21}. One study¹⁷ found no significant association between maternal intake of folic acid during pregnancy and asthma or allergic rhinitis in childhood. Two studies showed no significant association between maternal intake of folic acid and wheeze^{15,20} (pooled OR=0.91, 95% CI=0.49-1.68, I² 48%) and three studies showed no relation with eczema^{17,20,21} (pooled OR=0.91, 95% CI=0.76-1.09, I² 0%). Two studies^{17,20} showed no significant association between maternal intake of vitamin B2 during pregnancy and childhood eczema (pooled OR=0.86, 95%CI 0.74-1.01, I² 0%). In another cohort study²⁰, there was no significant association between maternal dietary intake of vitamin B12 or vitamin B6 during pregnancy and childhood wheeze or eczema.

Vitamin C

Six cohort studies examined maternal dietary intake of vitamin C and asthma or atopic conditions^{13,14,15,16,17,18}. Four studies^{13,15,16,18} found no significant association between maternal intake of vitamin C during pregnancy and childhood wheeze (pooled OR=0.99, 95% CI= 0.48-2.04, I²=79%). Five studies^{13,14,16,17,18} found no significant association between prenatal dietary intake of vitamin C and childhood eczema (pooled OR=0.95, 95% CI=0.69-1.31, I²=56%). Two studies^{14,17} showed no significant association between maternal dietary intake of vitamin C during pregnancy and asthma or other atopic outcomes (allergic rhinitis, food allergy or STR to allergens)¹⁸.

Vitamin D

Five cohort studies reported on maternal dietary intake of vitamin D during pregnancy and asthma or atopic diseases^{22,23,24,25,26}. Maternal vitamin D intake during pregnancy was associated with a reduction in the odds of childhood wheeze^{22,23,25,26} (pooled OR=0.58, 95%CI=0.38-0.88, $I^2=59\%$; **Figure 2**). We found no significant association between maternal dietary intake of vitamin D during pregnancy and asthma^{24,25}, eczema^{22,24,25,26} (pooled OR=0.76, 95% CI=0.46-1.26, $I^2=72\%$) or allergic rhinitis²⁴. Our findings thus support an inverse association between maternal vitamin D intake during pregnancy and childhood wheeze. However, our results do not support an association between maternal vitamin D intake during pregnancy and asthma or other atopic diseases during childhood.

Vitamin E CAN

Seven cohort studies examined maternal dietary intake of vitamin E and atopic conditions during childhood^{13,14,15,16,17,18,19}. Maternal vitamin E intake was associated with a 46% reduction in the odds of childhood wheeze^{13,14,15,16,18} (pooled OR=0.54, 95% CI=0.41-0.71, $I^2=0\%$ [**Figure 3**]). There were no significant results for asthma^{14,17,19}, eczema^{13,16,17,18} (pooled OR=0.87, 95% CI=0.65-1.18, $I^2=48\%$), allergic rhinitis¹⁷, food allergy¹⁸ or STR to allergens¹⁸. Hence, our results are consistent with an inverse association between maternal dietary intake of vitamin E during pregnancy and childhood wheeze. We found no supportive evidence of an association between maternal vitamin E intake during pregnancy and asthma or any atopic condition during childhood.

Zinc

Six cohort studies examined the relation between maternal dietary intake of zinc during pregnancy and the outcomes of interest^{13,14,15,16,17,18}. Maternal zinc intake during pregnancy was associated with a reduction in the odds of childhood wheeze^{15,16,18} (pooled OR=0.57, 95% CI=0.40-0.81, $I^2=0\%$ [Figure 4]). There were no significant results for asthma^{14,17}, eczema^{13,14,16,17,18} (pooled OR=1.00, 95% CI=0.69-1.45, $I^2=63\%$), allergic rhinitis¹⁷, food allergy¹⁸ or STR to allergens¹⁸. Thus, our results support an inverse association between maternal zinc intake during pregnancy and childhood wheeze. On the other hand, our findings do not support an association between prenatal intake of zinc and asthma or allergies in childhood.

Copper

Of the five cohort studies that examined maternal dietary intake of copper during pregnancy and asthma or allergic diseases during childhood^{13,14,15,17,18}, one¹⁸ found an inverse association between maternal copper intake and food allergy in their offspring (OR 0.38, 95%CI=0.15-0.95), but none found a significant association with asthma^{14,17} wheeze^{13,14,15,18}, eczema^{13,14,17,18}, allergic rhinitis^{14,17} or STR to allergens^{14,18} during childhood. Thus, our findings offer limited support for an inverse association between maternal copper intake during pregnancy and food allergy during childhood.

Calcium

Three cohort studies reported the association between maternal dietary intake of calcium during pregnancy and asthma or other allergic diseases during childhood^{17,25,26}. We found no significant association between maternal calcium intake and asthma^{17,26}, wheeze^{25,26}, eczema^{17,25,26} (pooled

OR=0.94, 95% CI=0.82-1.08, $I^2=0\%$) or allergic rhinitis¹⁷ during childhood.

Magnesium

Three cohort studies examined maternal dietary intake of magnesium and atopic conditions during childhood^{13,14,17}. Although maternal magnesium intake during pregnancy was significantly associated with eczema in one study¹⁷ (OR=0.78, 95% CI=0.62-0.97), our analysis yielded negative results for asthma¹⁷, wheeze¹³ and allergic rhinitis^{14,17}. Thus, there is limited evidence of an inverse association between maternal magnesium intake during pregnancy and childhood eczema.

Manganese and Selenium

Two cohort studies found no significant association between maternal dietary intake of manganese during pregnancy and asthma¹⁷ or any atopic condition (including wheeze¹³, eczema¹⁷ and allergic rhinitis¹⁷) during childhood. Two cohort studies found no significant evidence of an association between maternal dietary intake of selenium during pregnancy and asthma¹⁷, wheeze¹³, eczema¹⁷ or allergic rhinitis¹⁷ during childhood.

Vegetables

Eight prospective cohort studies^{16,27,28,29,30,31,32,33} and one retrospective cohort study³⁴ assessed maternal consumption of vegetables during pregnancy and atopic conditions in childhood. Maternal vegetable intake during pregnancy was associated with a reduction in the odds of eczema using data from two studies with similar design^{16,31} (pooled OR=0.71, 95% CI=0.53-0.96, $I^2=0\%$), but not in a third study³³ (RR=1.21, 95% CI= 0.97-1.51). The later study did not

report OR for any results, so these results were not used to calculate the pooled OR. There were negative findings for wheeze^{16,29,30,32} (pooled OR=0.90, 95% CI= 0.69-1.18, I²=62%), asthma^{31,32}, allergic rhinitis³² or STR to allergens²⁹ in childhood.

Fruits

Six prospective cohort studies^{16,27,28,30,32,33} and one retrospective cohort study³⁴ evaluated the relation between maternal intake of fruits during pregnancy and the outcomes of interest. We found no significant association between maternal fruit intake during pregnancy and asthma³², wheeze^{16,30,32} (pooled OR=1.03, 95% CI=0.81-1.31, I²=56%), eczema^{16,33} or allergic rhinitis³² in childhood.

Fish

Eleven prospective cohort studies^{27,28,29,30,31,32,33,35,36,37,38} and one retrospective cohort study³⁴ examined maternal fish consumption during pregnancy and asthma or allergic diseases. We found no significant association between maternal fish intake during pregnancy and wheeze^{30,32,35,36,38} (pooled OR=1.00, 95% CI=0.89-1.13, I²=26%), asthma^{31,32}, eczema^{27,28,31,35,36,38} (pooled OR=0.84, 95% CI=0.68-1.05, I²=56%), allergic rhinitis^{28,32}, or sensitization to allergens (assessed by specific IgEs^{27,35} or allergy skin testing³⁵) in childhood.

Meat

Six cohort studies evaluated the association between maternal meat intake during pregnancy and atopic conditions^{28,32,33,36,37,38} and so did one retrospective study³⁴. There was no relation between maternal meat intake and wheezing^{32,36,38} (pooled OR 1.16, 95%CI 0.97-1.40, I² 35%),

asthma³², eczema^{36,38} or allergic rhinitis³².

Dairy

Six prospective cohort studies^{25,26,27,28,30,32,33} and one retrospective cohort study³⁴ examined maternal consumption of dairy products during pregnancy and atopic diseases in childhood. Maternal dairy intake during pregnancy was associated with a 17% reduction in the risk of wheeze in one cohort³³ (95% CI for RR=0.72-0.96) but not in other four studies^{25,30,32,26} (pooled OR=0.88, 95% CI=0.69-1.13, $I^2=62\%$). There were negative findings for asthma^{26,32}, eczema^{25,26,27,32}, allergic rhinitis³² or an increased total IgE²⁷ during childhood.

Fats

Six cohort studies investigated the association between maternal intake of (any type of) fats during pregnancy and asthma or atopic outcomes in childhood^{32,33,36,37,38,39}. Maternal consumption of saturated fatty acids (SFAs) was associated with decreased odds of wheeze in three studies^{36,38,39} (pooled OR=0.80, 95% CI=0.65-0.97, $I^2=0\%$), but not in a fourth study³³ (RR=1.08, 95% CI=0.93-1.26). There were negative findings for childhood eczema^{36,38,39} or allergic rhinitis³⁹.

Maternal dietary intake of monounsaturated fatty acids (MUFAs) during pregnancy was not associated with wheeze^{33,36,38,39}, eczema^{33,36,38,39} or allergic rhinitis³⁹ in childhood. Similarly, there was no significant association between maternal consumption of n3 or n6 polyunsaturated fatty acids (PUFAs) during pregnancy and wheeze^{36,38,39}, eczema^{36,38,39} or allergic rhinitis³⁹ in childhood. We also found no significant association between maternal intake of total fats during

pregnancy and asthma³², eczema^{36,38,39} or allergic rhinitis^{32,39} in childhood.

In summary, we found limited and inconsistent evidence of an inverse association between maternal intake of SFAs during pregnancy and wheeze. Maternal consumption of any other type of fats (MUFAs, n3 PUFAs, n6 PUFAs) or total fat during pregnancy was not significantly associated with asthma or any other atopic conditions.

Mediterranean Diet and Other Dietary Patterns

Two prospective cohort studies^{29,33} and two retrospective cohort studies^{40,41} examined the relation between a maternal “Mediterranean diet” during pregnancy and asthma or other atopic conditions. Whereas two studies^{29,40} found a significant inverse association between a maternal “Mediterranean diet” and atopic conditions (including persistent wheeze²⁹, atopic wheeze²⁹, STR to allergens²⁹, asthma ever⁴⁰, and rhinitis ever⁴⁰) at ages 6-7 years, the other two studies yielded negative findings for wheeze or eczema^{33 41}.

Two cohort studies analyzed the association between specific maternal dietary patterns during pregnancy and the outcomes of interest^{42,43}. In the first study⁴², none of five maternal dietary patterns (health conscious, traditional, processed, vegetarian and confectionery) was significantly associated with asthma, wheeze, allergic rhinitis or eczema in childhood. The second study⁴³ examined three maternal dietary patterns during pregnancy (healthy, western and Japanese). In that study, a maternal western diet during pregnancy was inversely associated with wheeze (OR=0.59, 95% CI=0.35-0.98) but not with eczema in childhood. Neither a healthy diet nor a Japanese diet during pregnancy was associated with wheeze or eczema in childhood.

In summary, there is inconsistent or non-significant evidence for an association between maternal dietary patterns during pregnancy and asthma or other atopic conditions in childhood.

Sweetened Beverages

A single cohort study assessed maternal consumption of artificially sweetened soft drinks during pregnancy and asthma or other atopic conditions⁴⁴. In that study, maternal intake of ≥ 1 soft drink per day was associated with increased odds of asthma (OR=1.23, 95% CI= 1.13-1.33) but not with allergic rhinitis.

DISCUSSION

Because *in utero* exposures likely affect the development of asthma and other common allergic illnesses (food allergy, allergic rhinitis and eczema), identifying and modifying such exposures could have a substantial impact on public health. After conducting a meta-analysis of recently published studies on maternal dietary intake (as specific nutrients or dietary patterns) and asthma or atopic conditions in childhood, we found suggestive evidence of a protective effect of maternal dietary intake of each of three vitamins or nutrients (vitamin D, vitamin E and zinc) against wheeze during childhood. However, none of these nutrients was consistently associated with asthma *per se* or other atopic conditions, and thus there is inconclusive evidence for either a true cause-effect relationship or the mechanisms underlying our findings.

In a pooled analysis of four studies (including ~4,570 mother-child pairs), we found that higher maternal intake of vitamin D intake during pregnancy was significantly associated with reduced odds of wheeze. Our findings for vitamin D are consistent with those of previous reviews^{45,46}

which concluded that maternal dietary intake of vitamin D may protect against childhood wheeze. On the other hand, we and others have found inconsistent or non-significant evidence for an association between maternal vitamin D intake during pregnancy and atopy or asthma. Since wheeze in early childhood may be “transient” or “viral-induced” and not due to “true asthma”, these findings suggest that the maternal vitamin D intake during pregnancy may not be immune modulatory but either affect lung development or protect against viral infections (a theory supported by both experimental and human studies showing potential effects of vitamin D against severe asthma exacerbations in children, which are often induced by viral illnesses^{46,47}). Given that childhood wheeze causes significant suffering and is a major public health problem (regardless of its etiology), there is ample justification for ongoing clinical trials of prenatal vitamin D supplementation to prevent asthma or wheeze⁴⁸. Long-term follow up of children participating in these trials should help determine not only whether vitamin D supplementation during pregnancy helps prevent wheeze, but also whether such supplementation reduces the risk of asthma or other atopic conditions.

We also found that higher maternal intake of vitamin E or zinc during pregnancy is associated with reduced risk of childhood wheeze. Although the total sample size of the studies pooled in our meta-analysis was smaller for zinc than for vitamin E, all studies included in the meta-analysis of each nutrient had results in the same direction and there was very low heterogeneity ($I^2=0\%$).

Oxidative stress, an imbalance between pro-oxidant and antioxidant, can cause dysfunction in cell signaling and arachidonic acid metabolism, and increase airway and systemic

inflammation⁴⁹. Because vitamin E is an antioxidant and may inhibit secretion of IL-4 by T cells⁵⁰, there has been interest in the role of vitamin E supplementation to prevent or treat asthma in children and adults, particularly in children with asthma exposed to outdoor pollution⁵¹. However, a recent Cochrane review of five randomized controlled trials concluded that additional research is needed to establish whether vitamin E supplementation (in pre- or post-natal life) protects against inception or morbidity from asthma (including that induced by air pollution) or particular asthma endotypes (e.g. exercise-induced bronchoconstriction)⁵².

Zinc deficiency has been associated with impaired fetal lung growth in rats⁵³, and *ADAM33* (a gene implicated in asthma susceptibility) is a zinc-dependent metalloproteinase⁵⁴. At least one randomized controlled trial has reported that zinc supplementation, together with usual therapy, leads to improvement in lung function and respiratory symptoms in subjects with asthma⁵⁵.

Even though we found an inverse relation between maternal SFA and a western diet pattern intake with wheeze, this data come from a limited number of studies, so it should be cautiously interpreted.

We recognize limitations of our findings. Firstly, our results for maternal dietary patterns or certain nutrients during pregnancy must be cautiously interpreted due to the paucity of studies available for meta-analysis. In contrast to our results, a Mediterranean-style post-natal diet was significantly associated with lower risk of asthma in a recent meta-analysis that included almost 40,000 children in eight studies⁵⁶. Moreover, a recent large meta-analysis found that post-natal dietary intake of vegetables and/or fruits was associated with a reduced risk of wheeze or asthma

in children and adults⁵⁷. Secondly, we may have failed to identify some studies in spite of using a systematic approach, and thus the studies included may not represent the entire body of knowledge on this topic. Also, the pooled estimates are only adjusted for the covariates included in the original studies, and thus we cannot exclude residual confounding by unmeasured factors. Finally, all studies recorded food consumption using a food frequency questionnaire, not a daily register which would be more accurate. Some included cohorts previously performed validation of their questionnaires, comparing this instrument with a three-day diet record and serum biomarkers^{58,59}, finding a good correlation between all these methods.

In summary, there is insufficient evidence to recommend any dietary pattern or nutrient supplementation to prevent or treat asthma or allergies. Our findings fully support ongoing clinical trials of vitamin D supplementation during pregnancy to prevent wheeze or asthma.

On the other hand, additional research studies (either observational longitudinal studies or clinical trials of post-natal supplementation) are needed to determine whether a randomized clinical trial of prenatal supplementation of certain nutrients (e.g. vitamin E or its isoforms, zinc) is warranted, particularly given potential and long-lasting adverse effects (e.g. in cancer in adults supplemented with vitamin E). A more fundamental question in this field is whether a “whole-diet” approach is preferable to supplementation with individual nutrients or vitamins, particularly when such nutrients or vitamins are largely derived from dietary intake. Intake of single nutrients from food is generally correlated with the type of diet e.g. individuals with high intake of vegetables generally have high intake of e.g. both vitamin C and beta-carotene. We did not adjust each particular vitamin by other nutrients or dietary pattern. Ongoing and future

studies (both observational and interventional) should help answer this important question over the next few years.

REFERENCES

1. Moorman JE, Akinbami LJ, Bailey CM, et al. *National Surveillance of Asthma: United States, 2001-2010.*; 2012:1-67. <http://www.ncbi.nlm.nih.gov/pubmed/24252609>.
2. Akinbami LJ, Moorman JE, Bailey C, et al. Trends in asthma prevalence, health care use, and mortality in the United States, 2001-2010. *NCHS Data Brief.* 2012;(94):1-8. <http://www.ncbi.nlm.nih.gov/pubmed/22617340>.
3. Seaton a, Godden DJ, Brown K. Increase in asthma: a more toxic environment or a more susceptible population? *Thorax.* 1994;49:171-174. doi:10.1136/thx.49.2.171.
4. Consumption of selected household foods, 1942 to 1996. Food Statistics Branch, National Food Survey. 1996.
5. Devereux G, Seaton A. Diet as a risk factor for atopy and asthma. *J Allergy Clin Immunol.* 2005;115:1109-1118. doi:10.1016/j.jaci.2004.12.1139.
6. Leone L, Nocerino R, Canani RB. Mechanisms of the Immune System. 2014:4706-4719. doi:10.3390/nu6114706.
7. Netting M, Middleton P, Makrides M. Does maternal diet during pregnancy and lactation affect allergy outcomes in their offspring ? A systematic review of food based approaches. *Clin Transl Allergy.* 2013;3(11-12):O19. doi:10.1186/2045-7022-3-S3-O19.
8. Miles E a., Calder PC. Maternal diet and its influence on the development of allergic disease. *Clin Exp Allergy.* 2015;45:63-74. doi:10.1111/cea.12453.
9. Devereux G. Asthma and Allergy : Food for Thought. *Nature.* 2006;6(November):869-875.
10. Stroup DF, Berlin JA, Morton SC, Olkin I, Williamson GD, Rennie D, Moher D, Becker BJ, Sipe TA, Thacker SB. Meta-analysis of observational studies in epidemiology: a proposal for reporting. Meta-analysis Of Observational Studies in Epidemiology (MOOSE) group. *JAMA.* 2000 Apr 19;283(15):2008-12.
11. Liberati A, Altman DG, Tetzlaff J, et al. Annals of Internal Medicine Academia and Clinic The PRISMA Statement for Reporting Systematic Reviews and Meta-Analyses of Studies That Evaluate Health Care Interventions : *Ann Intern Med.* 2009;151(4):W65-W94. doi:10.1371/journal.pmed.1000100.
12. Wells GA, Shea B, O'Connell D, Peterson J, Welch V, et al. (2011) The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomized studies in meta-analysis. Available: www.ohri.ca/programs/clinical_epidemiology/oxford.asp. 2011:2011.
13. Martindale S, McNeill G, Devereux G, Campbell D, Russell G, Seaton A. Antioxidant intake in pregnancy in relation to wheeze and eczema in the first two years of life. *Am J Respir Crit Care Med.* 2005;171(39):121-128. doi:10.1164/rccm.200402-220OC.
14. Devereux G, Turner SW, Craig LC a, et al. Low maternal vitamin E intake during pregnancy is associated with asthma in 5-year-old children. *Am J Respir Crit Care Med.* 2006;174:499-507. doi:10.1164/rccm.200512-1946OC.

15. Litonjua AA, Rifas-Shiman SL, Ly NP, Tantisira KG, Rich-Edwards JW, Camargo CA Jr, Weiss ST, Gillman MW, Gold DR. Maternal antioxidant intake in pregnancy and wheezing illnesses in children at 2 y of age. *Am J Clin Nutr*. 2006 Oct;84(4):903-11.
16. Miyake Y, Sasaki S, Tanaka K, Hirota Y. Consumption of vegetables, fruit, and antioxidants during pregnancy and wheeze and eczema in infants. *Allergy Eur J Allergy Clin Immunol*. 2010;65:758-765. doi:10.1111/j.1398-9995.2009.02267.x.
17. Nwaru BI, Erkkola M, Ahonen S, et al. Intake of antioxidants during pregnancy and the risk of allergies and asthma in the offspring. *Eur J Clin Nutr*. 2011;65:937-943. doi:10.1038/ejcn.2011.67.
18. West CE, Dunstan J, McCarthy S, et al. Associations between maternal antioxidant intakes in pregnancy and infant allergic outcomes. *Nutrients*. 2012;4:1747-1758. doi:10.3390/nu4111747.
19. Maslova E, Hansen S, Strom M, Halldorsson TI, Olsen SF. Maternal intake of vitamins A, E and K in pregnancy and child allergic disease: a longitudinal study from the Danish National Birth Cohort. *Br J Nutr*. 2014;111:1096-1108. doi:10.1017/s0007114513003395.
20. Miyake Y, Sasaki S, Tanaka K, Hirota Y. Maternal B vitamin intake during pregnancy and wheeze and eczema in Japanese infants aged 16-24 months: the Osaka Maternal and Child Health Study. *Pediatr Allergy Immunol*. 2011;22(4):69-74. doi:10.1111/j.1399-3038.2010.01081.x.
21. Dunstan J a., West C, McCarthy S, et al. The relationship between maternal folate status in pregnancy, cord blood folate levels, and allergic outcomes in early childhood. *Allergy Eur J Allergy Clin Immunol*. 2012;67(3):50-57. doi:10.1111/j.1398-9995.2011.02714.x.
22. Camargo CA, Jr., Rifas-Shiman SL, Litonjua AA et al. Maternal intake of vitamin D during pregnancy and risk of recurrent wheeze in children at 3 y of age. *Am J Clin Nutr* 2007; 85: 788-795.
23. Devereux G, Litonjua A a, Turner SW, et al. Maternal vitamin D intake during pregnancy and early childhood wheezing. *Am J Clin Nutr*. 2007;85:853-859.
24. Erkkola M, Kaila M, Nwaru BI, et al. Maternal vitamin D intake during pregnancy is inversely associated with asthma and allergic rhinitis in 5-year-old children. *Clin Exp Allergy*. 2009;39:875-882. doi:10.1111/j.1365-2222.2009.03234.x.
25. Miyake Y, Sasaki S, Tanaka K, Hirota Y. Dairy food, calcium and vitamin D intake in pregnancy, and wheeze and eczema in infants. *Eur Respir J*. 2010;35(6):1228-1234. doi:10.1183/09031936.00100609.
26. Miyake Y, Tanaka K, Okubo H, Sasaki S, Arakawa M. Maternal consumption of dairy products, calcium, and vitamin D during pregnancy and infantile allergic disorders. *Ann Allergy, Asthma Immunol*. 2014;113(1):82-87. doi:10.1016/j.anai.2014.04.023.
27. Sausenthaler S, Koletzko S, Schaaf B, et al. Maternal diet during pregnancy in relation to eczema and allergic sensitization in the offspring at 2 y of age. *Am J Clin Nutr*. 2007;85(2):530-537.
28. Willers SM, Devereux G, Craig LC a, et al. Maternal food consumption during pregnancy and asthma, respiratory and atopic symptoms in 5-year-old children. *Thorax*. 2007;62:773-779. doi:10.1136/thx.2006.074187.
29. Chatzi L, Torrent M, Romieu I, et al. Mediterranean diet in pregnancy is protective for wheeze and atopy in childhood. *Thorax*. 2008;63:507-513. doi:10.1136/thx.2007.081745.

30. Willers SM, Wijga AH, Brunekreef B, et al. Maternal food consumption during pregnancy and the longitudinal development of childhood asthma. *Am J Respir Crit Care Med*. 2008;178:124-131. doi:10.1164/rccm.200710-1544OC.
31. Oien T, Storrø O, Johnsen R. Do early intake of fish and fish oil protect against eczema and doctor-diagnosed asthma at 2 years of age? A cohort study. *J Epidemiol Community Health*. 2010;64:124-129. doi:10.1136/jech.2008.084921.
32. Erkkola M, Nwaru BI, Kaila M, et al. Risk of asthma and allergic outcomes in the offspring in relation to maternal food consumption during pregnancy: A Finnish birth cohort study. *Pediatr Allergy Immunol*. 2012;23:186-194. doi:10.1111/j.1399-3038.2012.01272.x.
33. Chatzi L, Garcia R, Roumeliotaki T, Basterrechea M, Begiristain H. Mediterranean diet adherence during pregnancy and risk of wheeze and eczema in the first year of life : INMA (Spain) and RHEA (Greece) mother – child cohort studies British Journal of Nutrition. 2013;2058-2068. doi:10.1017/S0007114513001426.
34. Ushiyama Y, Matsumoto K, Shinohara M, Wakiguchi H, Sakai K, Komatsu T, Yamamoto S. Nutrition during pregnancy may be associated with allergic diseases in infants. *J Nutr Sci Vitaminol (Tokyo)*. 2002 Oct;48(5):345-51.
35. Romieu I, Torrent M, Garcia-Esteban R, et al. Maternal fish intake during pregnancy and atopy and asthma in infancy. *Clin Exp Allergy*. 2007;37:518-525. doi:10.1111/j.1365-2222.2007.02685.x.
36. Miyake Y, Sasaki S, Tanaka K, Ohfuji S, Hirota Y. Maternal fat consumption during pregnancy and risk of wheeze and eczema in Japanese infants aged 16-24months: the Osaka Maternal and Child Health Study. *Thorax*. 2009 Sep;64(9):815-21. doi: 10.1136/thx.2009.115931.
37. Saito K, Yokoyama T, Miyake Y, et al. Maternal meat and fat consumption during pregnancy and suspected atopic eczema in Japanese infants aged 3-4 months: The Osaka Maternal and Child Health Study. *Pediatr Allergy Immunol*. 2010;21(2):38-46. doi:10.1111/j.1399-3038.2009.00897.x.
38. Miyake Y, Tanaka K, Okubo H, Sasaki S, Arakawa M. Maternal fat intake during pregnancy and wheeze and eczema in Japanese infants: The Kyushu Okinawa Maternal and Child Health Study. *Ann Epidemiol*. 2013;23(11):674-680. doi:10.1016/j.annepidem.2013.08.004.
39. Nwaru BI, Erkkola M, Lumia M, et al. Maternal intake of fatty acids during pregnancy and allergies in the offspring. *Br J Nutr*. 2012;108:720-732. doi:10.1017/S0007114511005940.
40. De Batlle J, Garcia-Aymerich J, Barraza-Villarreal a, Antó JM, Romieu I. Mediterranean diet is associated with reduced asthma and rhinitis in Mexican children. *Allergy*. 2008;63:1310-1316. doi:10.1111/j.1398-9995.2008.01722.x.
41. Castro-Rodriguez J a., Garcia-Marcos L, Sanchez-Solis M, Pérez-Fernández V, Martinez-Torres A, Mallol J. Olive oil during pregnancy is associated with reduced wheezing during the first year of life of the offspring. *Pediatr Pulmonol*. 2010;45(March):395-402. doi:10.1002/ppul.21205.
42. Shaheen SO, Northstone K, Newson RB, Emmett PM, Sherriff a, Henderson a J. Dietary patterns in pregnancy and respiratory and atopic outcomes in childhood. *Thorax*. 2009;64:411-417. doi:10.1136/thx.2008.104703.

43. Miyake Y, Okubo H, Sasaki S, Tanaka K, Hirota Y. Maternal dietary patterns during pregnancy and risk of wheeze and eczema in Japanese infants aged 16-24months: The Osaka Maternal and Child Health Study. *Pediatr Allergy Immunol*. 2011;22(1):734-741. doi:10.1111/j.1399-3038.2011.01176.x.
44. Maslova E, Strøm M, Olsen SF, Halldorsson TI. Consumption of Artificially-Sweetened Soft Drinks in Pregnancy and Risk of Child Asthma and Allergic Rhinitis. *PLoS One*. 2013;8(2). doi:10.1371/journal.pone.0057261.
45. Han YY, Blatter J, Brehm JM, Forno E, Litonjua AA, Celedón JC. Diet and asthma: vitamins and methyl donors. *Lancet Respir Med*. 2013 Dec;1(10):813-22. doi: 10.1016/S2213-2600(13)70126-7.
46. Paul G, Brehm JM, Alcorn JF, Holguín F, Aujla SJ, Celedón JC. Vitamin D and asthma. *Am J Respir Crit Care Med*. 2012 Jan 15;185(2):124-32. doi: 10.1164/rccm.201108-1502CI.
47. Brehm JM, Acosta-Pérez E, Klei L, et al. Vitamin D insufficiency and severe asthma exacerbations in Puerto Rican children. *Am J Respir Crit Care Med*. 2012;186(10):140-146. doi:10.1164/rccm.201203-0431OC.
48. Litonjua A a., Lange NE, Carey VJ, et al. The Vitamin D Antenatal Asthma Reduction Trial (VDAART): Rationale, design, and methods of a randomized, controlled trial of vitamin D supplementation in pregnancy for the primary prevention of asthma and allergies in children. *Contemp Clin Trials*. 2014;38:37-50. doi:10.1016/j.cct.2014.02.006.
49. Moreno-Macias H, Romieu I. Effects of antioxidant supplements and nutrients on patients with asthma and allergies. *J Allergy Clin Immunol*. 2014;133(5):1237-1244. doi:10.1016/j.jaci.2014.03.020.
50. Li-Weber M, Giaisi M, Treiber M, Krammer P. Vitamin E inhibits IL-4 gene expression in peripheral blood T cells. *Eur J Immunol*. 2002;32:-. doi:10.1002/1521-4141(200209)32:9<2401::aid-immu2401>3.0.co;2-s.
51. Romieu I, Sienra-Monge JJ, Ramírez-Aguilar M, et al. Antioxidant supplementation and lung functions among children with asthma exposed to high levels of air pollutants. *Am J Respir Crit Care Med*. 2002;166(October 1998):703-709. doi:10.1164/rccm.2112074.
52. Wilkinson M, Hart A, Milan SJ, Sugumar K. Vitamins C and E for asthma and exercise-induced bronchoconstriction. *Cochrane Database Syst Rev*. 2014 Jun 17;6:CD010749.
53. Vojnik C, Hurley LS. Abnormal prenatal lung development resulting from maternal zinc deficiency in rats. *J Nutr*. 1977;107:862-872.
54. Powell RM, Hamilton LM, Holgate ST, Davies DE, Holloway JW. ADAM33: a novel therapeutic target for asthma. *Expert Opin Ther Targets*. 2003;7(4):485-494. doi:10.1517/14728222.7.4.485.
55. Ghaffari J, Khalilian a, Salehifar E, Khorasani E, Rezaii MS. Effect of zinc supplementation in children with asthma: a randomized, placebo-controlled trial in northern Islamic Republic of Iran. *East Mediterr Health J*. 2014;20:391-396. <http://www.ncbi.nlm.nih.gov/pubmed/24960516>.
56. Garcia-Marcos L, Castro-Rodriguez J a, Weinmayr G, Panagiotakos DB, Priftis KN, Nagel G. Influence of Mediterranean diet on asthma in children: a systematic review and meta-analysis. *Pediatr Allergy Immunol*. 2013;24(3):330-338. doi:10.1111/pai.12071.
57. Seyedrezazadeh E, Pour Moghaddam M, Ansarin K, Reza Vafa M, Sharma S, Kolahdooz F. Fruit and vegetable intake and risk of wheezing and asthma: A systematic review and meta-analysis. *Nutr Rev*. 2014;72(7):411-428. doi:10.1111/nure.12121.

58. Sasaki S, Ushio F, Amano K, et al. Serum biomarker-based validation of a self-administered diet history questionnaire for Japanese subjects. *J Nutr Sci Vitaminol (Tokyo)*. 2000;46(6):285-296. doi:10.3177/jnsv.46.285.
59. Sasaki S, Yanagibori R, Amano K. Self-administered diet history questionnaire developed for health education: a relative validation of the test-version by comparison with 3-day diet record in women. *J Epidemiol*. 1998;8(4):203-215. doi:10.2188/jea.8.203.

FIGURE LEGENDS

Figure 1.

Flowchart for identification of studies used.

Figure 2.

Pooled difference for maternal dietary vitamin D intake during pregnancy and childhood wheeze.

Figure 3.

Pooled difference for maternal dietary vitamin E intake during pregnancy and childhood wheeze.

Figure 4.

Pooled difference for maternal dietary zinc intake during pregnancy and childhood wheeze

Author	Geographic area	Design	Included participants	Dietary component measured	Outcomes	Statistical methods for potential confounder factor adjustments
Ushiyama 2002 ³⁴	Japan	Retrospective cohort	2070 mother-offspring pairs	FFQ of multiple nutrients intake including fish, meat, egg, dairy, vegetables, fruits, sugar, fatty food, total energy during pregnancy (obtained at recruitment time)	Atopic eczema, food allergies and asthma during recruitment time (Mean 7.6 ± 2.8 months after birth)	Logistic regression
Martindale 2005 ¹³	United Kingdom	Prospective Cohort	1924 mother-offspring pairs 1300 pairs at end of follow up (68%)	FFQ of antioxidante intake including vitamin E, C, β-carotene, selenium, magnesium, manganese, copper and zinc (obtained during pregnancy)	Wheeze and eczema, at 6, 12 and 24 months after birth	Logistic regression
Devereux 2006 ¹⁴	United Kingdom	Prospective Cohort	1924 mother-offspring pairs 1253 pairs at end of follow up (65%)	FFQ of vitamin E , C, β-carotene, selenium, magnesium, copper, iron and zinc intake (obtained during pregnancy)	Wheeze, eczema and asthma at 5 years of age	Logistic regression
Litonjua 2006 ¹⁵	United States	Prospective Cohort	1863 mother-offspring pairs 1290 pairs at end of follow up (69%)	FFQ of antioxidant intake including vitamin C, E, folic acid, α and β-carotene, copper and zinc (obtained during pregnancy)	Recurrent wheeze and eczema, at 6, 12 and 24 months after birth	Bivariate and multivariate logistic regression

Camargo 2007 ²²	United States	Prospective Cohort	1863 mother-offspring binomials 1194 pairs at end of follow up (64%)	FFQ of vitamin D intake (obtained during pregnancy)	Recurrent wheeze and eczema, at 6, 12, 24 and 36 months after birth	Multivariate logistic regression
Devereux 2007 ²³	United Kingdom	Prospective Cohort	1924 mother-offspring pairs 1253 pairs at end of follow up (65%)	FFQ of vitamin D intake (obtained during pregnancy)	Wheeze and asthma at 2 and 5 years of age Atopic sensitization, spirometry and FE _{NO} at 5 years of age	Multivariate logistic regression
Romieu 2007 ³⁵	Spain	Prospective Cohort	483 mother-offspring pairs 462 pairs at end of follow up (96%)	FFQ of dietary fish intake (obtained 3 months after delivery)	Eczema, atopy, persistent wheeze at 6.5 years of age	Multivariate logistic regression
Sausenthaler 2007 ²⁷	Germany	Prospective Cohort	3097 mother-offspring pairs 2641 pairs at end of follow up (85%)	FFQ of multiple nutrients intake including fish, egg, dairy, and sorts of vegetables and fruits (obtained few days after delivery)	Eczema, at 6, 12, 18 and 24 months after birth Allergic sensitization at 2 years of age	Logistic regression
Willers 2007 ²⁸	United Kingdom	Prospective Cohort	1924 mother-offspring pairs 1253 pairs at end of follow up (65%)	FFQ of multiple nutrients intake including fish, meat, egg, dairy, vegetables, fruits, sugar, fatty food, total energy	Asthma, wheeze, eczema and hayfever at 5 years of age Atopic sensitization, spirometry and FE _{NO} at 5 years of age	Multivariate logistic regression

				(obtained during pregnancy)		
Chatzi 2008 ²⁹	Spain	Prospective Cohort	482 mother-offspring pairs 468 pairs at end of follow up (97%)	FFQ evaluating mediterranean diet consumption (MD) and MD components (obtained 3 months after delivery)	Atopy and persistent wheeze at 6.5 years of age	Multivariate logistic regression
de Batlle 2008 ⁴⁰	Mexico	Retrospective cohort	1476 mother-offspring pairs	FFQ evaluating mediterranean diet consumption (MD) during pregnancy (obtained at recruitment time)	Asthma, wheeze, allergic rhinitis, during recruitment time (6-7 years of age)	Logistic regression
Willers 2008 ³⁰	The Netherlands	Prospective Cohort	4146 mother-offspring pairs 3320 pairs at end of follow up (80%)	FFQ of multiple nutrients intake including fish, egg, dairy, vegetables, fruits, sugar, fatty food, total energy during pregnancy (obtained during pregnancy)	Wheeze and asthma symptoms at 3 months after birth and then yearly from 1 to 8 years of age	Multiple logistic regression
Miyake Y 2009 ³⁶	Japan	Prospective Cohort	1002 mother-offspring pairs 763 pairs at end of follow up (76%)	FFQ of fat intake including fish and meat (obtained during pregnancy)	Wheeze and eczema at 2 to 9 and 16 to 24 months after birth	Multiple logistic regression
Erkkola 2009 ²⁴	Finland	Prospective Cohort	1669 pairs at end of follow up	FFQ evaluating vitamin D intake (obtained until 3 months after delivery)	Asthma, allergic rhinitis and eczema at 5 years of age (children with HLA-DQB1 susceptibility for type 1 diabetes)	Logistic regression

Shaheen 2009 ⁴²	United Kingdom	Prospective Cohort	14062 mother-offspring pairs at beginning, no description of number of pairs at end of follow up	FFQ evaluating 5 dietary patterns (obtained during pregnancy)	Wheeze, asthma, allergic rhinitis, eczema and atop at 7-8 years of age	Logistic regression
Castro-Rodriguez 2010 ⁴¹	Spain	Retrospective cohort	1409 mother-offspring binomials	FFQ of mediterranean diet and olive oil intake during pregnancy (obtained at recruitment time)	Wheeze during first year of live (15-18 months of age at recruitment)	Multivariate logistic regression
Miyake 2010 ²⁵	Japan	Prospective Cohort	1002 mother-offspring pairs 763 pairs at end of follow up (76%)	FFQ of dairy, calcium and vitamin D intake (obtained during pregnancy)	Wheeze and eczema at 16 to 24 months after birth	Multiple logistic regression
Miyake 2010 ¹⁶	Japan	Prospective Cohort	1002 mother-offspring pairs 763 pairs at end of follow up (76%)	FFQ of vegetables, fruits and several antioxidants, including vitamin C, vitamin E, α and β -carotene and zinc (obtained during pregnancy)	Wheeze and eczema at 16 to 24 months after birth	Multiple logistic regression
Oien 2010 ³¹	Norway	Prospective	3086 pairs at end of	FFQ of vegetables, fish and cod liver oil (obtained 12 months after	Asthma and eczema at 24 months	Multivariate logistic

		Cohort	follow up	delivery)	after birth	regression
Saito 2010 ³⁷	Japan	Prospective Cohort	1002 mother-offspring pairs 771 pairs at end of follow up (77%)	FFQ of fat intake including fish and meat (obtained during pregnancy)	Eczema at 3 to 4 months of age	Multiple logistic regression
Miyake 2011 ⁴³	Japan	Prospective Cohort	1002 mother-offspring pairs 763 pairs at end of follow up (76%)	FFQ evaluating 3 dietary patterns (obtained during pregnancy)	Wheeze and eczema at 16 to 24 months after birth	Multiple logistic regression
Miyake 2011 ²⁰	Japan	Prospective Cohort	1002 mother-offspring pairs 763 pairs at end of follow up (76%)	FFQ evaluating folic acid and vitamins B12, B6 and B2 (obtained during pregnancy)	Wheeze and eczema at 16 to 24 months after birth	Multiple logistic regression

Nwaru 2011 ¹⁷	Finland	Prospective Cohort	2441 pairs at end of follow up	FFQ of antioxidante intake including vitamin A, C, E, zinc, cooper, selenium, magnesium, manganese, calcium, folate, riboflavin (B2) (obtained until 3 months after delivery)	Asthma, allergic rhinitis and eczema at 5 years of age (children with HLA-DQB1 susceptibility for type 1 diabetes)	Logistic regression
Dunstan 2012 ²¹	Australia	Prospective Cohort	628 mother-offspring pairs 484 pairs at end of follow up (77%)	FFQ of folate intake (obtained during pregnancy)	Eczema, food allergy and allergic sensitization at 12 months of age	Logistic regression
Erkkola 2012 ³²	Finland	Prospective Cohort	2441 pairs at end of follow up	FFQ of multiple nutrients intake including fats, fish, meat, dairy, vegetables and fruits during pregnancy (obtained until 3 months after delivery)	Asthma, allergic rhinitis and wheezing at 5 years of age (children with HLA-DQB1 susceptibility for type 1 diabetes)	Logistic regression
Nwaru 2012 ³⁹	Finland	Prospective Cohort	2441 pairs at end of follow up	FFQ of dietary fats during pregnancy (obtained until 3 months after	Wheezing, allergic rhinitis and eczema at 5 years of age (children with HLA-DQB1 susceptibility for type 1 diabetes)	Logistic regression

				delivery)		
West 2012 ¹⁸	Australia	Prospective Cohort	450 mother-offspring pairs 300 pairs at end of follow up (67%)	FFQ of vitamin C, vitamin E, β -carotene, copper and zinc (obtained during pregnancy)	Wheeze, eczema, food allergy and allergic sensitization at 12 months of age	Multivariate logistic regression
Maslova 2013 ⁴⁴	Denmark	Prospective Cohort	60466 mother-offspring pairs 38149 pairs at end of follow up (63%)	FFQ of artificially sweetened soft drinks (obtained during pregnancy)	Asthma at 18 months of age and asthma and allergic rhinitis at 7 years of age	Logistic regression
Chatzi 2013 ³³	Spain and Greece	Prospective Cohort	Spain: 1784 pairs at end of follow up Greece: 1317 mother-offspring pairs 798 pairs at end of follow up (45%)	FFQ about mediterranean diet, vegetables, fruits, fish, meat and dairy products intake (obtained during pregnancy)	Wheeze and eczema at 12 months of age	Multivariate log-binominal regression

Miyake 2013 ³⁸	Japan	Prospective Cohort	1590 mother-offspring pairs 1362 pairs at end of follow up (86%)	FFQ of fat intake including fish and meat (obtained during pregnancy)	Wheeze and eczema at 4, 12 and 24 months after birth	Multiple logistic regression
Maslova 2014 ¹⁹	Denmark	Prospective Cohort	60466 mother-offspring pairs 28678 pairs at end of follow up (47%)	FFQ about vitamin A, E and K intake (obtained during pregnancy)	Asthma at 18 months of age and asthma and allergic rhinitis at 7 years of age	Multivariate logistic regression
Miyake 2014 ²⁶	Japan	Prospective Cohort	1590 mother-offspring pairs 1362 pairs at end of follow up (86%)	FFQ of dairy, calcium and vitamin D intake (obtained during pregnancy)	Wheeze and eczema at 4, 12 and 24 months after birth	Multiple logistic regression

FFQ: food frequency questionnaire

	Selection				Comparability	Outcome		
	Representative- ness of the exposed cohort	Selection of the non exposed cohort	Ascertainment of exposure	Demonstration that outcome of interest was not present at start of study	Comparability of cohorts on the basis of the design or analysis	Assessment of outcome	Follow-up long enough for outcomes to occur	Adequacy of follow up
Ushiyama 2002 ³⁴	*	*		*	*		*	
Martindale 2005 ¹³	*	*	*	*	*		*	
Devereux 2006 ¹⁴	*	*	*	*	*		*	
Litonjua 2006 ¹⁵	*	*	*	*	*		*	
Camargo 2007 ²²	*	*	*	*	*		*	
Devereux 2007 ²³	*	*	*	*	*		*	
Romieu 2007 ³⁵	*	*	*	*	*		*	*

Sausenthaler 2007 ²⁷	*	*	*	*	*		*	*
Willers 2007 ²⁸	*	*	*	*	*		*	
Chatzi 2008 ²⁹	*	*	*	*	*		*	*
de Batlle 2008 ⁴⁰	*	*		*	*		*	
Willers 2008 ³⁰	*	*	*	*	*		*	*
Miyake Y 2009 ³⁶	*	*	*	*	*		*	*
Erkkola 2009 ²⁴	*	*	*	*	*		*	
Shaheen 2009 ⁴²	*	*	*	*	*		*	
Castro- Rodriguez 2010 ⁴¹	*	*		*	*		*	

Miyake 2010 ²⁵	*	*	*	*	*		*	*
Miyake 2010 ¹⁶	*	*	*	*	*		*	*
Oien 2010 ³¹	*	*	*	*	*		*	
Saito 2010 ³⁷	*	*	*	*	*		*	*
Miyake 2011 ⁴³	*	*	*	*	*		*	*
Miyake 2011 ²⁰	*	*	*	*	*		*	*
Nwaru 2011 ¹⁷	*	*	*	*	*		*	
Dunstan 2012 ²¹		*	*	*	*		*	*
Erkkola 2012 ³²	*	*	*	*	*		*	

Nwaru 2012 ³⁹	*	*	*	*	*		*	
West 2012 ¹⁸		*	*	*	*		*	
Maslova 2013 ⁴⁴	*	*	*	*	*	*	*	
Chatzi 2013 ³³	*	*	*	*	*	*	*	
Miyake 2013 ³⁸	*	*	*	*	*		*	*
Maslova 2014 ¹⁹	*	*	*	*	*	*	*	
Miyake 2014 ²⁶	*	*	*	*	*		*	*

Selection

1) Representativeness of the exposed cohort

- a) truly representative *
- b) somewhat representative*
- c) selected group of users eg nurses, volunteers
- d) no description of the derivation of the cohort

2) Selection of the non exposed cohort

- a) drawn from the same community as the exposed cohort *
- b) drawn from a different source
- c) no description of the derivation of the non exposed cohort

3) Ascertainment of exposure

- a) secure record (eg surgical records) *
- b) structured interview*
- c) written self report
- d) no description

4) Demonstration that outcome of interest was not present at start of study

- a) yes *

b) no

Comparability

1) Comparability of cohorts on the basis of the design or analysis

a) study controls for the studied conditions *

b) study controls for any additional factor *

Outcome

1) Assessment of outcome

a) independent blind assessment *

b) record linkage *

c) self report

d) no description

2) Was follow-up long enough for outcomes to occur

a) yes *

b) no

3) Adequacy of follow up of cohorts

a) complete follow up *

- b) subjects lost to follow up unlikely to introduce bias , with follow up > 75 % or description provided of those lost *
- c) follow up rate < 75% and no description of those lost
- d) no statement

	Wheezing		Asthma		Eczema		Allergic Rhinitis		Food Allergy		Atopic Sensitization	
Exposure	Studies	Results	Studies	Results	Studies	Results	Studies	Results	Studies	Results	Studies	Results
Vitamin A												
α carotene	2	OR=1.16 [0.79-1.69]	1	HR=1.20 [0.92-1.56]	2	OR=0.95 [0.68-1.35]						
β carotene	3	OR=0.97 [0.68-1.37]	1	RR=1.02 [0.93-1.11]	3	OR=0.76 [0.46-1.26]			1	OR=0.38 [0.11-1.27]	1	OR=0.73 [0.33-1.61]
Vitamin B2	1	OR=1.27 [0.64-2.52]			2	OR=0.86 [0.74-1.01]						
Vitamin B6	1	OR=0.92 [0.47-1.77]			1	OR=1.06 [0.52-2.15]						
Vitamin B9	2		1	HR=1.04 [0.71-1.52]	2		1	HR=0.95 [0.79-1.15]				
Vitamin B12	1	OR=0.91 [0.49-1.68] OR=1.27 [0.59-2.73]			1	OR=0.91 [0.76-1.09] OR=0.85 [0.39-1.84]						
Vitamin C	4	OR=0.99 [0.48-2.04]	1	HR=0.92 [0.70-1.22]	4	OR=0.95 [0.69-1.31]	1	HR=0.93 [0.82-1.07]	1	OR=0.46 [0.16-1.36]	1	OR=0.76 [0.34-1.67]
Vitamin D	4	OR=0.58 [0.38-0.88]	1 1	OR=0.59 [0.27-1.29] HR=0.81 [0.45-1.47]	4	OR=0.99 [0.72-1.35]	1	HR=0.98 [0.69-1.37]				
Vitamin E	5	OR=0.54 [0.41-0.71]	1 1	OR=0.47 [0.24-0.92]	4	OR=0.87 [0.65-1.18]	1	HR=0.99 [0.83-1.18]	1	OR=0.57 [0.19-1.72]	1	OR=0.64 [0.29-1.41]

			1	RR=1.01 [0.91-1.12] HR=1.02 [0.70-1.47]								
Zinc	3	OR=0.57 [0.40-0.81]	1 1	OR=0.51 [0.27-0.97] HR=1.15 [0.77-1.72]	4	OR=1.00 [0.69-1.45]	1	HR=1.08 [0.86-1.35]	1	OR=0.52 [0.16-1.73]	1	OR=0.87 [0.39-1.96]
Copper	2	OR=0.55 [0.19-1.58]	1	HR=1.00 [0.71-1.41]	2	OR=0.91 [0.76-1.10]	1	HR=1.08 [0.89-1.31]	1	OR=1.08 [0.89-1.31]	1	OR=0.64 [0.30-1.37]
Calcium	2	OR=0.90 [0.39-2.08]	1 1	OR=0.72 [0.31-1.67] HR=1.14 [0.89-1.48]	3	OR=0.94 [0.82-1.08]	1	OR=0.97 [0.83-1.14]				
Magnesium			1	HR=0.91 [0.59-1.41]	1	OR=0.78 [0.62-0.97]	1	HR=0.95 [0.76-1.19]				
Manganese			1	HR=0.94 [0.73-1.20]	1	OR=0.87 [0.76-1.00]	1	OR=1.00 [0.87-1.15]				
Selenium			1	HR=1.37 [0.89-2.11]	1	OR=0.99 [0.81-1.21]	1	HR=0.99 [0.80-1.22]				
Vegetables	4 1	OR=0.90 [0.69-1.18] RR=0.94 [0.79-1.12]	2 1	OR=1.02 [0.58-1.81] HR=0.75 [0.46-1.22]	2 1	OR=0.71 [0.53-0.96] RR=1.21 [0.97-1.51]	1	HR=1.07 [0.78-1.47]			1	OR=1.09 [0.76-1.57]
Fruits	3 1	OR=1.03 [0.81-1.31] RR=0.94 [0.81-1.09]	1	HR=0.97 [0.60-1.58]	1 1	OR=0.78 [0.45-1.35] RR=0.82 [0.64-1.05]	1	HR=0.95 [0.68-1.34]				

Fish	6	OR=1.00 [0.89-1.13]	2 1	OR=0.99 [0.72-1.37] HR=1.09 [0.66-1.78]	6 1	OR=0.84 [0.68-1.05] RR=1.07 [0.86-1.33]	1 1	OR=0.28 [0.06-1.19] HR=0.92 [0.67-1.28]			2	OR=0.99 [0.75-1.30]
Meat	3 1	OR=1.16 [0.97-1.40] RR=1.22 [1.00-1.49]	1	HR=0.79 [0.48-1.32]	2 1	OR=1.28 [0.92-1.78] RR=0.96 [0.76-1.20]	1	HR=0.95 [0.67-1.33]				
Dairy	4 1	OR=0.88 [0.69-1.13] RR=0.83 [0.72-0.96]	1 1	OR=0.82 [0.35-1.88] HR=1.00 [0.59-1.69]	4 1	OR=0.95 [0.81-1.11] RR=0.95 [0.76-1.18]	1	HR=0.85 [0.61-1.19]			1	OR=0.93 [0.67-1.28]
Fats												
Total fats	3	OR=0.87 [0.71-1.06]	1	HR=0.92 [0.54-1.56]	3	OR=0.97 [0.80-1.17]	2	HR=0.98 [0.79-1.22]				
SFA	3 1	OR=0.80 [0.65-0.97] RR=1.08 [0.93-1.26]			3 1	OR=1.03 [0.86-1.24] RR=1.13 [0.88-1.45]	1	HR=1.06 [0.82-1.37]				
MUFA	3 1	OR=0.93 [0.77-1.14]			3 1	OR=0.99 [0.82-1.19]	1	HR=0.96 [0.74-1.24]				
n-6 PUFA	3	RR=1.06 [0.79-1.40]			3	RR=1.25 [0.96-1.62]	1					
n-3 PUFA	3	OR=1.06 [0.87-1.30]			3	OR=1.24 [0.87-1.76]	1	HR=0.84 [0.64-1.11]				

		OR=0.86 [0.58-1.26]				OR=0.94 [0.78-1.14]		HR=0.82 [0.63-1.08]				
--	--	---------------------	--	--	--	---------------------	--	---------------------	--	--	--	--

HR: Hazard Ratio- MUFA: Monounsaturated Fatty Acids- OR: Odds Ratio- PUFA: Polyunsaturated Fatty Acids- RR: Risk Ratio- SFA: Saturated Fatty Acids

Table 3: Main results of the studies included in the meta-analysis of maternal nutrition during pregnancy and risk of asthma, wheeze and atopic diseases during childhood.



