

Accepted Manuscript

Epidemiologic Patterns of In-Hospital Anaphylaxis in Pediatric Surgical Patients

Sara A. Sun, BA, Xiaoyue Ma, BS, Guohua Li, MD, DrPH, Caleb Ing, MD, MS

PII: S0091-6749(17)32918-4

DOI: [10.1016/j.jaci.2017.11.030](https://doi.org/10.1016/j.jaci.2017.11.030)

Reference: YMAI 13178

To appear in: *Journal of Allergy and Clinical Immunology*


Received Date: 8 September 2017

Revised Date: 31 October 2017

Accepted Date: 24 November 2017

Please cite this article as: Sun SA, Ma X, Li G, Ing C, Epidemiologic Patterns of In-Hospital Anaphylaxis in Pediatric Surgical Patients, *Journal of Allergy and Clinical Immunology* (2018), doi: 10.1016/j.jaci.2017.11.030.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Epidemiologic Patterns of In-Hospital Anaphylaxis in Pediatric Surgical Patients

Authors: Sara A. Sun, BA¹; Xiaoyue Ma, BS¹; Guohua Li, MD, DrPH^{1,2}; Caleb Ing, MD, MS^{1,2}

Affiliations: ¹Department of Anesthesiology, Columbia University College of Physicians and Surgeons, New York, NY, ²Department of Epidemiology, Mailman School of Public Health, New York, NY.

Capsule Summary

The incidence of in-hospital anaphylaxis in pediatric surgical patients has more than doubled between 2003 and 2012. Patients with hematological and myeloproliferative disorders, particularly when admitted for bone marrow transplant, are at especially heightened risk.

Key Words

Anaphylaxis; children; pediatric; surgical; inpatients; United States; bone marrow transplant

Abbreviations

KID: Kids' Inpatient Database; ICD-9: International Classification of Diseases, Ninth Revision;
DRG: diagnosis-related group; MDC: major diagnostic category; PPV: positive predictive value;
OR: odds ratio; CI: confidence interval

Sources of Funding

Foundation for Anesthesia Education & Research
Agency for Healthcare Research and Quality: K08HS022941
Centers for Disease Control and Prevention: R49 CE002096
Herbert Irving Scholar Award

Corresponding Author:

Caleb Ing, MD, MS
Email: ci2119@cumc.columbia.edu
Affiliation: Departments of Anesthesiology and Epidemiology, Columbia University College of
Physicians and Surgeons and Mailman School of Public Health, New York, NY
Address: 622 W. 168th St. BHN 4-440
New York, NY 10032, USA
Phone: 212-305-2413
Fax: 212-305-2395

To the Editor:

While several population-based epidemiological studies have evaluated the incidence of anaphylaxis in children and its change over time, few have focused specifically on anaphylaxis occurring during hospitalization. The purpose of this study is to assess the incidence and risk factors of in-hospital anaphylaxis in children admitted for surgery.

Data for this study came from the Kids' Inpatient Database (KID), which accounts for over 95% of all pediatric inpatient hospital admissions in the United States.^{1,2} The KID is released every three years. This study was based on the 2003, 2006, 2009, and 2012 KID datasets. In-hospital birth admissions and patients aged 18 years or older at admission were excluded from analysis. Children were identified as having had anaphylaxis if they received a primary or secondary diagnosis with International Classification of Diseases, Ninth Revision (ICD-9) codes for anaphylaxis or anaphylactic shock (995.0) or for an anaphylactic reaction to serum (999.4). These two codes were chosen because they have shown a high positive predictive value (PPV) for anaphylaxis when evaluated in inpatient samples (77.4%).³ Medical admissions were identified by a medical diagnosis-related group (DRG) code and surgical admissions by a surgical DRG.⁴ Since children who present to the hospital with anaphylaxis are unlikely to be admitted for a surgical procedure, diagnoses of anaphylaxis during surgical admissions were interpreted as having developed during hospitalization, and likely iatrogenic in nature.

Descriptive statistics and regression analyses in this study were calculated using weighted data.⁵ Using all available years of data, a total of 315 surgical admissions were found to have a diagnosis of anaphylaxis, while 1,750,151 did not. A total of 3,320 medical admissions had a diagnosis of anaphylaxis, while 7,817,791 did not. Of anaphylaxis diagnoses, the ICD-9 code

995.0 was used in 96.4% of surgical admissions and 97.0% of medical admissions, while the code 999.4 was used in the remaining admissions. The overall incidence of anaphylaxis in medical and surgical admissions more than doubled between 2003 and 2012 (Figure 1). Of anaphylactic episodes that occurred during medical admissions, depending on the year, 64-68% were coded as the primary diagnosis, whereas 97-100% of anaphylaxis diagnoses in surgical admissions were coded as a secondary diagnosis.

The overall in-hospital mortality ratio for children was 0.38% (0.75% for surgical admissions and 0.43% for medical admissions). The in-hospital mortality ratio for children who developed anaphylaxis during surgical admission was 2.47%, significantly higher than the in-hospital mortality ratio for pediatric surgical patients who did not develop anaphylaxis (0.75%, $p = 0.0004$). Yet, the in-hospital mortality ratio for children with any diagnosis of anaphylaxis during medical admission, 0.29%, was not significantly different from the in-hospital mortality ratio for pediatric medical patients without a diagnosis of anaphylaxis (0.43%, $p = 0.21$).

The characteristics of children with and without anaphylaxis during a surgical admission were evaluated. Of the 315 pediatric surgical patients with anaphylaxis, 15.6% were admitted for hematological and myeloproliferative disorders, compared to 1.5% of all pediatric surgical patients (see ETable I in Online Repository). All characteristics were included as covariates in a multivariable logistic regression model and the independent association between each covariate and the diagnosis of anaphylaxis was evaluated. Records with missing covariate data were excluded from analysis. To accommodate small cell sizes, some major diagnostic categories (MDCs) were merged together for regression analysis (see ETable II in Online Repository). Factors associated with significantly increased odds ratios of anaphylaxis included older age, larger hospital size, teaching hospital status, Western region of the United States, and admissions

for respiratory and cardiovascular diseases (Table I). The strongest association, however, was with hematological and myeloproliferative disorders, with an estimated odds ratio of 13.47 (95% confidence interval, 8.99-20.19) compared to digestive, hepatobiliary, and endocrine disorders, the most common reason for pediatric surgical admissions. Of the children admitted for hematological and myeloproliferative disorders, bone marrow transplant admissions accounted for 68.2% of admissions with an anaphylactic event, but only 21.4% of admissions without anaphylaxis. The incidence of anaphylaxis in children undergoing bone marrow transplant was 485.9 per 100,000 admissions, 27 times the incidence in the general pediatric surgical population (18.1 events per 100,000 surgical admissions). Overall, children who underwent bone marrow transplant accounted for 13.3% of the anaphylaxis incidence in surgical admissions.

In this study, we found anaphylaxis during pediatric surgical admissions to be associated with several patient and hospital variables, as well as an increased risk for mortality not seen in medical admissions. A particularly strong association with anaphylaxis was found with admissions for hematological and myeloproliferative disorders, especially for bone marrow transplants. While it is not possible to determine the triggering agent for anaphylaxis using this dataset, chemotherapeutic agents and more recent monoclonal antibody treatments have been recognized causes of immediate hypersensitivity and anaphylaxis in bone marrow transplant patients, but these events have typically only been documented in case reports or as adverse events in clinical trials.⁶⁻⁸ The exact mechanism for hypersensitivity to these agents is unclear, and while mild to moderate reactions are relatively common, severe hypersensitivity reactions are thought to be rare.⁹ Since severe hypersensitivity after treatment with chemotherapy has not been well described in a population-based study, its incidence may be underestimated, resulting in inadequate preparation for responding to these events.

Despite its relatively rare occurrence, in-hospital anaphylaxis is a serious event that can lead to significant morbidity and mortality. It is worth noting that while the PPV for anaphylaxis diagnostic codes 995.0 and 999.4 in inpatient records is relatively high, the sensitivity of these codes is unclear. Thus, it is likely that these codes still underestimate the true incidence of anaphylaxis in the pediatric population. Given the potentially life-threatening nature of an anaphylactic episode, our findings of a strong association between bone marrow transplant and anaphylaxis may direct future studies to evaluate reasons for this association and methods to reduce its occurrence.

Sara A. Sun, BA¹

Xiaoyue Ma, MS¹

Guohua Li, MD, DrPH^{1,2}

Caleb Ing, MD, MS^{1,2}

From the ¹Department of Anesthesiology, Columbia University College of Physicians and Surgeons, New York, NY, ²Department of Epidemiology, Mailman School of Public Health, New York, NY. E-mail: ci2119@cumc.columbia.edu

Sara Sun was supported by the Foundation for Anesthesia Education and Research (FAER). Dr. Caleb Ing is supported by a Herbert Irving Scholars Award as well as the Agency for Healthcare Research and Quality (AHRQ) under award number K08HS022941. Dr. Guohua Li is supported by the Centers for Disease Control and Prevention (CDC) under award number R49 CE002096.

The content of this publication is solely the responsibility of the authors and does not necessarily represent the official views of the AHRQ or CDC.

References

1. HCUP. HCUP Databases: HCUP-US KID Overview. 2017; <https://www.hcup-us.ahrq.gov/kidoverview.jsp>.
2. HCUP. *Introduction to the HCUP Kids' Inpatient Database (KID)*. U.S. Agency for Healthcare Research and Quality;2012.
3. Walsh KE, Cutrona SL, Foy S, Baker MA, Forrow S, Shoaibi A, et al. Validation of anaphylaxis in the Food and Drug Administration's Mini-Sentinel. *Pharmacoepidemiol Drug Saf*. 2013;22(11):1205-1213.
4. HCUP. DRG Formats Program. *HCUP Tools: Labels and Formats* 2017; https://www.hcup-us.ahrq.gov/db/tools/DRG_Formats.TXT.
5. Chu B, Houchens R, Elixhauser A, Ross D. *Using the Kids' Inpatient Database (KID) to Estimate Trends: HCUP Methods Series Report #2007-02*. U.S. Agency for Healthcare Research and Quality;2007.
6. Neugut AI, Ghatak AT, Miller RL. Anaphylaxis in the United States: an investigation into its epidemiology. *Arch Intern Med*. 2001;161(1):15-21.
7. Wickner PG, Hong D. Immediate Drug Hypersensitivity. *Curr Allergy Asthma Rep*. 2016;16(7):49.
8. Woods D, Winchester K, Towerman A, Gettinger K, Carey C, Timmermann K, et al. From the Children's Oncology Group: Evidence-Based Recommendations for PEG-Asparaginase Nurse Monitoring, Hypersensitivity Reaction Management, and Patient/Family Education. *J Pediatr Oncol Nurs*. 2017:1043454217713455.

- 147 9. Lenz HJ. Management and preparedness for infusion and hypersensitivity reactions. *Oncologist*.
148 2007;12(5):601-609.

149


ACCEPTED MANUSCRIPT

150 **Figure 1.** Incidence of Anaphylaxis as a Primary or Secondary Diagnosis in Surgical and
151 Medical Admissions.

Table I. Multivariable Logistic Regression including Risk Factors for Anaphylaxis in Surgical Admissions.

Risk Factors	OR Estimate (95% CI)
Age	1.05 (1.03-1.07)
Sex	
Male	(Reference)
Female	1.07 (0.85-1.35)
Admission type	
Elective	(Reference)
Emergent	0.83 (0.65-1.05)
Hospital bed size	
Large	(Reference)
Medium	0.66 (0.50-0.87)
Small	0.63 (0.43-0.93)
Hospital location	
Urban	(Reference)
Rural	0.79 (0.36-1.71)
Hospital teaching status	
Non-teaching	(Reference)
Teaching	2.86 (1.87-4.37)
Hospital region	
South	(Reference)
Northeast	0.95 (0.66-1.36)
Midwest	1.34 (0.97-1.86)
West	1.72 (1.28-2.31)
Major diagnostic category	
Digestive, hepatobiliary, endocrine disorders	(Reference)
Nervous system disorders	1.20 (0.73-1.98)
Eye, ear, nose, throat disorders	0.68 (0.37-1.26)
Respiratory disorders	3.00 (1.78-5.07)
Cardiovascular disorders	2.43 (1.55-3.80)
Musculoskeletal and skin disorders	1.18 (0.81-1.72)
Hematological and myeloproliferative disorders	13.47 (8.99-20.19)
Other disorders	1.17 (0.78-1.76)

OR = odds ratio, CI = confidence interval, (Reference) = reference value for each variable



ETable I. Demographic and Hospital Characteristics of Pediatric Surgical Admissions.

	Surgical Admissions	
	Reference (n = 1,750,151)	Anaphylaxis (n = 315)
Age, median (IQR)	10 (2 – 15)	11 (5 – 15)
Sex, n (%)		
Male	929,726 (53.1)	165 (52.3)
Female	795,820 (45.5)	151 (47.7)
Missing	24,606 (1.4)	0 (0.0)
Admission type, n (%)		
Elective	694,649 (39.7)	154 (48.8)
Emergent	1,047,051 (59.8)	160 (50.8)
Missing	8,451 (0.5)	*
Hospital bed size, n (%)		
Large	996,221 (56.9)	198 (62.9)
Medium	456,532 (26.1)	68 (21.4)
Small	228,134 (13.0)	31 (9.8)
Missing	69,264 (4.0)	19 (5.9)
Hospital location, n (%)		
Urban	1,574,911 (90.0)	290 (91.8)
Rural	105,976 (6.1)	*
Missing	69,264 (4.0)	19 (5.9)
Hospital teaching status, n (%)		
Non-teaching	418,680 (23.9)	28 (8.9)
Teaching	1,262,208 (72.1)	269 (85.2)
Missing	69,264 (4.0)	19 (5.9)
Hospital region, n (%)		
South	609,057 (34.8)	93 (29.4)
Northeast	300,090 (17.2)	45 (15.4)
Midwest	380,653 (21.8)	76 (24.2)
West	460,351 (26.3)	101 (32.1)
Missing	0 (0.0)	0 (0.0)
Major diagnostic category, n (%)		
Digestive, hepatobiliary, endocrine disorders	535,723 (30.6)	61 (19.5)
Nervous system disorders	144,133 (8.2)	26 (8.2)
Eye, ear, nose, throat disorders	163,382 (9.3)	15 (4.8)
Respiratory disorders	53,088 (3.0)	20 (6.2)
Cardiovascular disorders	125,033 (7.1)	38 (11.9)
Musculoskeletal and skin disorders	396,848 (22.7)	62 (19.5)
Hematological and myeloproliferative disorders	26,541 (1.5)	49 (15.6)
Kidney and reproductive disorders	110,283 (6.3)	12 (3.7)
Other disorders	195,118 (11.2)	33 (10.6)

IQR = interquartile range

*These data are not presented as the report of data for any subgroup with 10 or fewer subjects is prohibited by the Healthcare Cost and Utilization Project Data Use Agreement.

Note: Absolute numbers presented are based on weighted data. Total percentages for some variables may not sum exactly to 100% due to rounding in the presented values.

ETable II. Merged Major Diagnostic Categories (MDCs) used in Demographic and Regression Analyses.

Merged Disease Categories Used in Analyses	MDC Codes	Associated Major Diagnostic Category
Nervous system disorders	1	Diseases and disorders of the nervous system
Eye, ear, nose, throat disorders	2	Diseases and disorders of the eye
	3	Diseases and disorders of the ear, nose, mouth, and throat
Respiratory disorders	4	Diseases and disorders of the respiratory system
Cardiovascular disorders	5	Diseases and disorders of the circulatory system
Digestive, hepatobiliary, endocrine disorders	6	Diseases and disorders of the digestive system
	7	Diseases and disorders of the hepatobiliary system and pancreas
	10	Diseases and disorders of the endocrine, nutritional, and metabolic system
Musculoskeletal and skin disorders	8	Diseases and disorders of the musculoskeletal system and connective tissue
	9	Diseases and disorders of the skin, subcutaneous tissue, and breast
Kidney and reproductive disorders	11	Diseases and disorders of the kidney and urinary tract
	12	Diseases and disorders of the male reproductive system
	13	Diseases and disorders of the female reproductive system
Hematological and myeloproliferative disorders	16	Diseases and disorders of the blood and blood-forming organs and immunological disorders
	17	Myeloproliferative diseases and disorders (Poorly differentiated neoplasms)
Other disorders	14	Pregnancy, childbirth, and puerperium
	15	Newborn and other neonates (Perinatal period)
	18	Infectious and parasitic diseases and disorders (Systemic or unspecified sites)
	19	Mental diseases and disorders
	20	Alcohol/drug use or induced mental disorders
	21	Injuries, poison, and toxic effect of drugs

- 22 Burns
 - 23 Factors influencing health status and other
 - contacts with health services
 - 24 Multiple significant trauma
 - 25 Human Immunodeficiency Virus (HIV)
 - infection
-