

Flow cytometry *with vision*

Introducing FlowSight

- > Capable
- > Intuitive
- > Affordable



amnis[®]
www.amnis.com



Basophils, IgE, and Autoantibody-Mediated Kidney Disease

Xavier Bosch, Francisco Lozano, Ricard Cervera, Manuel Ramos-Casals and Booki Min

This information is current as of May 19, 2011

J Immunol 2011;186;6083-6090

doi:10.4049/jimmunol.1002648

<http://www.jimmunol.org/content/186/11/6083>

-
- | | |
|----------------------|---|
| References | This article cites 85 articles , 33 of which can be accessed free at:
http://www.jimmunol.org/content/186/11/6083.full.html#ref-list-1 |
| Subscriptions | Information about subscribing to <i>The Journal of Immunology</i> is online at
http://www.jimmunol.org/subscriptions |
| Permissions | Submit copyright permission requests at
http://www.aai.org/ji/copyright.html |
| Email Alerts | Receive free email-alerts when new articles cite this article. Sign up at
http://www.jimmunol.org/etoc/subscriptions.shtml/ |



Basophils, IgE, and Autoantibody-Mediated Kidney Disease

Xavier Bosch,* Francisco Lozano,[†] Ricard Cervera,[‡] Manuel Ramos-Casals,[‡] and Booki Min[§]

Basophils are of interest in immunology due to their ability to produce a Th2-signature cytokine, IL-4, following activation. A new understanding of the role of basophils in immunity shows novel functions at a cellular level through which basophils influence adaptive immunity. This review summarizes new advances in basophil biology and discusses new roles for basophils in human disease, especially in the mediation of the pathogenesis of lupus nephritis. Recently, basophils have been shown to contribute to self-reactive Ab production in systemic lupus erythematosus and may enhance pre-existing loss of B cell tolerance, suggesting that basophils, IL-4, and IgE mediate the pathogenesis of lupus nephritis by promoting the Th2 environment and activating autoreactive B cells. In addition to envisaging exciting therapeutic prospects, these novel findings open the way for the study of basophils in other autoimmune and renal diseases. *The Journal of Immunology*, 2011, 186: 6083–6090.

Basophils are bone marrow-derived basophilic granulocytes that contain large cytoplasmic granules and two- to three-lobed nuclei. They constitute <0.3% of the total nucleated cells in the bone marrow and are the least abundant granulocytes found in the circulation (1). Since the discovery of basophils >130 y ago, they have been implicated in many types of allergic inflammatory responses. Earlier studies using a guinea pig model of cutaneous hypersensitivity found that basophils reach tissue sites, where they are triggered by Ags to release mediators (2). Basophils also have been shown to confer resistance against ectoparasite infections (3). During *Amblyomma americanum* tick infection in calves, significant decreases in feeding are associated with cutaneous basophil infiltration and blood basophilia (4). Transfer of anti-basophil serum abolishes resistance to tick infection in guinea pigs (5). Nevertheless, a tool to identify or selectively

purify basophils has not been available until recently. Studies in the last 7 y have made significant progress in identifying the biologic functions of basophils, a list that continues to grow (see below for details). Surface phenotypes to analyze and purify basophils have been characterized extensively: FcεRI⁺ CD49b⁺ CD203c⁺ Thy1⁺ c-Kit⁺ FcγR⁺ CD45^{intermediate} (6). Animal models to study basophil functions in vivo have been developed. Most importantly, genetically engineered basophil-deficient animals have been generated recently (7, 8). We now are beginning to understand novel functions of basophils at the cellular level through which they influence adaptive immunity. This review summarizes new advances in basophil biology and novel roles of basophils in human disease, particularly their mediation in the pathogenesis of lupus nephritis (LN).

IL-3 is a key factor for basophil development

IL-3 is involved in the differentiation of early precursors into fully mature basophils in the bone marrow and spleen (9, 10). Because basophil differentiation is controlled at a transcriptional level by differential expression of two key transcription factors, C/EBPα and GATA2 (11), IL-3 may regulate expression of these factors, influencing fate decision during early development. IL-3 deficiency results in significant defects in the generation of basophils in the course of Th2 immunity (12, 13). Nonetheless, the basal production of basophils in naive IL-3-deficient mice is similar to that in wild-type mice (12). Therefore, the role of IL-3 in basophil development during steady-state conditions and immune responses is likely to differ. Basophil differentiation in steady-state conditions could be IL-3-independent; however, IL-3 becomes the major factor enhancing basophil generation in the bone marrow under Th2-associated inflammatory conditions (13).

Factors involved in basophil activation

The list of factors that positively and negatively regulate basophil activation continues to grow (Fig. 1). The primary action of IL-3 on mature basophils is to enhance IL-4 pro-

*Department of Internal Medicine, Hospital Clinic, Institut d'Investigacions Biomèdiques August Pi i Sunyer, University of Barcelona, 08036 Barcelona, Spain; [†]Department of Immunology, Hospital Clinic, Institut d'Investigacions Biomèdiques August Pi i Sunyer, University of Barcelona, 08036 Barcelona, Spain; [‡]Department of Autoimmune Diseases, Hospital Clinic, Institut d'Investigacions Biomèdiques August Pi i Sunyer, University of Barcelona, 08036 Barcelona, Spain; and [§]Department of Immunology, Lerner Research Institute, Cleveland Clinic Foundation, Cleveland, OH 44195

Received for publication November 10, 2010. Accepted for publication March 9, 2011.

This work was supported by National Institutes of Health Grant AI080908 (to B.M.).

Address correspondence and reprint requests to Dr. Xavier Bosch, Department of Internal Medicine, Hospital Clinic, Institut d'Investigacions Biomèdiques August Pi i

Sunyer, University of Barcelona, 08036 Barcelona, Spain. E-mail address: xavbosch@clinic.ub.es

Abbreviations used in this article: ACPA, anti-citrullinated protein Ab; ANA, anti-nuclear Ab; CIC, circulating immune complex; DC, dendritic cell; DPGN, diffuse proliferative glomerulonephritis; IRF2, IFN regulatory factor 2; LN, lupus nephritis; MGN, membranous glomerulonephritis; MHC II, MHC class II; RA, rheumatoid arthritis; SLE, systemic lupus erythematosus.

Copyright © 2011 by The American Association of Immunologists, Inc. 0022-1767/11/\$16.00




FIGURE 1. Potential functions of basophils stimulated via multiple pathways are illustrated. LT, leukotriene; PAMP, pathogen-associated molecular pattern; TSLP, thymic stromal lymphopoietin; IPSE.

duction (14–16). IL-3 also primes basophils to augment IL-4 production in response to other stimuli (15). Ig receptor (FcR) cross-linking induces both IL-4 and IL-13 production and histamine release in basophils (17); preincubation of basophils with IL-3 or the addition of IL-3 during stimulation significantly enhances cytokine production (15, 18). Both IL-18 and IL-33, members of the IL-1 family of cytokines, stimulate IL-4 and IL-13 production independently of FcR cross-linking (18–20). Notably, basophil survival significantly increases when stimulated with IL-18 or IL-33 by a PI3K/Akt-dependent pathway (21). IL-3 enhances basophil survival in vitro, although no in vivo effect on survival was observed (13). IL-25, a member of the IL-17 family of cytokines, plays a crucial role in developing Th2 immune responses (22). Basophils from patients with seasonal allergic rhinitis express high levels of IL-17RB, a receptor for IL-25 (23). In fact, IL-25 stimulation inhibits basophil apoptosis and enhances IgE-mediated degranulation, indicating the possible involvement of IL-25 in basophil activation and Th2 immunity (23).

Basophils express receptors involved in innate immunity, including TLR2 and TLR4 (24–26). Stimulation of basophils with LPS increases cytokine production through FcR-dependent or independent pathways (26). Ags associated with parasites are known to have basophil-activating properties. IPSE/ α -1, a glycoprotein from *Schistosoma* egg Ags, induces IL-4 production in basophils (27). Protease Ags, such as papain, and house dust mite Ags also activate basophil cytokine production (28, 29), although the underlying mechanism is not clear. Ags from pathogens also modulate basophil functions. *Helicobacter pylori*-derived cecropin-like peptide induces basophil chemotaxis (30). HIV glycoprotein gp120 induces IL-4 and IL-13 release from human basophils through interaction with the V_H3 region of IgE (31). Complement C5a induces strong IgE-independent production of cytokines and mediators from basophils (32).

Cytoplasmic signaling molecules and transcription factors that downregulate basophil activation have been reported. IFN regulatory factor 2 (IRF2) inhibits basophil expansion. As a result, mice deficient in IRF2 show marked basophilia and

spontaneous Th2 differentiation (33). Lyn, a Src kinase family member, dampens basophil expression of GATA3 (34). Like IRF2^{-/-} mice, *lyn*^{-/-} mice present basophilia and constitutive Th2 effector CD4 T cells (34). SHIP, a negative regulator of the PI3K pathway, plays a key role in suppressing IL-4 production from basophils (35). Notably, Th2-prone immune responses found in these mice require both basophils and IL-4, further highlighting the importance of IL-4-dependent immune modulation by basophils. The surface Ags that negatively regulate basophil activation remain to be determined.

Basophil functions in vivo

Th1 CD4 T cells mainly produce IFN- γ , protecting hosts from intracellular pathogens, and factors inducing Th1 differentiation have been identified, including IL-12, IFN- γ , and transcription factor T-bet (36). Th2 CD4 T cells produce IL-4, IL-5, and IL-13, protecting hosts from extracellular pathogens. IL-4 and transcription factor GATA3 are key Th2-inducing factors. Unlike Th1 immunity, in which activated Ag-presenting dendritic cells (DCs) provide IL-12, which then program activated naive CD4 T cells to Th1 lineage cells, the source of initial IL-4 necessary to initiate Th2 differentiation remains elusive. Studies using cytokine-reporter mice (4get or G4 mice) identified basophils as potential candidates for the role of Th2 inducers (37, 38). In vitro, the stimulation of naive CD4 T cells in the presence of basophils promotes Th2 differentiation without exogenous addition of IL-4 (39). A key mechanism of basophils that supports Th2 differentiation operates via IL-4, because basophils deficient in IL-4 or IL-4 neutralization lack or abolish basophil-mediated Th2 differentiation (Fig. 1).

Unlike CD4 T cells, CD8 T cells stimulated in the presence of basophils produce high levels of IL-10, which is not observed when T cells are stimulated by other APCs (40). Because basophils express MHC class I molecules, they directly interact with and promote IL-10 production in CD8 T cells. IL-4 neutralization abolishes basophil-mediated IL-10 expression; however, rIL-4 alone does not induce IL-10 in CD8 T cells, suggesting an additional factor in this process. IL-6 has a synergistic effect on CD8 T cell IL-10 expression when cocultured with IL-4, although IL-6 alone had no effect (40).

Basophils also can enhance B cell IgE production in vitro through the production of Th2 cytokines and CD40L expression (41, 42). It was reported recently that basophils efficiently capture intact Ags and enhance B cell activation, particularly B cell proliferation and Ig production (43). Both IL-4 and IL-6 produced by basophils are key cytokines that optimize CD4 T cell-dependent B cell help (43). In vivo, basophil-mediated effects on B cells are critical to mount protective humoral immunity against bacterial infection (44). In humans, circulating IgD binds to basophils and induces B cell-stimulating factors, such as BAFF (also known as B lymphocyte stimulator), IL-1, and IL-4, together with antimicrobial and opsonizing factors (44). Whether the IgD-mediated basophil activation pathway also operates in mice remains to be determined.

Basophil function as APCs and the controversy

Studies show that basophils play critical roles as APCs during Th2-type responses (Fig. 1). Basophils uptake exogenous

protein Ags, process, and present them to naive CD4 T cells via surface MHC class II (MHC II) (45). Mice immunized with OVA protein plus the protease papain developed OVA-specific Th2-type immune responses. Because papain injection recruits circulating basophils into the draining lymph nodes where basophils are found within T cell areas, it is suggested that basophils provide key signals (MHC II–Ag complexes plus IL-4) through which Ag-specific Th2 differentiation is optimized (45). Papain-mediated Th2 immune responses were not altered by CD11c⁺ DC depletion. In fact, the transfer of Ag-pulsed basophils induces Th2 differentiation within CIITA-deficient mice, where resident APCs, including DCs, lack functional MHC II molecules (45). Basophils incubated with IL-3 plus Ag or IgE–Ag complexes induce the development of Ag-specific Th2 cells and Ag-specific B cell responses in naive mice, and the administration of IgE–Ag complexes induces Th2 differentiation only in the presence of endogenous basophils, suggesting that basophils could function as potent APCs by capturing IgE complexes (24). A *Trichuris muris* infection model also demonstrated that DC-mediated immune priming is insufficient for the development of parasite-specific Th2 immunity using MHC II–CD11c transgenic mice in which MHC II expression is restricted to CD11c⁺ DCs (46). *T. muris*-specific Th2 immunity was impaired substantially in these mice, although Th1 differentiation was intact. Basophil depletion significantly impaired Th2 cytokine responses and parasite expulsion (46).

However, the role of basophils as Th2-inducing APCs was challenged recently by three independent studies demonstrating that basophils are dispensable for Th2 induction (47–49). CD11c⁺ DC depletion dramatically reduced Th2 induction in mice challenged with *Schistosoma* egg Ags or infected with *Schistosoma mansoni*, but treatment with MAR-1 Ab had little or no effect on Th2 responses (47). Th2 cell differentiation was not impaired after papain/OVA immunization using basophil-deficient mice (49). In response to house dust mite allergens, basophils play no role in Th2 immunity, but FcεRI⁺CD11c⁺ DC subsets recruited into the draining lymph nodes were responsible for Th2 immunity (48).

The discrepancies in these studies require explanation. Experimental models may determine the contribution of basophils to subsequent T cell immunity. Basophil-independent (and DC-dependent) Th2 immunity could arise from more complicated immunogens derived from parasites or multiple Ags, including house dust mite allergens. In contrast, papain could act directly on basophils via cysteine protease activity, which then activates basophils, including Ag-presenting machineries. However, Th2 immunity raised from papain immunization requires DC–basophil cooperation (45). Alternatively, DC subsets expressing FcεRI may have been overlooked in studies using MAR-1 Ab. Thus, the defects observed following MAR-1 Ab could be due to DC depletion.

Basophil-deficient animals: new functions revealed

A transgenic mouse, Mcpt8Cre, was generated by expressing the Cre recombinase gene under the control of the *Mcpt8* gene. Basophil numbers were constitutively lower in this animal, while other leukocytes, such as peritoneal mast cells, remained intact. In vivo Th2 responses, including alum-induced allergic lung inflammation, alum-induced humoral immune responses,

and Ig-mediated passive systemic anaphylaxis, were induced normally. Furthermore, primary Th2 immune responses against infection with the intestinal nematode *Nippostrongylus brasiliensis* were not impaired without basophils (49), which is consistent with our recent report (50). In contrast, protective memory immune responses against secondary *N. brasiliensis* infection were impaired significantly (49). Interestingly, lung infiltration of Th2-type CD4 T cells and eosinophils following *N. brasiliensis* infection was not altered; instead, eosinophil recruitment into the draining mediastinal lymph node was diminished greatly without basophils, suggesting that they may contribute to eosinophil recruitment (51). Karasuyama and colleagues (7) independently generated Mcpt8^{DTR} mice in which basophils are depleted conditionally by transgenic expression of the human diphtheria toxin receptor under the control of the mast cell protease 8 (Mcpt8) promoter. A single injection of diphtheria toxin resulted in transient (yet highly efficient) depletion of basophils from the bone marrow, circulation, and spleen, while other cell types including mast cells were not affected. Using a tick infection model, they demonstrated that basophil depletion resulted in the loss of acquired tick resistance. Because Ig receptor expression on basophils, but not on mast cells, was necessary to confer protection, these results provide strong evidence supporting Ab-mediated acquired immunity by basophils against tick infection (Fig. 1).

Basophil mobilization: a key step for basophil function in vivo

In steady-state conditions, the primary locations of basophils are the bone marrow and the circulation, while basophils are found very rarely in secondary lymphoid tissues. A key step that allows basophils to contribute to Th2 immune responses is their transient mobilization into secondary lymphoid tissues. Basophils within draining lymph nodes are observed in many in vivo immune responses: immunization with protease Ags, such as papain (29) and house dust mite Ags (48), and infection with *N. brasiliensis* (50), *S. mansoni* (47), *T. muris* (46), and ticks (7). Recruitment is found only in draining lymphoid tissues, suggesting active immune responses. IL-3 plays a crucial role in this recruitment process (50). Because IL-3 is produced primarily by activated T cells (13), Ag-induced T cell activation within local lymphoid tissues seems to be the primary underlying factor. In addition, recruitment is transient, only found between days 3 and 4 post-Ag challenge (29, 50). It is unclear how IL-3–derived basophil recruitment is tightly controlled, although IL-3–responsive cells have been shown to be bone marrow-derived cells (50). Likewise, recruited basophils are in close contact with T cells within the T cell zone (29). Basophils may function as APCs (24, 45, 46) and/or produce IL-4, creating an optimal microenvironment favoring Th2 differentiation. Basophils express chemokine receptors, including CCR1, CCR2, CCR3, CXCR1, CXCR3, and CXCR4 (52). A microarray study of basophils isolated from the lungs of *N. brasiliensis*-infected mice showed that CCR2 is highly expressed on basophils (38). Whether IL-3–dependent basophil recruitment is chemokine-dependent and, if so, which chemokines are involved in basophil recruitment remains to be determined.

Basophils: from allergy to autoimmunity

Because basophils play an immunomodulatory role, provoking Th2 cell differentiation in vivo, it could be suggested that they

also might play a leading role in immune diseases other than allergies. In addition to the finding of peripheral basophilia and constitutive Th2 skewing of CD4 T cells in *lyn*^{-/-} mice, basophil involvement also is reported in autoimmune urticaria, where autoantibodies to the high-affinity receptor FcεRIα induce basophil activation (53, 54). A Th2 component is well-described in autoimmune urticaria, and basophils contribute to the development of autoimmune symptoms through their immunomodulatory and effector responses. Reports show that, in some murine models, mast cells contribute to autoimmune arthritis (55). In this and other autoimmune diseases, mast cell effector and immunomodulatory functions initiate inflammation, whereas the enhancement of inflammation may be due to Th1 and Th17 cell responses. These CD4 T cell subsets are thought to be the primary effector cells involved in chronic inflammation in experimental autoimmune encephalomyelitis, a model of human multiple sclerosis, experimental autoimmune uveitis, a model of human posterior uveitis, and collagen-induced arthritis, a murine model of human rheumatoid arthritis (RA) (56).

Although IgE is recognized as a human Th2 marker, some reports describe increased levels of IgE in the setting of Th1 and Th17 responses in patients with autoimmune diseases (57–59). Although the significance of Th1- and Th17-driven responses in the development of autoimmunity is evident, it seems improbable that Th2 responses do not contribute to the disease. Furthermore, allergy and autoimmunity share various features, such as manifestations associated with autoantibody production and circulating immune complex (CIC) formation (60). Studies report a common structure of some allergens and autoantigens (61) and suggest that numerous environmental allergens have common structural conformations with self-antigens, which could explain the development of autoimmune diseases, such as chronic urticaria. However, as discussed later, increased IgE levels are not associated with increased allergic disease in other autoimmune diseases, such as systemic lupus erythematosus (SLE) (57).

The recent finding that IgD-mediated basophil activation by a yet unidentified receptor is relevant to some chronic auto-inflammatory diseases, such as hyper IgD syndrome, TNFR-associated periodic syndrome, and Muckle-Wells syndrome (44), suggests that this activation could be involved in B cell homeostasis and the production of Abs.

SLE as a paradigm of autoimmune disease

SLE is characterized by functional changes in B and T cells, monocytes, plasmacytoid DCs, and granulocytes, autoantibody deposition at inflammation sites, and increased circulating cytokine and chemokine levels. In LN, immune complexes deposited in glomeruli include IgG, IgM, and IgA autoantibodies, which are directed against ubiquitous nuclear Ags, especially dsDNA, possibly leading to kidney failure and death (62). Mechanisms contributing to LN pathogenesis include those mediated by Th1 and Th17 cells (63).

Although humoral responses produced by self-reactive Abs act as effector mediators of LN and IgE autoantibodies have been observed in some cases, the manner of Th2 cell and B cell activation in LN pathogenesis remains unclear (62).

Role of T cells and cytokines in the pathogenesis of SLE and LN

In SLE, T cells provide “excessive help” to B cells and mount inflammatory responses while producing low levels of IL-2

(63). Defective IL-2 production in T cells could explain the known reduction in cytotoxic activity, defective regulatory T cell function, and decreased activation-induced cell death in SLE patients (64). Helper CD4 T cells are vital components of immune responses in SLE; after differentiation, CD4 T cell effector functions correlate with the profiles of the cytokines that they secrete (65).

Amplified Th1, Th2, Th17, and follicular helper T cell responses are related to autoantibody production in murine and human SLE, and their effector cytokines may mediate target-organ inflammation. The cytokine response might reflect the primary activation pathway, which could be governed by genetic factors, and/or a specific environmental trigger (65, 66).

As mentioned, studies show a reduction in the amount and/or function of regulatory T cells in SLE patients and lupus-prone mice (67). Regulatory T cells have a reduced capacity to suppress CD4 T cell proliferation in active lupus compared with those from inactive lupus or healthy controls (68).

In addition to Th1, Th17, and regulatory T cells, reports suggest a role for Th2 cells in SLE (68–70). The strong humoral response in SLE indicates that there may be a Th2 component, with reports showing increased IgE and the presence of self-reactive IgE in the serum from some SLE patients, with no increase in allergy (71, 72). However, it is unclear whether Th2 cytokines, such as IL-4, and IgE contribute to SLE and which cell type may be implicated.

The presentation of LN is phenotypically and histologically heterogeneous. Diffuse proliferative glomerulonephritis (DPGN) and membranous glomerulonephritis (MGN) are, morphologically, complete opposites (73). DPGN pathogenesis is associated with a predominance of Th1 cytokines, suggesting that it occurs in a Th1-dominant immune response. Renal tissue from DPGN patients shows increased Th1 cytokine levels, including IFN-γ, and the IFN-γ/IL-4 production ratio correlates with the nephritic histological activity index. In contrast, the ratio is reduced in PBLs of MGN patients, suggesting that Th2-dominant cytokine responses are associated with MGN pathogenesis (74).

IgE in SLE and LN

Studies show that SLE patients without allergy have elevated total IgE and anti-nuclear IgE autoantibody levels, which appear to be related to disease severity (57). Likewise, the presence of IgE in glomerular immune complex deposits in renal biopsies of SLE patients and a murine model of systemic autoimmune disease suggest that they play a role in LN (75). A study of anti-nuclear IgE Ab specificity and cytokine involvement suggests that IgE Abs against cell autoantigens implicated in protein expression, cell proliferation, and apoptosis are observed in SLE patients and that IL-10 may downregulate IgE autoimmune responses in SLE (75).

Lyn, basophils, and Th2

Lyn is one of various Src tyrosine kinases involved in B cell activation. Aged *lyn*^{-/-} mice develop an autoimmune disease resembling human SLE (76). *lyn*^{-/-} mice have circulating autoantibodies to dsDNA and anti-nuclear Abs (ANAs). Glomerular CIC deposition in these mice results in renal damage and death.

lyn^{-/-} mice develop early strong, constitutive Th2 skewing and exacerbated responses to Th2 challenges (77, 78). In addition, B cells from some SLE patients express reduced levels of Lyn kinase (79). Therefore, *lyn*^{-/-} mice are a likely model to study the influence of a Th2 environment on lupus-like nephritis.

The cell types causing Th2 cell differentiation are not clear. Identification of the cell types and molecules responsible for Th2 cell response dysregulation could help to control these responses (34). Although important, factors controlling basophil-mediated Th2 cell differentiation remain unknown. Lyn kinase dampens basophil expression of the transcription factor GATA3 and the initiation and extent of Th2 cell differentiation. Reportedly, *lyn*^{-/-} mice had marked basophilia, a constitutive Th2 cell skewing that was exacerbated by in vivo basophil challenge, produced Abs to normally inert Ags, and did not respond appropriately to a Th1 cell-inducing pathogen. Th2 cell skewing was dependent on basophils, IgE, and IL-4 but independent of mast cells, suggesting that basophil-expressed Lyn kinase exerts regulatory control on Th2 cell differentiation and function (34).

Basophils, IgE, Th2, and LN

Charles et al. (80) recently investigated whether Th2 skewing in *lyn*^{-/-} mice plays a role in the development of late-life lupus-like nephritis and whether similar characteristics exist in SLE patients. They found that the Th2 phenotype contributes to lupus-like nephritis in *lyn*^{-/-} mice and is associated with LN in human SLE. Basophils and self-reactive IgE were identified as key players in renal disease. In mice, basophil depletion or a lack of IL-4 or IgE led to reduced autoantibody production and preserved renal function. Therefore, without basophils, autoantibody levels might not be sufficient to cause renal disease.

Charles et al. (80) studied the role played by the Th2 environment in the development of an SLE-like phenotype using mice deficient in IgE and Lyn, IL-4 and Lyn, or mast cells and Lyn, respectively. As reported (77), IgE contributes to increased mast cell levels in *lyn*^{-/-} mice, suggesting that it plays a role in mast cell survival (81). However, neither IL-4 nor IgE was associated with basophilia in mice, and the lupus-like nephritis seen was, therefore, dependent on IgE and IL-4 but not on mast cells. Basophil depletion in *lyn*^{-/-} mice aged >32 wk substantially decreased ANA and reduced splenic plasma cells and the renal proinflammatory environment. Basophils supported splenic plasma cells and enhanced the production of autoantibodies in an IL-4- and IgE-dependent manner, resulting in kidney damage.

Charles et al. (80) also determined whether self-reactive IgE is observed in murine circulation, whether they activate FcεRI-bearing basophils, resulting in Th2 cytokine expression, and whether IgE or IgG immune complexes stimulate basophil IL-4 production. Although IgE immune complexes induced basophil IL-4 production, IgG immune complexes did not.

Circulating basophils had increased CD62L expression, which permits leukocyte homing to peripheral lymphoid tissues. A lack of IL-4 or IgE inhibited CD62L expression on circulating basophils. Large amounts of basophils were present in the spleen and lymph nodes, where the proportion of basophils was reduced markedly in IL-4- or IgE-deficient but

not in mast cell-deficient settings. Lymph node basophils expressed membrane-associated BAFF, showing a possible role for lymph node basophils in B cell survival and differentiation. In addition, basophils from mice lymph nodes and spleen had high expression of MHC II. Higher expression of MHC II and/or BAFF in the lymph nodes and spleen could permit communication with T and B cells.

SLE patients had high numbers of C1q-reactive CICs, which activate the classical complement pathway and were increased markedly in mild and active disease. SLE patients also showed high total IgE and self-reactive dsDNA-specific IgE levels, which were associated with active LN. IgGs directed against IgE also were found in sera in SLE patients, with significantly increased levels correlating with active disease.

Basophils from all of the SLE patients had high CD203c expression compared with that of healthy controls, indicating basophil activation. CD62L and HLA-DR expression also was elevated on SLE basophils and correlated with greater disease activity. The number of total circulating basophils was reduced in SLE patients and was associated with immunosuppressive therapy but not with basophil activation.

Interpretation and caveats

Charles et al. (80) showed that basophils contribute to self-reactive Ab production in SLE and could enhance pre-existing loss of B cell tolerance. The development of lupus-like disease in *lyn*^{-/-} mice depended on the presence of IL-4 and IgE, whose absence reversed the Th2 skewing seen in these mice. IgG anti-dsDNA and ANA titers and CIC levels also were reduced in mice deficient for IgE and Lyn, and IL-4 and Lyn, respectively, demonstrating that autoantibody production in *lyn*^{-/-} mice was, at least partly, reliant on the Th2 environment. Therefore, these results demonstrate the participation of basophils, IgE, and IL-4 in the maintenance of autoantibody production in *lyn*^{-/-} mice and indicate that the Th2 environment contributes to the autoimmunity. They also show that, in some settings, basophils are strong inducers of Th2 cell differentiation, thus connecting the "allergic" environment to the development of autoimmune diseases, such as SLE, although the suggestion that basophils are necessary for Th2 cell differentiation must be inferred with caution. Research is limited on this topic, and the crucial test of challenging mice with a Th2 Ag in the complete absence of basophils remains to be done.

Although this study by Charles et al. (80) increases the understanding of LN pathogenesis, it has some limitations. IgE autoantibodies are found only in one third of SLE patients (75). Therefore, it is unlikely that the *lyn*^{-/-} mouse models SLE in most patients. However, it is known that IFN-γ, the primary Th1 cytokine, can negatively regulate both basophils and IL-4 production; therefore, it remains to be determined whether the 30% of SLE patients with high IgE ANA levels would respond to therapeutic basophil modulation, as suggested by this study in *lyn*^{-/-} mice (65).

It is also unclear whether the reported upregulation of BAFF on basophils depends on FcεRI engagement or needs concomitant TLR9 engagement or, alternatively, BAFF produces basophil-mediated class switching and Ig production (44). Activated basophils migrate to peripheral lymphoid organs, but the manner and site of interaction with T or B cells remain to be elucidated.

Finally, Charles et al. (80) observed a significant decrease in the number of circulating basophils in SLE patients, which was associated with increased disease activity, findings consistent with basophil homing to the secondary lymphoid tissues. A reduced number of circulating basophils was not observed in *lyn*^{-/-} mice, although the mice had peripheral basophilia, which was not observed in patients. In *lyn*^{-/-} mice, dysregulation of basophil proliferation probably covers circulating basophil depletion after basophil homing to the secondary lymphoid tissues.

Therapeutic implications

This study suggests that reducing circulating levels of self-reactive IgE or dampening basophil activity could have therapeutic benefits in LN. Therefore, determining whether basophil reduction or inactivation early in the disease course could defer or rescue the early development of LN is of substantial interest.

A recent phase 3 multicenter, placebo-controlled trial in patients with active SLE found that belimumab, a mAb against BAFF, is safe and efficacious (62, 82). Belimumab, besides affecting B cells, might deplete basophils (which contain membrane-bound BAFF), suggesting that it could be of help in some LN patients, perhaps combined with treatments suppressing IgE production.

The study showed that IgE immune complexes activate basophils and that removing self-reactive IgEs forming functional CICs prevents renal disease. As circulating IgE levels are decreased, omalizumab, a mAb binding to human IgE and decreasing FcεRI expression on basophils (83), may provide therapeutic benefits in SLE patients with increased levels of IgE. This approach could be tested easily, because omalizumab is used currently to treat asthma and allergic rhinitis and seems to be well tolerated (80, 81).

Future directions

The variety of basophil immunoregulatory activities explains why their pharmacological modulation by specific inhibitors or regulators is actively pursued, especially in allergic diseases. This approach may be used fairly soon in other pathologic conditions, including autoimmunity, as illustrated by the recent finding of basophil activation by anti-citrullinated protein Abs (ACPAs) in the serum of RA patients (84). Schuerwegh et al. (84) showed that IgE-bearing basophils from ACPA-positive RA patients are activated directly on exposure to citrullinated Ags, probably reflecting the presence of FcεRI-bound IgE-ACPAs on basophil surfaces. Identification of this activation pathway mediated by IgE-ACPAs in basophils could lead to innovative therapeutic schemes in RA.

The role of basophils also should be tested in immune-mediated renal disorders other than LN. A memory-type immune response prevails in transplantation and various autoimmune diseases with the constant presence of alloantigens/autoantigens. Together with specific Abs, these Ags may activate basophils, which may enhance ongoing humoral immune responses and further boost Ig production (85).

Although studies suggest that basophils may play a role in transplant rejection, studies using consistent methods of detecting basophils and determining their activation in some renal disorders are required. Basophils should be studied in minimal-change disease and focal segmental glomerulo-

sclerosis, where T cells and allergy/atopy are believed to play a major role, and Th2 skewed cytokine profiles have been recognized (85). Basophils, as inducers of Th2 immune responses and an important source of IL-4, could initiate disease and play a role in sustaining Th2-immune responses. Basophils also should be assessed in diseases driven by Ig directed against defined kidney structures (e.g., anti-glomerular basement membrane disease) or deposited in the glomeruli (e.g., membranous nephropathy) or driven by systemic vasculitis (e.g., anti-neutrophil cytoplasmic Ab-associated vasculitis), in which basophils may contribute to the generation and maintenance of high levels of specific Ig and, eventually, to disease progression.

Conclusions

Although our knowledge of the immunomodulatory and effector functions of basophils remains imperfect, it can be stated that, in addition to their role in allergy and host defense against parasites, basophils can promote Th2 cell differentiation and amplify the humoral immune response. Their role in autoimmune diseases is not clear, but the studies reported here provide a persuasive argument for a role in the amplification of LN rather than disease initiation. Essential requirements are autoreactive IgE-dependent basophil activation, IL-4 production by basophils, and promotion of Th2 cell differentiation. This does not negate the proven roles of Th1, Th17, and the loss of regulatory T cell function in the development of autoimmunity but rather suggests that the robust humoral constituent of autoimmune diseases, such as SLE, requires basophils to reach the autoantibody production levels that produce renal disease. These novel findings open the way for the study of basophils in other autoimmune and renal diseases. Future studies also may show whether basophil-mediated immune activation is linked to other SLE-prone mouse models in addition to *lyn*^{-/-} mice.

Disclosures

The authors have no financial conflicts of interest.

References

1. Falcone, F. H., D. Zillikens, and B. F. Gibbs. 2006. The 21st century renaissance of the basophil? Current insights into its role in allergic responses and innate immunity. *Exp. Dermatol.* 15: 855–864.
2. Askenase, P. W. 1979. Mechanisms of hypersensitivity: cellular interactions. Basophil arrival and function in tissue hypersensitivity reactions. *J. Allergy Clin. Immunol.* 64: 79–89.
3. Brown, S. J., F. M. Graziano, and P. W. Askenase. 1982. Immune serum transfer of cutaneous basophil-associated resistance to ticks: mediation by 7SIgG1 antibodies. *J. Immunol.* 129: 2407–2412.
4. Brown, S. J., R. W. Barker, and P. W. Askenase. 1984. Bovine resistance to *Amblyomma americanum* ticks: an acquired immune response characterized by cutaneous basophil infiltrates. *Vet. Parasitol.* 16: 147–165.
5. Brown, S. J., S. J. Galli, G. J. Gleich, and P. W. Askenase. 1982. Ablation of immunity to *Amblyomma americanum* by anti-basophil serum: cooperation between basophils and eosinophils in expression of immunity to ectoparasites (ticks) in guinea pigs. *J. Immunol.* 129: 790–796.
6. Min, B. 2008. Basophils: what they 'can do' versus what they 'actually do'. *Nat. Immunol.* 9: 1333–1339.
7. Wada, T., K. Ishiwata, H. Koseki, T. Ishikura, T. Ugajin, N. Ohnuma, K. Obata, R. Ishikawa, S. Yoshikawa, K. Mukai, et al. 2010. Selective ablation of basophils in mice reveals their nonredundant role in acquired immunity against ticks. *J. Clin. Invest.* 120: 2867–2875.
8. Ohnmacht, C., C. Schwartz, M. Panzer, I. Schiedewitz, R. Naumann, and D. Voehringer. 2010. Basophils orchestrate chronic allergic dermatitis and protective immunity against helminths. *Immunity* 33: 364–374.
9. Ohmori, K., Y. Luo, Y. Jia, J. Nishida, Z. Wang, K. D. Bunting, D. Wang, and H. Huang. 2009. IL-3 induces basophil expansion in vivo by directing granulocyte-monocyte progenitors to differentiate into basophil lineage-restricted progenitors in

- the bone marrow and by increasing the number of basophil/mast cell progenitors in the spleen. *J. Immunol.* 182: 2835–2841.
10. Arinobu, Y., H. Iwasaki, M. F. Gurish, S. Mizuno, H. Shigematsu, H. Ozawa, D. G. Tenen, K. F. Austen, and K. Akashi. 2005. Developmental checkpoints of the basophil/mast cell lineages in adult murine hematopoiesis. *Proc. Natl. Acad. Sci. USA* 102: 18105–18110.
 11. Arinobu, Y., H. Iwasaki, and K. Akashi. 2009. Origin of basophils and mast cells. *Allergol. Int.* 58: 21–28.
 12. Lantz, C. S., J. Boesiger, C. H. Song, N. Mach, T. Kobayashi, R. C. Mulligan, Y. Nawa, G. Dranoff, and S. J. Galli. 1998. Role for interleukin-3 in mast-cell and basophil development and in immunity to parasites. *Nature* 392: 90–93.
 13. Shen, T., S. Kim, J. S. Do, L. Wang, C. Lantz, J. F. Urban, G. Le Gros, and B. Min. 2008. T cell-derived IL-3 plays key role in parasite infection-induced basophil production but is dispensable for in vivo basophil survival. *Int. Immunol.* 20: 1201–1209.
 14. Lantz, C. S., B. Min, M. Tsai, D. Chatterjea, G. Dranoff, and S. J. Galli. 2008. IL-3 is required for increases in blood basophils in nematode infection in mice and can enhance IgE-dependent IL-4 production by basophils in vitro. *Lab. Invest.* 88: 1134–1142.
 15. Le Gros, G., S. Z. Ben-Sasson, D. H. Conrad, I. Clark-Lewis, F. D. Finkelman, M. Plaut, and W. E. Paul. 1990. IL-3 promotes production of IL-4 by splenic non-B, non-T cells in response to Fc receptor cross-linkage. *J. Immunol.* 145: 2500–2506.
 16. Hida, S., S. Yamasaki, Y. Sakamoto, M. Takamoto, K. Obata, T. Takai, H. Karasuyama, K. Sugane, T. Saito, and S. Taki. 2009. Fc receptor gamma-chain, a constitutive component of the IL-3 receptor, is required for IL-3-induced IL-4 production in basophils. *Nat. Immunol.* 10: 214–222.
 17. Ben-Sasson, S. Z., G. Le Gros, D. H. Conrad, F. D. Finkelman, and W. E. Paul. 1990. Cross-linking Fc receptors stimulate splenic non-B, non-T cells to secrete interleukin 4 and other lymphokines. *Proc. Natl. Acad. Sci. USA* 87: 1421–1425.
 18. Yoshimoto, T., H. Tsutsui, K. Tominaga, K. Hoshino, H. Okamura, S. Akira, W. E. Paul, and K. Nakanishi. 1999. IL-18, although antiallergic when administered with IL-12, stimulates IL-4 and histamine release by basophils. *Proc. Natl. Acad. Sci. USA* 96: 13962–13966.
 19. Schneider, E., A. F. Petit-Bertron, R. Bricard, M. Levasseur, A. Ramadan, J. P. Girard, A. Herbelin, and M. Dy. 2009. IL-33 activates unprimed murine basophils directly in vitro and induces their in vivo expansion indirectly by promoting hematopoietic growth factor production. *J. Immunol.* 183: 3591–3597.
 20. Smithgall, M. D., M. R. Comeau, B. R. Yoon, D. Kaufman, R. Armitage, and D. E. Smith. 2008. IL-33 amplifies both Th1- and Th2-type responses through its activity on human basophils, allergen-reactive Th2 cells, iNKT and NK cells. *Int. Immunol.* 20: 1019–1030.
 21. Kroeger, K. M., B. M. Sullivan, and R. M. Locksley. 2009. IL-18 and IL-33 elicit Th2 cytokines from basophils via a MyD88- and p38alpha-dependent pathway. *J. Leukoc. Biol.* 86: 769–778.
 22. Fort, M. M., J. Cheung, D. Yen, J. Li, S. M. Zurawski, S. Lo, S. Menon, T. Clifford, B. Hunte, R. Lesley, et al. 2001. IL-25 induces IL-4, IL-5, and IL-13 and Th2-associated pathologies in vivo. *Immunity* 15: 985–995.
 23. Wang, H., R. Mobini, Y. Fang, F. Barrenäs, H. Zhang, Z. Xiang, and M. Benson. 2010. Allergic challenge of peripheral blood mononuclear cells from patients with seasonal allergic rhinitis increases IL-17RB, which regulates basophil apoptosis and degranulation. *Clin. Exp. Allergy* 40: 1194–1202.
 24. Yoshimoto, T., K. Yasuda, H. Tanaka, M. Nakahira, Y. Imai, Y. Fujimori, and K. Nakanishi. 2009. Basophils contribute to T(H)2-IgE responses in vivo via IL-4 production and presentation of peptide-MHC class II complexes to CD4+ T cells. *Nat. Immunol.* 10: 706–712.
 25. Sabroe, I., E. C. Jones, L. R. Usher, M. K. Whyte, and S. K. Dower. 2002. Toll-like receptor (TLR)2 and TLR4 in human peripheral blood granulocytes: a critical role for monocytes in leukocyte lipopolysaccharide responses. *J. Immunol.* 168: 4701–4710.
 26. Bieneman, A. P., K. L. Chichester, Y. H. Chen, and J. T. Schroeder. 2005. Toll-like receptor 2 ligands activate human basophils for both IgE-dependent and IgE-independent secretion. *J. Allergy Clin. Immunol.* 115: 295–301.
 27. Schramm, G., K. Mohrs, M. Wodrich, M. J. Doenhoff, E. J. Pearce, H. Haas, and M. Mohrs. 2007. Cutting edge: IPSE/alpha-1, a glycoprotein from *Schistosoma mansoni* eggs, induces IgE-dependent, antigen-independent IL-4 production by murine basophils in vivo. *J. Immunol.* 178: 6023–6027.
 28. Phillips, C., W. R. Coward, D. I. Pritchard, and C. R. Hewitt. 2003. Basophils express a type 2 cytokine profile on exposure to proteases from helminths and house dust mites. *J. Leukoc. Biol.* 73: 165–171.
 29. Sokol, C. L., G. M. Barton, A. G. Farr, and R. Medzhitov. 2008. A mechanism for the initiation of allergen-induced T helper type 2 responses. *Nat. Immunol.* 9: 310–318.
 30. de Paulis, A., N. Prevete, I. Fiorentino, A. F. Walls, M. Curto, A. Petraroli, V. Castaldo, P. Ceppa, R. Fiocca, and G. Marone. 2004. Basophils infiltrate human gastric mucosa at sites of *Helicobacter pylori* infection, and exhibit chemotaxis in response to H. pylori-derived peptide Hp(2-20). *J. Immunol.* 172: 7734–7743.
 31. Patella, V., G. Florio, A. Petraroli, and G. Marone. 2000. HIV-1 gp120 induces IL-4 and IL-13 release from human Fc epsilon RI+ cells through interaction with the VH3 region of IgE. *J. Immunol.* 164: 589–595.
 32. Kurimoto, Y., A. L. de Weck, and C. A. Dahinden. 1989. Interleukin 3-dependent mediator release in basophils triggered by C5a. *J. Exp. Med.* 170: 467–479.
 33. Hida, S., M. Tadachi, T. Saito, and S. Taki. 2005. Negative control of basophil expansion by IRF-2 critical for the regulation of Th1/Th2 balance. *Blood* 106: 2011–2017.
 34. Charles, N., W. T. Watford, H. L. Ramos, L. Hellman, H. C. Oettgen, G. Gomez, J. J. Ryan, J. J. O'Shea, and J. Rivera. 2009. Lyn kinase controls basophil GATA-3 transcription factor expression and induction of Th2 cell differentiation. *Immunity* 30: 533–543.
 35. Kuroda, E., V. Ho, J. Ruschmann, F. Antignano, M. Hamilton, M. J. Rauh, A. Antov, R. A. Flavell, L. M. Sly, and G. Krystal. 2009. SHIP represses the generation of IL-3-induced M2 macrophages by inhibiting IL-4 production from basophils. *J. Immunol.* 183: 3652–3660.
 36. Zhu, J., H. Yamane, and W. E. Paul. 2010. Differentiation of effector CD4 T cell populations (*). *Annu. Rev. Immunol.* 28: 445–489.
 37. Min, B., M. Prout, J. Hu-Li, J. Zhu, D. Jankovic, E. S. Morgan, J. F. Urban Jr., A. M. Dvorak, F. D. Finkelman, G. LeGros, and W. E. Paul. 2004. Basophils produce IL-4 and accumulate in tissues after infection with a Th2-inducing parasite. *J. Exp. Med.* 200: 507–517.
 38. Voehringer, D., K. Shinkai, and R. M. Locksley. 2004. Type 2 immunity reflects orchestrated recruitment of cells committed to IL-4 production. *Immunity* 20: 267–277.
 39. Oh, K., T. Shen, G. Le Gros, and B. Min. 2007. Induction of Th2 type immunity in a mouse system reveals a novel immunoregulatory role of basophils. *Blood* 109: 2921–2927.
 40. Kim, S., T. Shen, and B. Min. 2009. Basophils can directly present or cross-present antigen to CD8 lymphocytes and alter CD8 T cell differentiation into IL-10-producing phenotypes. *J. Immunol.* 183: 3033–3039.
 41. Gauchat, J. F., S. Henchoz, G. Mazzei, J. P. Aubry, T. Brunner, H. Blasey, P. Life, D. Talabot, L. Flores-Romo, J. Thompson, et al. 1993. Induction of human IgE synthesis in B cells by mast cells and basophils. *Nature* 365: 340–343.
 42. Yanagihara, Y., K. Kajiwara, Y. Basaki, K. Ikizawa, M. Ebisawa, C. Ra, H. Tachimoto, and H. Saito. 1998. Cultured basophils but not cultured mast cells induce human IgE synthesis in B cells after immunologic stimulation. *Clin. Exp. Immunol.* 111: 136–143.
 43. Denzel, A., U. A. Maus, M. Rodriguez Gomez, C. Moll, M. Niedermeier, C. Winter, R. Maus, S. Hollingshead, D. E. Briles, L. A. Kunz-Schughart, et al. 2008. Basophils enhance immunological memory responses. *Nat. Immunol.* 9: 733–742.
 44. Chen, K., W. Xu, M. Wilson, B. He, N. W. Miller, E. Bengtén, E. S. Edholm, P. A. Santini, P. Rath, A. Chiu, et al. 2009. Immunoglobulin D enhances immune surveillance by activating antimicrobial, proinflammatory and B cell-stimulating programs in basophils. *Nat. Immunol.* 10: 889–898.
 45. Sokol, C. L., N. Q. Chu, S. Yu, S. A. Nish, T. M. Laufer, and R. Medzhitov. 2009. Basophils function as antigen-presenting cells for an allergen-induced T helper type 2 response. *Nat. Immunol.* 10: 713–720.
 46. Perrigoue, J. G., S. A. Saenz, M. C. Siracusa, E. J. Allenspach, B. C. Taylor, P. R. Giacomin, M. G. Nair, Y. Du, C. Zaph, N. van Rooijen, et al. 2009. MHC class II-dependent basophil-CD4+ T cell interactions promote T(H)2 cytokine-dependent immunity. *Nat. Immunol.* 10: 697–705.
 47. Phythian-Adams, A. T., P. C. Cook, R. J. Lundie, L. H. Jones, K. A. Smith, T. A. Barr, K. Hochweller, S. M. Anderton, G. J. Hämmerling, R. M. Maizels, and A. S. MacDonald. 2010. CD11c depletion severely disrupts Th2 induction and development in vivo. *J. Exp. Med.* 207: 2089–2096.
 48. Hammad, H., M. Plantinga, K. Deswarte, P. Pouliot, M. A. Willart, M. Kool, F. Muskens, and B. N. Lambrecht. 2010. Inflammatory dendritic cells—not basophils—are necessary and sufficient for induction of Th2 immunity to inhaled house dust mite allergen. *J. Exp. Med.* 207: 2097–2111.
 49. Ohnmacht, C., C. Schwartz, M. Panzer, I. Schiedewitz, R. Naumann, and D. Voehringer. 2010. Basophils orchestrate chronic allergic dermatitis and protective immunity against helminths. *Immunity* 33: 364–374.
 50. Kim, S., M. Prout, H. Ramshaw, A. F. Lopez, G. LeGros, and B. Min. 2010. Cutting edge: basophils are transiently recruited into the draining lymph nodes during helminth infection via IL-3, but infection-induced Th2 immunity can develop without basophil lymph node recruitment or IL-3. *J. Immunol.* 184: 1143–1147.
 51. Mukai, K., K. Matsuoka, C. Taya, H. Suzuki, H. Yokozeki, K. Nishioka, K. Hirokawa, M. Etori, M. Yamashita, T. Kubota, et al. 2005. Basophils play a critical role in the development of IgE-mediated chronic allergic inflammation independently of T cells and mast cells. *Immunity* 23: 191–202.
 52. Iikura, M., M. Miyamasu, M. Yamaguchi, H. Kawasaki, K. Matsushima, M. Kitauro, Y. Morita, O. Yoshie, K. Yamamoto, and K. Hirai. 2001. Chemokine receptors in human basophils: inducible expression of functional CXCR4. *J. Leukoc. Biol.* 70: 113–120.
 53. Grattan, C. E. 2001. Basophils in chronic urticaria. *J. Investig. Dermatol. Symp. Proc.* 6: 139–140.
 54. Bischoff, S. C., R. Zwahlen, M. Stucki, G. Müllner, A. L. de Weck, B. M. Stadler, and C. A. Dahinden. 1996. Basophil histamine release and leukotriene production in response to anti-IgE and anti-IgE receptor antibodies. Comparison of normal subjects and patients with urticaria, atopic dermatitis or bronchial asthma. *Int. Arch. Allergy Immunol.* 110: 261–271.
 55. Nigrovic, P. A., and D. M. Lee. 2007. Synovial mast cells: role in acute and chronic arthritis. *Immunol. Rev.* 217: 19–37.
 56. Damsker, J. M., A. M. Hansen, and R. R. Caspi. 2010. Th1 and Th17 cells: adversaries and collaborators. *Ann. N. Y. Acad. Sci.* 1183: 211–221.
 57. Atta, A. M., C. P. Sousa, E. M. Carvalho, and M. L. Sousa-Atta. 2004. Immunoglobulin E and systemic lupus erythematosus. *Braz. J. Med. Biol. Res.* 37: 1497–1501.
 58. Sekigawa, I., T. Yoshiike, N. Iida, H. Hashimoto, and H. Ogawa. 2003. Allergic diseases in systemic lupus erythematosus: prevalence and immunological considerations. *Clin. Exp. Rheumatol.* 21: 117–121.
 59. Veldhoen, M. 2009. The role of T helper subsets in autoimmunity and allergy. *Curr. Opin. Immunol.* 21: 606–611.
 60. Martin, F., and A. C. Chan. 2004. Pathogenic roles of B cells in human autoimmunity: insights from the clinic. *Immunity* 20: 517–527.
 61. Valenta, R., S. Seiberler, S. Natter, V. Mahler, R. Mossabeh, J. Ring, and G. Stingl. 2000. Autoallergy: a pathogenetic factor in atopic dermatitis? *J. Allergy Clin. Immunol.* 105: 432–437.

62. Kaveri, S. V., L. Mouthon, and J. Bayry. 2010. Basophils and nephritis in lupus. *N. Engl. J. Med.* 363: 1080–1082.
63. Crispin, J. C., S. N. Liossis, K. Kis-Toth, L. A. Lieberman, V. C. Kyttaris, Y. T. Juang, and G. C. Tsokos. 2010. Pathogenesis of human systemic lupus erythematosus: recent advances. *Trends Mol. Med.* 16: 47–57.
64. Crispin, J. C., V. C. Kyttaris, Y. T. Juang, and G. C. Tsokos. 2008. How signaling and gene transcription aberrations dictate the systemic lupus erythematosus T cell phenotype. *Trends Immunol.* 29: 110–115.
65. Davidson, A., and B. Diamond. 2010. Activated basophils give lupus a booster shot. *Nat. Med.* 16: 635–636.
66. Singh, R. R., V. Saxena, S. Zang, L. Li, F. D. Finkelman, D. P. Witte, and C. O. Jacob. 2003. Differential contribution of IL-4 and STAT6 vs STAT4 to the development of lupus nephritis. *J. Immunol.* 170: 4818–4825.
67. Rahman, A., and D. A. Isenberg. 2008. Systemic lupus erythematosus. *N. Engl. J. Med.* 358: 929–939.
68. Valencia, X., C. Yarboro, G. Illei, and P. E. Lipsky. 2007. Deficient CD4+CD25high T regulatory cell function in patients with active systemic lupus erythematosus. *J. Immunol.* 178: 2579–2588.
69. Zeng, D., Y. Liu, S. Sidobre, M. Kronenberg, and S. Strober. 2003. Activation of natural killer T cells in NZB/W mice induces Th1-type immune responses exacerbating lupus. *J. Clin. Invest.* 112: 1211–1222.
70. Pernis, A. B. 2009. Th17 cells in rheumatoid arthritis and systemic lupus erythematosus. *J. Intern. Med.* 265: 644–652.
71. Tiller, T., M. Tsuiji, S. Yurasov, K. Velinzon, M. C. Nussenzweig, and H. Wardemann. 2007. Autoreactivity in human IgG+ memory B cells. *Immunity* 26: 205–213.
72. Tsuiji, M., S. Yurasov, K. Velinzon, S. Thomas, M. C. Nussenzweig, and H. Wardemann. 2006. A checkpoint for autoreactivity in human IgM+ memory B cell development. *J. Exp. Med.* 203: 393–400.
73. Shimizu, S., N. Sugiyama, K. Masutani, A. Sadanaga, Y. Miyazaki, Y. Inoue, M. Akahoshi, R. Katafuchi, H. Hirakata, M. Harada, et al. 2005. Membranous glomerulonephritis development with Th2-type immune deviations in MRL/lpr mice deficient for IL-27 receptor (WSX-1). *J. Immunol.* 175: 7185–7192.
74. Masutani, K., M. Akahoshi, K. Tsuruya, M. Tokumoto, T. Ninomiya, T. Kohsaka, K. Fukuda, H. Kanai, H. Nakashima, T. Otsuka, and H. Hirakata. 2001. Predominance of Th1 immune response in diffuse proliferative lupus nephritis. *Arthritis Rheum.* 44: 2097–2106.
75. Atta, A. M., M. B. Santiago, F. G. Guerra, M. M. Pereira, and M. L. Sousa Atta. 2010. Autoimmune response of IgE antibodies to cellular self-antigens in systemic Lupus Erythematosus. *Int. Arch. Allergy Immunol.* 152: 401–406.
76. Yu, C. C., T. S. Yen, C. A. Lowell, and A. L. DeFranco. 2001. Lupus-like kidney disease in mice deficient in the Src family tyrosine kinases Lyn and Fyn. *Curr. Biol.* 11: 34–38.
77. Odom, S., G. Gomez, M. Kovarova, Y. Furumoto, J. J. Ryan, H. V. Wright, C. Gonzalez-Espinosa, M. L. Hibbs, K. W. Harder, and J. Rivera. 2004. Negative regulation of immunoglobulin E-dependent allergic responses by Lyn kinase. *J. Exp. Med.* 199: 1491–1502.
78. Beavitt, S. J., K. W. Harder, J. M. Kemp, J. Jones, C. Quilici, F. Casagrande, E. Lam, D. Turner, S. Brennan, P. D. Sly, et al. 2005. Lyn-deficient mice develop severe, persistent asthma: Lyn is a critical negative regulator of Th2 immunity. *J. Immunol.* 175: 1867–1875.
79. Liossis, S. N., E. E. Solomou, M. A. Dimopoulos, P. Panayiotidis, M. M. Mavrikakis, and P. P. Sfikakis. 2001. B-cell kinase lyn deficiency in patients with systemic lupus erythematosus. *J. Invest. Med.* 49: 157–165.
80. Charles, N., D. Hardwick, E. Daugas, G. G. Illei, and J. Rivera. 2010. Basophils and the T helper 2 environment can promote the development of lupus nephritis. *Nat. Med.* 16: 701–707.
81. Asai, K., J. Kitaura, Y. Kawakami, N. Yamagata, M. Tsai, D. P. Carbone, F. T. Liu, S. J. Galli, and T. Kawakami. 2001. Regulation of mast cell survival by IgE. *Immunity* 14: 791–800.
82. Navarra, S. V., R. M. Guzmán, A. E. Gallacher, S. Hall, R. A. Levy, R. E. Jimenez, E. K. Li, M. Thomas, H. Y. Kim, M. G. León, et al. 2011. Efficacy and safety of belimumab in patients with active systemic lupus erythematosus: a randomised, placebo-controlled, phase 3 trial. *Lancet* 377: 721–731.
83. Lin, H., K. M. Boesel, D. T. Griffith, C. Prussin, B. Foster, F. A. Romero, R. Townley, and T. B. Casale. 2004. Omalizumab rapidly decreases nasal allergic response and FcεRI on basophils. *J. Allergy Clin. Immunol.* 113: 297–302.
84. Schuerwegh, A. J., A. Ioan-Facsinay, A. L. Dorjée, J. Roos, I. M. Bajema, E. I. van der Voort, T. W. Huizinga, and R. E. Toes. 2010. Evidence for a functional role of IgE anticitrullinated protein antibodies in rheumatoid arthritis. *Proc. Natl. Acad. Sci. USA* 107: 2586–2591.
85. Mack, M., and A. R. Rosenkranz. 2009. Basophils and mast cells in renal injury. *Kidney Int.* 76: 1142–1147.