



Clonal deletion of T cell repertoires with specific T cell receptor V β chains by two endogenous superantigens in NC/Nga mice

Kozo Ohkusu-Tsukada, Teruyo Tsukada & Kimimasa Takahashi

To cite this article: Kozo Ohkusu-Tsukada, Teruyo Tsukada & Kimimasa Takahashi (2017): Clonal deletion of T cell repertoires with specific T cell receptor V β chains by two endogenous superantigens in NC/Nga mice, Bioscience, Biotechnology, and Biochemistry

To link to this article: <http://dx.doi.org/10.1080/09168451.2017.1374829>

 View supplementary material 

 Published online: 22 Sep 2017.

 Submit your article to this journal 

 View related articles 

 View Crossmark data 



Note

Clonal deletion of T cell repertoires with specific T cell receptor V β chains by two endogenous superantigens in NC/Nga miceKozo Ohkusu-Tsukada^{1,*} , Teruyo Tsukada² and Kimimasa Takahashi¹¹Department of Veterinary Pathology, Nippon Veterinary & Life-science University, Tokyo, Japan; ²Radiation Biology Team, Nishina Center for Accelerator-based Science, RIKEN, Saitama, Japan

Received June 13, 2017; accepted August 25, 2017

<https://doi.org/10.1080/09168451.2017.1374829>

Superantigens (SAGs) are powerful T-cell stimulatory proteins. Because an atopic dermatitis (AD) model NC/Nga mice had two endogenous SAGs, namely *minor lymphocyte-stimulating locus-1^a* (*Mls-1^a*) and *mouse mammary tumor virus (MMTV)*(*SHN*), SAG-responsive T-cells bearing V β 5.1, V β 6, V β 8.1, V β 8.2, V β 8.3, V β 9, and V β 11 should be endogenously deleted. Here, we discuss that the endogenous SAGs-expression may be involved in AD-sensitivity in NC/Nga mice.

Key words: Superantigen; TCR V β ; NC/Nga mice; *Mls-1^a*; *MMTV*(*SHN*)

Superantigens (SAGs) bind to major histocompatibility complex (MHC) class II molecules and activate both CD4⁺ and CD8⁺ T-cells bearing variable regions of specific T-cell receptor β chains (TCR V β) domain.¹⁾ Endogenous SAG of *mouse mammary tumor virus* (MMTV or *Mtv*) causes tolerance or the clonal deletion of T cells bearing a specific V β element, thus allowing proliferation of infected B cells and transmission of the virus to offspring.¹⁾ Of the murine endogenous SAGs encoded by *Mtv*, the *minor lymphocyte-stimulating locus-1^a* (*Mls-1^a*), encoded by an open reading frame in the 3' long terminal repeat of the *Mtv*-7 provirus, is expressed in AKR, CBA/J, C58, DBA/2, and NZB mice, where it induces tolerance or clonal deletion of T-cells bearing TCR V β 6, V β 8.1, and V β 9.¹⁾ TCR V β 7 is also weakly stimulated by *Mls-1^a*.²⁾ In contrast, *human endogenous retroviruses type K* (*HERV-K*), *HIV*, and *human T cell leukemia virus (HTLV)* are complex retroviruses with features similar to *MMTV*.³⁾ The *HERV-K* family also contains endogenous SAGs, suggesting their potential involvement in autoimmunity.^{3,4)} *HERV-K* family members (particularly *HERV-K10* and *HERV-K18*), *HTLV-1-related endogenous sequence (HRES)-1*, *endogenous retrovirus (ERV)-3*, and *HERV-E4-1* may also function in the pathogenesis of systemic lupus erythematosus.⁵⁾ Although *HERV-K18* is recognized by human T-cells bearing TCR V β 7 and

V β 13,^{3,4)} it is unknown whether other *HERV* are recognized by other T-cells bearing a specific V β element.

NC/Nga mice were established as an inbred strain by Dr K. Kondo in 1957 based on the Japanese fancy mice (Nishiki-Nezumi).⁶⁾ This strain has been reported to have the following biological characteristics: liver and kidney esterase isozyme patterns similar to those in DBA/2 mice and high susceptibility to X ray-irradiation and to anaphylactic shock from ovalbumin (OVA).⁶⁾ NC/Nga mice spontaneously develop atopic dermatitis (AD)-like disease under conventional conditions, but not under specific pathogen-free conditions.⁷⁾ Here, we show that the clonal deletion of T-cell repertoire with a specific TCR V β is due to the expression of two endogenous SAGs in NC/Nga mice, and discuss the role of endogenous SAGs in developing spontaneous AD-like disease in NC/Nga mice.

Animal; NC/Nga (H-2^{nc}, previously unknown),⁸⁾ BALB/c (H-2^d, *Mls-1^b*), DBA/2 (H-2^d, *Mls-1^a*), and B10.BR (H-2^k, *Mls-1^b*) (male) (Japan SLC, Hamamatsu, Japan) were maintained in SPF conditions, and used at 8–10 weeks of age. All mice were maintained in our full-barrier animal facility under controlled temperature, humidity, and a 12-hour light/dark regimen. All experiments were approved by the Institutional Review Board (IRB) for Animal Studies of the Nippon Veterinary and Life-science University (NVLU), and were performed following the guidelines provided by the committee.

Flow cytometry; Peripheral blood mononucleated cells (PBMCs) were isolated from mouse blood using Lymphoprep (ProGen, Heidelberg, Germany). For FACS analysis, PBMCs suspended in PBS were stained with a mixture of FITC or PE-conjugated CD4 (clone; GK1.5) or CD8 (clone; 53-6.7) and FITC-conjugated V β 8.1 8.2 (clone; KJ16-133.18), PE-conjugated V β 5.1 5.2 (clone; MR9-4), PE-conjugated V β 6 (clone; RR4-7)-specific mAbs, PE-conjugated V β 9 (clone; MR10-2)-specific mAbs, or PE-conjugated V β 11 (clone; KT11)-specific mAbs (BioLegend, St. Louis,

*Corresponding author. Email: tkd-oks@nvlu.ac.jp

Abbreviations: Sags, superantigens; TCR V β , variable regions of T cell receptor β chains; APCs, antigen-presenting cells; MHC, major histocompatibility complex; AD, atopic dermatitis; Mls, minor lymphocyte-stimulating locus; MMTV, mouse mammary tumor virus; HERV, human endogenous retroviruses; mAbs, monoclonal antibodies; SEB, Staphylococcal enterotoxin B.

MO). Two-color analysis was conducted by FACS (FACSCalibur, Nippon Becton Dickinson, Japan).

RT-PCR; CD4⁺ T-cells, CD8⁺ T-cells, B-cells (3×10^5 /well), or myeloid cells were enriched from LNs using anti-mouse CD4, anti-mouse CD8, anti-mouse B220, or anti-mouse CD11b Magnetic Particles-DM (BD Biosciences). Antigen-presenting cells (APCs) were used as matured myeloid cells cultured with treatment of 20 ng/ml recombinant (r)GM-CSF (Pepro-Tech, Rocky Hill, NJ) and 20 ng/ml rIL-4 (PeproTech) for 72 h at 37°C in 5% CO₂. These RNA were extracted using ISOGEN (Nippon Gene, Tokyo, Japan). First strand cDNA was synthesized using 1 µg of isolated RNA template, M-MLV reverse transcriptase, Oligo (dT)₁₅ primer, RNase inhibitor and dNTP according to the manufacturer's protocol (Nippon Gene). The primers (20 nM) for PCR were sense (5'-GTCAA-GAACAGGTGCAAGGAC-3') and antisense (5'-AAGGGATCGAAGCCAACGCG-3') for *Mls-1^a* (443 bp), sense (5'-TCTGCGCACAAACGGATGAG-3') and antisense (5'-AAGGGGGCATCTGTTGGTCT-3') for *MMTV(SHN)* (572 bp) (GenBank: X78590.1), sense (5'-ATGGCTTCTGTGGCTACAGACC-3') for Vβ2, sense (5'-AAGGACAAAAAGCAAAGATGAGG-3') for Vβ3, sense (5'-AACACTGCCTTCCCTGACCC-3') for Vβ5.1, sense (5'-CAAAACTGACCTTGAAATGT CAA-3') for Vβ6, sense (5'-AGAATGTTTTGCTG-GAATGTGGA-3') for Vβ7, sense (5'-GAAAGGTGA-CATTGAGCTGTCAC-3') for Vβ8.1, sense (5'-GGAAAGGTGACATTGAGCTGTAAT-3') for Vβ8.2, sense (5'-GAAAGGTGACATTGAGCTGTCAC-3') for Vβ8.3, sense (5'-CTTCTGTCTTCTTGCAGCCACTT-3') for Vβ9, sense (5'-TGCCTCTTGGAATAGGCC-3') for Vβ10, sense (5'-TGCTTCTTGAGAGCAGAAC-CAA-3') for Vβ11, antisense (5'-GCAATCTCTGC TTTTGATGGCT-3') for Cβ,⁹⁾ and sense (5'-ACCA-CAGTCCATGCCATCAC-3'), and antisense (5'-TCCA CCACCTGTGCTGTA-3') for GAPDH (981 bp). The PCR conditions of *Mls-1^a*, *MMTV(SHN)*, Vβ5.1, and Vβ6 using *Taq* DNA polymerase (BioAcademia, Osaka, Japan) were as follows: 36 cycles at 98 °C for 10 s, 56.5 °C for 30 s, and 72 °C for 70 s, then the annealing temperature for Vβ2, Vβ3, Vβ7, and Vβ8.1 are 61 °C, and the annealing temperature for Vβ8.2, Vβ8.3, Vβ9, Vβ10, and Vβ11 are 58.5 °C. The products were diluted five-fold with loading buffer consisting of xylene cyanol and bromophenol blue dyes and were electrophoresed on 2% agarose gel.

Statistical analysis; Statistical analysis was performed by ANOVA using Excel (Microsoft) and Stat-Plus (AnalystSoft, Alexandria, VA). A *p*-value < 0.05 was considered significant.

The *Mls-1^a* and *MMTV(SHN)* genes were detected in mRNA of NC/Nga mice B-cells or antigen-presenting cells (APCs) using RT-PCR (Fig. 1(A) and (B)). Although the clonal deletion of TCR Vβ6 and Vβ9 by *Mls-1^a* expression in NC/Nga mice was assumed to be the same as in control DBA/2 mice (*Mtv-1*, *Mtv-6*, *Mtv-7*, and *Mtv-13*), other T-cells bearing TCR Vβ5.1, Vβ8.1, Vβ8.2, Vβ8.3, and Vβ11 should be also deleted in NC/Nga mice CD4⁺ T-cells (Fig. 1(c)). Then, when the expression of Vβ chains on also thymocytes and




Fig. 1. The expression of TCR Vβ repertoires and endogenous superantigens in NC/Nga mice. (A) Endogenous superantigen *Mls-1^a*-specific expression was measured by RT-PCR using APCs purified from BALB/c and NC/Nga mice and B-cells purified from BALB/c (*Mls-1^b*), NC/Nga (unknown), and DBA/2 (*Mls-1^a*) mice, (B) The mRNA expression of *MMTV(SHN)* in B-cells purified from BALB/c, B10.BR, and NC/Nga mice were measured using RT-PCR, and (c) The mRNA expression of TCR Vβ repertoires (Vβ2, Vβ3, Vβ5.1, Vβ6, Vβ7, Vβ8.1, Vβ8.2, Vβ8.3, Vβ9, Vβ10, and Vβ11) in CD4⁺ T-cells purified from BALB/c, DBA/2, and NC/Nga mice was measured using RT-PCR. Data are representative of two independent experiments.

LN cells in NC/Nga strain was compared, the results indicated before and after clonal deletion of peripheral T cells in NC/Nga mice. (Supplemental Fig. S1). Because *MMTV(SHN)* mRNA was expressed in B10.BR or NC/Nga mice but not BALB/c mice (Fig. 1(B)), the clonal deletion of a specific TCR Vβ by *MMTV(SHN)* should also be induced. Besides, the PCR products of *Mls-1^a* and *MMTV(SHN)* were confirmed by the sequence analysis (Supplemental Fig. S2). Furthermore, TCR Vβ11, but not Vβ5.1, was deleted in BALB/c mice (*Mtv-6*, *Mtv-8*, and *Mtv-9*), although it is reported that the expression of *Mls-1^a* in BALB/c mice induced the deletion or tolerance of T-cells bearing TCR Vβ5.1, Vβ11, and Vβ12.¹⁰⁾

RT-PCR results were then confirmed using FACS analysis. T-cell repertoires of TCR Vβ6, Vβ8.1, Vβ8.2, Vβ9, and Vβ11 were clearly deleted in NC/Nga mice. However the deletion of Vβ5.1 in NC/Nga mice was not confirmed by FACS analysis because the population of CD4⁺ T cells was detected by mAb specific to Vβ5.1 5.2 (data not shown). Vβ11 was deleted in also BALB/c mice (*p* < 0.05, ANOVA). Vβ6 and Vβ9 were deleted in DBA/2 mice having *Mls-1^a* (*p* < 0.05, ANOVA; Fig. 2(A) and (B)).

Because physiological barrier dysfunctions of the skin, high numbers of *Staphylococcus aureus* among the skin surface bacterial flora, and an increased hypersensitivity to itch were reported in AD patients,¹¹⁾ the effects of applying exogenous SAg *Staphylococcal enterotoxin B (SEB)* to the skin of AD model NC/Nga mice have been investigated.¹²⁾ However, the effect of




Fig. 2. FACS analysis of the expression of TCR V β 6, V β 8.1–8.2, V β 9, and V β 11. (A) Samples were stained using a mixture of mAbs specific to CD4 and TCR V β 6, V β 8.1–8.2, V β 9, or V β 11 in PBMCs of NC/Nga, BALB/c, B10.BR, and DBA/2 mice and analyzed using FACS. An example of a two-color panel is shown. (B) The percentages of TCR V β 6, V β 8.1–8.2, V β 9, or V β 11-positive populations in CD4⁺ cells (dark gray) or CD8⁺ cells (light gray) of NC/Nga, BALB/c, B10.BR, and DBA/2 mice ($n = 4$) are shown. Significant downregulation (down arrow) by clonal deletion is indicated.

Notes: * $p < 0.02$; ANOVA, when compared with the respective NC/Nga mice. ** $p < 0.05$, ANOVA, when compared with respective *-marks. Bars represent mean $\pm$ SE of data. Data are representative of two independent experiments.

SEB on NC/Nga mice may be reviewed because it was identified that endogenous SAg *Mls-1^a* and *MMTV (SHN)* were expressed in these mice. Although it was reported that TCR V β 8⁺ T-cells, but not V β 2.1⁺ and V β 7.1⁺ T-cells, were absent in NC/Nga mice,¹²⁾ it is considered to be caused by the endogenous clonal deletion by *Mls-1^a*. The expression of the *Mtv-7 provirus* product, *Mls-1^a*, induces the deletion or tolerance of TCR V β 6, V β 8.1, and V β 9.¹⁾ Our results indicated NC/Nga mice lack T-cells bearing TCR V β 5.1, V β 6, V β 8.1, V β 8.2, V β 8.3, V β 9, and V β 11 and contain *MMTV(SHN)* in addition to *Mls-1^a*. Because the expression of *Mtv-8*, *Mtv-9*, and *Mtv-11 proviruses* product, *Mls^f*, induces the deletion or tolerance of TCR V β 5.1, V β 11, and V β 12,¹⁰⁾ and the expression of *Mtv-4 provirus* product, *MMTV(SHN)*, induces the deletion or tolerance of TCR V β 7, V β 8.1, V β 8.2, and V β 8.3,¹³⁾ we suggest that NC/Nga mice also contain *Mls^f* in addition to *Mls-1^a* and *MMTV(SHN)*. In contrast, *SEB* stimulates V β 3⁺, V β 7⁺, V β 8.1⁺, V β 8.2⁺, V β 8.3⁺, or V β 17⁺ T-cells in mice.¹²⁾ Interestingly, *SEB*-induced autoreactive T-cell responses were blocked by IL-10 production from tolerized V β 8⁺ T-cells in CBA/J mice (*Mls-1^a*).¹⁴⁾ A similar phenomenon may also occur in NC/Nga mice. *SEB*-induced Th1 dominant state mediated by IL-12 or IL-18 was inhibited by the absence of V β 8⁺ T-cells in NC/Nga mice, resulting in a Th2 dominant state.¹²⁾ Recent studies conducted on the pathogenesis, biomarkers, and itch in AD have reported the potential role of *Staphylococcus aureus*, IL-18, nerve growth factor (NGF), and semaphorin 3A in the pathogenesis and treatment of AD.¹⁵⁾ To investigate this, NC/Nga mice have been used as an AD model. Therefore, the correct characterization of NC/Nga mice is important when interpreting the *in vivo* responses.

Author contribution

K. O-T. designed the experiments, and performed the experiments. The manuscript was written by K. O-T. All authors participated in the discussion of the data and in the production of the final version of the manuscript.

Acknowledgment

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors. The authors would like to thank Enago for the English language review.

Disclosure statement

No potential conflict of interest was reported by the authors.

Supplemental material

The supplemental material for this paper is available at <https://doi.org/10.1080/09168451.2017.1374829>.

ORCID

Kozo Ohkusu-Tsukada  <http://orcid.org/0000-0002-4998-8895>

References

- [1] Acha-Orbea H, Held W, Waanders GA, et al. Exogenous and endogenous mouse mammary tumor virus superantigens. *Immunol Rev.* 1993;131:5–25.

- [2] Waanders GA, McDonald HR. Hierarchy of responsiveness *in vivo* and *in vitro* among T cells expressing distinct *Mls-I^a*-reactive V beta domains. *Eur J Immunol.* 1992;22:291–293.
- [3] Conrad B, Weissmahr RN, Böni J, et al. A human endogenous retroviral superantigen as candidate autoimmune gene in type I diabetes. *Cell.* 1997;90:303–313.
- [4] Sutkowski N, Conrad B, Thorley-Lawson DA, et al. Epstein-barr virus transactivates the human endogenous retrovirus HERV-K18 that encodes a superantigen. *Immunity.* 2001;15:579–589.
- [5] Nelson P, Rylance P, Roden D, et al. Viruses as potential pathogenic agents in systemic lupus erythematosus. *Lupus.* 2014;23:596–605.
- [6] Kondo K, Nagami T, Tadokoro S. Comparative cellular and species radiosensitivity. In: Bond PV, Sugawara T, editors. Tokyo: Igakushoin; 1969. p. 20.
- [7] Matsuda H, Watanabe N, Geba GP, et al. Development of atopic dermatitis-like skin lesion with IgE hyperproduction in NC/Nga mice. *Int Immunol.* 1997;9:461–466.
- [8] Ohkusu-Tsukada K, Yamashita T, Tsukada T, et al. Low expression of a D^{dm7}/L^{dm7}-hybrid mutant (D/L^{dm7}) in the novel haplotype H-2^{nc} identified in atopic dermatitis model NC/Nga mice. *Gene Immun.* Forthcoming.
- [9] Wettstein P, Strausbauch M, Therneau T, et al. The application of real-time PCR to the analysis of T cell repertoires. *Nucleic Acids Res.* 2008;36:e140.
- [10] Foo-Phillips M, Kozak CA, Principato MA, et al. Characterization of the Mls^f system. II. Identification of mouse mammary tumor virus proviruses involved in the clonal deletion of self-Mlsf-reactive T cells. *J Immunol.* 1992;149:3440–3447.
- [11] Stalder JF, Fleury M, Sourisse M, et al. Local steroid therapy and bacterial skin flora in atopic dermatitis. *Br J Dermatol.* 1994;131:536–540.
- [12] Habu Y, Seki S, Takayama E, et al. The mechanism of a defective IFN-gamma response to bacterial toxins in an atopic dermatitis model, NC/Nga mice, and the therapeutic effect of IFN-gamma, IL-12, or IL-18 on dermatitis. *J Immunol.* 2001;166:5439–5447.
- [13] Luther S, Shakhov AN, Xenarios I, et al. New infectious mammary tumor virus superantigen with V beta-specificity identical to staphylococcal enterotoxin B (SEB). *Eur J Immunol.* 1994;24:1757–1764.
- [14] Ito K, Takaishi H, Jin Y, et al. Staphylococcal enterotoxin B stimulates expansion of autoreactive T cells that induce apoptosis in intestinal epithelial cells: regulation of autoreactive responses by IL-10. *J Immunol.* 2000;164:2994–3001.
- [15] Ikezawa Z, Komori J, Ikezawa Y, et al. A role of *Staphyococcus aureus*, interleukin-18, nerve growth factor and semaphorin 3A, an axon guidance molecule, in pathogenesis and treatment of atopic dermatitis. *Allergy Asthma Immunol Res.* 2010;2:235–246.