

**Galectin 3 modulates immune and inflammatory responses during
helminthic infection: impact of galectin 3 deficiency
on the functions of dendritic cells**

**Laetitia Breuilh,^{1,2,3,5, ‡} François Vanhoutte,^{1,2,3, ‡} Josette Fontaine,^{1,2,3} Caroline M.W. van
Stijn,⁴ Isabelle Tillie-Leblond,^{1,2,5} Monique Capron,^{1,2,3} Christelle Faveeuw,^{1,2,3}
Thierry Jouault,^{2,6} Irma van Die,⁴ Philippe Gosset,^{1,2,5} and François Trottein^{1,2,3,*}**

Institut National de la Recherche Médicale, U547,¹ Institut Pasteur de Lille, Institut Fédératif
de Recherche 142,² Université de Lille 2, Lille, F-59019 France,³ VU University Medical
Center, 1081 BT Amsterdam, The Netherlands,⁴ Inserm, U774, Lille, F-59019 France,⁵
Inserm U799, Lille, F-59037 France⁶

[‡] Contribute equally to the work

Running title : Galectin-3 during schistosomiasis

* Corresponding author. Mailing address: Unité Inserm U547, Institut Pasteur de Lille, 1 rue
du Professeur Calmette, 59019 Lille Cedex, France. Phone: (333) 2087-7885. Fax: (333)
2087-7888. E-mail: françois.trottein@pasteur-lille.fr.

ABSTRACT

Galectin-3 (Gal-3) is a multifunctional β -galactoside-binding lectin that senses self and microbial glycoconjugates. Although important in immune reactions and host defence in some experimental models, the function of Gal-3 during helminthic diseases (e.g. schistosomiasis) is still elusive. We show that, compared to wild-type *S. mansoni*-infected mice, infected Gal-3^{-/-} mice have a reduced number of T and B lymphocytes in the spleen, develop reduced liver granulomas 7 wk (acute phase) and 14 wk (chronic phase) post-infection and mount a biased cellular and humoral Th1 response. In an attempt to understand this latter phenomenon, we studied the role of endogenous Gal-3 in dendritic cells (DCs), the most potent APC, both *in vitro* and *in vivo*. Although Gal-3 deficiency in DCs does not impact on their differentiation and maturation processes, it greatly influences the strength (but not the nature) of the adaptive immune response that they trigger, suggesting that Gal-3 deficiency in some other cell types may be important during murine schistosomiasis. As a whole, this study implies Gal-3 as a modulator of the immune/inflammatory responses during helminthic infection and reveals for the first time that Gal-3 expression in DCs is pivotal to control the magnitude of T lymphocyte priming.

INTRODUCTION

Mammalian lectins display important functions on the innate/acquired immune system by sensing self and/or pathogen-derived glycoconjugates. Among these, galectins represent an emerging β -galactoside-binding lectin family involved in the regulation of host immune responses during physiological and pathological conditions, including inflammation, cancer and infection (for reviews, (2, 32, 33, 41-43)). Although they lack a classical signal sequence, galectins are present in extracellular fluid and on the cell surface and are also located inside the cells (for review, (22)). Extracellular and intracellular galectins display numerous functions including cell adhesion, signalling, proliferation, differentiation, survival and apoptosis (for review, (32)). Among the 15 galectin members, galectin-3 (Gal-3, previously known as Mac-2) is expressed in many immunocompetent/inflammatory cells, including monocytes, dendritic cells (DCs), macrophages, eosinophils, mast cells, NK cells and activated T and B cells (for reviews, (8, 16, 48)). Through its ability to bind polylactosamine (poly-LN) structures ($\text{Gal}\beta 1\text{-4GlcNAc}_n$) on endogenous ligands (20, 28), Gal-3 intervenes in many cellular processes *in vitro*. In particular, it favours cell-cell (for instance DC/T cell) and cell-matrix glycoprotein adhesion (37), exerts chemotactic effects (47), controls cell proliferation (13, 58), and promotes phagocytosis by macrophages (19, 46, 54). Gal-3 also takes part in the control of T cell and monocyte survival and activation. For instance, extracellular Gal-3, by associating with *N*-glycans on T cell receptor (TCR) has been shown to down-modulate TCR responsiveness and to regulate the production of Th1 and Th2 cytokines by differentiated T cells (13, 35). On the other hand, intracellular Gal-3 can positively or negatively impact on intracellular signaling pathways by regulating the activities of various kinases, including protein kinases C, mitogen-activated protein kinase and phosphatidylinositol 3-kinase (8). The role of Gal-3 in the promotion/control of inflammation and in the regulation of the immune response has been recently evaluated in different models.

Gal-3 has been proposed to be a powerful pro-inflammatory signal *in vitro* (24, 27, 47, 57) and studies of Gal-3 deficient mice have provided support for the pro-inflammatory role of this lectin (10, 21, 59). For instance, the lack of Gal-3 protects mice against asthma reactions, an effect associated with an enhanced Th1 response in challenged animals (59). On the other hand, administration of Gal-3 (by mean of gene therapy) inhibits asthmatic reactions (12, 34). These studies reveal opposite effects of the endogenous and exogenous Gal-3 and suggest that, according to its location (extracellular vs intracellular), Gal-3 differentially controls immune/inflammatory cell survival, migration and cytokine release.

Along with its role in inflammation and immune responses in non-infectious conditions, Gal-3 can also sense certain microorganisms. Recently, it has been demonstrated that Gal-3 binds to multimeric LN and LDN (LacdiNAc, GalNac β 1-4GlcNAc) motifs (54) as well as to other carbohydrate structures on glycoproteins and glycolipids (including lipopolysachcharides, LPS) from many pathogens such as (myco)bacteria, protozoan parasites and yeasts (36). Through this activity, Gal-3 participates in the phagocytosis of some microorganisms by macrophages and triggers host responses to pathogens, at least *in vitro* (26, 46). Whether or not it also contributes to signalling events in accessory cells during *in vivo* infection is unclear. Likewise, its role in signalling pathways triggered by Toll-like receptors (TLRs), which represent key sensors in innate cells (for review, (23)), is still elusive. The *in vivo* role of Gal-3 during infection has not been extensively studied. It appears to clear late *Mycobacterium tuberculosis* infection (5), to contribute to neutrophil recruitment at the site of *Streptococcal pneumonia* infection (49) and to exert an important role in innate immunity during *Toxoplasma gondii* infection (6). In the present report, to further investigate the role of Gal-3 in infection and innate/acquired immunity, we studied its role during murine schistosomiasis, an helminthic disease characterized by a dominant Th2 response triggered by

the egg stage of the parasite (for review, (39)). Recently, van den Berg *et al.* demonstrated that LDN motifs, which are expressed by *Schistosoma mansoni* eggs (29, 50), bind Gal-3 and suggested that this interaction may play an important role in the Th2-mediated inflammatory response that occurs in the liver (54). Support for this hypothesis has been recently provided by Van de Vijver and co-workers who showed that beads that carry glycoconjugates with terminal LN or LDN elements gave rise to granulomas similar to the granulomatous lesions caused by schistosome eggs in a natural infection (53). Recently, although conflicting, two studies have addressed the role of endogenous Gal-3 during *Schistosoma* infection (7, 38). Here, in agreement with (38), we show that, relative to wild-type (WT) mice, *S. mansoni*-infected Gal-3 deficient mice develop reduced granuloma formation and have a dramatic decreased number of total lymphocytes in the spleen. Moreover, Gal-3 deficiency results in a more pronounced Th1 response in infected mice. In an attempt to understand this latter phenomenon, we studied the role of endogenous Gal-3 in DCs both *in vitro* and *in vivo*. We report that Gal-3 deficiency in DCs is important to control the priming of T lymphocytes but has no role in the Th1/Th2 balance of the immune response suggesting that Gal-3 deficiency in some other cell types may be important during murine schistosomiasis.

MATERIALS AND METHODS

Animals. Female 129 mice were purchased from Iffa-Credo (I'Arbresle, France). The generation of Gal-3 deficient (Gal-3^{-/-}) 129 mice has been described by Colnot et al. (9). These animals show no morphological abnormalities and have no apparent phenotypes under standard pathogen-free conditions.

Reagents and Abs. mAbs against CD3 (unconjugated or FITC-conjugated), CD4 (unconjugated), Mac-1 (unconjugated), CD62L (biotin-conjugated), CD11c (biotin or APC-conjugated), CD19 (APC-conjugated), CD4 (FITC-conjugated), CD25 (APC-conjugated), and B220 (PE-conjugated) were purchased from Becton Dickinson (Le Pont de Claix, France). The 120G8 mAb was a gift from Dr Asselin-Paturel (3). APC-conjugated streptavidin was from Clinisciences (Montrouge, France) and CD4⁺ T cell isolation kit and streptavidin-conjugated microbeads used for MACS purification were from Miltenyi Biotec (Paris, France). Biotinylated mAbs against murine IgG2b and IgG1 were purchased from Southern Biotechnologies (Birmingham, USA). Anti-Gal-3 mAb (clone SC-32790, PE-conjugated or not) was purchased from Santa Cruz Biotechnology (Santa Cruz, California) and anti- β -actin Ab and HRP-conjugated anti-mouse IgG were from Sigma Aldrich (Saint Quentin Fallavier, France). Pam₃CSK₄ was purchased from EMC Microcollections (Tuebingen, Germany) and polyinosine-polycytidylic acid (poly(I:C)) and ultrapure lipopolysaccharide (LPS) (from *Escherichia coli* serotype 0111:B4) were from Cayla (Toulouse, France). Polyacrylamide (PAA) coupled glycoconjugates LDN-PAA, LN-PAA and glucitol-PAA (~ 20% substitution) (FITC-labelled or not) were from Lectinity (Lappeenranta, Finland). Lactose was from Sigma Aldrich.

Parasites and Ag preparation. *Schistosoma mansoni* (Puerto Rican strain) cercariae were obtained from infected *Biomphalaria glabrata* snails (44). Schistosome eggs and soluble egg Ag (52) were prepared as previously described (17).

Infection of mice and analysis of the parasitological and immunological parameters.

Eight-week old female WT and Gal-3^{-/-} mice were infected with 45 *S. mansoni* cercariae via the percutaneous route. Worm burdens were measured by liver perfusion 7 wk post infection (p.i.). At the time of perfusion, the small intestines and the livers were also collected for measurement of egg numbers deposited in these organs. Tissues were digested in 4% KOH as previously described (15). Records of the organ weights allowed the calculation of total tissue eggs per organ. Spleens and mesenteric lymph nodes (MLNs) were harvested 7 and 14 wk p.i. and cells were stimulated with increasing doses of SEA for 3 days at 37°C. During the last 18 h, 0.5 µCi of ³H-thymidine/well was added. Production of IFN-γ, IL-4, IL-5, IL-10 and IL-13 was measured by ELISA (R&D Systems, Abingdon, UK). To study the humoral response, mice were bled 7 and 14 wk p.i. and the anti-SEA IgG1 and IgG2b titrations were determined for each mouse by ELISA (17). Serum titers are defined as the dilution which gives an optical density reading at least two fold higher than the mean background of non-infected mouse serum.

Analysis of the egg-associated liver pathology in infected mice. The egg-associated pathology was analyzed on the livers of mice 7 and 14 wk p.i. in a double blind fashion. Briefly, a small piece of liver from each mouse was fixed in Immunohistofix (Genta Molecular Products, Brussels, Belgium) for 2-3 days and embedded in wax. Then, 5 µm liver sections were prepared for Rojkind coloration to evaluate collagen deposition. The staining mix is made of a saturated solution of picric acid in distilled water containing 0.1% Fast

Green FCF (Sigma-Aldrich, Saint Quentin Fallavier, France) and 0.1% Sirius Red. After staining, slides were washed in physiological water, and mounted in Acrytol (Surgipath, Labonord, France) after dehydration. The diameters of granulomas surrounding single, mature, and viable eggs were measured using the Metaview computing software (Roper Scientific, Evry, France), and the volume of each granuloma was calculated assuming a spherical shape. The proportion of eosinophils was determined by May Grunwald Giemsa staining and that of monocytes/macrophages and T lymphocytes by immunohistochemistry using anti-Mac-1 and anti-CD4 mAbs, respectively. Collagen deposition in granulomas was visualized by Sirius red staining and fibrosis was quantified using an arbitrary scale from 0 (no collagen deposition) to 3 (strong and tight collagen deposition all over the granuloma). For each parameters, ~ 5 granulomas/mouse were examined.

Preparation and analysis of BM-DCs. Bone-marrow (BM)-DCs were generated from the BMs of WT or Gal-3^{-/-} 129 mice by culture in GM-CSF-conditioned medium as previously described (1). DCs were used on day 14 of culture (>95% pure, as assessed by CD11c immunostaining). Gal-3 expression in BM-DCs was analyzed by Western blot using anti-galectin-3 mAb (0.4μg/ml). Phenotypic analysis was performed by flow cytometry using standard procedures. For endocytosis analysis, immature WT and Gal-3^{-/-} BM-DCs (2 x 10⁵ cells/well) were incubated in the presence of FITC-labelled glucitol-, LDN- or LN-conjugated PAA (50μg/ml) or FITC-labelled dextran (10μg/ml) for 2 h at 4°C or at 37°C, to evaluate extracellular binding and specific endocytosis. Glycan endocytosis was then assessed by flow cytometry. In activation experiments, BM-DCs (1 x 10⁶ cells/ml) were cultured with live eggs (1/200 cells), Pam₃CSK₄ (500ng/mL) plus poly(I:C) (2μg/mL) or with LPS (100ng/mL), or were left untreated. In some experiments, LDN- or glucitol-conjugated PAA (50μg/ml) were added 30 min before TLR agonists. After 18-20 h, culture supernatants were collected and

TNF- α , IL-6, IP-10/CXCL10 and MIG/CXCL9 concentrations were measured by ELISA (R&D Systems).

Mixed Leucocyte Reaction (MLR). WT and Gal-3^{-/-} BM-DCs (3×10^6 cells/plate), either stimulated or not, were treated with mitomycin ($25\mu\text{g/ml}$) for 30 min at 37°C , extensively washed and co-cultured at different ratio with purified naïve CD4⁺ T cells in a flat-bottom 96-well plate (1×10^5 T cells/well). Naïve CD4⁺ T cells were obtained from Balb/c mice spleens by MACS purification, using a CD4⁺ T cell isolation kit, followed by anti-CD62L mAb treatment and streptavidin-conjugated microbead purification (CD4⁺ CD62L⁺, purity > 95%). Five days later, cocultures were transferred into anti-CD3 mAb-bound plates for 48 h at 37°C . Culture supernatants were then harvested and cytokine production was determined by ELISA. Proliferation was assessed by Blue Alamar addition ($20\mu\text{l/well}$), and the OD was measured 24 h later, according to the manufacturer's protocol (Serotec, Cergy Saint-Christophe, France).

Immunization protocol and analysis of the immune response. After overnight culture, Keyhole Limpet haemocyanin (KLH)-pulsed DCs, cultured in the presence or absence of schistosome eggs (1:200 DCs), were collected, washed and administered at a dose of 1.5×10^6 cells i.v. in syngenic naïve mice. Seven days later, splenic cells (5×10^5 cells/well in flat-bottom 96-well plate) were harvested and restimulated with KLH or SEA for 72 h at 37°C .

Statistical analysis. Results are expressed as the mean $\pm$ SD. All other p values were determined by using the Student's t test. Values of $p < 0.05$ were considered as significant.

RESULTS

***Schistosoma mansoni*-infected Gal-3^{-/-} mice have an impaired number of splenic B and T cells, relative to WT infected mice.**

Gal-3^{-/-} and WT mice were percutaneously infected with *S. mansoni* and 7 wk later, the parasitological parameters were analyzed. As shown in Table I, worm burdens (both males and females, not shown) as well as tissue (liver and intestine) egg numbers were similar in the two groups, indicating that the lack of Gal-3 does not influence parasite survival and fecundity. This is consistent with recent results reporting no role of Gal-3 in the establishment of *Schistosoma* infection in mice (7, 38). Having established this, we next determined the frequency and the total number of B and T lymphocytes, as well as DCs, in the spleens of WT and Gal-3^{-/-} infected mice. Relative to uninfected WT animals, we noticed no variations in the frequency or in the number of splenic B cells (CD3⁻ CD19⁺) and T cells (CD3⁺ CD19⁻), including CD4⁺ CD25⁺ T cells in uninfected Gal-3^{-/-} mice (Table II). In *S. mansoni*-infected mice, although the frequency of these cells remained unchanged, except for T cells (decreased proportion compared to uninfected mice), their total number increased dramatically in WT-, but less in Gal-3^{-/-}-infected animals (~ 40% reduction). These data are in agreement with those from Oliveira et al. who reported a reduction in total lymphocyte counts in the spleen of *S. mansoni*-infected Gal-3 deficient mice (38). Analysis of both myeloid (CD11c⁺ 120G8⁻) and plasmacytoid (CD11c⁺ 120G8⁺) DCs did not show any significant variation between the two uninfected animal groups. Interestingly, in WT and Gal-3^{-/-}-infected mice, we observed a dramatic enhancement (~ 2.2 fold to 3.5 fold) in the frequency and number (x9) of myeloid DCs. On the other hand, the number of plasmacytoid DCs remained unchanged during infection and was similar in both WT and Gal-3^{-/-} mice. Thus, Gal-3 deficiency affects the number, but not the proportion, of both splenic T and B cells in *S. mansoni*-infected mice whereas it does not modulate those of DCs.

Infected Gal-3^{-/-} mice develop a biased Th1 response compared to WT controls. We next determined the nature of the cellular immune response in infected WT and Gal-3^{-/-} mice. To this end, spleens and MLNs were removed 7 wk p.i. and cells were restimulated with various doses of SEA. As seen in Fig. 1A, upon *in vitro* SEA challenge, splenic or MLN cells from WT and Gal-3^{-/-} mice proliferate in a comparable manner. In contrast, major differences were observed in terms of cytokine synthesis. Interestingly, the production of the Th1 cytokine IFN- γ by SEA-stimulated splenic and MLN cells from Gal-3^{-/-} infected mice was dramatically increased (~ 4 to 6-fold), compared to WT mice (Fig. 1B). On the other hand, the production of the Th2-type cytokines IL-4, IL-5 and IL-13 (not shown) was similar between Gal-3^{-/-} and WT mice. Notably, although no significant differences were detected in splenic cells, the level of IL-10 was decreased in the supernatant of Gal-3^{-/-} MLN cells.

We then determined the titers of SEA-specific IgG1 (a marker of a Th2 response) and IgG2b (a marker of a Th1 response) in infected mice. Although the overall Ig levels were identical in both animal groups (not shown), Gal-3^{-/-}-infected mice produced increased SEA-specific IgG2b (Fig. 2) and IgG2a (not shown), compared to control mice, in agreement with the enhanced production of IFN- γ seen in Fig. 1A. Although the level of Th2-type cytokines was not affected in these mice, Gal-3^{-/-}-infected mice produced less SEA-specific IgG1 (Fig. 2) compared to WT animals. Overall, this suggests that Gal-3 is important in the Th1/Th2 balance of the immune response during infection.

Gal-3 influences the development of liver granulomas in *S. mansoni*-infected mice.

We then evaluated the influence of Gal-3 deficiency on the formation of liver granulomas during the acute (7 wk) and chronic (14 wk) phases of the disease. In Gal-3^{-/-}-infected mice, and relative to WT animals, the size of the granulomas surrounding eggs was significantly decreased at wk 7 (Fig. 3A, left panel). However, analysis of the cellular composition of the

granulomas indicated no significant difference in the frequency of eosinophils (~75%), monocytes/macrophages (~2%) and T lymphocytes (~2%) between the two animal groups (not shown). Thus, during the acute phase of the disease, Gal-3 appears to contribute, in a quantitative but not qualitative manner, in the recruitment of immune/inflammatory cells into granulomas. Rojkind staining of liver sections showed no significant difference in the deposition of collagenous material and in fibrosis between the two animal groups 7 wk p.i. (Fig. 3B and C).

We next investigated the impact of Gal-3 deficiency during the chronic phase of the disease. The average size of granulomas in the Gal-3^{-/-}-infected mice was also significantly reduced at 14 wk p.i, relative to that in WT animals (Fig. 3A, right panel). As observed after 7 wk p.i., no difference in the proportion of eosinophils, monocytes/macrophages and T lymphocytes within the granulomas was noticed 14 wk p.i. (not shown). Finally, as visualized by Rojkind staining, the collagenous network was less compact in Gal-3 deficient mice relative to WT animals (Fig. 3B and C). As a whole, Gal-3^{-/-}-infected mice develop smaller lesions at 7 and 14 wk p.i and less severe hepatic fibrosis at 14 wk p.i. than do WT animals.

Gal-3 is functional in DCs but does not influence their differentiation or their maturation. Since DCs play a key role in orchestrating immune responses (4), we next investigated the possibility that Gal-3 deficiency in DCs could, at least in part, explain the dysregulated immune response observed in Gal-3^{-/-}-infected mice. No difference in the yield of DCs, either generated from WT or Gal-3^{-/-} BM progenitors was found, suggesting that the lack of Gal-3 has no major consequences on cell death/survival during the differentiation process. As seen in Fig. 4A (left panel), Western blot analysis confirmed the presence of Gal-3 protein in DCs generated from WT, but not Gal-3^{-/-}, BMs. Immunostaining revealed that Gal-3 is expressed on the surface of immature DCs (Fig. 4A, right panel). Of note, no

substantial changes in Gal-3 expression were observed in DCs matured with the canonical TLR4 agonist LPS (Fig. 4A, and not shown). We next investigated whether the lack of Gal-3 in DCs could impact on their phenotype and functions. FACS analysis revealed similar levels of mannose receptor, MHC class II, CD1d, CD54 and CD80 expression in immature WT and Gal-3^{-/-} DCs (not shown). To study the impact of Gal-3 deficiency on endocytosis, a key function of immature DCs, WT or Gal3^{-/-} BM-DCs were exposed to FITC-labelled dextran and internalization was followed by FACS analysis. As seen in Fig. 4B, Gal-3^{-/-} DCs internalized dextran (at 37°C) as efficiently as WT counterparts. To assess Gal-3-mediated endocytosis, FITC-labelled LDN-PAA and LN-PAA were utilized. FITC-labelled glucitol-PAA was used as a negative control. Compared to Gal-3 competent cells, Gal-3^{-/-} DCs failed to internalize LDN-PAA and LN-PAA (Fig. 4B). Furthermore, pre-treatment of WT DCs with lactose, a competitive inhibitor of Gal-3 binding, prevented LDN-PAA (Fig. 4C) and LN-PAA internalization (not shown). These data indicate that Gal-3 expression on DCs mediates endocytosis of canonical Gal-3 ligands. On the other hand, Gal-3 deficiency does not influence the differentiation, the phenotype or the Gal-3-independent endocytic functions of immature DCs.

Next, the ability of WT and Gal-3^{-/-} DCs to mature in response to different stimuli, including whole parasite eggs and canonical TLR agonists, was studied. Since schistosome eggs have been shown to induce DC maturation through TLR-2 and TLR-3 (1), a combination of Pam₃CSK₄ (a TLR2 agonist) and poly(I:C) (a TLR3 agonist) was used, although this binary activation does not recapitulate the total stimulatory effects of the eggs. The TLR4 agonist LPS was also included in this study. Whatever the mode of stimulation, and compared to WT cells, Gal-3^{-/-} DCs did not show any defect in their ability to up-regulate MHC class II, CD86 and CD40 (not shown). Similarly, as seen in Fig. 4D (left panel) (and data not shown), mature WT and Gal-3^{-/-} DCs produced equal amounts of the inflammatory cytokines TNF-α

and IL-6 as well as the chemokine IP-10/CXCL10 and MIG/CXCL9. Finally, we investigated the possibility that engagement of Gal-3 could modulate the maturation of DCs. To address this issue, WT DCs were pre-treated with LDN-PAA (or glucitol-PAA as a control) 30 min before Pam₃CSK₄/poly(I:C) or LPS stimulation. LDN-PAA did not modulate the production of inflammatory cytokines or chemokines (Fig. 4D, right panel and data not shown). In addition, LDN-PAA alone (without TLR agonists) did not trigger cytokine/chemokine production by DCs. Overall, Gal-3 deficiency in DCs or Gal-3 ligation does not cooperate nor interfere with TLR signalling pathways.

Expression of Gal-3 in DCs is necessary to control, but not to polarize, the cytokine response of T lymphocytes. Next, we studied the impact of Gal-3 deficiency in DCs on the strength and on the nature of the T cell response. To this end, we first compared the ability of Gal-3^{-/-} and WT DCs, either immature or mature, to induce an MLR response. For this, DCs stimulated or not (medium) with schistosome eggs, Pam₃CSK₄/poly(I:C) or LPS were co-cultured at different ratio with purified naïve T cells isolated from WT Balb/c mice. As expected, compared to immature DCs, mature DCs induced a higher T cell proliferation (Fig. 5A). Interestingly, relative to WT counterparts, mature Gal-3^{-/-} DCs induced an increased proliferation as well as an enhanced production of both IFN- γ and IL-4 by T cells (Fig. 5A) (as well as IL-5 and IL-10, not shown). Analysis of the IFN- γ /IL-4 ratio (not shown) revealed that Gal-3^{-/-} DCs do not skew the immune response in either a Th1 or Th2 direction.

To investigate whether Gal-3^{-/-} DCs prime T cells more efficiently than WT DCs *in vivo*, WT or Gal-3^{-/-} DCs were pulsed overnight with KLH, in the presence or absence of schistosome eggs, and were then injected i.v into naive syngenic WT mice. Seven days later, whole splenic cells were re-stimulated with KLH or egg Ag (52). Of note, a higher number of splenic T lymphocytes was detected in mice that received Gal-3 deficient DCs (~ 75%

enhancement compared to mice that received WT DCs), whilst no difference in T cell survival was observed (not shown). After restimulation, spleen cells from mice that received Gal-3^{-/-} DCs proliferated more vigorously (not shown) and produced an increased level of cytokines (Fig. 5B). For instance, after KLH re-stimulation, administration of Gal-3^{-/-} DCs (in particular those sensitized with parasite eggs) induced higher amounts of IFN- γ and IL-4 (as well as IL-5 and IL-10, not shown), compared to WT DCs. Finally, after SEA re-stimulation, Gal-3^{-/-} DCs cultured with whole eggs induced higher levels of both IFN- γ and IL-4 by T cells. Notably, although all groups of mice contained an identical frequency of IFN- γ positive T cells in the spleen (after CD3 or PMA/ionomycin re-stimulation), the mean fluorescent intensity of the IFN- γ staining tended to be higher in mice that received Gal-3 deficient DCs (not shown). Thus, it is likely that the increased production of cytokines in these mice is due to the increased number of T cells and to an enhanced cytokine production per cell number. To conclude, Gal-3 deficiency in DCs increases the T cell cytokine response both *in vitro* and *in vivo*, without biasing the immune response towards a Th1 or Th2 direction.

DISCUSSION

The role of Gal-3 in the Th1/Th2 immune and inflammatory responses appears to vary according to the experimental models used (administration of Gal-3 vs Gal-3 null animals) (10, 13, 21, 59). To further investigate its role in these settings, we studied Gal-3 deficient mice with regard to their response to *Schistosoma mansoni* infection, which is characterized by a Th2 polarized immune response and inflammation in the liver. Because DCs are extremely potent in dictating the outcome of innate and acquired immune responses, and since the functions of endogenous Gal-3 in DCs are presently unknown, we also attempted to determine its potential role in their maturation process and in their immunostimulatory and immunoregulatory properties.

Analysis of the parasitological parameters in infected WT and Gal-3^{-/-} mice indicated no variation in the number of adult worms nor in the number of eggs deposited in the tissues, confirming that parasites develop normally in infected Gal-3^{-/-} mice (7, 38). This is in contrast to the Gal-3 induced direct anti-microbial (fungicidal) activity recently described (30). Analysis of spleen cell composition and numbers revealed a dramatic reduction in the number (but not in the frequency) of B and T cells (but not of DCs) in infected Gal-3^{-/-} mice, compared to WT controls. This effect may be due to differences in cell recruitment or proliferation in this immunological site between the two animal groups. Gal-3 has indeed been shown to exert potent chemotactic activities on many immune cell types and to play a part in their survival (8, 16), although in our setting we found no difference in the rate of T cell survival between the different animal groups (not shown). Analysis of the immune responses during the acute phase of the disease (7 wk p.i.) clearly shows that during *S. mansoni* infection, Gal-3^{-/-} mice mount an enhanced Th1-biased response, manifested by a higher level of IFN- γ and a decreased ratio of *S. mansoni*-specific IgG1 to IgG2b in the sera

(Fig. 2). Interestingly enough, this enhanced Th1-biased response persisted in Gal-3^{-/-}-infected mice at a more chronic time point (14 wk p.i.) (not shown). These data are in accordance with a recent study showing that Gal-3 deficiency leads to the development of a heightened Th1 response (and a decreased inflammation) in mice infected with *Toxoplasma gondii* (6). These authors postulated that these effects are mediated, at least in part, by an enhanced IL-12 production by DCs. During murine schistosomiasis, IL-12 production in DCs is difficult to evaluate and we were unable to assess this parameter. Moreover, our data are also in line with those from Zuberi et al. (59) reporting that, in a murine model of asthma, Gal-3^{-/-} mice develop a higher Th1 response (and a diminished inflammation in the lungs) relative to WT animals. We next investigated whether Gal-3 deficiency could lead to modulation of liver inflammation in infected mice. Recently, Van de Vijver *et al.* showed that, when coupled to inert beads, LN- and LDN-containing glycoconjugates induce granuloma formation *in vivo*, with similar features to schistosome egg-induced granulomas (53). Consistent with this finding, and in agreement with Oliveira and co-workers (38), we here demonstrate that Gal-3 deficiency affects the size of liver granulomas but does not influence the proportion of immune cell types within the inflammatory foci. During the chronic phase of the disease, the liver of Gal-3 deficient mice were less damaged (not shown) and contain smaller lesions with a more diffuse network of collagen. Thus, endogenous Gal-3 plays a role in the pathology (lesion size and hepatic fibrosis) during murine schistosomiasis although whether this is due to endogenous and/or parasite-derived glycoconjugates is still unknown. Similarly, although we tend to favour the hypothesis that the reduced granuloma formation in Gal-3^{-/-} animals is rather due to a more general defect in immune/inflammatory cell recruitment in inflamed sites in these mice, it is also possible that the Th1 biased response developed in these mice might play a part in the pathology (56).

We next investigated whether the observed amplified Th1 response in infected Gal-3^{-/-} mice could be attributed to Gal-3 deficiency in DCs. The expression and the role of galectins in DCs have not been extensively studied and their functions have been mainly investigated by treating cells with exogenously added recombinant protein. For instance, exogenous addition of Gal-1 (18, 31) or Gal-9 (11) promotes DC maturation and increases their immunostimulatory functions (increased Th1 response). Although artificial Gal-1 overexpression in DCs also results in similar effects (40), no study has addressed the role of endogenous galectins in DCs, and whether or not they could exert activities on DCs, through a paracrine/autocrine manner, is unknown. Similarly, no studies have addressed the function of Gal-3, which is known to exert many antagonistic functions compared to Gal-1 and Gal-9 (32, 45), in DCs. Gal-3 has been shown to be expressed in both human and mouse DCs (14, 51, 55). Here, we show that Gal-3 is expressed on the surface of BM-DCs and that it mediates the internalization of LN- and LDN-conjugated molecules in immature cells. Phenotypic analysis indicates that immature WT and Gal-3^{-/-} BM-DCs display identical features, thus eliminating Gal-3 as a key factor in DC differentiation. This was confirmed *in vivo* by analyzing the number and the phenotype of splenic DCs in WT and Gal-3^{-/-} mice (Table II and not shown). Moreover, we showed that Gal-3 deficiency does not impact, in either a positive or a negative manner, on the TLR-induced maturation of DCs. Moreover, Gal-3 ligation in the absence of other stimulation did not induce DC maturation in terms of phenotype (not shown) and cytokine production (Fig. 4D), indicating that Gal-3 does not act as a danger signal in DCs. We next addressed the question of whether Gal-3 expression in DCs could modulate the priming and/or the differentiation of naïve T cells. Previous studies have shown that Gal-3 exerts potent effects on T cell proliferation (25), apoptosis and homeostasis (58). On the other hand, Gal-3 has also been reported to associate with specific N-glycans on the TCR complex to control TCR-mediated intracellular pathways in T lymphocytes (13, 35). This process

appears to maintain the development of immune responses, but whether this is mediated through Gal-3 expression by DCs or by other cells (including T cells themselves) is unknown. Here we show for the first time that Gal-3 expression in DCs is essential and sufficient to control the magnitude of T cell priming both *in vitro* and *in vivo*, since Gal-3^{-/-} DCs enhanced the proliferation and the cytokine release by T lymphocytes. *In vivo*, this effect was not due to an enhanced migration of DCs into the spleen or to an increased survival of T cells in this organ (not shown). Importantly, although it controls cytokine release by T cells, loss of Gal-3 expression by DCs does not modulate the Th1/Th2 balance of the immune response. This is in line with the non-essential role of Gal-3 in DC maturation (Fig. 4D), a complex process influencing the quality of the ensuing immune response. In our experimental infectious model, it is difficult to address the exact mechanisms behind the more pronounced Th1 bias observed in Gal-3 deficient animals. Although our data show that the lack of Gal-3 in DCs enhances the T cell response (Fig. 5), it cannot fully explain our observations (similar Th2 response in WT and Gal-3^{-/-} mice). Thus, it is likely that Gal-3 deficiency in other immune cell types is also important in this phenomenon.

Taken as a whole, this study reveals for the first time that Gal-3 expression in DCs plays part in the control of the T cell response but does not regulate the differentiation or maturation of DCs. This study also confirms, in an infectious model, that Gal-3 acts as an important regulator of immune and inflammatory responses.

ACKNOWLEDGMENTS

We gratefully acknowledge C. Vendeville for her outstanding technical assistance. We thank Dr R. Pierce (Inserm U547, Institut Pasteur de Lille) for the critical reading of the manuscript. This work was supported by the Institut National de la Santé et de la Recherche Médicale, the Pasteur Institute of Lille and the University of Lille 2. This project was also supported by the Conseil Régional Nord Pas de Calais / ARCir (Actions de Recherche Concertées d'initiative régionale) and FEDER (Fonds Européen de Développement Régional). LB was supported by a grant from the Ministère de l'Education Nationale de la Recherche et Technique and FV was the recipient of a doctoral fellowship from l'Ecole Normale Supérieure de Lyon. CV, PG TJ and CF are supported by the Inserm and FT by the Centre National de la Recherche Scientifique.

FIGURE LEGENDS

FIG. 1. Analysis of the cellular response in *S. mansoni*-infected WT and Gal-3^{-/-} mice. A, Proliferative response (A) and cytokine production (B) of spleen or MLN cells recovered from *S. mansoni*-infected WT or Gal-3^{-/-} mice, 7 wk p.i.. Cells were restimulated with increasing doses of SEA (shown are 2.5 and 10µg/ml) or were left unstimulated. Cytokine production and proliferation were measured after 3 days of culture. Results represent the mean of triplicate cultures ± SD (6 to 8 mice per group). One representative experiment out of two is shown. Significant differences are designated by ** (p<0.01), *** (p<0.005).

FIG. 2. Analysis of the humoral response in *S. mansoni*-infected WT and Gal-3^{-/-} mice. SEA-specific Ab isotype responses by *S. mansoni*-infected WT or Gal-3^{-/-} mice, 7 wk p.i.. Individual IgG1 and IgG2b titers and the IgG1/IgG2b ratio are shown. The mean titers ± SD (6 to 7 mice per group) are presented. One representative experiment out of two is shown. Significant differences are designated by ** (p<0.01), *** (p<0.005).

FIG. 3. Analysis of the granulomatous response in *S. mansoni*-infected WT and Gal-3^{-/-} mice. Rojkind staining of liver sections from *S. mansoni*-infected WT or Gal-3^{-/-} mice, 7 wk and 14 p.i.. A, The volume of the granulomas are indicated at wk 7 and wk 14 p.i. for each animal group. Significant differences are designated by ** (p<0.01) and *** (p<0.005). B, Representative staining obtained at 7 wk or 14 wk p.i. (x 100 magnification). C, The fibrosis, visualized on Sirius red-stained sections, was quantified using an arbitrary scale from 0 to 3. ***, p<0.005.

FIG. 4. A, Expression of Gal-3 in immature BM-DCs. Left panel, Western blot analysis of Gal-3 protein expression in DCs (5 x 10⁵ cells/lane). Right panel, gated CD11c⁺ DCs were

analyzed for Gal-3 expression (bold line). The isotype control is indicated (thin line). *B*, Endocytic function of Gal-3 in immature BM-DCs was assessed by flow cytometry. WT or Gal-3^{-/-} BM-DCs were incubated with FITC-labelled dextran, LDN-PAA, LN-PAA or glucitol-PAA for 2 hrs at 4°C (thin line) or 37°C (bold line). Endocytosis is displayed as an increase of fluorescence intensity at 37°C compared with 4°C. *C*, Lactose (5 mM) was added or not to WT DCs 30 minutes before adjunction of FITC-labelled LDN-PAA. *D*, Effects of Gal-3 deficiency on DC maturation (left panel). WT or Gal-3^{-/-} DCs were incubated overnight with schistosome eggs, a combination of Pam₃CSK₄ plus poly(I:C) or LPS. Right panel, LDN-PAA or glucitol-PAA (50µg/ml) was added to WT DCs 30 min before or not Pam₃CSK₄ plus poly(I:C) or LPS stimulation. After 18 hrs stimulation, quantification of secreted cytokines was performed by ELISA. The mean value ± SD are presented (4 experiments performed).

FIG. 5. Effects of Gal-3 deficiency in DCs on the priming of T cells. *A*, WT or Gal-3^{-/-} BM-DCs were incubated for 18 hrs with medium, schistosome eggs, a combination of Pam₃CSK₄ plus poly(I:C) or LPS. MLR was carried out by coculturing DCs with naïve T cells prepared from the spleens of Balb/c mice. Different DC:T cell ratios were tested. For clarity, SD values are not shown. One representative experiment out of three is shown. Significant differences are designated by * (p<0.05). *B*, WT or Gal-3^{-/-} BM-DCs were incubated overnight with KLH in the presence or not of schistosome eggs. Cells were then injected i.v. into syngenic mice and 7 days later, splenic cells were restimulated with KLH or SEA. Supernatants were harvested 72 hrs later for cytokine production. In the absence of restimulation, no proliferation or cytokine production was observed. Results represent the mean of triplicates cultures ± SD. One representative experiment out of two is shown. Significant differences are designated by * (p<0.05), ** (p<0.01), *** (p<0.005).

Key words: Rodent; Galectins; Dendritic Cells; Parasitic-Helminth; Th1/Th2 response

Abbreviations used in this paper: Gal-3, galectin-3 ; DCs, dendritic cells ; LN, LacNAc ; LDN, LacdiNAc ; WT, wild type ; PAA, polyacrylamide ; SEA, soluble egg antigen ; MLN, mesenteric lymph nodes ; BM-DCs, bone-marrow derived DCs ; KLH , Keyhole Limpet haemocyanin.

ACCEPTED

REFERENCES

1. Aksoy, E., C. S. Zouain, F. Vanhoutte, J. Fontaine, N. Pavelka, N. Thieblemont, F. Willems, P. Ricciardi-Castagnoli, M. Goldman, M. Capron, B. Ryffel, and F. Trottein. 2005. Double-stranded RNAs from the helminth parasite *Schistosoma* activate TLR3 in dendritic cells. *J Biol Chem* **280**:277-283.
2. Almkvist, J., and A. Karlsson. 2004. Galectins as inflammatory mediators. *Glycoconj J* **19**:575-581.
3. Asselin-Paturel, C., G. Brizard, J. J. Pin, F. Briere, and G. Trinchieri. 2003. Mouse strain differences in plasmacytoid dendritic cell frequency and function revealed by a novel monoclonal antibody. *J Immunol* **171**:6466-6477.
4. Banchereau, J., and R. M. Steinman. 1998. Dendritic cells and the control of immunity. *Nature* **392**:245-252.
5. Beatty, W. L., E. R. Rhoades, D. K. Hsu, F. T. Liu, and D. G. Russell. 2002. Association of a macrophage galactoside-binding protein with *Mycobacterium*-containing phagosomes. *Cell Microbiol* **4**:167-176.
6. Bernardes, E. S., N. M. Silva, L. P. Ruas, J. R. Mineo, A. M. Loyola, D. K. Hsu, F. T. Liu, R. Chammas, and M. C. Roque-Barreira. 2006. *Toxoplasma gondii* infection reveals a novel regulatory role for galectin-3 in the interface of innate and adaptive immunity. *Am J Pathol* **168**:1910-1920.
7. Bickle, Q., and H. Helmby. 2007. Lack of galectin-3 involvement in murine intestinal nematode and schistosome infection. *Parasite Immunol* **29**:93-100.
8. Chen, H. Y., F. T. Liu, and R. Y. Yang. 2005. Roles of galectin-3 in immune responses. *Arch Immunol Ther Exp (Warsz)* **53**:497-504.
9. Colnot, C., D. Fowles, M. A. Ripoche, I. Bouchaert, and F. Poirier. 1998. Embryonic implantation in galectin 1/galectin 3 double mutant mice. *Dev Dyn* **211**:306-313.
10. Colnot, C., M. A. Ripoche, G. Milon, X. Montagutelli, P. R. Crocker, and F. Poirier. 1998. Maintenance of granulocyte numbers during acute peritonitis is defective in galectin-3-null mutant mice. *Immunology* **94**:290-296.
11. Dai, S. Y., R. Nakagawa, A. Itoh, H. Murakami, Y. Kashio, H. Abe, S. Katoh, K. Kontani, M. Kihara, S. L. Zhang, T. Hata, T. Nakamura, A. Yamauchi, and M. Hirashima. 2005. Galectin-9 induces maturation of human monocyte-derived dendritic cells. *J Immunol* **175**:2974-2981.
12. del Pozo, V., M. Rojo, M. L. Rubio, I. Cortegano, B. Cardaba, S. Gallardo, M. Ortega, E. Civantos, E. Lopez, C. Martin-Mosquero, G. Peces-Barba, P. Palomino, N. Gonzalez-Mangado, and C. Lahoz. 2002. Gene therapy with galectin-3 inhibits bronchial obstruction and inflammation in antigen-challenged rats through interleukin-5 gene downregulation. *Am J Respir Crit Care Med* **166**:732-737.
13. Demetriou, M., M. Granovsky, S. Quaggin, and J. W. Dennis. 2001. Negative regulation of T-cell activation and autoimmunity by Mgat5 N-glycosylation. *Nature* **409**:733-739.
14. Dietz, A. B., P. A. Bulur, G. J. Knutson, R. Matasic, and S. Vuk-Pavlovic. 2000. Maturation of human monocyte-derived dendritic cells studied by microarray hybridization. *Biochem Biophys Res Commun* **275**:731-738.
15. Doenhoff, M., R. Musallam, J. Bain, and A. McGregor. 1978. Studies on the host-parasite relationship in *Schistosoma mansoni*-infected mice: the immunological dependence of parasite egg excretion. *Immunology* **35**:771-778.
16. Dunic, J., S. Dabelic, and M. Flogel. 2006. Galectin-3: an open-ended story. *Biochim Biophys Acta* **1760**:616-635.

17. **Faveeuw, C., V. Angeli, J. Fontaine, C. Maliszewski, A. Capron, L. Van Kaer, M. Moser, M. Capron, and F. Trottein.** 2002. Antigen presentation by CD1d contributes to the amplification of Th2 responses to *Schistosoma mansoni* glycoconjugates in mice. *J Immunol* **169**:906-912.
18. **Fulcher, J. A., S. T. Hashimi, E. L. Levroney, M. Pang, K. B. Gurney, L. G. Baum, and B. Lee.** 2006. Galectin-1-matured human monocyte-derived dendritic cells have enhanced migration through extracellular matrix. *J Immunol* **177**:216-226.
19. **Furtak, V., F. Hatcher, and J. Ochieng.** 2001. Galectin-3 mediates the endocytosis of beta-1 integrins by breast carcinoma cells. *Biochem Biophys Res Commun* **289**:845-850.
20. **Hirabayashi, J., T. Hashidate, Y. Arata, N. Nishi, T. Nakamura, M. Hirashima, T. Urashima, T. Oka, M. Futai, W. E. Muller, F. Yagi, and K. Kasai.** 2002. Oligosaccharide specificity of galectins: a search by frontal affinity chromatography. *Biochim Biophys Acta* **1572**:232-254.
21. **Hsu, D. K., R. Y. Yang, Z. Pan, L. Yu, D. R. Salomon, W. P. Fung-Leung, and F. T. Liu.** 2000. Targeted disruption of the galectin-3 gene results in attenuated peritoneal inflammatory responses. *Am J Pathol* **156**:1073-1083.
22. **Hughes, R. C.** 1999. Secretion of the galectin family of mammalian carbohydrate-binding proteins. *Biochim Biophys Acta* **1473**:172-185.
23. **Janeway, C. A., Jr., and R. Medzhitov.** 2002. Innate immune recognition. *Annu Rev Immunol* **20**:197-216.
24. **Jeng, K. C., L. G. Frigeri, and F. T. Liu.** 1994. An endogenous lectin, galectin-3 (epsilon BP/Mac-2), potentiates IL-1 production by human monocytes. *Immunol Lett* **42**:113-116.
25. **Joo, H. G., P. S. Goedegebuure, N. Sadanaga, M. Nagoshi, W. von Bernstorff, and T. J. Eberlein.** 2001. Expression and function of galectin-3, a beta-galactoside-binding protein in activated T lymphocytes. *J Leukoc Biol* **69**:555-564.
26. **Jouault, T., M. El Abed-El Behi, M. Martinez-Esparza, L. Breuilh, P. A. Trinel, M. Chamaillard, F. Trottein, and D. Poulain.** 2006. Specific Recognition of *Candida albicans* by Macrophages Requires Galectin-3 to Discriminate *Saccharomyces cerevisiae* and Needs Association with TLR2 for Signaling. *J Immunol* **177**:4679-4687.
27. **Karlsson, A., P. Follin, H. Leffler, and C. Dahlgren.** 1998. Galectin-3 activates the NADPH-oxidase in exudated but not peripheral blood neutrophils. *Blood* **91**:3430-3438.
28. **Kasai, K., and J. Hirabayashi.** 1996. Galectins: a family of animal lectins that decipher glycodes. *J Biochem (Tokyo)* **119**:1-8.
29. **Khoo, K. H., and A. Dell.** 2001. Glycoconjugates from parasitic helminths: structure diversity and immunobiological implications. *Adv Exp Med Biol* **491**:185-205.
30. **Kohatsu, L., D. K. Hsu, A. G. Jegalian, F. T. Liu, and L. G. Baum.** 2006. Galectin-3 Induces Death of *Candida* Species Expressing Specific beta-1,2-Linked Mannans. *J Immunol* **177**:4718-4726.
31. **Levroney, E. L., H. C. Aguilar, J. A. Fulcher, L. Kohatsu, K. E. Pace, M. Pang, K. B. Gurney, L. G. Baum, and B. Lee.** 2005. Novel innate immune functions for galectin-1: galectin-1 inhibits cell fusion by Nipah virus envelope glycoproteins and augments dendritic cell secretion of proinflammatory cytokines. *J Immunol* **175**:413-420.
32. **Liu, F. T.** 2000. Galectins: a new family of regulators of inflammation. *Clin Immunol* **97**:79-88.

33. **Liu, F. T., and G. A. Rabinovich.** 2005. Galectins as modulators of tumour progression. *Nat Rev Cancer* **5**:29-41.
34. **Lopez, E., V. del Pozo, T. Miguel, B. Sastre, C. Seoane, E. Civantos, E. Llanes, M. L. Baeza, P. Palomino, B. Cardaba, S. Gallardo, F. Manzarbeitia, J. M. Zubeldia, and C. Lahoz.** 2006. Inhibition of chronic airway inflammation and remodeling by galectin-3 gene therapy in a murine model. *J Immunol* **176**:1943-1950.
35. **Morgan, R., G. Gao, J. Pawling, J. W. Dennis, M. Demetriou, and B. Li.** 2004. N-acetylglucosaminyltransferase V (Mgat5)-mediated N-glycosylation negatively regulates Th1 cytokine production by T cells. *J Immunol* **173**:7200-7208.
36. **Ochieng, J., V. Furtak, and P. Lukyanov.** 2004. Extracellular functions of galectin-3. *Glycoconj J* **19**:527-535.
37. **Ochieng, J., M. L. Leite-Browning, and P. Warfield.** 1998. Regulation of cellular adhesion to extracellular matrix proteins by galectin-3. *Biochem Biophys Res Commun* **246**:788-791.
38. **Oliveira, F. L., P. Frazao, R. Chammas, D. K. Hsu, F. T. Liu, R. Borojevic, C. M. Takiya, and M. C. El-Cheikh.** 2007. Kinetics of mobilization and differentiation of lymphohematopoietic cells during experimental murine schistosomiasis in galectin-3-/- mice. *J Leukoc Biol.*
39. **Pearce, E. J., and A. S. MacDonald.** 2002. The immunobiology of schistosomiasis. *Nat Rev Immunol* **2**:499-511.
40. **Perone, M. J., A. T. Larregina, W. J. Shufesky, G. D. Papworth, M. L. Sullivan, A. F. Zahorchak, D. B. Stolz, L. G. Baum, S. C. Watkins, A. W. Thomson, and A. E. Morelli.** 2006. Transgenic galectin-1 induces maturation of dendritic cells that elicit contrasting responses in naive and activated T cells. *J Immunol* **176**:7207-7220.
41. **Rabinovich, G. A., L. G. Baum, N. Tinari, R. Paganelli, C. Natoli, F. T. Liu, and S. Iacobelli.** 2002. Galectins and their ligands: amplifiers, silencers or tuners of the inflammatory response? *Trends Immunol* **23**:313-320.
42. **Rabinovich, G. A., and A. Gruppi.** 2005. Galectins as immunoregulators during infectious processes: from microbial invasion to the resolution of the disease. *Parasite Immunol* **27**:103-114.
43. **Rabinovich, G. A., N. Rubinstein, and M. A. Toscano.** 2002. Role of galectins in inflammatory and immunomodulatory processes. *Biochim Biophys Acta* **1572**:274-284.
44. **Ramalho-Pinto, F. J., G. Gazzinelli, R. E. Howells, T. A. Mota-Santos, E. A. Figueiredo, and J. Pellegrino.** 1974. *Schistosoma mansoni*: defined system for stepwise transformation of cercaria to schistosomule in vitro. *Exp Parasitol* **36**:360-372.
45. **Rubinstein, N., J. M. Ilarregui, M. A. Toscano, and G. A. Rabinovich.** 2004. The role of galectins in the initiation, amplification and resolution of the inflammatory response. *Tissue Antigens* **64**:1-12.
46. **Sano, H., D. K. Hsu, J. R. Apgar, L. Yu, B. B. Sharma, I. Kuwabara, S. Izui, and F. T. Liu.** 2003. Critical role of galectin-3 in phagocytosis by macrophages. *J Clin Invest* **112**:389-397.
47. **Sano, H., D. K. Hsu, L. Yu, J. R. Apgar, I. Kuwabara, T. Yamanaka, M. Hirashima, and F. T. Liu.** 2000. Human galectin-3 is a novel chemoattractant for monocytes and macrophages. *J Immunol* **165**:2156-2164.
48. **Sato, S., and J. Nieminen.** 2004. Seeing strangers or announcing "danger": galectin-3 in two models of innate immunity. *Glycoconj J* **19**:583-591.

49. **Sato, S., N. Ouellet, I. Pelletier, M. Simard, A. Rancourt, and M. G. Bergeron.** 2002. Role of galectin-3 as an adhesion molecule for neutrophil extravasation during streptococcal pneumonia. *J Immunol* **168**:1813-1822.
50. **Srivatsan, J., D. F. Smith, and R. D. Cummings.** 1992. *Schistosoma mansoni* synthesizes novel biantennary Asn-linked oligosaccharides containing terminal beta-linked N-acetylgalactosamine. *Glycobiology* **2**:445-452.
51. **Thery, C., M. Boussac, P. Veron, P. Ricciardi-Castagnoli, G. Raposo, J. Garin, and S. Amigorena.** 2001. Proteomic analysis of dendritic cell-derived exosomes: a secreted subcellular compartment distinct from apoptotic vesicles. *J Immunol* **166**:7309-7318.
52. **Truong, M. J., V. Gruart, J. P. Kusnierz, J. P. Papin, S. Loiseau, A. Capron, and M. Capron.** 1993. Human neutrophils express immunoglobulin E (IgE)-binding proteins (Mac-2/epsilon BP) of the S-type lectin family: role in IgE-dependent activation. *J Exp Med* **177**:243-248.
53. **Van de Vijver, K. K., A. M. Deelder, W. Jacobs, E. A. Van Marck, and C. H. Hokke.** 2006. LacdiNAc- and LacNAc-containing glycans induce granulomas in an in vivo model for schistosome egg-induced hepatic granuloma formation. *Glycobiology* **16**:237-243.
54. **van den Berg, T. K., H. Honing, N. Franke, A. van Remoortere, W. E. Schiphorst, F. T. Liu, A. M. Deelder, R. D. Cummings, C. H. Hokke, and I. van Die.** 2004. LacdiNAc-glycans constitute a parasite pattern for galectin-3-mediated immune recognition. *J Immunol* **173**:1902-1907.
55. **Vray, B., I. Camby, V. Vercruysse, T. Mijatovic, N. V. Bovin, P. Ricciardi-Castagnoli, H. Kaltner, I. Salmon, H. J. Gabius, and R. Kiss.** 2004. Up-regulation of galectin-3 and its ligands by *Trypanosoma cruzi* infection with modulation of adhesion and migration of murine dendritic cells. *Glycobiology* **14**:647-657.
56. **Wynn, T. A., R. W. Thompson, A. W. Cheever, and M. M. Mentink-Kane.** 2004. Immunopathogenesis of schistosomiasis. *Immunol Rev* **201**:156-167.
57. **Yamaoka, A., I. Kuwabara, L. G. Frigeri, and F. T. Liu.** 1995. A human lectin, galectin-3 (epsilon bp/Mac-2), stimulates superoxide production by neutrophils. *J Immunol* **154**:3479-3487.
58. **Yang, R. Y., D. K. Hsu, and F. T. Liu.** 1996. Expression of galectin-3 modulates T-cell growth and apoptosis. *Proc Natl Acad Sci U S A* **93**:6737-6742.
59. **Zuberi, R. I., D. K. Hsu, O. Kalayci, H. Y. Chen, H. K. Sheldon, L. Yu, J. R. Apgar, T. Kawakami, C. M. Lilly, and F. T. Liu.** 2004. Critical role for galectin-3 in airway inflammation and bronchial hyperresponsiveness in a murine model of asthma. *Am J Pathol* **165**:2045-2053.

1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394

Table I. Analysis of the worm burden and of the total tissue eggs in WT and Gal-3^{-/-} infected mice 7 wk p.i. ^a

	WT	Gal-3 ^{-/-}
Worm Burden	17.1 ± 7.5	16.7 ± 2.8
Eggs/g of liver/female	776.8 ± 222.7	796.2 ± 126.1
Eggs/g of intestine/female	959.3 ± 309.0	1063.5 ± 484.6

^a Results represent the mean number ± SD (8 mice per group). One representative experiment out of two is shown.

1395

1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406

Table II. Analysis of the percentage and number of splenic cells in WT and Gal-3^{-/-} infected mice 7 wk p.i.^a

		uninfected		<i>S. mansoni</i> infected	
		WT	Gal-3 ^{-/-}	WT	Gal-3 ^{-/-}
CD3 ⁺ CD19 ⁺	%	39.56 ± 1.54	35.14 ± 4.25	39.75 ± 4.33	36.30 ± 2.51
	N	20.56 ± 8.19	21.79 ± 5.34	91.54 ± 7.36	57.29 ± 4.45 ***
CD3 ⁺ CD19 ⁻	%	45.04 ± 5.69	44.23 ± 4.61	30.44 ± 2.49	28.24 ± 3.70
	N	22.58 ± 6.49	27.42 ± 5.22	70.46 ± 8.17	44.48 ± 5.33 **
CD4 ⁺ CD25 ⁺	%	3.23 ± 0.58	2.95 ± 0.53	2.62 ± 0.31	2.21 ± 0.42
	N	1.73 ± 0.90	1.83 ± 0.72	6.03 ± 0.65	3.46 ± 0.21 ***
CD11c ⁺ 120G8 ⁻	%	0.82 ± 0.04	0.64 ± 0.23	1.80 ± 0.44	2.28 ± 0.4
	N	0.45 ± 0.12	0.40 ± 0.05	4.11 ± 0.78	3.58 ± 0.43
CD11c ⁺ 120G8 ⁺	%	0.83 ± 0.19	0.55 ± 0.31	0.24 ± 0.06	0.23 ± 0.08
	N	0.45 ± 0.03	0.34 ± 0.25	0.55 ± 0.13	0.37 ± 0.15

^a Results represent the mean percentage or number ± SD (4 mice per group). One representative experiment out of two is shown. N : number of cells × 10⁶/spleen. Significant differences between WT and Gal-3^{-/-} infected mice are represented by ** (p < 0.01) or *** (p < 0.005).

1407
1408

Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5

