

IgA MAb blocks SARS-CoV-2 Spike-ACE2 interaction providing mucosal immunity

Monir Ejemel^{#a}, Qi Li^{#a}, Shurong Hou^{b#}, Zachary A. Schiller^{a#}, Aaron L. Wallace^a, Alla Amcheslavsky^a, Nese Kurt Yilmaz^b, Jacqueline R. Toomey^a, Ryan Schneider^a, Brianna J. Close^c, Da-Yuan Chen^c, Hasahn L. Conway^c, Saeed Mohsan^c, Lisa A. Cavacini^{a*}, Mark S. Klempner^{a*}, Celia A. Schiffer^{b*}, Yang Wang^{a*}

^aMassBiologics of the University of Massachusetts Medical School, Boston, Massachusetts, USA

^bBiochemistry and Molecular Pharmacology, University of Massachusetts Medical School, Worcester, Massachusetts, USA

^cNational Emerging Infectious Diseases Laboratories, Boston University, Boston, Massachusetts, USA

[#]Co-first authors

^{*}Co-corresponding authors

17 **Summary:**

18 COVID-19 caused by SARS-CoV-2 has become a global pandemic requiring the development of
 19 interventions for the prevention or treatment to curtail mortality and morbidity. No vaccine to
 20 boost mucosal immunity or as a therapeutic has yet been developed to SARS-CoV-2. In this
 21 study we discover and characterize a cross-reactive human IgA monoclonal antibody,
 22 MAb362. MAb362 binds to both SARS-CoV and SARS-CoV-2 spike proteins and
 23 competitively blocks hACE2 receptor binding, by completely overlapping the hACE2 structural
 24 binding epitope. Furthermore, MAb362 IgA neutralizes both pseudotyped SARS-CoV and
 25 SARS-CoV-2 in human epithelial cells expressing hACE2. SARS-CoV-2 specific IgA
 26 antibodies, such as MAb362, may provide effective immunity against SARS-CoV-2 by inducing
 27 mucosal immunity within the respiratory system, a potentially critical feature of an effective
 28 vaccine.

Introduction:

In December 2019, a novel coronavirus (SARS-CoV-2) was identified as the cause of an outbreak of acute respiratory infections that emerged in Wuhan, China. The coronavirus disease 2019 (COVID-19) ranges from *mild* to *severe acute respiratory* infection, with a fatality rate estimated *to range* from 2 to 3%¹⁻⁴. Within three months of the first report cases, COVID-19 rapidly disseminated through the human population and had become a global pandemic by March 2020. Phylogenetic analysis has classified SARS-CoV-2 within the sarbecoviruses subgenus, the β lineage that also contains SARS-CoV, sharing proximately 79.6% sequence identity⁴.

Interventions for the prevention or treatment of COVID-19 are crucial for the ongoing outbreak. Pre- or post-exposure immunotherapies with neutralizing antibodies, would be of great use by providing immediate mucosal immunity against SARS-CoV-2. Although concerns, as occurred with SARS-CoV^{5,6}, that vaccines may cause disease enhancement will need to be addressed. The feasibility of human monoclonal antibodies (MAbs) as immunoprophylaxis or therapy against coronaviruses including SARS-CoV⁷⁻¹⁰ and MERS-CoV¹¹ has been demonstrated. These anti-coronavirus MAbs primarily target the viral spike (S) glycoprotein, a type I transmembrane glycoprotein that produces recognizable crown-like spike structures on the virus surface. The receptor-binding domain (RBD) of the S protein facilitates viral entry into human cells through human angiotensin-converting enzyme 2 (hACE2) receptor binding leveraging a similar mechanism as SARS-CoV¹²⁻¹⁴.

Most current anti-SARS-CoV MAbs neutralize virus by binding to epitopes on the spike protein RBD of SARS-CoV¹⁵. We and others have demonstrated that neutralizing MAbs that block RBD-hACE2 binding could confer potent protection against SARS-CoV as both prophylaxis and treatment in various animal models^{7,9,10}. Several anti-SARS-COV MAbs have demonstrated cross-neutralizing activities against the S protein of SARS-CoV-2^{16,17}. However, to date no antibody has directly bound to the hACE2 RBD interface of SARS-CoV-2 competitively blocking the SARS-CoV-2 spike:hACE2 complex.

Antibody-dependent enhancement of viral infections are major *hurdles* in the development of effective vaccines. This enhancement is likely facilitated by the Fc domain of IgG but not for its isotype variant IgA¹⁸. The avidity of mucosal IgA, in comparison with IgG, due to the multimeric structure, enhances the antibody binding with antigens. In addition, the diverse, high level of glycosylation of IgA antibodies, further protects the mucosal surface with non-specific interference. In animal models, high titers of mucosal IgA in the lung is correlated with reduced pathology upon viral challenge with SARS-CoV¹⁹. How precisely which isotype may protect the mucosa from SARS-CoV-2 infection remains an open question.

In the current study we describe the discovery of a cross-neutralizing human IgA monoclonal antibody, MAb362 IgA. This IgA antibody binds to SARS-CoV-2 RBD with high affinity directly competing at the hACE2 binding interface by blocking interactions with the receptor. MAb362 IgA neutralizes both pseudotyped SARS-CoV and SARS-CoV-2 in human epithelial cells expressing ACE2. Our results demonstrate that IgA isotype, plays a critical role in SARS-CoV-2 neutralization.

Results:

Selection of MAb binding to RBD of SARS-CoV-2 in ELISA

We have previously developed and characterized a panel of human MAbs that targets the RBD of the SARS-CoV S glycoprotein, isolated from transgenic mice expressing human immunoglobulin genes^{9,10}. These 36 hybridomas were recently screened against the SARS-CoV-2 Spike protein for potential cross-binding activity. MAb362 was identified with cross-binding activity against both the S1 subunit of the SARS-CoV S₁₋₅₉₀ and SARS-CoV-2 S₁₋₆₀₄ proteins (Extended Data Table 1).

While both IgG and IgA are expressed at the mucosa, IgA is more effective on a molar basis and thus the natural choice for mucosal passive immunization as we recently demonstrated in other mucosal infectious disease^{20,21}. To further characterize the functionality of MAb362, variable sequences of MAb362 were cloned into expression vectors as either IgG or IgA1 isotypes. Both MAb362 IgG and IgA were assessed in ELISA binding assays against the receptor-binding domain (RBD) of the S1 subunit for SARS-CoV S₂₇₀₋₅₁₀ and SARS-CoV-2 S₃₁₉₋₅₄₁ (Figure 1a). MAb362 IgA showed better binding activities, compared to its IgG counterpart against with SARS-CoV-2 S₃₁₉₋₅₄₁ (Figure 1b). Assessment of the binding kinetics was consistent with the ELISA binding trends, the binding affinity of IgA with RBD of SARS-CoV-2 is significantly higher (0.3 nM) than that of IgG (13 nM) due to a much slower dissociation rate as an IgA ($K_{off} = 1.13 \times 10^{-3} \pm 1.06 \times 10^{-4}$) compared to an IgG ($K_{off} = 7.75 \times 10^{-5} \pm 5.46 \times 10^{-5}$) (Figure 1c-f).

Structural modeling MAb362 binding to the core domain of RBD and competes for hACE2

binding

To define the antibody-binding epitope, known co-crystal and cryo-EM complexes from SARS-CoV and MERS spike protein in complex with neutralizing antibodies were evaluated for their potential to competitively block hACE2 binding, based on the structural interface of hACE2-SARS-CoV-2-RBD (PDB ID- 6VW1)²². The 80R-SARS-CoV-RBD complex (PDB ID- 2GHW)²³, a crystal structure of SARS-CoV RBD in complex with a neutralizing antibody, 80R was found most closely to have these characteristics. When the sequence was evaluated, we ascertained that the two antibodies, MAb362 and 80R had frameworks with *striking 90% amino acid sequence identity*. Thus, the crystal structure 2GHW provided an outstanding scaffold to build a highly accurate atomic homology model of MAb362. This structure permitted the modeling with the superposition of the hACE2:SARS-CoV-2-RBD (PDB ID-6VWI) for the modeling of MAb362:SARS-CoV-2-RBD (**Figure 2a**).

The interface between the MAb362:SARS-CoV-2-RBD complex is predicted to form extensive van der Waals contacts (**Figure 2b**). The CDRs of both the heavy and light chain make extensive interactions with SARS-CoV-2-RBD (**Figure 2c**), with the heavy chain of CDR-3 having the most extensive interaction. The binding interface of MAb362 is predicted to form 32 extensive contacts with residues on SARS-CoV-2-RBD (12 of which vary in sequence relative to SARS-CoV-RBD shown in red font) (**Figure 2d**). Seventeen of these contacts also are major points of contact between hACE2 on the SARS-CoV-2-RBD (**Figure 2e**). Thus MAb362 appears to directly compete for SARS-CoV-2 binding with hACE2.

Point mutations were engineered into the SARS-CoV-2-RBD based on this model and the overlap with the hACE2-RBD binding interface to further validate this model (**Figure 2f**). A combination of alanine and lysine mutations showed that charge mutations at the periphery (L455K) or outside the interface (N478K) had no impact, while sites that formed more extensive interactions Y449A, F456A and Y489A caused dramatic loss of binding affinity, only N501A retained affinity in a way that suggest a water mediated interaction may preserve this site. Both Y449 and Y489 are conserved with SARS-CoV while F456 is a Leucine (**Extended Data Figure 1-3**). Interestingly examination of the complex structure shows the close stacking of F456 against Y489 (**Figure 2g**) that together forming a combined extensive interface with light chain of MAb362, specifically the hydroxyl Y489 forms both extensive van der Waals interactions and hydrogen network at the interface. F456 forms less direct interactions, but structurally stabilizes the interactions of Y489, which explains the strong impact of the two alanine mutations F456A and Y489A. Y449A also forms extensive interactions (**Figures 2h**). Thus the loss of binding interaction from these site mutation validates our model of this complex as being biologically relevant complex

Complementing the mutational analysis, to correlate the epitope binding with functionality, MAb362 IgG and IgA were tested in a receptor-blocking assay with hACE2 expressing Vero E6 cells. The result suggested that both MAb362 IgG and IgA block SARS-CoV-2 RBD binding to receptors in a concentration dependent manner starting at ~ 4 nM (**Figure 2i, Extended Data Figure 4**). Thus confirming that the MAb362 epitope is directly competing for the hACE2 binding epitope on SARS-CoV-2 Spike.

MAb362 structural epitope

The epitope of MAb362 is in fact very different from the other recently reported MAb complexes to the SARS-CoV-2-RBD, such as CR3022¹⁶ or 309¹⁷. MAb362 overlaps entirely with the hACE2 epitope on the RBD (**Figure 3a**). This contrasts with CR3022 and 309 that bind to epitopes further away from the receptor-binding interface (**Figure 3b**). This finding was consistent with the unique activity of MAb362 of compromising RBD-receptor interaction. As with the binding of hACE2, the MAb362 binding epitope can only be exposed if the RBD was in the open or up conformation in the trimer (**Figure 3c**). In the closed conformation, this epitope would not be accessible to MAb362 without major steric clashes. However, unlike CR3022, MAb362 could access the hACE2 binding epitope(s) if one or more of the trimers is in this open conformation, potentially accounting for the added neutralizing activity.

MAb362 IgA1 neutralizes SARS-CoV and SARS-CoV-2 better than IgG

To evaluate the neutralization potency of cross-reactive MAb362, we performed a pseudovirus assay using lentiviral pseudovirions on 293 cells expressing hACE2 receptor²⁴. Both MAb362 IgG and IgA showed potent neutralization activity against SARS-CoV (**Figure 4a**). MAb362 IgG weakly neutralized SARS-CoV-2 pseudovirus despite its activities to block receptor binding. Interestingly, isotype switch to MAb362 IgA1 resulted in significantly enhanced neutralization potency, compared to its IgG subclass variant (**Figure 4b**). Monomeric MAb362 IgA1 was also co-expressed with J chain to produce dimeric IgA, which further improved neutralization (**Figure 4b**). This is consistent with our prior study showing isotype switch to IgA1 lead to improved antibody neutralization of HIV infection²⁵. Our data extends this observation to coronavirus, suggesting that IgA may play an important role in SARS-CoV-2 neutralization.

Discussion:

This study is the first report of a cross-reactive epitope within the core receptor-binding interface of the S protein of both SARS-CoV and SARS-CoV-2. MAb362 IgA neutralizes the virus via directly competing S protein binding to hACE2 receptors. Interestingly, our results show that despite the same blocking of spike interaction with hACE2, MAb362 IgG weakly neutralizes SARS-CoV-2 while its IgA1 isotype variant and its dimeric form showed significantly enhanced neutralization potency. Crystal structure studies demonstrated that IgA1 has a lengthy hinge region with a 13-a.a. insertion and a relaxed “T” like structure as compared to the more rigid “Y” like structure in IgG^{26,27}. Thus, the increase flexibility of IgA1 would likely afford a greater reach towards its epitopes on the target and decrease steric hindrance. MAb362 IgA binds when Spike protein (trimer) is in open form. The longer IgA1 hinge may allow two Fabs to reach two RBDs of the trimer at the same time without clashes, which may not be achieved by the shorter hinge in IgG. Our results suggest that compared to IgG, SARS-CoV-2-specific IgA antibody may play an important independent role in providing protective mucosal immunity.

Other recent structure studies have characterized antibodies targeting the RBD domain but distal from the receptor binding core interface of SARS-CoV-2, thus lack the characteristics of how MAb362 blocks the hACE2 binding epitope. Furthermore, these neutralizing IgGs, 47D11 and 309, neutralize SARS-CoV-2 with high potency, but do not block receptor binding to hACE2^{17,28}. Potentially hACE2 may not be the sole receptor for SARS-CoV-2, similar to SARS-CoV²⁹, or these antibodies may prevent a conformational change necessary for viral entry. Further study of the interaction between MAb362, and other receptor blocking and neutralizing antibodies against SARS-CoV-2 will provide insight into the design of vaccine and

prophylactic/therapeutic antibodies against future emerging infections caused by this viral family.

References:

- 1 Li, Q. *et al.* Early Transmission Dynamics in Wuhan, China, of Novel Coronavirus-Infected Pneumonia. *N Engl J Med*, doi:10.1056/NEJMoa2001316 (2020).
- 2 Jiang, S., Du, L. & Shi, Z. An emerging coronavirus causing pneumonia outbreak in Wuhan, China: calling for developing therapeutic and prophylactic strategies. *Emerg Microbes Infect* **9**, 275-277, doi:10.1080/22221751.2020.1723441 (2020).
- 3 Huang, C. *et al.* Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *The Lancet*, doi:10.1016/s0140-6736(20)30183-5 (2020).
- 4 Zhou, P. *et al.* A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature*, doi:10.1038/s41586-020-2012-7 (2020).
- 5 Czub, M., Weingartl, H., Czub, S., He, R. & Cao, J. Evaluation of modified vaccinia virus Ankara based recombinant SARS vaccine in ferrets. *Vaccine* **23**, 2273-2279, doi:10.1016/j.vaccine.2005.01.033 (2005).
- 6 Liu, L. *et al.* Anti-spike IgG causes severe acute lung injury by skewing macrophage responses during acute SARS-CoV infection. *JCI Insight* **4**, doi:10.1172/jci.insight.123158 (2019).
- 7 Zhu, Z. *et al.* Potent cross-reactive neutralization of SARS coronavirus isolates by human monoclonal antibodies. *Proc Natl Acad Sci U S A* **104**, 12123-12128, doi:10.1073/pnas.0701000104 (2007).
- 8 Sui, J. *et al.* Potent neutralization of severe acute respiratory syndrome (SARS) coronavirus by a human mAb to S1 protein that blocks receptor association. *Proc Natl Acad Sci U S A* **101**, 2536-2541, doi:10.1073/pnas.0307140101 (2004).
- 9 Greenough, T. C. *et al.* Development and characterization of a severe acute respiratory syndrome-associated coronavirus-neutralizing human monoclonal antibody that provides effective immunoprophylaxis in mice. *J Infect Dis* **191**, 507-514, doi:10.1086/427242 (2005).
- 10 Roberts, A. *et al.* Therapy with a severe acute respiratory syndrome-associated coronavirus-neutralizing human monoclonal antibody reduces disease severity and viral burden in golden Syrian hamsters. *J Infect Dis* **193**, 685-692, doi:10.1086/500143 (2006).
- 11 Du, L. *et al.* MERS-CoV spike protein: a key target for antivirals. *Expert Opin Ther Targets* **21**, 131-143, doi:10.1080/14728222.2017.1271415 (2017).
- 12 Letko, M. & Munster, V. Functional assessment of cell entry and receptor usage for lineage B betacoronaviruses, including 2019-nCoV. doi:10.1101/2020.01.22.915660 (2020).
- 13 Li, W. *et al.* Angiotensin-converting enzyme 2 is a functional receptor for the SARS coronavirus. *Nature* **426**, 450-454, doi:10.1038/nature02145 (2003).
- 14 Becker, M. M. *et al.* Synthetic recombinant bat SARS-like coronavirus is infectious in cultured cells and in mice. *Proc Natl Acad Sci U S A* **105**, 19944-19949, doi:10.1073/pnas.0808116105 (2008).
- 15 Jiang, S., Hillyer, C. & Du, L. Neutralizing Antibodies against SARS-CoV-2 and Other Human Coronaviruses. *Trends Immunol*, doi:10.1016/j.it.2020.03.007 (2020).
- 16 Yuan, M. *et al.* A highly conserved cryptic epitope in the receptor-binding domains of SARS-CoV-2 and SARS-CoV. *Science*, doi:10.1126/science.abb7269 (2020).

230 17 Pinto, D. *et al.* Structural and functional analysis of a potent sarbecovirus neutralizing antibody.
231 *bioRxiv*, doi:10.1101/2020.04.07.023903 (2020).

232 18 Bakema, J. E. & van Egmond, M. Immunoglobulin A: A next generation of therapeutic
233 antibodies? *MAbs* **3**, 352-361, doi:10.4161/mabs.3.4.16092 (2011).

234 19 Du, L. *et al.* The spike protein of SARS-CoV--a target for vaccine and therapeutic development.
235 *Nat Rev Microbiol* **7**, 226-236, doi:10.1038/nrmicro2090 (2009).

236 20 Stoppato, M. *et al.* Oral administration of an anti-CfaE secretory IgA antibody protects against
237 Enterotoxigenic Escherichia coli diarrheal disease in a nonhuman primate model. *Vaccine* **38**,
238 2333-2339, doi:10.1016/j.vaccine.2020.01.064 (2020).

239 21 Hu, Y. *et al.* Preformulation Characterization and Stability Assessments of Secretory IgA
240 Monoclonal Antibodies as Potential Candidates for Passive Immunization by Oral
241 Administration. *J Pharm Sci* **109**, 407-421, doi:10.1016/j.xphs.2019.07.018 (2020).

242 22 Shang, J. *et al.* Structural basis of receptor recognition by SARS-CoV-2. *Nature*,
243 doi:10.1038/s41586-020-2179-y (2020).

244 23 Hwang, W. C. *et al.* Structural basis of neutralization by a human anti-severe acute respiratory
245 syndrome spike protein antibody, 80R. *J Biol Chem* **281**, 34610-34616,
246 doi:10.1074/jbc.M603275200 (2006).

247 24 Ou, X. *et al.* Characterization of spike glycoprotein of SARS-CoV-2 on virus entry and its immune
248 cross-reactivity with SARS-CoV. *Nat Commun* **11**, 1620, doi:10.1038/s41467-020-15562-9 (2020).

249 25 Yu, X. *et al.* Impact of IgA constant domain on HIV-1 neutralizing function of monoclonal
250 antibody F425A1g8. *J Immunol* **190**, 205-210, doi:10.4049/jimmunol.1201469 (2013).

251 26 Boehm, M. K., Woof, J. M., Kerr, M. A. & Perkins, S. J. The Fab and Fc fragments of IgA1 exhibit a
252 different arrangement from that in IgG: a study by X-ray and neutron solution scattering and
253 homology modelling. *J Mol Biol* **286**, 1421-1447, doi:10.1006/jmbi.1998.2556 (1999).

254 27 Woof, J. M. & Burton, D. R. Human antibody-Fc receptor interactions illuminated by crystal
255 structures. *Nat Rev Immunol* **4**, 89-99, doi:10.1038/nri1266 (2004).

256 28 Wang, C. *et al.* A human monoclonal antibody blocking SARS-CoV-2 infection. *bioRxiv*,
257 doi:10.1101/2020.03.11.987958 (2020).

258 29 Jeffers, S. A. *et al.* CD209L (L-SIGN) is a receptor for severe acute respiratory syndrome
259 coronavirus. *Proc Natl Acad Sci U S A* **101**, 15748-15753, doi:10.1073/pnas.0403812101 (2004).

260 30 Walls, A. C. *et al.* Structure, Function, and Antigenicity of the SARS-CoV-2 Spike Glycoprotein.
261 *Cell*, doi:10.1016/j.cell.2020.02.058 (2020).

262 31 Wrapp, D. *et al.* Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation.
263 *Science* **367**, 1260-1263, doi:10.1126/science.abb2507 (2020).

264 32 Giuntini, S. *et al.* Identification and Characterization of Human Monoclonal Antibodies for
265 Immunoprophylaxis against Enterotoxigenic Escherichia coli Infection. *Infect Immun* **86**,
266 doi:10.1128/IAI.00355-18 (2018).

267 33 Wang, Y. *et al.* G glycoprotein amino acid residues required for human monoclonal antibody
268 RAB1 neutralization are conserved in rabies virus street isolates. *Antiviral Res* **91**, 187-194,
269 doi:10.1016/j.antiviral.2011.06.002 (2011).

270 34 Li, F., Li, W., Farzan, M. & Harrison, S. C. Structure of SARS coronavirus spike receptor-binding
271 domain complexed with receptor. *Science* **309**, 1864-1868, doi:10.1126/science.1116480 (2005).

272 35 Harder, E. *et al.* OPLS3: A Force Field Providing Broad Coverage of Drug-like Small Molecules and
273 Proteins. *J Chem Theory Comput* **12**, 281-296, doi:10.1021/acs.jctc.5b00864 (2016).

274 36 Leidner, F., Kurt Yilmaz, N., Paulsen, J., Muller, Y. A. & Schiffer, C. A. Hydration Structure and
275 Dynamics of Inhibitor-Bound HIV-1 Protease. *J Chem Theory Comput* **14**, 2784-2796,
276 doi:10.1021/acs.jctc.8b00097 (2018).

Acknowledgements:

We thank Dr. Julia Tree at Medical Interventions Group, National Infection Service at Public Health England for support and advice on antibody screen. We thank Dr. Robert Finberg at UMMS for helpful advice on functional assays. The initial mouse immunization work in 2005 was supported by NIH Contract NO1-AI-65315. S.H., N.Y.K. and C.A.S. were supported by NIH R01AI150478

Author Contributions:

M.E., Q.L. cloned, expressed and purified MAbs, S proteins, truncations, and variants with assistance from A.W., A.A., J.T., and R.S. M.E. and Q.L. designed and performed affinity and binding assays and flow cytometry with assistance from Y.W., L.A.C., and Z.A.S. S.H., carried out structural modeling and analyses with assistance from Q.L. N.Y.K. Y.W. and C.A.S. M.E. and Q.L. conducted pseudovirus neutralization with assistance from B.J.C., D-Y.C., H.L.C., S.M., L.A.C., and Y.W. Z.A.S. conducted data and statistical analysis with assistance from M.E., Q.L., and Y.W. Z.A.S. and Y.W. wrote the paper with assistance from M.E., Q.L., L.A.C., M.S.K., and C.A.S. M.S.K., L.A.C., and Y.W. supervised the project.

Competing interests

A patent application has been filed on 5 May 2020 on monoclonal antibodies targeting SARS-CoV-2 (U.S. Patent and Trademark Office patent application no. 63/020,483; patent applicants: Y.W., M.E., Q.L., and M.K., University of Massachusetts Medical School).




Figure 1. Binding of MAb362 IgG and IgA1 to the RBD of SARS-CoV and SARS-CoV-2

MAb362 IgG and IgA1 bind to purified RBD truncations of the S glycoprotein of SARS-CoV (S₂₇₀₋₅₁₀) (a) and SARS-CoV-2 (S₃₁₉₋₅₄₁) (b). Affinity measurements of MAb362 IgG (c, d) and IgA1 (e, f) against the RBD truncations of S glycoprotein of SARS-CoV (S₂₇₀₋₅₁₀) and SARS-CoV-2 (S₃₁₉₋₅₄₁) were conducted using bio-layer interferometry and demonstrate nano and sub-nanomolar affinities. Data is plotted as the average ± SD from two independent experiments.




Figure 2. Structural modeling MAb362 binding to the core domain of RBD and competes for hACE2 binding

MAb362 is predicted to have an extensive complementary interface with SARS-CoV-2 RBD that extensively overlaps with the hACE2 binding interface. **(a)** Surface representation of the complex, MAb362 in green, with SARS-CoV-2 RBD in violet. **(b)** The complex is separated and rotated by ~45° to highlight the extensive van der Waals buried surface areas on each protein, the spectrum of color coding represents the extent of the predicted van der Wall contact, with red being the most extensive contact and dark blue the least. **(c)** Detailed van der Waals of MAb362, residues from all the CDR's from both heavy and light chains pack against the SARS-CoV-2 RBD. **(d)** The binding interface on SARS-CoV-2 RBD with MAb362. **(e)** The binding

317 interface on SARS-CoV-2 RBD with hACE2. Residues names shown in red are those that differ
 318 in sequence between SARS-CoV and SARS-CoV-2.
 319
 320 **(f)** Binding affinity of wildtype and six SARS-CoV-2 RBD variants validated critical residues
 321 identified by SARS-CoV-2 RBD modeling. Data is plotted as the average $\pm$ SD from three
 322 independent experiments. **(g)** MAb362 IgG and IgA1 dose dependent reduction of S₁ truncation
 323 of SARS-CoV-2 (S₁₋₆₀₄) binding to hACE2 receptors on transfected Vero E6 cells. Data is
 324 plotted as the average $\pm$ SD from two independent experiments. **(h)** Interactions of SARS-CoV-2
 325 RBD (violet) Y489 and F456 (in pink) with MAb362 (green) dotted lines indicate potential
 326 hydrogen bonds. **(i)** Interactions of SARS-CoV-2 RBD (violet) Y449 (pink) with MAb362
 327 (green).




Figure 3. MAb362 structural epitope

(a) Superposition of the space filling structures of MAb362 (green) on the complex of hACE2 (orange) -SARS-CoV-2 RBD (violet) (**6VW1**) two views are rotated 180°. (b) Positioning of MAb362 on SARS-CoV-2 RBD (violet) relative to the binding of MAb CR3022¹⁶ (cream) or 309¹⁷ epitope (pink). (c) MAb362 modeled on the Spike trimer in open conformation with one RBD domain exposed **6VYB**³⁰. Note that the binding has no steric clashes with the other monomers.




Figure 4. IgA isotype switch enhances MAb362 neutralization of SARS-CoV-2

MAb362 antibody-mediated neutralization of luciferase-encoding pseudovirions with spike proteins of SARS-CoV (a) and SARS-CoV-2 (b). SARS-CoV and SARS-CoV-2 pseudovirions pre-incubated with fivefold dilutions of MAb362 (0.32 to 200 nM) were used to infect 293 cells expressing hACE2 receptor. Pseudoviral transduction was measured by luciferase activities in cell lysates 48 hrs post transduction to calculate neutralization (%) relative to non-antibody-treated controls. Data is plotted as the average $\pm$ SD from two independent experiments.

Materials and Methods:

S glycoprotein expression and purification

SARS-CoV and SARS-CoV-2 S glycoproteins were expressed and purified as previously described⁹. Briefly, the amino acid sequence of the SARS-CoV S glycoprotein (Urbani strain, National Center for Biotechnology Information [strain no. AAP13441]) and SARS-CoV-2 S glycoprotein sequence (GeneBank: MN908947) were used to design a codon-optimized version of the gene encoding the ectodomain of the S glycoproteins a.a. 1–1255 [S₁₋₁₁₉₀] for SARS-CoV and a.a. 1-1273 [S₁₋₁₂₅₅] for SARS-CoV-2, as described elsewhere³¹. The synthetic gene was cloned into pcDNA3.1 Myc/His in-frame with c-Myc and 6-histidine epitope tags that enabled detection and purification. Truncated soluble S glycoproteins were generated by polymerase chain reaction (PCR) amplification of the desired fragments from the vectors encoding S₁₂₅₅ and S₁₂₇₃. The SARS-CoV-2 RBD constructs carrying point mutation were generated by following the standard protocol from QuikChange® II XL Kit (Agilent). The cloned genes were sequenced to confirm that no errors had accumulated during the PCR process. All constructs were transfected into Expi293 cells using ExpiFectamine™ 293 Transfection Kit (Thermo Fisher). Antibodies were purified by immobilized metal chelate affinity chromatography using nickel-nitrilotriacetic acid (Ni-NTA) agarose beads. Antibodies were eluted from the columns using 250 mmol/L imidazole and then dialyzed into phosphate buffered saline (PBS), pH 7.2.

Generation of MAbs

Previously generated frozen hybridomas of anti-SARS-CoV MAbs⁹ were recovered and scaled up. Hybridoma supernatants were screened for reactivity to the SARS-CoV-2 S protein. Positive cell clones were selected for antibody sequencing. The heavy chain and light chain variable

regions were amplified from hybridoma cells and cloned into an immunoglobulin G1 (IgG) expression vector. Isotype switching was conducted using primers designed to amplify the variable heavy chain of the IgG antibody. Products were digested and ligated into a pcDNA 3.1 vector containing the heavy constant IgA1 chain. The vector was transformed in NEB5- α competent cells, and sequences were verified ahead of transient transfection. IgG and IgA1 antibodies were transfected in Expi293 cells and purified as previously described³². For dimeric IgA (dIgA), the heavy and light chain vectors were cotransfected with pcDNA-containing DNA for the connecting J chain. Purified antibodies were dialyzed against PBS and then concentrated and quality tested by SDS-PAGE.

ELISA

Dilutions of purified MAbs were tested in ELISA for reactivity against recombinant S protein. Briefly, 96-well plates were coated with S proteins followed by incubation overnight at 4°C. The plates were blocked with 1% BSA with 0.05% Tween 20 in PBS. Hybridoma supernatant or purified antibody diluted in 1× PBS plus 0.1% Tween 20 and added to the 96-well plates and incubated for 1 hour at room temperature. The plates were stained with horseradish peroxidase-conjugated anti-kappa (1:2,000) for 1 h and developed using 3,3',5,5'-tetramethylbenzidine. Absorbance at an optical density at 450 nm (OD₄₅₀) was measured on an Emax precision plate reader (Molecular Devices) using Softmax software.

Flow cytometry-based receptor binding inhibition assay

Vero E6 cells were harvested with PBS containing 5 mM EDTA and aliquoted to 1×10⁶ cells per reaction. Cells were pelleted then resuspended in PBS containing 10% FBS. Before mixing with

the cells, Myc-tagged SARS-CoV S₁₋₅₉₀ or SARS-CoV-2 S₁₋₆₀₄ was incubated with the MAb at varying concentrations for 1 hour at room temperature, then the S protein was added to the Vero cells to a final concentration of 10 nM. The cells-S protein mixture was incubated for 1 h at room temperature. After incubation, the cell pellets were washed and then resuspended in PBS with 2% FBS and incubated with 10 µg/mL of anti-Myc antibody for 1 hour at 4°C. Pellets were washed again then subsequently incubated with a Phycoerythrin-conjugated anti-mouse IgG (Jackson Immuno Research) for 40 minutes at 4°C. Cells were washed twice then subjected to flow cytometric analysis using a MACSquant Flow Cytometer (Miltenyi Biotec) and analyzed by FlowJo (version 10). Binding was expressed as relative to cells incubated with S proteins only.

Pseudotyped virus neutralization assay

Production of pseudotyped SARS-CoV and SARS-CoV-2 was performed as previously described³³. Pseudovirus was generated employing an HIV backbone that contained a mutation to prevent HIV envelope glycoprotein expression and a luciferase gene to direct luciferase expression in target cells (pNL4-3.Luc.R-E-, obtained from Dr. Nathaniel Landau, NIH). SARS-S and SARS2-S spike protein was provided in *trans* by co-transfection of 293T cells with pcDNA-G with pNL4-3.Luc.R-E-. Supernatant containing virus particles was harvested 48–72 h post-transfection, concentrated using Centricon 70 concentrators, aliquoted and stored frozen at -80 degree. Before assessing antibody neutralization, the 293T cells were transiently transfected with 100ng pcDNA-hACE2 each well in 96 well plates, and the cells were used for the pseudovirus infection 24 hours after transfection. A titration of pseudovirus was performed on 293T cells transiently transfected with hACE2 receptor to determine the volume of virus need to generate 50,000 counts per second (cps) in the infection assay. The appropriate volume of

pseudovirus was pre-incubated with varying concentrations of MAbs for 1 h at room temperature before adding to 293T cells expressing hACE2. 24 hours after the infection, the pseudovirus was replaced by the fresh complete media, and 24 hours after media changing the infection was quantified by luciferase detection with BrightGlo luciferase assay (Promega) and read in a Victor3 plate reader (Perkin Elmer) for light production.

Structural modeling and analyses.

Three crystal structures, 2GHW the complex of 80R:SARS-CoV-RBD²³, 2AJF the complex of hACE2:SARS-CoV-RBD³⁴ and 6VW1 the complex of hACE2:SARS-CoV-2-RBD²² were used as initial scaffolds in the determinations of the models of MAb362:SARS-CoV-RBD and MAb362:SARS-CoV-2-RBD. The amino acid sequence of MAb362 was aligned to the amino acid sequences of 80R. The molecular modeling of MAb362 was performed through the program Modeller 9.15 using the basic modeling and forming the initial MAb362:SARS-CoV RBD complex. This structure was further refined using iterative energy minimization by Desmond as previously described^{35,36}. The MAb362:SARS-CoV-2-RBD complex was made by replacing the SARS-CoV-2-RBD from 2AJF on the SARS-CoV-RBD structure and further optimized. The structural model of MAb362 binds to the SARS-CoV-2 Spike trimer was based on 6VYB³⁰. All figures were made within the PyMOL package. Hydrogen bonds were determined for pairs of eligible donor/acceptor atoms using criteria set by Schrodinger (Schrodinger, LLC, The PyMOL Molecular Graphics System, Version 1.3r1. 2010.). The residue van der Waals potential between the various complexes was extracted from the structures energies using the energy potential within Desmond. Epitope residues predicted by modeling were individually mutated using BioXp™ 3200 System (SGI-DNA). The genes were cloned into

RBD expression vectors and RBD proteins were purified as described above. Mutant RBD were confirmed intact expression on proteins gels, and the same amount of proteins were coated on the plate for ELISA assays. ELISAs assay was performed to determine binding of the MAbs to the mutant proteins compared to the wild-type.

Affinity determination

Bio-layer interferometry (BLI) with an Octet HTX (PALL/ForteBio) was used to determine the affinity of MAb362 IgG and IgA1 to the RBD of SARS-CoV and SARS-CoV-2 S protein. MAbs were added to 96 wells plates at 1000 nM and titrated 1:2 to 62 nM using PBS. RBD from SARS-CoV and SARS-CoV-2 were biotinylated (Thermo Fisher) and immobilized on Streptavidin (SA) Biosensors (ForteBio) for 120 seconds at 1600nM concentration. After a baseline step, MAb362-RBD binding rate was determined when the biosensors with immobilized RBD were exposed to MAb362 IgG or IgA1 at different concentrations for 120 seconds. Following association, the MAb362-RBD complex was exposed to PBS and the rate of the MAb362 dissociation from RBD was measured. Each assay was performed in triplicate. c

Statistical Analysis

Statistical calculations were performed using Prism version 7.03 (GraphPad Software, La Jolla, CA).

Clone	Epitope (A.A.)	SARS-CoV S1 S ₁₋₅₀₉	SARS-CoV RBD S ₂₇₀₋₅₁₀	SARS-CoV-2 S1 S ₁₋₆₀₄	SARS-CoV-2 RBD S ₃₁₉₋₅₄₁
7-508-395	90-190	>0.5	-	-	-
7-508-16	130-150	>0.5	-	-	-
7-508-39	130-150	>0.5	-	-	-
7-508-68	130-150	>0.5	-	-	-
7-508-104	130-150	>0.5	-	-	-
7-508-415	130-150	>0.5	-	-	-
7-508-466	370-510	>0.5	>0.5	-	-
12-28-1	490-510	>0.5	>0.5	-	-
7-73-121	490-510	>0.5	>0.5	-	-
7-508-201	490-510	>0.5	>0.5	-	-
7-508-669	490-510	>0.5	>0.5	-	-
7-508-362	N/A	>0.5	>0.5	>0.5	>0.5
7-508-94	N/A	>0.5	>0.5	-	-
7-508-411	N/A	>0.5	>0.5	-	-
7-508-478	N/A	>0.5	>0.5	-	-
7-508-568	N/A	>0.5	>0.5	-	-
7-509-2	N/A	>0.5	>0.5	-	-
7-508-73	N/A	>0.5	-	-	-
7-508-19	N/A	>0.5	-	-	-
7-508-165	N/A	>0.5	-	-	-
7-508-254	N/A	>0.5	-	-	-
7-508-223	N/A	>0.5	-	-	-
7-508-315	N/A	>0.5	-	-	-
7-508-528	N/A	>0.5	-	-	-
7-508-592	N/A	>0.5	-	-	-
7-512-9	N/A	>0.5	-	-	-
7-508-143	N/A	-	0.1 - 0.5	-	-
7-508-198	N/A	0.1 - 0.5	0.1 - 0.5	-	-
7-508-380	N/A	0.1 - 0.5	>0.5	-	-
12-28-4	N/A	-	-	-	-
7-73-57	N/A	-	-	-	-
7-508-448	N/A	-	-	-	-
7-508-573	N/A	-	-	-	-
7-508-646	N/A	-	-	-	-
7-508-680	N/A	-	-	-	-
7-508-694	N/A	-	-	-	-

Extended Data Table 1. Screening of a Panel of SARS-CoV MAbs for Cross-Binding

Activity

A panel of 36 previously generated frozen hybridomas of anti-SARS-CoV MAbs were recovered and scaled up. Hybridoma supernatants were screened for reactivity to the SARS-CoV-2 S

- 5 protein. Binding for each truncation is reported as absorbance at 450 nm for hybridoma
- 6 supernatant. Positive cell clones were selected for antibody sequencing.


RBD Core
Point mutation
F456 and Y489

SARS-CoV-1	384	ADSFVVKGDVVRQIAPGQTGVIADYNYKLPDDFMGCVLAWNTRNIDATST	433
		:: : . . : :: : :...	
SARS-CoV-2	397	ADSFVIRGDEVVRQIAPGQTGKIADYNYKLPDDFTGCVIAWNSNNLDSKVG	446
SARS-CoV-1	434	GNYNKYRYLRHGKLRPFERDISNVPFSPDGKPCT-PPALNCYWPLNDYG	482
		: ...:..... : ..	
SARS-CoV-2	447	GNYNLYRLERKSNLKPFERDISTEIQAGSTPCNGVEGFNCYEPLOSYG	496
SARS-CoV-1	483	FYTTTGIGYQPYRVVLSFELLNAPATVCGPKLSTDLIKNCVNFNFNGL	532
		.. . : : . : : : : :	
SARS-CoV-2	497	FQPTNGVGYQPYRVVLSFELLHAPATVCGPKKSTNLVKNKCVNFNFNGL	546

Extended Data Figure 1. Sequence alignment of SARS-CoV and SARS-CoV-2 S protein


RBDs

Sequence alignment show the identified critical residues (Y449, F456, and Y489) reside within the core domain of the SARS-CoV and SARS-CoV-2 S protein RBD.




Extended Data Figure 2. hACE2 - SARS-CoV-2 RBD complex.

MAb362 is predicted to have an extensive complementary interface with SARS-CoV-2 RBD that extensively overlaps with the hACE2 binding interface. **(a)** Surface representation of the complex, hACE2 in orange, with SARS-CoV-2 RBD in violet. **(b)** The complex is separated and rotated by ~45° to highlight the extensive van der Waals buried surface areas on each protein, the spectrum of color coding represents the extent of the predicted van der Waals contact, with red being the most extensive contact and dark blue the least. **(c)** Detailed van der Waals of hACE2. **(d)** The binding interface on SARS-CoV-2 RBD with hACE2. Residues names shown in red are those that differ in sequence between SARS-CoV and SARS-CoV-2.



Extended Figure 3. MAb362 - SARS-CoV RBD complex

MAb362 is predicted to have an extensive complementary interface with SARS-CoV RBD that extensively overlaps with the hACE2 binding interface. **(a)** Surface representation of the complex, MAb362 in green, with SARS-CoV RBD in violet. **(b)** The complex is separated and rotated by ~45° to highlight the extensive van der Waals buried surface areas on each protein, the spectrum of color coding represents the extent of the predicted van der Wall contact, with red being the most extensive contact and dark blue the least. **(c)** Detailed van der Waals of MAb362, residues from all the CDR's from both heavy and light chains pack against the SARS-CoV RBD. **(d)** The binding interface on SARS-CoV RBD with MAb362. **(e)** The binding interface on SARS-CoV RBD with hACE2. Residues names shown in red are those that differ in sequence between SARS-CoV and SARS-CoV-2.



Extended Data Figure 4. MAb362 blocks SARS-CoV and SARS-CoV-2 binding to hACE2 receptor

Flow cytometry of MAb362 IgG and IgA1 blocking binding of S₁ truncations of the SARS-CoV (S₁₋₅₉₀) and SARS-CoV-2 (S₁₋₆₀₄) binding to cell surface receptor hACE2 on Vero E6 cells.