

DR. JULIA FRICKE (Orcid ID : 0000-0002-7625-474X)

MS. THERESA KELLER (Orcid ID : 0000-0002-3603-534X)

DR. KARSTEN WELLER (Orcid ID : 0000-0003-4437-0313)

DR. SUSANNE LAU (Orcid ID : 0000-0002-5189-4265)

PROF. MARCUS MAURER (Orcid ID : 0000-0002-4121-481X)

PROF. TORSTEN ZUBERBIER (Orcid ID : 0000-0002-1466-8875)

Article type : Original Article: Epidemiology and Genetics

Prevalence of chronic urticaria in children and adults across the globe: systematic review with meta-analysis

Dr. med. Julia Fricke¹, MPH; Gabriela Ávila^{1,2}, MD, MSc; Dr. rer. medic. Theresa Keller^{1,3}, MSc; PD Dr. med. Karsten Weller⁴; Prof. Dr. med. Susanne Lau⁵; Prof. Dr. med. Marcus Maurer⁴; Prof. Dr. med. Torsten Zuberbier⁴; Prof. Dr. med. Thomas Keil^{1,6,7}, MSc

Affiliations:

¹Institute for Social Medicine, Epidemiology and Health Economics, Charité - Universitätsmedizin Berlin, Berlin, Germany

²Institute of Medical Informatics, Biometry and Epidemiology (IBE), Ludwig-Maximilians-Universität Munich, Germany

³Institute of Biometry and Clinical Epidemiology, Charité - Universitätsmedizin Berlin, Berlin, Germany

⁴Department of Dermatology and Allergy, Comprehensive Allergy Centre Charité, Charité - Universitätsmedizin Berlin, Germany

⁵Department for Pediatric Pneumology and Immunology, Charité - Universitätsmedizin Berlin, Berlin, Germany

⁶Institute of Clinical Epidemiology and Biometry, University of Wuerzburg, Wuerzburg, Germany

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/all.14037

This article is protected by copyright. All rights reserved.

⁷Institute for Health Resort Medicine and Health Promotion, Bavarian Health and Food Safety Authority, Bad Kissingen, Germany

Address of correspondence:

Dr. med. Julia Fricke, MPH

Institute for Social Medicine, Epidemiology and Health Economics,

Charité-Universitätsmedizin Berlin,

Luisenstr. 57, 10117 Berlin

Germany

Phone: +49-30450 529 015

julia.fricke@charite.de

Abstract:

Background and objectives: Urticaria is a frequent skin condition, but reliable prevalence estimates from population studies particularly of the chronic form are scarce. The objective of this study was to systematically evaluate and summarize the prevalence of chronic urticaria by evaluating population-based studies worldwide.

Methods: We performed a systematic search in PUBMED and EMBASE for population-based studies of cross-sectional or cohort design and studies based on health insurance/system databases. Risk of bias was assessed using a specific tool for prevalence studies. For meta-analysis we used a random effects model.

Results: 18 studies were included in the systematic evaluation and 11 in the meta-analysis including data from over 86,000,000 participants. Risk of bias was mainly moderate, whereas the statistical heterogeneity (I^2) between the studies was high. Asian studies combined showed a higher point prevalence of chronic urticaria (1.4%, 95%-CI 0.5-2.9) than those from Europe (0.5%, 0.2-1.0) and Northern American (0.1%, 0.1-0.1). Women were slightly more affected than men, whereas in children <15 years we did not find a sex-specific difference in the prevalence. The four studies that examined time trends indicated an increasing prevalence of chronic urticaria over time.

Conclusions: On a global level, the prevalence of chronic urticaria showed considerable regional differences. There is a need to obtain more sex-specific population-based and standardized international data particularly for children and adolescents, different chronic urticaria subtypes and potential risk and protective factors.

Key words:

Chronic urticaria; Prevalence; Systematic Review; Meta-analysis; Sex differences

Word count: 3538 (incl. Conflict of interest disclosure and Statement of contribution by each author)

Introduction

Urticaria is considered by clinicians as a relatively common skin condition, characterized by the development of wheals (hives), angioedema, or both [1]. By contrast there is a paucity of studies assessing urticaria prevalence and usually they do not distinguish between acute and chronic forms [2-5]. Epidemiological data is lacking especially for chronic urticaria (CU), defined as the recurrence of wheals, angioedema, or both for longer than 6 weeks [1].

Addressing previous inconsistencies, the updated EAACI/GA²LEN/EDF/WAO guideline for urticaria now clearly distinguishes two subtypes of CU: chronic spontaneous urticaria and chronic inducible urticaria – the latter including, for example, cold urticaria, cholinergic urticaria, and symptomatic dermographism [1].

CU carries a substantial burden not only for affected patients but also for health care systems. In most patients, CU markedly impairs quality of life with significant impact on sleep, work performance and social interactions [6-8]. In addition, patients with CU often present mental health problems [9-11]. Health care systems are facing high costs for managing patients with CU – including frequent health care visits, pharmacotherapy, absences from work, and loss of productivity [12, 13].

To the best of our knowledge, there is no comprehensive systematic review of population-based studies assessing the prevalence of CU worldwide. Therefore, the primary aim of this systematic review was to examine the prevalence of CU by assessing the evidence from population-based studies worldwide. As secondary outcome, the incidence of CU was investigated, if assessed within the same studies. As subgroup analyses we planned a stratification of prevalence estimates by age, sex and world region.

Material and methods

This systematic review is reported in accordance with Preferred Reporting Items for Systematic reviews and Meta-Analyses (PRISMA) guidelines [14]. The protocol of the present review was published on PROSPERO (https://www.crd.york.ac.uk/prospero/display_record.asp?ID=CRD42017073948), an international prospective register of systematic reviews.

Search strategy

We performed a systematic search in the major medical databases PUBMED and EMBASE. No date and language restrictions were applied. In addition to the electronic search, reference lists of selected articles were hand-searched. The results were managed using Endnote X7®. The search strategy is provided in the electronic supplement (table e1).

Eligibility criteria

The inclusion criteria were the following: (1) population-based studies of cross-sectional or cohort design, register studies or studies based on health insurance or physician databases, (2) studies in which prevalence and/or incidence data for CU can be extracted or calculated (self-reported/parent-reported or diagnosed by physician).

The following studies were excluded: (1) intervention studies, (2) case reports/series, (3) ecological studies, (4) hospital/out-patient studies, (5) studies with participants based on their occupations, and (6) studies with participants of only male or only female sex. Abstracts, editorials, notes, letters, and reviews were also not eligible for inclusion in the review.

Study selection

The selection of studies was conducted in a three-stage process by two independent reviewers (J.F., G.A.). First, each reviewer scanned the identified publications by title and classified each publication as “include”, “exclude” or “unclear”. Second, every publication categorized as “include” or “unclear” was reviewed using the abstract. Third, full text of publications rated as “include” or “unclear” were assessed according to the eligibility criteria. Any disagreements were discussed and if necessary referred to a third reviewer (T.Keil) for final decision.

Data was extracted independently by two reviewers (J.F., G.A.). Disagreements were solved by consensus. If necessary, a third reviewer was involved (T.Keller). The following data were extracted: continent, country, study design, sample size, database, response rate, observation period, age of participants, definition of disease, method of data collection, type of urticaria, prevalence and incidence data. In case data were missing, authors of the article were contacted by e-mail. Only data of authors who responded within two weeks were included in the meta-analysis. Additional data from authors was checked if it was in line with the original publication.

Risk of bias of individual studies was assessed by two reviewers (J.F., G.A.) using a tool that was designed for population-based prevalence studies and recently modified [15]. The tool includes 10 items (external validity: representation, sampling, random selection; internal validity: non-response bias, data collection, case definition, reliability/validity of tool, method of data collection, prevalence period, numerator(s) and denominator(s)) and a summary assessment (low risk, moderate risk and high risk of bias). The scope of the tool was to identify whether studies had attempted to minimize bias and not to assess an overall numeric rating of risk of bias. The modified tool has shown to have a high interrater agreement and had been adopted previously in several studies [16, 17]. Disagreements were resolved by consensus. Studies rated with a high risk of bias were not included in the meta-analysis.

Quantitative synthesis

We extracted the prevalence of CU for each study and then calculated the pooled prevalence estimates using the arcsine transformation for variance stabilization [18] with a 95% confidence interval using random effects meta-analysis due to the expected heterogeneity of the studies. For subgroup analyses we planned a stratification by world region, sex and age. I^2 was calculated to quantify this heterogeneity. Statistical analyses were done with R (R Foundation for Statistical Computing, Vienna, Austria).

Results

Characteristics of included studies

A total of 4,844 records were identified, and finally full texts of 55 records were assessed for eligibility. 18 studies were included in the systematic evaluation and 11 in the meta-analysis. Selection process is shown in figure 1.

Six of the 18 studies were carried out in Europe [19-24], two in Latin America [25, 26], three in Northern America [27-29], six in Asia [30-35] and one in Africa [36]. 16 studies were published after 2003. Five studies comprised only adults and three only children. Ten were cross-sectional studies of primary data and eight based on cross-sectional analyses of secondary data (health insurance or national health system database). Seven studies used self-reported data, ten studies data based on physician diagnosis and one used a 3 step assessment (self-reporting questionnaire, than telephone interview, than examination with physician diagnosis). Data on incidence was only available from three studies, but these were too heterogeneous for meta-analysis. Two authors sent additional information and data of one author was included. A detailed summary of each study is included in table e2 and table e3.

Most studies were rated as having a moderate risk of bias. Only Hellgren 1972 [19] had a high risk of bias, and this study was not included in the meta-analysis. It should be noted that for secondary health analysis studies or survey panels response rate was not specified. Risk of bias was found most often for the external validity items representation and sampling frame. Within the internal validity section risk of bias was more often found for the appropriateness of numerators/denominators. In case calculation or reporting of numerators/denominators contained errors, we re-calculated them or, if this was not possible, excluded them from the meta-analysis. Another concern was the validity/reliability of the study instrument. It should be pointed out that validated instruments, like the widely used ISAAC questions for assessing asthma, rhinitis and atopic eczema on a population level [2], are lacking for the assessment of CU in population-based studies.

Four studies [23, 24, 31, 34] provided prevalence estimates, but the published information was incomplete and therefore not included in the meta-analysis. Furthermore, El-Khateeb et al 2014 [36] was included in the evaluation but excluded from the meta-analysis because it assessed only the one-day prevalence. Similarly, Ohmi et al 1984 [30] was excluded because it assessed only the prevalence of one subtype of chronic inducible urticaria [37]. Finally, pooled estimates of prevalence were determined from 11 studies. In total, data from n=86,632,267 persons was considered for

meta-analysis (study samples: 2,613 [minimum]-50,316,384 [maximum]). Due to the expected heterogeneity of the studies a random effects model was chosen for meta-analysis instead of a fixed effects model. In a random effects model the weights of the single studies are more balanced and the size of the individual boxes for each study in the forest plots are therefore not substantially different.

Lifetime prevalence of chronic urticaria

Five studies [20, 21, 25, 26, 29] reported the overall lifetime prevalence of CU, which was on average 4.4% (95%CI 1.6-8.4). Excluding one study with an unusually high lifetime prevalence [25] in relation to the other studies, the adjusted lifetime prevalence was 1.4% (95%CI 0.8-2.2, $I^2=98.9\%$, figure 2).

Point prevalence of chronic urticaria

Based on eleven studies [20-22, 25-29, 32-34] the overall point prevalence of CU was 0.7% (95%CI 0.2-1.4, $I^2=100\%$) (figure 3). Most studies presented 12-month-prevalence estimates, however three studies assessed the prevalence of current treatment for CU [22, 26, 29] and one study the combination of one-week-prevalence plus current treatment [20].

Point prevalence of chronic urticaria by world region, sex and age

The regions with the highest point prevalence estimates were Latin America and Asia: 1.5% (95%CI 0.0-6.0) and 1.4% (95%CI 0.5-2.9) respectively. The region with the lowest prevalence was Northern America: 0.1%; 95%CI 0.1-0.1 (figure 4).

Seven studies [20, 21, 25, 26, 28, 32, 33] presented data stratified by sex. The point prevalence estimate for women was slightly higher than for men with overlapping corresponding 95%-CIs based on random effect meta-analysis : 1.3% (95%CI 0.1-2.2) versus 0.8% (95%CI 0.2-3.2) (figure 5).

Stratifying by world regions we found that this difference was only statistically significant in Europe and Northern America whereas in the Asian studies we did not detect such sex-specific difference in the prevalence of CU (figure e1-e3).

Nine studies [20-22, 25, 26, 28, 29, 32, 33] presented data stratified by age. Only one of the studies included in meta-analysis assessed only data on children (4-13 years) [32]. Based on the available studies, we decided to calculate prevalence estimates for children (0-19 years) and adults (> 19 years). For this age-specific analysis we excluded one study that presented age strata, which did not correspond with the age strata of the other studies (≤ 11 years, 12-24 years, ≥ 25 years [28]). The summary point prevalence estimate for children was slightly higher than for adults (figure 6).

Looking at sex-specific differences in children < 15 years [28, 32, 33] a subgroup analysis yielded a point prevalence of 1.0% (95%CI 0.0-3.4) for girls and of 1.1% (95%CI 0.0-3.9) for boys (figure e4).

Additional analysis

Studies assessing point prevalence based on self-reported questionnaires (cross-sectional studies) (0.7%, 95%CI 0.4-1.0) yielded similar prevalence estimates as studies assessing point prevalence based on physician diagnosis (secondary data) (0.6%, 95% CI 0.0-1.8) (figure e5).

Regarding possible time trends we identified four studies from the systematic review that assessed the point prevalence at different time points in the same region with the same methods. All four studies [23, 24, 33, 34] showed an increasing point prevalence over time (table e3).

Discussion

Main findings

Our systematic review with meta-analysis showed that CU affects a considerable part of the population around the globe with overall lifetime and point prevalence rates of 1.4% and 0.7%, respectively. CU seems to be more prevalent in Asia than in Europe and Northern America. Regarding sex-specific analyses, women seemed to be more affected than men. In children < 15 years we did not find a sex-specific difference. Varying time trend evaluations showed an increasing prevalence of CU in recent years.

Comparison with other studies

Based on four studies only, the overall lifetime prevalence of 1.4% needs to be interpreted with caution. It may be an underestimation because several studies did not ask specifically for symptoms of chronic inducible urticaria and participants may not be aware of having it because symptoms can be mild. In a selected sample of students, Zuberbier et al. assessed a prevalence of 11.2% for cholinergic urticaria after presenting urticaria-specific symptoms in slides to them [38]. On the other hand, a recently published study from Poland with similar methods yielded a lifetime prevalence of CU of only 0.6% [39].

The overall point prevalence of less than 1% may be also an underestimation because the definition of CU in several of the included studies required current treatment, but not all CU patients are receiving treatment continuously. In fact, only less than 50% of patients seem to be responsive to the first-line treatment with antihistamines [37]. Another reason may be that three studies [27, 28, 34] assessed only chronic spontaneous urticaria.

Latin America and Asia showed a higher point prevalence of CU than other regions. However the estimate for Latin America, based on two studies only, yielded a very wide confidence interval. The study from Mexico [25] assessed with 3.4% a specifically higher point prevalence in relation to the other studies. The authors of this study pointed out that their study was undertaken in a region of Mexico that is according to their assessment predestined for allergic diseases due to climatic reasons [25]. Opposite to the data from Mexico, a recently published study from Argentina yielded a point prevalence of only 0.29% among a highly selective sample in Buenos Aires, i.e. members of a private health maintenance organization [40]. Northern America showed the lowest point prevalence. This may be explainable by the fact that two of the three studies – covering 98.5% of the Northern America sample - assessed only chronic spontaneous urticaria and not chronic inducible urticaria. More population-based studies from Latin and Northern America assessing both chronic spontaneous and chronic inducible urticaria are needed to further elucidate the prevalence patterns.

The point prevalence estimate for women was slightly higher than for men, confirming results from studies in samples of patients with chronic spontaneous urticaria [41]. Interestingly, one study from Asia found no sex-specific difference and it was the study including only children (4-13 years) [32]. Evaluating sex-specific differences in children < 15 years in three studies [28, 32, 33] we found again no sex-specific difference. A sex-related prevalence shift from childhood to adulthood has been identified for allergies, like allergic rhinitis in adolescence [42, 43] and asthma after the onset of puberty [44, 45]. Maybe the sex-specific difference in CU that has been shown for patients [41] results because in the transition to adulthood more females are developing urticaria? A recent study from South Korea assessed a predominance of women for new-onset urticaria only for the age group 20–64 years [46], being a possible explanation for the preponderance of female patients with CU in this group.

The point prevalence estimate of CU for children was slightly higher than for adults, although the limited number of childhood studies is hampering valid comparisons. A recently published study among physicians from five European countries yielded a point prevalence of 1.4% among pediatric patients [47], which is within the range of the population-based prevalences for children in our meta-analysis. These results indicate that more studies are needed to validly assess prevalence data in children, preferably differentiating between childhood, adolescence and early adulthood.

Prevalence rates of CU appear to be increasing. This should be confirmed and characterized in more detail in future longitudinal studies.

Strengths and Limitations

To the best of our knowledge the current review has been the first to determine systematically the prevalence of CU worldwide. Our meta-analysis included large datasets with a total of more than 86,000,000 participants of population-based studies worldwide. We searched the two major medical databases, PUBMED and EMBASE, which cover most of the available medical literature. Although we did not restrict our search to specific languages, most publications were in English. We cannot completely rule out that we may have missed studies, however, to minimize such bias we conducted manual searches in addition to the systematic search.

Unfortunately not all world regions were represented in this present evaluation. For Oceania, no study was identified and the only study from Africa had to be excluded from meta-analysis, but was included in the systematic review. Three of the most recent studies included in meta-analysis were from Asia, adding new evidence to the epidemiology of CU worldwide.

For some of our planned subgroup analyses there were only few studies eligible hampering comparisons. This concerns especially the estimates for overall lifetime prevalence and point prevalence for children.

There was a great difference in sample size between cross-sectional studies and studies assessing complete health insurance/national health system databases. This may be one reason for the considerable heterogeneity in all meta-analyses as indicated by the Higgins' I^2 tests. By applying the random effects model, studies with larger samples had a smaller and studies with smaller samples had a bigger effect. Another reason for heterogeneity may be having taking into account different data collection modes like cross sectional studies using self-reported questionnaires as well as

secondary data analyses assessing physician diagnoses. More possible reasons are the inclusion of different regions and age strata.

Risk of bias assessment showed that the included studies were mainly rated as moderate risk of bias. One concern was the validity/reliability of the instruments used in the studies. Interestingly, the point prevalence did not considerably differ when comparing studies using physician diagnosis with studies administering self-reported questionnaires. So even without including a validated questionnaire for the assessment of CU, the questionnaire-based studies did not seem to over- or underestimate notably the CU prevalence.

Most studies did not distinguish between relevant CU subtypes. Future studies should address this issue and aim to assess both types of CU separately. Only a few studies assessed incidence data, which was insufficient to conduct a meta-analysis. Future cohort studies should start early in life and try to collect incidence data from childhood onwards to get a broader picture of the development of this disease and assess potential predictors, risk and protective factors. Another potential determinant to be investigated may be urban and rural setting as rural residence has been suggested as a risk factor for urticaria [46].

Conclusions

This first systematic review on CU prevalence showed considerable regional differences. CU seems to be more prevalent in Asia than in Europe and Northern America. Women seemed to be more affected than men, whereas in children < 15 years there was no sex-specific difference in the prevalence of CU. Temporal analysis showed an increasing prevalence over time. Our quality assessment showed that the risk of bias was mainly moderate, whereas the statistical heterogeneity (I^2) between the studies included in the meta-analysis was rather high.

There is a need to conduct further properly performed population-based studies on the prevalence of CU, especially regarding certain age groups, sex-specific differences and regions. Further research should focus specifically on children and adolescents and different CU subtypes. Prospective investigations are required to examine the incidence and potential protective and risk factors of CU in order to develop preventive strategies. The need for global studies may be facilitated by the global network of urticaria centers of reference and excellence (UCARE) [48].

Conflict of interest:

J. Fricke, G. Ávila, T. Keller, T. Keil have nothing to disclose.

K. Weller reports personal fees from Novartis, personal fees from MOXIE, personal fees from Uriach, personal fees from UCB, personal fees from Dr. Pfleger, outside the submitted work; .

S. Lau reports personal fees from Sanofi Genzyme, personal fees from DBV, personal fees from ALK, personal fees from Allergopharma, outside the submitted work; .

Dr. Maurer reports grants and personal fees from Allakos, personal fees from Aralez, grants from AstraZeneca, grants and personal fees from FAES, grants and personal fees from Genentech, grants and personal fees from Menarini, grants and personal fees from Novartis, grants and personal fees from Moxie, grants and personal fees from MSD, grants and personal fees from Roche, grants and personal fees from Sanofi, grants and personal fees from UCB, grants and personal fees from Uriach, outside the submitted work; .

T. Zuberbier reports personal fees from Bayer Health Care, personal fees from FAES, personal fees from Novartis, personal fees from Henkel, from Novartis, from Henkel, personal fees from AstraZeneca Fee for talk, personal fees from AbbVie Fee for talk, personal fees from ALK Fee for talk, personal fees from Almirall Fee for talk, personal fees from Astellas Fee for talk, personal fees from Bayer Health Care Fee for talk, personal fees from Bencard Fee for talk, personal fees from Berlin Chemie Fee for talk, personal fees from FAES Fee for talk, personal fees from HAL Fee for talk, personal fees from Leti Fee for talk, personal fees from Meda Fee for talk, personal fees from Menarini Fee for talk, personal fees from Merck Fee for talk, personal fees from MSD Fee for talk, personal fees from Novartis Fee for talk, personal fees from Pfizer Fee for talk, personal fees from Sanofi Fee for talk, personal fees from Stallergenes Fee for talk, personal fees from Takeda Fee for talk, personal fees from Teva Fee for talk, personal fees from UCB Fee for talk, personal fees from Henkel Fee for talk, personal fees from Kryolan Fee for talk, personal fees from L'Oréal Fee for talk, outside the submitted work; .

Statement of contribution by each author:

All the authors conceived and designed the study project. J. Fricke and G. Ávila performed the literature search, assessed study details and evaluated the study quality supported by T. Keil and T. Keller. T. Keller performed the statistical analyses. J. Fricke wrote the first draft of the paper with the support of G. Ávila, which was critically revised by all the other authors. All the authors gave final approval of the version to be submitted and agreed to be accountable for the whole paper.

References

1. Zuberbier T, Aberer W, Asero R et al. The EAACI/GA²LEN/EDF/WAO Guideline for the Definition, Classification, Diagnosis and Management of Urticaria. The 2017 Revision and Update . Allergy 2018 Jul;73(7):1393-1414
2. Bakke P, Gulsvik A, Eide GE. Hay fever, eczema and urticaria in southwest Norway. Lifetime prevalences and association with sex, age, smoking habits, occupational airborne exposures and respiratory symptoms. Allergy. 1990 Oct;45(7):515-22.
3. Popescu R, Popescu CM, Williams HC, Forsea D. The prevalence of skin conditions in Romanian school children. Br J Dermatol. 1999 May;140(5):891-6.
4. Campos A, Alamar R, Castelló JV, Enguñados, M. Proyecto orba: 1983-1993. Changes in the prevalence of asthma, rhinitis, urticaria and drug allergy in Valencia (Spanish). Alergol Inmunol Clin 2002; 17: 151-158

5. Brüske I, Standl M, Weidinger S et al. Epidemiology of urticaria in infants and young children in Germany--results from the German LISApplus and GINIplus Birth Cohort Studies. *Pediatr Allergy Immunol*. 2014 Feb;25(1):36-42.
6. O'Donnell BF, Lawlor F, Simpson J, Morgan M, Greaves MW. The impact of chronic urticaria on the quality of life. *Br J Dermatol* 1997;136(2):197-201.
7. Staubach P, Eckhardt-Henn A, Dechene M et al. Quality of life in patients with chronic urticaria is differentially impaired and determined by psychiatric comorbidity. *Br J Dermatol* 2006;154(2):294-8.
8. Dias GA, Pires GV, Valle SO et al. Impact of chronic urticaria on the quality of life of patients followed up at a university hospital. *An Bras Dermatol* 2016;91(6):754-9.
9. Ozkan M, Oflaz SB, Kocaman N et al. Psychiatric morbidity and quality of life in patients with chronic idiopathic urticaria. *Ann Allergy Asthma Immunol* 2007;99(1):29-33.
10. Engin B, Uguz F, Yilmaz E, Ozdemir M, Mevlitoglu I. The levels of depression, anxiety and quality of life in patients with chronic idiopathic urticaria. *J Eur Acad Dermatol Venereol* 2008;22(1):36-40.
11. Chung MC, Symons C, Gilliam J, Kaminski ER. Stress, psychiatric co-morbidity and coping in patients with chronic idiopathic urticaria. *Psychol Health* 2010;25(4):477-90.
12. DeLong LK, Culler SD, Saini SS, Beck LA, Chen SC. Annual direct and indirect health care costs of chronic idiopathic urticaria: a cost analysis of 50 nonimmunosuppressed patients. *Arch Dermatol* 2008;144(1):35-9.
13. Mendelson MH, Bernstein JA, Gabriel S et al. Patient-reported impact of chronic urticaria compared with psoriasis in the United States. *J Dermatolog Treat* 2017;28(3):229-36.
14. Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS medicine* 2009;6(7):e1000097.
15. Hoy D, Brooks P, Woolf A et al. Assessing risk of bias in prevalence studies: modification of an existing tool and evidence of interrater agreement. *J Clin Epidemiol* 2012;65(9):934-9.
16. Werfalli M, Musekiwa A, Engel ME, Ross I, Kengne AP, Levitt NS. The prevalence of type 2 diabetes mellitus among older people in Africa: a systematic review study protocol. *BMJ open* 2014;4(6):e004747.
17. Theroux J, Stomski N, Hodgetts CJ et al. Prevalence of low back pain in adolescents with idiopathic scoliosis: a systematic review. *Chiropr Man Therap* 2017; 25: 10. Published online 2017 Apr 20. doi: 10.1186/s12998-017-0143-1
18. Sokal RR, Rohlf FJ (1969). *Biometry: The principles and practice of statistics in biological research*. WH Freeman: San Francisco. xxi, 776 pp
19. Hellgren L. The prevalence of urticaria in the total population. *Acta Allergol* 1972;27(3):236-40.
20. Gaig P, Olona M, Munoz Lejarazu D et al. Epidemiology of urticaria in Spain. *J Investig Allergol Clin Immunol* 2004;14(3):214-20.

21. Zuberbier T, Balke M, Worm M, Edenharter G, Maurer M. Epidemiology of urticaria: a representative cross-sectional population survey. *Clin Exp Dermatol* 2010;35(8):869-73.
22. Balp MM, Vietri J, Tian H, Isherwood G. The impact of chronic urticaria from the patient's perspective: a survey in five european countries. *Patient* 2015;8(6):551-8.
23. Cantarutti A, Dona D, Visentin F et al. Epidemiology of frequently occurring skin diseases in italian children from 2006 to 2012: a retrospective, population-based study. *Pediatr Dermatol* 2015;32(5):668-78.
24. Lapi F, Cassano N, Pegoraro V et al. Epidemiology of chronic spontaneous urticaria: results from a nationwide, population-based study in Italy. *Br J Dermatol* 2016;174(5):996-1004.
25. Vázquez Nava F, Martínez Burnes J. Prevalence and factors associates to chronic urticaria. Analysis of the allergic rhinitis and symptoms related to asthma as a factors associated to chronic urticaria in a urban area of northeastern of Mexico. *Alergol Inmunol Clin* 2004(19):16-24.
26. Balp MM, Lopes da Silva N, Vietri J, Tian H, Ensina LF. The burden of chronic urticaria from Brazilian patients' perspective. *Dermatol Ther (Heidelb)* 2017 Dec; 7(4): 535–545.
27. Zazzali JL, Broder MS, Chang E, Chiu MW, Hogan DJ. Cost, utilization, and patterns of medication use associated with chronic idiopathic urticaria. *Ann Allergy Asthma Immunol* 2012 Feb;108(2):98-102.
28. Broder MS, Raimundo K, Antonova E, Chang E. Resource use and costs in an insured population of patients with chronic idiopathic/spontaneous urticaria. *Am J Clin Dermatol* 2015; 16(4): 313–321.
29. Vietri J, Turner SJ, Tian H, Isherwood G, Balp MM, Gabriel S. Effect of chronic urticaria on US patients: analysis of the National Health and Wellness Survey. *Ann Allergy Asthma Immunol* 2015;115(4):306-11.
30. Ohmi T, Mue S, Tamura G, Ishihara T, Taniguchi Y, Takishima T. An epidemiologic study of cholinergic urticaria in students of junior and senior high schools in miyagi prefecture. *Arerugi* 1984 Aug;33(8):447-53.
31. Confino-Cohen R, Chodick G, Shalev V, Leshno M, Kimhi O, Goldberg A. Chronic urticaria and autoimmunity: associations found in a large population study. *J Allergy Clin Immunol* 2012 May;129(5):1307-13.
32. Lee SJ, Ha EK, Jee HM et al. Prevalence and risk factors of urticaria with a focus on chronic urticaria in children. *Allergy Asthma Immunol Res* 2017;9(3):212-9.
33. Lee N, Lee JD, Lee HY, Kang DR, Ye YM. Epidemiology of chronic urticaria in Korea using the Korean health insurance database, 2010-2014. *Allergy Asthma Immunol Res* 2017;9(5):438-45.
34. Chu CY, Cho YT, Jiang JH, Lin EI, Tang CH. Epidemiology and comorbidities of patients with chronic urticaria in Taiwan: A nationwide population-based study. *J Dermatol Sci* 2017 Nov;88(2):192-198.

35. Shalom G, Magen E, Babaev M et al. Chronic urticaria and the metabolic syndrome: a cross-sectional community-based study of 11 261 patients. *J Eur Acad Dermatol Venereol*. 2018;32(2):276-281.
36. El-Khateeb EA, Lotfi RA, Abd Elaziz KM, El-Shiekh SE. Prevalences of skin diseases among primary schoolchildren in Damietta, Egypt. *Int J Dermatol* 2014 May;53(5):609-16.
37. Maurer M, Weller K, Bindslev-Jensen C et al. Unmet clinical needs in chronic spontaneous urticaria. A GA2LEN task force report. *Allergy* 2011;66:317–30.
38. Zuberbier T, Althaus C, Chantraine-Hess S, Czarnetzki BM. Prevalence of cholinergic urticaria in young adults. *J Am Acad Dermatol* 1994;31:978-981.
39. Raciborski F, Kłak A, Czarnecka-Operacz M et al. Epidemiology of urticaria in Poland - nationally representative survey results. *Postepy Dermatol Alergol*. 2018;35(1):67-73.
40. Parisi CA, Ritchie C, Petriz N et al. Chronic urticaria in a health maintenance organization of Buenos Aires, Argentina - new data that increase global knowledge of this disease. *An Bras Dermatol*. 2018;93(1):76-79.
41. Weller K, Altrichter S, Ardelean E, et al. Chronic urticaria. Prevalence, course, prognostic factors and impact. *Hautarzt* 2010 Sep;61(9):750-7.
42. Pinart M, Keller T, Reich A et al. Sex-related allergic rhinitis prevalence switch from childhood to adulthood: a systematic review and meta-Analysis. *Int Arch Allergy Immunol* 2017;172(4):224-235.
43. Fröhlich M, Pinart M, Keller T et al. Is there a sex-shift in prevalence of allergic rhinitis and comorbid asthma from childhood to adulthood? A meta-analysis. *Clin Transl Allergy* 2017 Dec 5;7:44.
44. Almqvist C, Worm M, Leynaert B; working group of GA2LEN WP 2.5 Gender. Impact of gender on asthma in childhood and adolescence: a GA2LEN review. *Allergy* 2008 Jan;63(1):47-57.
45. Keller T, Hohmann C, Standl M et al. The sex-shift in single disease and multimorbid asthma and rhinitis during puberty - a study by MedALL. *Allergy* 2018 Mar;73(3):602-614.
46. Eun SJ, Lee JY, Kim DY, Yoon HS. Natural course of new-onset urticaria: Results of a 10-year follow-up, nationwide, population-based study. *Allergol Int*. 2019 Jan;68(1):52-58.
47. Balp MM, Weller, K, Carboni V et al. Prevalence and clinical characteristics of chronic spontaneous urticaria in pediatric patients. *Pediatr Allergy Immunol* 2018 Sep;29(6):630-636.
48. Maurer M, Metz M, Bindslev-Jensen C et al. Definition, aims, and implementation of GA(2) LEN Urticaria Centers of Reference and Excellence. *Allergy*. 2016 Aug;71(8):1210-8.
49. Balke M. Epidemiology of Urticaria. PhD Thesis, Charité-Humboldt-University Berlin, 2003.

Figure and table legends:

Figure 1: PRISMA flow chart for the literature search

Figure 2: Overall lifetime prevalence of chronic urticaria

Figure 3: Overall point prevalence of chronic urticaria

Figure 4: Point prevalence of chronic urticaria by major global regions

Figure 5: Point prevalence of chronic urticaria by sex

Figure 6: Point prevalence of chronic urticaria by age group

Figure e1: Subgroup analysis: Point prevalence of chronic urticaria in Europe by sex

Figure e2: Subgroup analysis: Point prevalence of chronic urticaria in Asia by sex

Figure e3: Subgroup analysis: Point prevalence of chronic urticaria in Northern America by sex


Figure e4: Subgroup analysis: Point prevalence of chronic urticaria by children < 15 years and sex

Figure e5: Point prevalence of chronic urticaria by method of data collection

Table e1: Search strategy

Table e2: Descriptive data of individual studies

Table e3: Overview prevalence and incidence estimates of individual studies

Figure 1




Figure 2: Overall lifetime prevalence of chronic urticaria




Figure3: Overall point prevalence of chronic urticaria




Figure 4: Point prevalence of chronic urticaria by major global regions




Figure 5: Point prevalence of chronic urticaria by sex




Figure 6: Point prevalence of chronic urticaria by age group




Figure e1: Subgroup analysis: Point prevalence of chronic urticaria in Europe by sex




Figure e2: Subgroup analysis: Point prevalence of chronic urticaria in Asia by sex




Figure e3: Subgroup analysis: Point prevalence of chronic urticaria in Northern America by sex




Figure e4: Subgroup analysis: Point prevalence of chronic urticaria by children < 15 years and sex




Figure e5: Point prevalence of chronic urticaria by method of data collection

Table e1 Search strategy

Date: 7/06/2017		EMBASE	PUBMED
Terms	Search	Query	Query
Chronic Urticaria Term	#1	exp urticaria/	urticaria[MeSH Terms]
	#2	exp angioneurotic edema/	angioedema[MeSH Terms]
	#3	urticaria.ti.	chronic urticaria[Title/Abstract]
	#4	chronic urticaria.ti,ab.	chronic idiopathic urticaria[Title/Abstract]
	#5	chronic spontaneous urticaria.ti,ab.	chronic spontaneous urticaria[Title/Abstract]
	#6	chronic idiopathic urticaria.ti,ab.	hives[Title/Abstract]
	#7	hives.ti,ab.	nettle rash[Title/Abstract]
	#8	wheals.ti,ab.	weals[Title/Abstract]
	#9	weals.ti,ab.	wheals[Title/Abstract]
	#10	nettle rash.ti,ab.	urticaria[Title]
	#11	1 or 2 or 3 or 4 or 5 or 6 or 7 or 8 or 9 or 10	(((((((((urticaria[MeSH Terms]) OR angioedema[MeSH Terms]) OR chronic urticaria[Title/Abstract]) OR chronic idiopathic urticaria[Title/Abstract]) OR chronic spontaneous urticaria[Title/Abstract]) OR hives[Title/Abstract]) OR nettle rash[Title/Abstract]) OR weals[Title/Abstract]) OR wheals[Title/Abstract]) OR urticaria[Title]
Prevalence Term	#12	prevalence.m_titl.	prevalence[MeSH Terms]
	#13	incidence.m_titl.	incidence[MeSH Terms]
	#14	epidemiology.m_titl.	epidemiology[MeSH Terms]
	#15	exp prevalence/	prevalence[Title]
	#16	exp incidence/	incidence[Title]
	#17	exp epidemiology/	epidemiology[Title]
	#18	12 or 13 or 14 or 15 or 16 or 17	(((((prevalence[MeSH Terms]) OR incidence[MeSH Terms]) OR epidemiology[MeSH Terms]) OR prevalence[Title]) OR incidence[Title]) OR epidemiology[Title]
	#19	11 and 18	(((((((((((((urticaria[MeSH Terms]) OR angioedema[MeSH Terms]) OR chronic urticaria[Title/Abstract]) OR chronic idiopathic urticaria[Title/Abstract]) OR chronic spontaneous urticaria[Title/Abstract]) OR hives[Title/Abstract]) OR nettle rash[Title/Abstract]) OR weals[Title/Abstract]) OR wheals[Title/Abstract]) OR urticaria[Title]))) AND ((((((prevalence[MeSH Terms]) OR incidence[MeSH Terms]) OR epidemiology[MeSH Terms]) OR prevalence[Title]) OR incidence[Title]) OR epidemiology[Title])
Limits	#20	limit 19 to (human and (embase or medline) and (abstract report or article or article in press or reports or "review" or short survey) and (journal or major reference work or report))	

Table e2 Descriptive data of individual studies

Region	Authors/ Year of Publication	Study design	Country	Sample size	Database	Response rate	Observation period	Age of participants	Definition of disease	Method of data collection	Type of urticaria	Qualitative appraisal
EUROPE	Hellgren (1972) [19]	Cross-sectional study	Sweden	36,475	The total general population over the age of 7 of five major regions	80-90%	Not specified	≥8 years	Whitish or reddish, slightly elevated papules, macules, patches or wheals, often surrounded by a halo >6 weeks	Diagnosis by physician	CU (dermographism was excluded)	High risk of bias
	Gaig et al. (2004) [20]	Cross-sectional study	Spain	5,003	National telephone directory	99%	September - November 2001	≥ 18 years	Hives with or without angioedema daily for >6 weeks (+ current treatment)	Questionnaire by telephone interview	CU	Low risk of bias
	Zuberbier et al. (2010) [21]	Cross-sectional study	Germany	4,093	PANE study	30.8%	1998-2000	All ages	Weals and / or angioedema >6 weeks	3 step assessment (self-reporting questionnaire, telephone interview, diagnosis by physician)	CU.	Moderate risk of bias
	Balp et al. (2015) [22]	Cross-sectional study	France, Germany, Italy, Spain and UK	2010: 57,805 2011: 57,512 2013: 62,000	5EU National Health and Wellness Survey (NHWS)	Not specified (opt-in online survey panels)	2010, 2011, 2013	≥ 18 years	Experience, diagnosis, and treatment of chronic hives	Self-reporting Questionnaire	CU	Moderate risk of bias
	Cantarutti et al. (2015) [23]	Secondary data analysis	Italy	145,233	Pedianet database	Not specified	2006-2012	≤14 years	ICD-9 diagnosis (not specified in the article) or skin disorder names	Diagnosis by physician	CU	Low risk of bias
	Lapi et al. (2016) [24]	Secondary data analysis	Italy	~ 1,000,000	Health Search IMS Health Longitudinal Patient Database	Not specified	2002-2013	≥ 15 years	ICD-9 diagnosis t (algorithm)	Diagnosis by physician	CSU	Moderate risk of bias

LATIN AMERICA	Vazquez Nava and Martinez Burnes (2004) [25]	Cross-sectional study	Mexico	2,613	Not specified	87.1%	Not specified	20-50 Jahre	Hives with or without angioedema twice a week for >6 weeks	Self-reporting Questionnaire	CU	Moderate risk of bias
	Balp et al. (2017) [26]	Cross-sectional study	Brazil	36,000	National Health and Wellness Survey (NHWS) in Brazil	Not specified (opt-in online survey panels)	2011, 2012, 2015	≥ 18 years	Experience, diagnosis, and treatment of chronic hives	Self-reporting Questionnaire	CU	Moderate risk of bias
NORTHERN AMERICA	Zazzali et al. (2012) [27]	Secondary data analysis	USA	7,555,991	Health Insurance Portability and Accountability Act	Not specified	July 2007-June 2008	All ages	ICD-9 diagnosis (algorithm)	Diagnosis by physician	CSU	Low risk of bias
	Broder et al. (2015) [28]	Secondary data analysis	USA	5,802,466	US commercial health plan	Not specified	January-December 2012	All ages	ICD-9 diagnosis(algorithm)	Diagnosis by physician	CSU	Low risk of bias
	Vietri et al. (2015) [29]	Cross-sectional study	USA	197,463	US National Health and Wellness Survey	Not specified (opt-in online survey panels)	2010-2012	≥ 18 years	Experience, diagnosis, and treatment of chronic hives	Self-reporting Questionnaire	CU	Moderate risk of bias
ASIA	Ohmi et al. (1984) [30]	Cross-sectional study	Japan	113,630	Middle and high schools in Miyagi	78,90%	April to August 1982	Middle and High school students: ~12-17 years (age not specified)	Exercise induced cutaneous pruritus or whealing episode	Self-reporting Questionnaire	CIndU (cholinergic urticaria)	Moderate risk of bias
	Confino-Cohen et al. (2012) [31]	Secondary data analysis	Israel	~ 2,000,000	Maccabi Healthcare Services (Health maintenance organisation)	Not specified	January 1993-March 2010	All ages	ICD-9 CM diagnosis	Diagnosis by physician	CSU	Moderate risk of bias

	Lee SJ et al (2017) [32]	Cross-sectional study	South Korea	4,076	Seongnam Atopy Project in Korea (SAP2015)	85.3%	June to July of 2015	4 to 13 years	Burning, itching or a rash resembling a mosquito bite or swelling on the skin >6 weeks	Self-reporting Questionnaire	CU	Moderate risk of bias
	Lee N et al. (2017) [33]	Secondary data analysis	South Korea	Total Korean population in 2012 from Statistics Korea	Korean Health Insurance Review and Assessment Service (HIRA) database	Not specified Note: 98% of the Korean population are covered by NHIS	2010-2014	All ages	ICD-10 diagnosis (algorithm)	Diagnosis by physician	CU	Moderate risk of bias
	Chu et al. (2017) [34]	Secondary data analysis	Taiwan	2009: 22,282,479 2010: 22,336,029 2011: 22,537,021 2012: 22,532,255	National Health Insurance Research Database (NHIRD)	Not specified Note: System covers 99.9% of the total population of Taiwan	2009-2012	All ages	ICD-9 diagnosis (algorithm)	Diagnosis by physician	CSU	Low risk of bias
	Shalom et al.(2018) [35]	Secondary data analysis	Israel	~4,300,000	Clalit Health Service (Health maintenance organisation)	Not specified	January 2002- December 2012	All ages	8 urticaria diagnoses (with each two diagnoses registered within a period of 6 weeks)	Diagnosis by physician	CU	Moderate risk of bias
AFRICA	El-Khateeb et al. (2014) [36]	Cross-sectional study	Egypt	6,162	Primary schools in Damietta	Not specified	October 2011-March 2012	6 to 12 years	ICD-10 diagnosis Skin disorder	Diagnosis by physician	CIndU (Dermographism)	Low risk of bias

CU: chronic urticaria, CSU: chronic spontaneous urticaria, CIndU: chronic inducible urticaria

Table e3 Overview prevalence and incidence estimates of individual studies

Region	Authors/ Year	Sample size	Age of participants	Type of urticaria	Prevalence		Incidence	Comments
					Lifetime	Point		
EUROPE	Hellgren (1972) [19]	36,475	≥7 years	CU (dermographism was excluded)	Not reported	(One day) 0.08%	Not reported	
	Gaig et al (2004) [20]	5,003	≥18 years	CU	2.9% †	(1 week + current treatment) 0.6% (95%CI, 0.4%-0.8%)	Not reported	†95%CI not reported
	Zuberbier et al. (2010) [21]	4,093	All age groups	CU.	1.81%‡ (95% CI 1.4–2.3%)	(12 months) 0.81%‡ (95%CI, 0.6–1.1%)	0.1%‡	‡Data completed with data published in PhD thesis (Balke 2003 [49])
	Balp et al. (2015) [22]	2010: 57,805 2011: 57,512 2013: 62,000	≥18 years	CU	(diagnosis) 0.51%§ (95%CI 0.48-0.55) (experience) 0.63%§ (95%CI 0.59-0.67)	(diagnosis + current treatment) 0.23%§ (369/175,923)(95%CI, 0.20- 0.25)	Not reported	§Weighted
	Cantarutti et al. (2015) [23]	145,233	0 to 14 years	CU	Not reported	(12 months) Year 2006: 0.38% (95%CI, 0.35–0.4) Year 2012: 0.84% (95%CI, 0.8–0.9)	(per 1,000 person-years Year 2006: 2.1% (95%CI,1.9–2.4) Year 2012: Not reported	
	Lapi et al. (2016) [24]	~ 1,000,000	≥ 15 years	CSU	Not reported	(12 months) Year 2002: 0.02%¶ Year 2013: 0.38%¶	(per 1000 person-years) Year 2002 : 0.10%¶ Year 2013: 1.30%¶	¶Data from graph
LATIN AMERICA	Vazquez and Martinez (2004) [25]	2,613	20-50 years	CU	30.15%†	(12 months) 3.44%†	Not reported	†95%CI not reported
	Balp et al. (2017) [26]	36,000	≥18 years	CU	(diagnosis) 0.41%§ (95%CI, 0.33-0.49) (experience) 0.53%§ (95%CI,0.44-0.62)	(diagnosis + treatment) 0.21%§ (95%CI, 0.15-0.27%)	Not reported	§Weighted

NORTHERN AMERICA	Zazzali et al. (2012) [27]	7,555,991	Not reported	CSU	Not reported	(12 months) 0.08%†	Not reported	†95%CI not reported
	Broder et al. (2015) [28]	5,802,466	All age groups	CSU	Not reported	(12 months) 0.11%†	Not reported	†95%CI not reported
	Vietri et al. (2015) [29]	197,463	≥18 years	CU	(diagnosis) 0.40%§ (95%CI,0.37-0.43) (experience) 0.53%§ (95%CI,0.49-0.56)	(diagnosis + treatment) 0.13%§ (95%CI, 0.11-0.15)	Not reported	§Weighted
ASIA	Ohmi et al. (1984) [30]	113,630	Not reported	CIndU (cholinergic urticaria)	0.1%††	Not reported	Not reported	†† Calculated by us. Authors reported 0.13%
	Confino-Cohen et al. (2012) [31]	~2,000,000	All ages groups	CSU	Not reported (12,778 cases†† in a ~2-million-enrollee health organization database: ~0.64%)	Not reported	Not reported	††17 years period
	Lee SJ et al. (2017) [32]	4,076	4 to 13 years	CU	Not reported	(12 months) 1.4%††	Not reported	††Calculated by us. Authors reported 1.8%
	Lee N et al. (2017) [33]	Total Korean population in 2012 from Statistics Korea	All age groups	CU	Not reported	(12 months) Year 2010: 1.66% (per 100,000 person-years) Year 2014: 2.31% (per 100,000 person-years)	Not reported	
	Chu et al. (2017) [34]	2009: 22,282,479 2012: 22,532,255	All age groups	CSU	Not reported	(12 months) Year 2009: 0.69%† Year2012: 0.79%†	(Incidence rate) Year 2009: Not available Year 2012:0.50%¶¶	†95%CI not reported ¶¶Data from graph

	Shalom et al. (2018) [35]	~4,300,000	All age groups	CU	0.27%†	Not reported	Not reported	†95%CI not reported, 11 years period
AFRICA	El-Khateeb et al. (2014) [36]	6,162	6 to 12 years	CIndU (Dermographism)	Not reported	(One day) 0.03%†	Not reported	†95%CI not reported

CU: chronic urticaria, CSU: chronic spontaneous urticaria, CIndU: chronic inducible urticaria