

Birch pollen-related food allergy: Clinical aspects and the role of allergen-specific IgE and IgG₄ antibodies

Marija Geroldinger-Simic, MD,^a Thomas Zelniker, MD,^a Werner Aberer, MD,^c Christof Ebner, MD,^d Cornelia Egger, MD,^e Antonia Greiderer,^e Nicole Prem, MD,^c Jonas Lidholm, PhD,^f Barbara K. Ballmer-Weber, MD,^g Stefan Vieths, PhD,^h and Barbara Bohle, PhD^{a,b} Vienna, Graz, and Innsbruck, Austria, Uppsala, Sweden, Zurich, Switzerland, and Langen, Germany

Background: Patients with birch pollen allergy often develop allergic reactions to plant foods.

Objective: To evaluate the prevalence, main symptoms, and triggers of birch pollen-related food allergy and the role of food-specific IgG₄ antibodies in food tolerance.

Methods: Food-induced symptoms were evaluated in 225 individuals with birch pollen allergy by using a standardized questionnaire. IgE and IgG₄ levels specific for the major birch pollen allergen Bet v 1 and birch profilin Bet v 2 and the Bet v 1 homologs in apple (Mal d 1) and hazelnut (Cor a 1) were quantified by ImmunoCAP. Mock-treated and IgG-depleted sera from patients tolerating hazelnuts in food challenges were compared for their inhibitory activity for binding of Cor a 1-IgE complexes to B cells.

Results: In total, 73% of the study population experienced food allergy, which was perennial in 86% of the affected individuals. The oral allergy syndrome was the main clinical manifestation. However, more than 58% of the patients also experienced food-

induced rhinoconjunctivitis. Apples and hazelnuts were identified as the most frequent triggers. Food allergy correlated with IgE reactivity to Bet v 1 but not to Bet v 2. Mal d 1-specific and Cor a 1-specific IgG₄/IgE ratios were significantly higher in food-tolerant individuals than individuals with food allergy. Sera from IgG₄-positive food-tolerant patients possessed IgG-dependent IgE-inhibitory activity.

Conclusion: Birch pollen-related food allergy is highly prevalent and often perennial. High food allergen-specific IgG₄/IgE ratios seem associated with food tolerance, potentially because specific IgG₄ blocks IgE binding to food allergens. Thus, the presence of food allergen-specific IgG₄ antibodies is no diagnostic marker for birch pollen-related food allergy. (J Allergy Clin Immunol 2010;■■■■:■■■-■■■.)

Key words: Birch pollen allergy, food allergy, oral allergy syndrome, Bet v 1, IgE, blocking antibodies

From ^athe Department of Pathophysiology and Allergy Research and ^bthe Christian Doppler Laboratory for Immunomodulation, Medical University of Vienna; ^cthe University Clinic for Dermatology, Medical University of Graz; ^dthe Allergy Clinic Reumannplatz, Vienna; ^ethe Department of Dermatology, University Clinic Innsbruck; ^fthe Research and Development Department, Phadia AB, Uppsala; ^gthe Allergy Unit, Department of Dermatology, University Hospital Zurich; and ^hPaul-Ehrlich-Institut, Division of Allergology, Langen.

Supported by the Austrian Science Fund, project SFB-F1807-B13, the Christian Doppler Research Association, and Biomay AG, Vienna, Austria.

Disclosure of potential conflict of interest: B. Bohle receives research support from the Austrian Science Fund and the Christian Doppler Laboratory. S. Vieths is an associate for the Institute for Product Quality; has received honoraria for scientific presentations from Phadia; has received honoraria as an expert reviewer from the Food Allergy Resource and Research Program; has received research support from the European Union, the German Research Foundation, the German Federal Ministry of Education and Research, the Research Fund for the German Food Industry, Monsanto Co, Pioneer Hi-Bred International, Inc, the Food Allergy Resource and Research Program, and the European Directorate for the Quality of Medicines and Health Care; is chairman of the Food Allergy Interest Group and a member of the Executive Committee for EAACI; is chairman of the Allergen Standardization Subcommittee and secretary of the Allergen Nomenclature Subcommittee for the International Union of Immunological Studies; is a Registered Expert for the European Agency for the Evaluation of Medicinal Products and European Pharmacopoeia Commission; is chairman of Technical Committee 275 Working Group 12 "Food Allergens" for the European Committee for Standardization; and is a member of the Food Allergy Working Group for the German Society for Allergy and Clinical Immunology. The rest of the authors have declared that they have no conflict of interest.

Received for publication July 16, 2010; revised September 10, 2010; accepted for publication October 18, 2010.

Reprint requests: Barbara Bohle, PhD, Medical University of Vienna, Department of Pathophysiology and Allergy Research, Waehringer Guertel 18-20, AKH-3Q, A-1090 Wien, Austria. E-mail: barbara.bohle@meduniwien.ac.at.

0091-6749/\$36.00

© 2011 American Academy of Allergy, Asthma & Immunology

doi:10.1016/j.jaci.2010.10.027

Birch pollen is one of the most common causes of rhinoconjunctivitis and allergic asthma in Northern and Central Europe and North America.^{1,2} It has long been known that patients with birch pollen allergy may develop immediate reactions to fruits and vegetables in addition to seasonal respiratory symptoms.^{3,4} This birch-fruit-vegetable syndrome is characterized by local symptoms at the site of food contact such as itching of the lips, tongue, and throat, sometimes accompanied by swelling of the lips and tongue, referred to as the oral allergy syndrome.⁵ Occasionally, systemic and more severe IgE-mediated reactions such as urticaria, asthma, or anaphylactic shock may occur.⁶ Although birch pollen-related food allergy is often noticed in clinical practice, no recent data on the prevalence and main triggers of this type of food allergy are available.^{7,8}

Birch pollen-related food allergy is considered a consequence of immunologic cross-reactivity between ubiquitous birch pollen allergens and structurally related food proteins. IgE antibodies specific for the major birch pollen allergen, Bet v 1, have been shown to cross-react with homologous proteins identified in different stone-fruits, such as apple (Mal d 1), cherry (Pru av 1), and pear (Pyr c 1), as well as hazelnut (Cor a 1), celery (Api g 1), carrot (Dau c 1), soybean (Gly m 4), peanut (Ara h 8; see summary⁹), jackfruit,¹⁰ and kiwi (Act d 8).¹¹ These food allergens can also activate Bet v 1-specific T cells to proliferate and produce cytokines,¹² sometimes even resulting in T cell-mediated late-phase responses in target organs, such as flare-ups of atopic eczema in the skin.^{12,13} Bet v 2, the birch pollen profilin, is another allergen able to induce cross-reactive IgE antibodies.^{14,15} Bet v 2-specific IgE antibodies have been shown to recognize profilins in apple, banana, carrot, celery, cherry, hazelnut, pear, pineapple,

Abbreviations used

FAB: Facilitated allergen binding

r: Recombinant

SIT: Allergen-specific immunotherapy

potato, and tomato.¹⁶⁻²² IgE antibodies specific for the minor allergen Bet v 6, an isoflavone reductase-homologous protein of 35 kd, have been shown to cross-react with proteins of comparable size in apple, banana, carrot, lychee, mango, orange, and pear.^{23,24}

Often, patients with birch pollen allergy who display IgE reactivity to birch pollen-related allergens in foods do not develop clinical symptoms when consuming the foods.^{20,25} Recently, evidence accumulated that allergen-specific IgG, in particular IgG₄ antibodies, might play a role in food tolerance. For example, tolerance to cow's milk in atopic children and adults has been associated with elevated levels of specific IgG₄ in combination with low specific IgE.²⁶⁻²⁸ Furthermore, significant increases of food allergen-specific IgG₄ antibodies have been observed during specific oral immunotherapy with milk, egg, and peanut²⁹⁻³² and during sublingual immunotherapy with hazelnut and peach extract, respectively.^{33,34} These findings prompted us to investigate the role of allergen-specific IgG₄ antibodies in patients tolerant to birch pollen-related foods.

We included 225 patients with birch pollen allergy to analyze the prevalence, symptoms, and foods involved in birch pollen-related food allergy. Furthermore, the relevance of IgE reactivity to Bet v 1 and Bet v 2 for food-induced reactions was elucidated. Allergen-specific IgG₄ levels in patients tolerating and reacting to birch pollen-related foods were compared. Mock-treated and IgG-depleted sera from food-tolerant patients were investigated for their capacity to block IgE binding to Bet v 1-related food allergens.

METHODS

Study population and questionnaire

In total, 225 patients who attended Austrian Allergy Clinics in Innsbruck, Graz, and Vienna from May to June 2007 and were diagnosed as having birch pollen allergy were included. The diagnosis was made on the basis of clinical history, positive skin prick test responses to birch pollen extract, and specific IgE to birch pollen (≥ 0.35 kilo units [kU_A/L], ImmunoCAP; Phadia, Uppsala, Sweden). In 140 individuals, skin prick tests to grass pollen were performed and were positive in 65.7%. None of the individuals had undergone allergen-specific immunotherapy (SIT). All patients gave written informed consent. Patients were interviewed by using a standardized questionnaire in German adopted from Fernandez-Rivas et al³⁵ (see this article's Fig E1 in the Online Repository at www.jacionline.org) including 16 foods, namely fruits of the Rosaceae family (almond, apple, apricot, cherry, nectarine/peach, pear, plum), vegetables of the Apiaceae (carrot, celery) and Solanaceae families (potato peel, tomato), Leguminosae (peanuts, soybean), hazelnut, walnut, and kiwi.

Allergens

Recombinant (r) Bet v 1.0101 (termed Bet v 1 in this article), Bet v 2.0101 (termed Bet v 2), and Mal d 1.0108 (termed Mal d 1) were purchased from Biomay (Vienna, Austria). rCor a 1.0401 (termed Cor a 1) was kindly provided by Stephan Scheurer, Paul Ehrlich Institut, Langen, Germany.

Determination of allergen-specific IgE and IgG₄

IgE and IgG₄ levels specific for the different allergens were determined by ImmunoCAP (Phadia) and ELISA. For the latter, microtiter plates

TABLE I. Demographic data of the study population

Characteristics	Values
Sex, n (%)	
Female	136 (60.4)
Male	89 (39.6)
Age (y)	
Median	35.0
Range	6-76
Total IgE (kU/L)	
Median	143.0
Range	2.0->5000
Birch-specific IgE (kU _A /L)	
Median	20.0
Range	0.37->100

(Maxisorp, Nunc, Denmark) were coated with the recombinant allergens (2 μ g/mL each) in carbonate buffer (pH 9.6) overnight at 4°C. After washing and saturation, sera were incubated in duplicate overnight at 4°C. Bound IgE antibodies were detected by using alkaline phosphatase-conjugated antihuman IgE (BD Biosciences, San Jose, Calif). Bound IgG₄ was detected by mouse antihuman IgG₄ (BD Biosciences) followed by horseradish peroxidase-conjugated sheep antimouse IgG₄ (GE Healthcare, Buckinghamshire, United Kingdom). Sera from 10 subjects without allergy were used as negative controls in IgE-ELISA. The mean value of all negative controls plus 5 times the SD was set as the cutoff value for positive IgE reactivity.

Depletion of IgG antibodies

Sera from patients tolerating hazelnuts in double-blind placebo-controlled food challenges³⁶ were incubated with either protein G-Sepharose beads (Bio Vision Research Products, Mountain View, Calif) or Sepharose beads for 4 hours at room temperature. Thereafter, sera were centrifuged and incubated with fresh beads overnight at room temperature. These steps were repeated until ELISA results showed successful depletion of allergen-specific IgG₄. As controls, sera were used from 4 patients undergoing birch pollen SIT for 24 months who showed improved clinical symptoms to birch pollen and associated foods.

IgE-facilitated allergen binding inhibition assay

The facilitated allergen binding (FAB) inhibition assay measures the capacity of IgG antibodies to inhibit the formation of IgE-allergen complexes that bind to the low-affinity IgE receptor (CD23) on B cells.³⁷ In preliminary experiments, indicator sera with an ImmunoCAP >50 kU_A/L to rCor a 1 were incubated with different concentrations of rCor a 1 (1, 0.3, 0.1, and 0.03 μ g/mL) to form IgE-allergen complexes. We determined 0.03 μ g/mL as the optimum concentration. Thereafter, indicator serum (15 μ L) plus either mock-treated or IgG-depleted sera (20 μ L) were incubated with 5 μ L rCor a 1 (0.03 μ g/mL) for 1 hour at 37°C. Then 1×10^5 EBV-transformed B cells (5 μ L) were added and incubated for 1 hour at 4°C. Cells were washed, and bound complexes were detected by using a fluorescein isothiocyanate-labeled affinity purified goat-antihuman IgE antibody (KPL, Gaithersburg, Md) with a FACS Canto (BD Biosciences).

Statistical analysis

Statistical analysis was performed with SPSS 14.0 (SPSS Inc, Chicago, Ill). Differences of allergen-specific antibody levels between patients with food allergy and food-tolerant patients were evaluated by using the Wilcoxon signed-ranks test, and the Spearman rank test was used to determine correlation. Two-sided *P* values <.05 were considered significant.

TABLE II. Bet v 1 homologs and profilins in birch pollen-related foods

Food	Frequency of food hypersensitivity (%) [*]	Bet v 1 homolog	GenBank accession no.	Profilin	GenBank accession no.
Apple	80.0	Mal d 1	AJ417551	Mal d 4	AF129426 AF129427 AF129428
Hazelnut	59.4	Cor a 1	AF136945	Cor a 2	AF327622 AF327623
Nectarine/peach	50.9	Pru p 1	DQ251187	Pru p 4	AJ491881 AJ491882
Kiwi	47.9	Act d 8	AM489568	Act d 9	FG438715
Walnut	41.2	—	—	—	—
Carrot	34.5	Dau c 1	AF456481	Dau c 4	AF456482
Apricot	33.3	Pru ar 1	U93165	—	—
Cherry	32.1	Pru av 1	U66076	Pru av 4	AF129425
Pear	32.1	Pyr c 1	AF057030	Pyr c 4	AF129424
Almond	31.5	—	—	Pru du 4	AY081850 AY081852
Peanut	24.2	Ara h 8	AY328088	Ara h 5	AF059616
Plum	24.2	Pru d 1	—	Pru d 4	—
Tomato	21.2	—	—	Lyc e 1	AJ417553
Potato	18.8	—	—	—	—
Celery	15.8	Api g 1	Z48967	Api g 4	AF129423
Soybean	13.9	Gly m 4	X60043	Gly m 3	AJ223982

—, Not identified.

^{*}Percentage of positive reactions in 225 patients with birch pollen allergy.

RESULTS

Study population

Data on sex, age, and total and birch pollen-specific IgE levels of the 225 patients with birch pollen allergy are summarized in Table I. Testing all sera for IgE reactivity to rBet v 1 in ELISA revealed 205 positive patients (91%). In immunoblots with birch pollen extract 210 sera (93.3%) showed IgE reactivity to Bet v 1, and 73% of these patients were exclusively sensitized to the major birch pollen allergen (data not shown). The ImmunoCAP for rBet v 1 performed with 147 sera positively correlated with the respective ELISA results ($R = 0.827$; $P < .001$). ImmunoCAP results for rBet v 1-specific IgE also correlated highly with ImmunoCAP results for birch pollen ($R = 0.969$; $P < .001$). Of 147 patients, 22 (15%) were positive for rBet v 2 (ImmunoCAP ≥ 0.35 kU_A/L), and all but 1 were also sensitized to Bet v 1.

Main triggers and symptoms of birch pollen-related food allergy

In summary, 165 of 225 (73%) reported allergic reactions to at least 1 of the 16 foods included in the questionnaire. Among food-reactive patients, apple induced symptoms in 80% and hazelnut in 59.4%, followed by peach, kiwi, walnut, carrot, apricot, cherry, pear, almond, plum, peanut, tomato, potato peel, raw celery, and soybean (Table II). The most prevalent symptom was itching in the oral cavity (89.2%), followed by rhinitis (58.2%), conjunctivitis (57.6%), swelling of the lips (50.6%), tightness of the throat (50.8%), and itching in the ear (36.7%). Dyspnea (28.5%), cramps (18.4%), nausea (9.5%), urticaria (15.2%), and drop of blood pressure (8.9%) were less frequent. Of 121 food-reactive individuals who could comment on the onset of their food-induced symptoms, 85 (70.2%) reported that food allergy had started after birch pollinosis. Fifty-nine of 134

patients with food allergy (44%) experienced a worsening of food-induced symptoms during the birch pollen season, and 138 of 159 (86.8%) patients also had food allergy in winter, out of the birch pollen season.

Birch pollen-related food allergy is clearly associated with sensitization to Bet v 1

All foods included in the questionnaire contain Bet v 1 homologs and profilins (Table II). To evaluate the involvement of these birch pollen allergens in birch pollen-related food allergy, we compared the levels of total and birch pollen-specific IgE in 149 patients with food allergy and 56 food-tolerant patients, and Bet v 1-specific and Bet v 2-specific IgE levels in 115 patients with food allergy and 32 food-tolerant patients (Fig 1, A). No difference in total IgE levels between the groups was found (median, 163 kU_A/L, range, 11.2-5000 kU_A/L; and median, 140 kU_A/L, range, 2.0-3503 kU_A/L, respectively). Birch pollen-specific IgE levels were significantly higher in the group with food allergy (median, 19.1 kU_A/L; range, 0.6->100 kU_A/L) than in food-tolerant individuals (median, 6.1 kU_A/L; range, 0.4-68.4 kU_A/L; $P < .001$). Similarly, Bet v 1-specific IgE levels were significantly higher in patients with food allergy than tolerant patients (median, 26.7 kU_A/L, range, 0.9-381 kU_A/L; and median, 7.5 kU_A/L, range, 0.7-63.84 kU_A/L, respectively; $P < .001$). All patients with Bet v 1-specific IgE levels of >65 kU_A/L had food allergy. A significantly larger number of food-reactive individuals (96%) were sensitized to Bet v 1 than tolerant individuals (79%; $P = .046$; Fig 1, B). Fifteen of 115 (13.0%) of the individuals with food allergy and 7 of 32 (21.8%) of the tolerant individuals displayed IgE to Bet v 2 (Fig 1, B). Although all patients with higher Bet v 2-specific IgE levels (≥ 3.5 kU_A/L) belonged to the food-allergic group, these individuals were all concomitantly




FIG 1. IgE levels of patients with birch pollen allergy with and without food allergy. **A**, Levels of total IgE and IgE specific for birch pollen, Bet v 1, and Bet v 2 were compared in food-tolerant (open circles) and food-reactive (filled circles) patients. **B**, Percentage of Bet v 1-sensitized and Bet v 2-sensitized patients in tolerant (white bars) and food-reactive (gray bars) individuals. Horizontal lines indicate median values. Wilcoxon signed-ranks test, * $P < .05$, *** $P < .001$.

sensitized to Bet v 1. In total, 14 of 225 patients with birch pollen allergy showed no IgE reactivity to Bet v 1, of whom 7 individuals (50%) were tolerant to all foods asked about in the questionnaire. The IgE reactivity patterns to birch pollen allergens of the 7 Bet v 1-negative, food-reactive patients were analyzed in immunoblots (data not shown). Six individuals showed IgE reactivity to an allergen of 35 kd (presumably Bet v 6). Of these 6 patients, 2 showed IgE reactivity to Bet v 2 and 2 to an allergen of 9 kd (presumably Bet v 4). For 1 patient (birch pollen-specific IgE, 1.78 kU_A/L) who had reported allergic symptoms to stone-fruits, no IgE reactivity to any birch pollen allergen was detectable.

Food-tolerant patients show higher ratios of food allergen-specific IgG₄/IgE than patients with food allergy

Apple and hazelnut containing Mal d 1 and Cor a 1, respectively, were identified as the most frequent triggers for birch pollen-related food allergy. In patients with apple allergy, a strong positive correlation between IgE specific for Bet v 1 and Mal d 1 was found ($R = 0.811$; $P < .001$). Similarly, IgE levels specific for Bet v 1 and Cor a 1 highly correlated in patients with hazelnut allergy ($R = 0.927$; $P < .001$). Not all individuals who showed IgE reactivity to the Bet v 1-related food allergens also experienced food-induced allergic symptoms. Accordingly, 113 Mal d 1-sensitized patients were grouped into 85 individuals with apple allergy and 28 apple-tolerant individuals and 105 Cor a 1-sensitized patients into 62 patients with hazelnut allergy and 43 hazelnut-tolerant patients. IgG₄ levels specific for Mal d 1 and Cor a 1 were determined by ImmunoCAP (Fig 2). Apple-tolerant patients displayed significantly lower levels of Mal d 1-specific IgE ($P = .002$) and similar levels of IgG₄ compared with patients with apple allergy. In contrast, hazelnut-tolerant individuals displayed similar Cor a 1-specific IgE levels and significantly higher IgG₄ levels ($P = .03$) compared with patients with

hazelnut allergy (Fig 2). When IgG₄/IgE ratios were calculated for each individual, food-tolerant patients showed significantly higher IgG₄/IgE ratios than patients with food allergy (Fig 2). In apple-tolerant patients, the median of Mal d 1-specific IgG₄/IgE ratios was 27 (range, 6.1-202.7) compared with 14.5 (range, 1.5-426.4) in patients with apple allergy ($P = .005$). The median of Cor a 1-specific IgG₄/IgE ratios in hazelnut-tolerant patients was 4.73 (range, 0.43-212.7) compared with 2.49 (range, 0.14-72.8) in hazelnut-tolerant individuals ($P = .045$).

Specific IgG₄ antibodies in food-tolerant individuals block IgE binding to the allergen

We compared mock-treated and IgG-depleted sera from 6 patients with birch pollen allergy who tolerated hazelnuts in food challenges in the FAB inhibition assay.³⁷ All patients displayed Cor a 1-specific IgG₄ and IgE (≥ 4.8 kU_A/L), excluding that food tolerance is caused by very low allergen-specific IgE levels (Table III). In addition, we used sera from 4 patients with birch pollen allergy who had undergone successful birch pollen SIT and had developed Cor a 1-IgG₄ antibodies during therapy (Table III). Similar to SIT-treated individuals, sera from hazelnut-tolerant patients showed IgE inhibitory activity. Treatment with protein-G-Sepharose beads effectively removed allergen-specific IgG₄ antibodies in each serum (Fig 3, A). The depletion of IgG antibodies abolished the inhibitory activity of all sera (Fig 3, B). Sera collected from the same SIT-treated patients before the onset of therapy that contained no Cor a 1-specific IgG₄ showed no inhibitory activity (Fig 3, B).

DISCUSSION

We found that 73% of adult Austrian patients with birch pollen allergy experienced food allergy and identified apple and hazelnut as the most frequent triggers (Table II). These results match previous studies performed in Sweden,^{7,38} Italy,⁸ and Denmark.³⁹ Sensitization to Bet v 1 and/or Bet v 2 may cause




FIG 2. Specific IgE and IgG₄ levels in patients with and without food allergy. Patients with IgE >0.35 kU_A/L to Mal d 1 or Cor a 1 were grouped into food-tolerant (white) or food-reactive (gray). Food allergen-specific IgG₄ and IgG₄/IgE ratios were determined. For IgG₄/IgE ratios, box plots are shown. Wilcoxon signed-ranks test, **P* < .05.

food hypersensitivity.⁴⁰ In our study population, the frequency of Bet v 1-sensitized subjects as well as the levels of Bet v 1-specific IgE antibodies were significantly higher in food-reactive than food-tolerant patients (Fig 1). Moreover, 99 of 165 food-reactive patients (60%) showed exclusive IgE reactivity to Bet v 1 as determined by immunoblotting (data not shown). In contrast, neither the frequency nor the quantity of Bet v 2-specific IgE antibodies could be associated with food-induced reactions, and no patient with food allergy exclusively sensitized to Bet v 2 was found in our study population. In line with a previous study suggesting that Bet v 1-specific but not Bet v 2-specific IgE antibodies induced clinically relevant cross-reactivity to foods,⁴¹ our data underline the major role of Bet v 1 and the potential inferior role of Bet v 2 in birch pollen-related food allergy.

More than 90% of our study population displayed IgE antibodies to Bet v 1, but not all individuals had food allergy. To investigate a potential role of specific IgG₄ antibodies in the tolerance to birch pollen-related foods, we compared IgG₄ levels specific for Mal d 1 and Cor a 1 in patients tolerant or allergic to apple or hazelnut, respectively. No difference between Mal d 1-specific IgG₄ levels in apple-tolerant patients and patients with apple allergy was found (Fig 2). This observation may be explained by the fact that many patients with apple allergy continue to consume apple cultivars that induce only mild allergic reactions.⁴² This exposure to the food may raise specific IgG₄ antibody levels as observed in patients without allergy.⁴³ In contrast, hazelnuts were

TABLE III. Antibody levels in sera from patients tested in the FAB inhibition assay

Patient	Status*	Cor a 1-specific IgE		Cor a 1-specific IgG ₄	IgG ₄ /IgE ratio
		kU _A /L	μg/L	μg/L	
1	Tolerant	4.8	11.4	2538.3	222.1
2	Tolerant	13.4	32.2	6840.0	212.7
3	Tolerant	37.7	90.5	460.0	5.1
4	Tolerant	48.3	115.9	208.7	1.8
5	Tolerant	28.1	67.5	358.2	5.3
6	Tolerant	42.64	102.2	227.8	2.2
7	SIT-treated	0.0	0.0	1600	
8	SIT-treated	0.0	0.0	1800	
9	SIT-treated	36.5	87.6	1291	14.7
10	SIT-treated	49.7	119.4	847	7.1

*Patients 1 to 6 tolerated hazelnuts in food challenges, and patients 8 to 11 underwent birch pollen SIT.

strictly avoided by the majority of patients with hazelnut allergy, and patients tolerating hazelnut showed significantly higher Cor a 1-specific IgG₄ levels than patients with hazelnut allergy (Fig 2). When individual food-specific IgG₄/IgE ratios were calculated, patients tolerating either apple or hazelnut displayed significantly higher ratios than individuals with food allergy. Together, these results suggest that the individual specific IgG₄/IgE ratio might




FIG 3. IgG₄ in food-tolerant patients shows IgE inhibitory activity. Sera from Cor a 1-sensitized, hazelnut-tolerant patients (1-6) and patients who underwent birch pollen SIT (7-10; Table III) were included. **A**, Cor a 1-specific IgG₄ in mock-treated (white bars) and IgG-depleted (gray bars) sera assessed by ELISA. **B**, Serum inhibitory activity tested in FAB inhibition assays. Sera obtained from SIT-treated patients before therapy served as controls. Horizontal lines indicate median values.

be involved in tolerance to birch pollen-related foods as it has been postulated for tolerance to cow's milk in atopic children.²⁶⁻²⁸

In line with the EAACI Task Force report, our study confirms that the detection of food-specific IgG₄ antibodies is no diagnostic marker for food allergy.⁴³ Moreover, because of the high variation of the ratios between the individual patients, we do not consider IgG₄/IgE ratios suitable to predict or diagnose birch pollen-related food allergy.

Effective blocking of IgE binding to food allergens by specific IgG₄ may be one possible immune mechanism underlying food tolerance in patients with a clear food sensitization.⁴⁴ To evaluate this hypothesis, we used the FAB inhibition assay, a test system often used to test the blocking capacity of SIT-induced allergen-specific IgG antibodies.^{45,46} Accordingly, patients who underwent successful birch pollen SIT had developed IgG antibodies with IgE inhibitory capacity (Fig 3). Sera from 6 patients who tolerated hazelnuts in food challenges and displayed high Cor a 1-specific IgG₄/IgE ratios (Table III) also effectively inhibited the formation of Cor a 1-IgE complexes (Fig 3). The inhibitory activity of all sera could be attributed to the presence of IgG antibodies because depletion of IgG resulted in the loss of IgE-blocking capacity. Although based on a limited number of patients, these data provide the first evidence that naturally occurring food-specific IgG₄ antibodies block IgE binding to the respective allergens. To translate this blocking capacity into clinical food tolerance, the individual IgG₄/IgE ratio may be decisive.

Although often regarded as a nonsevere syndrome, birch pollen-related food allergy is highly prevalent and often perennial. This situation demands effective treatment strategies. However, it is debated whether successful birch pollen SIT concomitantly cures the associated food allergy. Therefore, the immune mechanisms underlying birch pollen-related food allergy should be further investigated to develop efficient treatment strategies for this important allergic disease.

We thank Michaela Gubesch, Renate Höppner, and Katja Parapatics for their excellent technical assistance.

Clinical implications: The vast majority of patients with birch pollen allergy develops allergy to apple and hazelnut. The detection of food-specific IgG₄ is no diagnostic marker for this food allergy.

REFERENCES

- Stevens WJ, Ebo DG, Hagendorens MM, Bridts CH, De Clerck LS. Is the prevalence of specific IgE to classical inhalant aeroallergens among patients with respiratory allergy changing? evidence from two surveys 15 years apart. *Acta Clin Belg* 2003;58:178-82.
- Lin RY, Clauss AE, Bennett ES. Hypersensitivity to common tree pollens in New York City patients. *Allergy Asthma Proc* 2002;23:253-8.
- Tuft L, Blumenstein GI. Studies in food allergy. Sensitization to fresh fruits: clinical and experimental observations. *J Allergy Clin Immunol* 1942;13:574-81.
- Hannuksela M, Lahti A. Immediate reactions to fruits and vegetables. *Contact Dermatitis* 1977;3:79-84.
- Mari A, Ballmer-Weber BK, Vieths S. The oral allergy syndrome: improved diagnostic and treatment methods. *Curr Opin Allergy Clin Immunol* 2005;5:267-73.
- Kleine-Tebbe J, Vogel L, Crowell DN, Hausteiner UF, Vieths S. Severe oral allergy syndrome and anaphylactic reactions caused by a Bet v 1-related PR-10 protein in soybean, SAM22. *J Allergy Clin Immunol* 2002;110:797-804.
- Eriksson NE, Formgren H, Svenonius E. Food hypersensitivity in patients with pollen allergy. *Allergy* 1982;37:437-43.
- Asero R, Massironi F, Velati C. Detection of prognostic factors for oral allergy syndrome in patients with birch pollen hypersensitivity. *J Allergy Clin Immunol* 1996;97:611-6.
- Bohle B. T-cell epitopes of food allergens. *Clin Rev Allergy Immunol* 2006;30:97-108.
- Bolhaar ST, Ree R, Bruijnzeel-Koomen CA, Knulst AC, Zuidmeer L. Allergy to jackfruit: a novel example of Bet v 1-related food allergy. *Allergy* 2004;59:1187-92.
- Oberhuber C, Bulley SM, Ballmer-Weber BK, Bublin M, Gaier S, DeWitt AM, et al. Characterization of Bet v 1-related allergens from kiwifruit relevant for patients with combined kiwifruit and birch pollen allergy. *Mol Nutr Food Res* 2008;52(suppl 2):S230-40.
- Jahn-Schmid B, Radakovics A, Luttkopf D, Scheurer S, Vieths S, Ebner C, et al. Bet v 1142-156 is the dominant T-cell epitope of the major birch pollen allergen

- and important for cross-reactivity with Bet v 1-related food allergens. *J Allergy Clin Immunol* 2005;116:213-9.
13. Bohle B, Zwolfer B, Heratizadeh A, Jahn-Schmid B, Antonia YD, Alter M, et al. Cooking birch pollen-related food: divergent consequences for IgE- and T cell-mediated reactivity in vitro and in vivo. *J Allergy Clin Immunol* 2006;118:242-9.
 14. Valenta R, Duchene M, Pettenburger K, Sillaber C, Valent P, Bettelheim P, et al. Identification of profilin as a novel pollen allergen; IgE autoreactivity in sensitized individuals. *Science* 1991;253:557-60.
 15. Radauer C, Willeroider M, Fuchs H, Hoffmann-Sommergruber K, Thalhamer J, Ferreira F, et al. Cross-reactive and species-specific immunoglobulin E epitopes of plant profilins: an experimental and structure-based analysis. *Clin Exp Allergy* 2006;36:920-9.
 16. Hirschwehr R, Valenta R, Ebner C, Ferreira F, Sperr WR, Valent P, et al. Identification of common allergenic structures in hazel pollen and hazelnuts: a possible explanation for sensitivity to hazelnuts in patients allergic to tree pollen. *J Allergy Clin Immunol* 1992;90:927-36.
 17. Ebner C, Hirschwehr R, Bauer L, Breiteneder H, Valenta R, Ebner H, et al. Identification of allergens in fruits and vegetables: IgE cross-reactivities with the important birch pollen allergens Bet v 1 and Bet v 2 (birch profilin). *J Allergy Clin Immunol* 1995;95:962-9.
 18. van Ree R, Fernandez-Rivas M, Cuevas M, van Wijngaarden M, Aalberse RC. Pollen-related allergy to peach and apple: an important role for profilin. *J Allergy Clin Immunol* 1995;95:726-34.
 19. Scheurer S, Wangorsch A, Hausteine D, Vieths S. Cloning of the minor allergen Api g 4 profilin from celery (*Apium graveolens*) and its cross-reactivity with birch pollen profilin Bet v 2. *J Chromatogr B Biomed Sci Appl* 2000;30:962-71.
 20. Scheurer S, Wangorsch A, Nerkamp J, Skov PS, Ballmer-Weber B, Wuthrich B, et al. Cross-reactivity within the profilin panallergen family investigated by comparison of recombinant profilins from pear (*Pyr c 4*), cherry (*Pru av 4*) and celery (*Api g 4*) with birch pollen profilin Bet v 2. *J Chromatogr B Biomed Sci Appl* 2001;756:315-25.
 21. Reindl J, Rihs HP, Scheurer S, Wangorsch A, Hausteine D, Vieths S. IgE reactivity to profilin in pollen-sensitized subjects with adverse reactions to banana and pineapple. *Int Arch Allergy Immunol* 2002;128:105-14.
 22. Westphal S, Kempf W, Foetisch K, Retzek M, Vieths S, Scheurer S. Tomato profilin Lyc e 1: IgE cross-reactivity and allergenic potency. *Allergy* 2004;59:526-32.
 23. Wellhausen A, Schoning B, Petersen A, Vieths S. IgE binding to a new cross-reactive structure: a 35 kDa protein in birch pollen, exotic fruit and other plant foods. *Z Ernahrungswiss* 1996;35:348-55.
 24. Karamloo F, Wangorsch A, Kasahara H, Davin LB, Hausteine D, Lewis NG, et al. Phenylcoumaran benzylic ether and isoflavonoid reductases are a new class of cross-reactive allergens in birch pollen, fruits and vegetables. *Eur J Biochem* 2001;268:5310-20.
 25. Bircher AJ, Van Melle G, Haller E, Curty B, Frei PC. IgE to food allergens are highly prevalent in patients allergic to pollens, with and without symptoms of food allergy. *Clin Exp Allergy* 1994;24:367-74.
 26. Ruiter B, Knol EF, van Neerven RJ, Garssen J, Bruijnzeel-Koomen CA, Knulst AC, et al. Maintenance of tolerance to cow's milk in atopic individuals is characterized by high levels of specific immunoglobulin G4. *Clin Exp Allergy* 2007;37:1103-10.
 27. Tomicic S, Norrman G, Falth-Magnusson K, Jenmalm MC, Devenney I, Bottcher MF. High levels of IgG4 antibodies to foods during infancy are associated with tolerance to corresponding foods later in life. *Pediatr Allergy Immunol* 2009;20:35-41.
 28. Savilahti EM, Rantanen V, Lin JS, Karinen S, Saarinen KM, Goldis M, et al. Early recovery from cow's milk allergy is associated with decreasing IgE and increasing IgG4 binding to cow's milk epitopes. *J Allergy Clin Immunol* 2010;125:1315-21, e9.
 29. Skripak JM, Nash SD, Rowley H, Brereton NH, Oh S, Hamilton RG, et al. A randomized, double-blind, placebo-controlled study of milk oral immunotherapy for cow's milk allergy. *J Allergy Clin Immunol* 2008;122:1154-60.
 30. Buchanan AD, Green TD, Jones SM, Scurlock AM, Christie L, Althage KA, et al. Egg oral immunotherapy in nonanaphylactic children with egg allergy. *J Allergy Clin Immunol* 2007;119:199-205.
 31. Jones SM, Pons L, Roberts JL, Scurlock AM, Perry TT, Kulis M, et al. Clinical efficacy and immune regulation with peanut oral immunotherapy. *J Allergy Clin Immunol* 2009;124:212-300.
 32. Blumchen K, Ulbricht H, Staden U, Dobberstein K, Beschoner J, de Oliveira LC, et al. Oral peanut immunotherapy in children with peanut anaphylaxis. *J Allergy Clin Immunol* 2010;126:83-91.
 33. Enrique E, Pineda F, Malek T, Bartra J, Basagana M, Tella R, et al. Sublingual immunotherapy for hazelnut food allergy: a randomized, double-blind, placebo-controlled study with a standardized hazelnut extract. *J Allergy Clin Immunol* 2005;116:1073-9.
 34. Fernandez-Rivas M, Garrido Fernandez S, Nadal JA, Diaz de Durana MD, Garcia BE, Gonzalez-Mancebo E, et al. Randomized double-blind, placebo-controlled trial of sublingual immunotherapy with a Pru p 3 quantified peach extract. *Allergy* 2009;64:876-83.
 35. Fernandez-Rivas M, Bolhaar S, Gonzalez-Mancebo E, Asero R, van Leeuwen A, Bohle B, et al. Apple allergy across Europe: how allergen sensitization profiles determine the clinical expression of allergies to plant foods. *J Allergy Clin Immunol* 2006;118:481-8.
 36. Hansen KS, Ballmer-Weber BK, Sastre J, Lidholm J, Andersson K, Oberhofer H, et al. Component-resolved in vitro diagnosis of hazelnut allergy in Europe. *J Allergy Clin Immunol* 2009;123:1134-41, 41 e1-e3.
 37. Shamji MH, Wilcock LK, Wachholz PA, Dearman RJ, Kimber I, Wurtzen PA, et al. The IgE-facilitated allergen binding (FAB) assay: validation of a novel flow-cytometric based method for the detection of inhibitory antibody responses. *J Immunol Methods* 2006;317:71-9.
 38. Ghunaim N, Gronlund H, Kronqvist M, Gronneberg R, Soderstrom L, Ahlstedt S, et al. Antibody profiles and self-reported symptoms to pollen-related food allergens in grass pollen-allergic patients from northern Europe. *Allergy* 2005;60:185-91.
 39. Osterballe M, Hansen TK, Mortz CG, Bindslev-Jensen C. The clinical relevance of sensitization to pollen-related fruits and vegetables in unselected pollen-sensitized adults. *Allergy* 2005;60:218-25.
 40. Asero R, Monsalve R, Barber D. Profilin sensitization detected in the office by skin prick test: a study of prevalence and clinical relevance of profilin as a plant food allergen. *Clin Exp Allergy* 2008;38:1033-7.
 41. Wensing M, Akkerdaas JH, van Leeuwen WA, Stapel SO, Bruijnzeel-Koomen CA, Aalberse RC, et al. IgE to Bet v 1 and profilin: cross-reactivity patterns and clinical relevance. *J Allergy Clin Immunol* 2002;110:435-42.
 42. Bolhaar ST, van de Weg WE, van Ree R, Gonzalez-Mancebo E, Zuidmeer L, Bruijnzeel-Koomen CA, et al. In vivo assessment with prick-to-prick testing and double-blind, placebo-controlled food challenge of allergenicity of apple cultivars. *J Allergy Clin Immunol* 2005;116:1080-6.
 43. Stapel SO, Asero R, Ballmer-Weber BK, Knol EF, Strobel S, Vieths S, et al. Testing for IgG4 against foods is not recommended as a diagnostic tool: EAACI Task Force Report. *Allergy* 2008;63:793-6.
 44. Aalberse RC, Stapel SO, Schuurman J, Rispens T. Immunoglobulin G4: an odd antibody. *Clin Exp Allergy* 2009;39:469-77.
 45. Wachholz PA, Soni NK, Till SJ, Durham SR. Inhibition of allergen-IgE binding to B cells by IgG antibodies after grass pollen immunotherapy. *J Allergy Clin Immunol* 2003;112:915-22.
 46. Durham SR, Emminger W, Kapp A, Colombo G, de Monchy JG, Rak S, et al. Long-term clinical efficacy in grass pollen-induced rhinoconjunctivitis after treatment with SQ-standardized grass allergy immunotherapy tablet. *J Allergy Clin Immunol* 2010;125:131-8, e1-e7.

Standardized questionnaire to evaluate the cause and symptoms of birch pollen-related food allergy**1. Do you have hay fever (allergic rhinoconjunctivitis) during the months March - May?**Yes ☐ No ☐**2. Have you noticed any reaction during and/or after consumption of:**

	Yes	No	Never tried
Hazelnuts			
Walnuts			
Peanuts			
Almonds			
Apple			
Pear			
Peach/ Nectarine			
Apricot			
Cherry			
Plum			
Celery (raw)			
Soy/Soymilk			
Kiwi			
Tomato			
Carrot (raw)			
Potato (raw, reaction during peeling)			
Others:			

3. What kind of reactions to these foods have you experienced?

	Yes	No
Itching in the mouth		
Itching in the ears		
Swelling of lips		
Lump in the throat		
Sneezing and/or running nose		
Streaming eyes		
Rush on the hands after contact		
Body rush after consumption		
Swelling of the face		
Difficulties in breathing		
Abdominal cramping		
Nausea and/or emesis		
Drop of the blood pressure		
Others:		

4. Are food-induced reactions worsening during the pollen season in spring?Yes ☐ No ☐**5. Do you experience food-induced reactions in winter as well?**Yes ☐ No ☐**6. Did you notice food-induced reactions before or after you have developed birch pollen allergy?**Before ☐ After ☐**7. Do you have neurodermitis?**Yes ☐ No ☐**FIG E1.** Standardized questionnaire to evaluate the cause and symptoms of birch pollen-related food allergy.