


Journal Pre-proof

Tick peptides evoke itch by activating MrgprC11/X1 to sensitize TRPV1 in pruriceptors

Xueke Li, MSc, Haifeng Yang, MSc, Yuewen Han, MSc, Shijin Yin, PhD, Bingzheng Shen, PhD, Yingliang Wu, PhD, Wenxin Li, PhD, Zhijian Cao, PhD



PII: S0091-6749(20)32413-1

DOI: <https://doi.org/10.1016/j.jaci.2020.12.626>

Reference: YMAI 14905

To appear in: *Journal of Allergy and Clinical Immunology*

Received Date: 4 May 2020

Revised Date: 21 November 2020


Accepted Date: 2 December 2020

Please cite this article as: Li X, Yang H, Han Y, Yin S, Shen B, Wu Y, Li W, Cao Z, Tick peptides evoke itch by activating MrgprC11/X1 to sensitize TRPV1 in pruriceptors, *Journal of Allergy and Clinical Immunology* (2021), doi: <https://doi.org/10.1016/j.jaci.2020.12.626>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Published by Elsevier Inc. on behalf of the American Academy of Allergy, Asthma & Immunology.

Tick peptides evoke itch by activating MrgprC11/X1 to sensitize TRPV1 in pruriceptors



1 Tick peptides evoke itch by activating MrgprC11/X1 to sensitize
2 TRPV1 in pruriceptors

3
4
5 Xueke Li, MSc^{1,§}, Haifeng Yang, MSc^{1,§}, Yuewen Han, MSc¹, Shijin Yin, PhD²,
6 Bingzheng Shen, PhD¹, Yingliang Wu, PhD¹, Wenxin Li, PhD¹, Zhijian Cao, PhD^{1,3,4,¶}
7

8
9
10 ¹ State Key Laboratory of Virology, College of Life Sciences, Wuhan University,
11 Wuhan 430072, P. R. China

12 ² School of Pharmaceutical Sciences, South-Central University for Nationalities,
13 Wuhan 430074, P. R. China.

14 ³ Bio-drug Research Center, Wuhan University, Wuhan 430072, P. R. China

15 ⁴ Hubei Province Engineering and Technology Research, Center for Fluorinated
16 Pharmaceuticals, Wuhan University, Wuhan 430072, P. R. China
17

18
19 Disclosure of potential conflict of interest:

20 The authors declare that they have no relevant conflicts of interest.
21
22

23 [§] These authors contributed equally

24 [¶] Correspondence: State Key Laboratory of Virology, College of Life Sciences,
25 Wuhan University, Wuhan 430072, P. R. China. Tel: +86-27-68752831, Fax:
26 +86-27-68756746. E-mail: zjcao@whu.edu.cn (Zhijian Cao)
27

Abstract

Background Tick bites severely threaten human health because they allow the transmission of many deadly pathogens, including viruses, bacteria, protozoa and helminths. Pruritus is a leading symptom of tick bites, but its molecular and neural bases remain elusive.

Objective To discover potent drugs and targets for the specific prevention and treatment of tick bite-induced pruritus and arthropod-related itch.

Methods We used live-cell calcium imaging, patch-clamp recordings, and genetic ablation and evaluated mouse behavior to investigate the molecular and neural bases of tick bite-induced pruritus.

Results We found that two tick salivary peptides, IPDef1 and IRDef2, induced itch in mice. IPDef1 was further revealed to have a stronger pruritogenic potential than IRDef2 and to induce pruritus in a histamine-independent manner. IPDef1 evoked itch by activating mouse MrgprC11 and human MrgprX1 on dorsal root ganglion (DRG) neurons. IPDef1-activated MrgprC11/X1 signaling sensitized downstream ion channel TRPV1 on DRG neurons. Moreover, IPDef1 also activated mouse MrgprB2 and its ortholog human MrgprX2 selectively expressed on mast cells, inducing the release of inflammatory cytokines and driving acute inflammation in mice, although mast cell activation did not contribute to IP-O-induced itch.

Conclusion Our study identifies tick salivary peptides as a new class of pruritogens that initiate itch through MrgprC11/X1-TRPV1 signaling in pruritoseptors. Our work will provide potential drug targets for the prevention and treatment of pruritus induced by the bites or stings of tick and maybe other arthropods.

Key words Tick; Peptide; Itch; Mrgprs; TRP channel

Abbreviations

IPDef1: IP defensin 1

IP-O: Oxidated form of IPDef1

IRDef2: IR defensin 2

MRGPR: Mas-related G protein coupled receptor

- 59 DRG: Dorsal root ganglion
60 TRPV1: Transient receptor potential vanilloid 1
61 RP-HPLC: Reverse-phase high-pressure liquid chromatography
62 α -CHCA: α -cyano-4-hydroxycinnamic acid
63 TFA: Trifluoroacetic acid
64 HEK293T: Human embryonic kidney 293T
65 MALDI-TOF MS: Matrix-assisted laser desorption ionization time-of-flight mass
66 spectrometry
67 CCTCC: China Center for Type Culture Collection
68 MIC: Minimum inhibitory concentration
69 CQ: Chloroquine
70 HBSS: Hank's balanced salt solution
71 HEPES: Hydroxyethyl piperazineethanesulfonic
72 Cap: Capsaicin
73 AITC: Allyl isothiocyanate
74 KO: Knockout
75 PBS: Phosphate buffer saline
76 SDS: Sodium dodecyl sulfate
77 SDS-PAGE: Sodium dodecyl sulfate-polyacrylamide gel electrophoresis
78 MCDM: Mast cell dissociation media
79 mSCF: Mouse stem cell factor
80 DMEM: Dulbecco's modified eagle medium
81 ELISA: Enzyme linked immune sorbent assay
82 MCP-1: Monocyte chemotactic protein 1
83 TNF- α : Tumor necrosis factor α
84 qRT-PCR: Quantitative real time PCR
85 GAPDH: Glyceraldehyde-3-phosphate dehydrogenase
86 SEM: Standard error of the mean
87 MW: Molecular weight
88 CS $\alpha\beta$: Cysteine-stabilized α -helical and β -sheet
89 GFP: Green fluorescent protein
90 EC₅₀: Concentration for 50% of maximal effect
91 CRISPR: Clustered regularly interspaced short palindromic repeats
92 WT: Wild-type

93 GPCR: G protein-coupled receptor
94 TRPA1: Transient receptor potential A1
95 5-HT: 5-hydroxy tryptamine
96 PMC: Peritoneal mast cell
97 CPA3: Carboxypeptidase A3
98 TLR4: Toll-like receptor 4

99

100 **Key Messages**

- 101 ● Two tick salivary peptides IPDef1 and IRDef2 induce itch in mice via a
102 histamine-independent pathway.
- 103 ● IPDef1 evokes itch by activating MrgprC11/X1 to sensitize downstream TRPV1
104 on DRG neurons.
- 105 ● IPDef1 activates MrgprB2/X2 on mast cells to cause acute inflammation.

106

107 **Capsule Summary**

108 Anti-histamine treatment does not improve the itching symptoms of patients bitten by
109 ticks. Tick salivary peptides induce itch via the MrgprC11/X1-TRPV1 signaling
110 pathway, providing potential drug targets for treatment of the disease.

111

Introduction

Pruritus is a dermatological symptom involving skin itching but no primary skin damage and is the most common clinical manifestation of skin diseases^{1, 2}. There are numerous factors that can induce pruritus, such as cold, warmth, chemical fibers, bites of ticks and insects, diabetes, liver disease and kidney disease. In addition to the H1/H4 receptor, Mas-related G protein coupled receptors (Mrgprs) and Piezo2 channels were recently identified as itch related membrane proteins^{3, 4}. However, itch is still an unmet clinic problem with no universal treatment because the molecular, cellular and neural circuit mechanisms of itch have not been fully understood. Therefore, it is important to identify new and specific pruritogens to dissect molecular and cellular mechanisms of itch as well as unravel the specificity and selectivity of itch receptors.

The bites or stings of many arthropods represent one of the most common causes of itching^{5, 6}. These arthropods mainly include insects and arachnids, such as fleas, mosquitoes, bedbugs, bees, wasps, mites and ticks. Each of them has thousands of species on the earth, constituting a large group of itch-related organisms. Ticks are small arachnids that belong to the order Ixodida of the class Arachnida. There are approximately 900 tick species in the world. Blood-sucking ticks attack various types of vertebrates. Moreover, some tick species carry pathogens such as viruses and rickettsiae, which can infect humans and animals^{7, 8}. Tick bites can lead to local lesions and systemic illness, referred to as tick toxicosis. Pruritus is a leading symptom of tick toxicosis. Patients bitten by lone star ticks exhibit skin manifestations, specifically a large number of pruritic papules⁹. Dogs bitten by the mouro tick *Ornithodoros brasiliensis* also present skin rash and itch symptoms¹⁰. However, the molecular and neural bases of tick bite-induced pruritus is largely unknown.

The saliva or venoms of the itch-inducing arthropods contain various toxic peptides used for prey and defense that exhibit extremely diverse primary sequences, spatial structures, targeting receptors and biological functions¹¹⁻¹⁵. It is possible that these arthropods may produce a class of common peptides that induce itch in humans and animals. Previous reports showed that the class of the ancient invertebrate defensin could serve as a common peptide component in the saliva or venoms of the itch-inducing arthropods^{16, 17}. We speculated that these ancient invertebrate defensin peptides may be potent candidate pruritogens.

In this study, two tick salivary defensin peptides, IPDef1 and IRDef2, were found to induce histamine-independent itch in mice while IPDef1 had a stronger activity. IPDef1 produced itch through directly activating dorsal root ganglion (DRG) neurons and triggering Ca^{2+} influx. Using live cell calcium imaging, patch-clamp recordings, co-immunoprecipitation and gene editing, mouse MrgprC11 and human MrgprX1 were identified as the main itchy receptors for IPDef1 on DRG neurons. The MrgprC11/X1-TRPV1 axis in DRG neurons was an important signaling pathway for IPDef1-induced itch. Interestingly, IPDef1 also activated mouse MrgprB2 and its human ortholog MrgprX2 selectively expressed on mast cells, thereby causing inflammatory cytokine release and inducing acute inflammation in mice. Unexpectedly, mast cell activation by IPDef1 did not contribute to its itch-inducing activity. Our study discloses the molecular and cellular basis of itch induced by the tick salivary peptides and provides potential drug targets for the prevention and treatment of pruritus induced by the bites or stings of arthropods such as ticks, mosquitoes and ants.

Methods

Oxidative refolding and homology modeling

Reduced IPDef1 and IRDef2 were synthesized by ChinaPeptides Co., Ltd. (China), and the purity of each peptide was greater than 97%. To form three disulfide linkages via intermolecular oxidative refolding, the reduced peptides (1 mg) were dissolved in 2 mL Tris-HCl buffer (0.1 M, pH 8.0) and incubated at 25 °C for 48 h with continuous shaking at 50 rpm. The oxidized peptides were centrifuged at 12,000 rpm for 10 min at 4 °C, and the supernatants were purified by reverse-phase high-pressure liquid chromatography (RP-HPLC) (Agilent, USA). The average molecular mass of each oxidized peptide was confirmed by matrix-assisted laser desorption ionization time-of-flight mass spectrometry (MALDI-TOF MS) (BiflexIII, Bruker, Daltonik GmbH, Bremen, Germany). Oxidized IPDef1 and IRDef2, which were desalinated and purified by RP-HPLC, were mixed with MALDI-matrix solution (1 mL, containing 10 mg/mL α -cyano-4-hydroxycinnamic acid (α -CHCA), 0.1% trifluoroacetic acid (TFA) and 45% acetonitrile). Then, 1 μL of each peptide sample mixture was spotted onto a MALDI target plate and left to air dry at room temperature. Mass spectrometry was performed with FlexControl software (Version 3.0, Bruker

Daltonics) for a mass range of m/z from 1,000 to 8,000 Da. The mass of the oxidized peptide was measured in positive-ion linear mode at an accelerating voltage of 25 kV. The secondary structure of each peptide was determined by circular dichroism (CD) spectroscopy using a JASCO J-810 spectrometer (JASCO International Co., Ltd., Japan). The peptides were dissolved in Milli-Q water at a concentration of approximately 200 $\mu\text{g/mL}$. CD spectra were obtained at wavelengths from 190 nm to 260 nm at room temperature (25 °C). The scanning speed was 50 nm/min, the resolution was 1 nm, and the response time was 2 sec. Each reading was repeated three times, and the results are shown as the mean residue molar ellipticity (θ). The 3D-structure prediction was determined using SWISS-MODEL work-space (<http://swissmodel.expasy.org>).

***In vitro* antimicrobial assays**

Reference strains of gram-positive bacteria and gram-negative bacteria were used to evaluate the *in vitro* antimicrobial activity of IPDef1 (IP-R and IP-O). *Staphylococcus aureus* AB94004, *S. aureus* ATCC25923, *S. aureus* ATCC6538, *Micrococcus luteus* AB93113, *Bacillus subtilis* AB91021, *Escherichia coli* AB94012 and *E. coli* ATCC25922 were purchased from the China Center of Type Culture Collection (CCTCC). The antimicrobial activities of IP-R and IP-O *in vitro* were evaluated by a two-fold serial dilution method as recommended by CLSI guidelines. The strains, which were stored in a refrigerator at -80 °C, were inoculated into solid medium plates and cultivated at 37 °C overnight in a thermostatic incubator. A single colony was selected and subcultured in liquid medium. After overnight culturing and activation, the test strains were diluted with medium to 10^4 - 10^6 CFU/mL. Then, 20 μL peptide at various concentrations was added to 80 μL diluted culture medium containing the test strains for a total volume of 100 μL . The 96-well microplates were incubated at 37 °C with continuous shaking at 100 rpm for 14-16 h, and the absorbance at 630 nm was measured to determine the minimum inhibitory concentration (MICs). The MIC was defined as the lowest peptide concentration that completely prevented growth and was measured with a microtiter optical plate reader. To monitor the validity and reproducibility of the assays, incubations were performed in triplicate with three parallel replicates.

Behavioral studies

Two- to 3-month-old male mice (C57BL/6, 20-30 g) were housed on a 12-h light-dark cycle at 24 °C. On the day of the experiment, the animals were allowed to acclimatize to the test chamber for 10 minutes prior to injection. A pruritic substance (i.e., IPDef1, IRDef2, histamine, CQ, PAMP9-20 or anti-IgE) was intradermally injected into the nape of the neck after acclimatization. A bout of scratching was defined as an episode in which a mouse lifted its paw and scratched directly at the area around the injection site continuously for any length of time and lasted until the paw was returned to the floor. The use of both forepaws was classified as grooming behavior and was not considered scratching. Scratching behavior was quantified by counting the number of scratching bouts during the 30-min observation period. An antagonist (i.e., cetirizine, JNJ7777120, AMG9810, or HC030031) or the mast cell stabilizer cromolyn sodium were intraperitoneally injected 30 min before the injection of the pruritic substances. All behavioral tests were performed by an experimenter blind to genotype. All experiments were performed under the policies and recommendations of the Institutional Animal Care and Use Committee of Wuhan University. Antagonists above were purchased from Targetmol.

Gene amplification

Since Mrgprs are expressed mainly in the peripheral and central nervous systems, we extracted total RNA from mouse DRG neurons using TRIzol reagent (BBI, Toronto, Canada). Then, total RNA was reverse-transcribed into the 1st strand cDNA using the First Strand cDNA Synthesis Kit (Thermo Scientific, USA). Subsequently, the synthesized cDNAs were used as templates for random primer p(dN)6 amplification by PCR. The mouse and human Mrgpr genes were amplified and characterized by PCR. Experiments involving tissue were also performed under the policies and recommendations of the Institutional Animal Care and Use Committee of Wuhan University.

DRG neuron culture

DRG neurons from all spinal levels were collected from 4- to 5-week-old mice, placed in cold HBSS and treated with enzyme solution at 37 °C. Briefly, neurons from sensory ganglia were dissected and incubated for 10 min in 1.4 mg/mL collagenase P (Roche) in Hanks calcium-free balanced salt solution. The neurons were then incubated in 0.25% standard trypsin (vol/vol) STV versene-EDTA solution for 3 min

with gentle agitation. After trituration and centrifugation, the cells were resuspended in media (Eagle's MEM with Earle's BSS medium supplemented with 10% horse serum (vol/vol), MEM vitamins, penicillin/streptomycin and l-glutamine), plated on glass coverslips coated with poly-D-lysine, cultured in an incubator at 37 °C, and used within 18 h. All results were also confirmed using neuronal cultures from adult mice.

HEK293T cell culture

HEK293T cells were cultured on poly-D-lysine-coated glass coverslips. The cells were transfected with 500 ng mouse *Mrgprs*, 500 ng human *Mrgprs*, 250 ng human *Trpa1*, or 250 ng human *Trpv1* plasmids with Tubofect (Invitrogen). The cells were replated on glass coverslips 20 h after transfection and used for Calcium imaging or patch-clamp recordings.

Calcium imaging

DRG neurons or HEK293T cells were loaded for 30-45 min in the dark with 10 μ M Fura-2AM (Yeasen Biotech Co., Ltd) supplemented with 0.01% Pluronic F-127 (wt/vol, Yeasen Biotech Co., Ltd) in physiological Ringer's solution containing 140 mM NaCl, 5 mM KCl, 10 mM HEPES, 2 mM CaCl_2 , 2 mM MgCl_2 and 10 mM d-(+)-glucose, pH 7.4.. After washing, the cells were imaged at an excitation wavelength of 340 and 380 nm to detect intracellular free calcium. Cells were considered to have exhibited a response if the $[\text{Ca}_2^+]_i$ rose by at least 30% for at least 10 sec allowing us to clearly distinguish ligand-induced responses from random flickering events. Each experiment was performed at least three times with at least 100 neurons or HEK293T cells were analyzed. For the assay of calcium response traces, each colored line represents an individual DRG neuron or HEK293T cell. For the assay of EC_{50} value or the response prevalence, calcium responses at each concentration or substance were normalized to the maximal response elicited subsequently. Each point represents data collected from an independent experiment. KCl (50 mM) was used to identify live cells. BAM8-22 (50 μ M), CQ (1 mM) or PAMP9-20 (20 μ M) was used as an agonist to identify the functional *MrgprC11*, *MrgprA3* or *MrgprB2*, respectively. Capsaicin (1 μ M) or allyl isothiocyanate (100 μ M) was used as a channel agonist to identify the functional TRPV1 or TRPA1,

respectively. Calcium imaging assays were performed by an experimenter blind to genotype or pretreatment.

Whole-cell patch-clamp recordings

HEK293T cells plated on coverslips were transferred to a chamber with extracellular solution. Patch pipettes with a resistance of 2-4 MU were used. For current-clamp recordings, action potentials were measured with an Axon 700B amplifier and the pCLAMP 9.2 software package (Axon Instruments). Cells were perfused with IP-O (10 μ M) for 30 s, and Cap (1 μ M) or AITC (100 μ M) was used as a positive control. All experiments were performed at room temperature (25 °C).

Knockout mice

The three knockout mouse strains we used were on the C57BL/6 background. MrgprC11 knockout mice were generated by the following strategy (Cyagen Biosciences). The MrgprC11 gene (NCBI Reference Sequence: NM_207540; Ensembl: ENSMUSG00000070552) is located on mouse chromosome 7. Two exons with the ATG start codon in exon 2 and the TGA stop codon in exon 2 were identified (Transcript: ENSMUST00000094390). Exon 2 was selected as the target site. Cas9 and gRNA were coinjected into fertilized eggs to produce KO mice. The pups were genotyped by PCR followed by sequencing analysis. Exon 2 starts from approximately 0.1% of the coding region. Exon 2 covers 100.0% of the coding region. The size of the effective KO region was 964 bp. The KO region did not contain any other known gene. *Trpa1*^{-/-} and *Trpv1*^{-/-} mice were obtained from Jackson Laboratory.

Immunoprecipitation

Cells transfected with the plamid pcDNA3.1 expressing N-flag-tagged Mrgpr receptors (mouse MrgprA3/C11 and human MrgprX1/X2/X3/X4) and TRP channels (TRPV1 and TRPA1) were washed with ice-cold PBS, and proteins were extracted according to the manufacturer's instructions. Con (control) represents HEK293T cells transfected with the plamid pcDNA3.1-Flag. After centrifugation, an adequate amount of soluble His-IP-O was added to the supernatant, and the mixture was incubated with rotation for 4 h at a temperature of 4 °C. Subsequently, 8 μ L protein G beads and 0.5 μ L anti-Flag antibody (Sigma) were added, and the samples were

incubated with rotation overnight. After overnight incubation, the Sepharose beads were washed three times with ice-cold modified NHG buffer and resuspended in 5 × SDS sample buffer. The samples were separated by SDS-PAGE and then transferred onto Immobilon-P membranes (Millipore) for western blot analysis.

Peritoneal mast cell purification and imaging

Three- to four-month-old adult C57BL/6 male and female mice were killed by CO₂ inhalation. Ice-cold mast cell dissociation media (MCDM; HBSS with 3% fetal bovine serum and 10 mM HEPES, pH 7.2) was used to perform two sequential peritoneal lavages; the media from these lavages were combined, and the cells were spun down at 200× *g*. The pellet from each mouse was resuspended in 2 mL MCDM, layered on top of 4 mL isotonic 70% Percoll suspension (2.8 mL Percoll, 320 μL 10 × HBSS, 40 μL 1 M HEPES, 830 μL MCDM), and spun down for 20 min with 500× *g* at 4 °C. Mast cells were recovered in the pellet. Purity was assayed by toluidine blue staining. Mast cells were resuspended at a concentration of 5×10^5 - 1×10^6 cells/mL in DMEM with 10% fetal bovine serum and 25 ng/mL recombinant mouse stem cell factor (Sigma) and allowed to recover for 2 h in a 37 °C incubator with 5% CO₂. The cells were then spun down, resuspended in HBSS, counted, and plated at a concentration of 300 cells/well in 75 μL HBSS in 96-well plates. After 2 h of incubation at 37 °C and 5% CO₂, the mast cells were used for calcium imaging according to the methods described above.

Hindpaw swelling and Evans blue extravasation

Adult male C57BL/6 mice were anesthetized by intraperitoneal injection of 0.2 mL chloral hydrate (3.5%). Fifteen minutes after induction of anesthesia, the mice were injected with 50 μL 12.5 mg/mL Evans blue (Sigma) in saline through the tail vein. Five minutes after Evans blue injection, the test substance (IP-O, 2 mg/mL, 5 μL) was intraplantarly injected into one hindpaw of each mouse, and saline (5 μL) was injected into the other hindpaw. Paw thickness was measured by Vernier calipers immediately after injection. Fifteen minutes later, paw thickness was measured again, and the mice were killed by decapitation. The hindpaws of the mice were imaged, and paw tissues were collected, dried for 24 h at 50 °C, and weighed. Evans blue was extracted by 24-hour incubation in formamide at 50 °C, and the OD values were read at 620 nm using a spectrophotometer. The concentration of Evans blue dye was determined

based on the corresponding standard curve and expressed as ng/mg of tissue weight. The mice were initially treated with PBS or cromolyn sodium (25 mg/kg) for 3 days by intraperitoneal injection. On the fourth day, the mice were subjected to the above experiment.

Enzyme linked immune sorbent assay (ELISA)

1×10^4 - 5×10^4 mast cells were incubated with test compound for 30 minutes before supernatant was collected. Supernatants were stored at -80 °C until used for ELISA. All data were normalized according to cell number. Histamine was detected with an ELISA Kit from Abcam according to manufacturer's instructions. Tryptase beta 2, serotonin, monocyte chemotactic protein 1 (MCP-1) and tumor necrosis factor- α (TNF- α) were analyzed with ELISA Kits from Cusa Bio. Each dot represents an independent biological replicates from PMCs isolated from >4 animals.

Quantitative real time PCR (qRT-PCR)

Total RNA was extracted from PMCs or hindpaw tissues using TRIzol reagent (Takara), and the first-strand cDNA was reversely transcribed by using the RevertAid First Strand cDNA Synthesis Kit (ThermoFisher Scientific). The cDNAs of the tested cytokines and chemokines were quantitated by qRT-PCR using the Bestar® SybrGreen qPCR master mix reagent (DBI® Bioscience). The shown data represent the relative abundance of the indicated RNA normalized to that of GAPDH. The nucleic acid stain (Super GelRed, no.: S-2001) was purchased from US Everbright Inc. The qRT-PCR primer sequences for the test cytokines and chemokines were shown in Table S1. All qRT-PCR experiments were performed on an ABI 7500 system according to the manufacturer's instructions.

Statistical analysis

The data are presented as the mean $\pm$ standard error of the mean (SEM). Statistical comparisons were made with unpaired Student's t test, and differences were considered significant at $P < 0.05$.

Results

Preparation, structural features and antimicrobial activity of the tick salivary

peptide IPDef1.

Since there have been some clinical cases of itch caused by the bites of hard ticks, we wondered whether the salivary peptide IPDef1 from the tick *Ixodes persulcatus* (Fig. 1A) can induce itching and scratching responses in mice. First, we chemically synthesized the reduced form of IPDef1 (IP-R) and then folded it by air oxidation in slightly alkaline Tris-HCl buffer. The oxidized product of IPDef1 (IP-O) was purified to homogeneity by reverse-phase high-performance liquid chromatography (RP-HPLC) and was eluted at a retention time (T_R) of 20.4 min, which was 1.4 min later than the reduced form was eluted (T_R of 19.0 min) (Fig. 1B), indicating that the reduced form and oxidized product of IPDef1 have different polarities. This difference suggests that the formation of an intramolecular disulfide bond can decrease the polarity of this peptide. To verify disulfide bond formation, we analyzed the reduced form and oxidized product of IPDef1 using matrix-assisted laser desorption/ionization time-of-flight mass spectrometry (MALDI-TOF MS). The results showed that the mass-to-charge ratios of IP-O and IP-R were 4195.1 and 4201.5, respectively (Fig. 1C). The measured molecular weight (MW) of the oxidized product (IP-O) was 4194.1 Da, which was 6.4 Da less than the MW of the reduced form (IP-R, 4200.5 Da). These data indicate that six hydrogen atoms of the cysteines of the reduced peptide were removed when three disulfide bridges were formed. Additionally, analysis of circular dichroism indicated that IP-O displayed a minimum at 208 nm and a maximum at 198 nm (Fig. S1A), demonstrating that the reduced form of IPDef1 folds into a native-like conformation similar to that of other peptides with cysteine-stabilized α -helical and β -sheet ($CS\alpha\beta$) structures. In contrast to that of IP-O, the secondary structure of IP-R was mainly dominated by irregular coils (Fig. S1A). The 3D structure of IPDef1 was modeled using the structure of the defensin MGD-1 as a template (PDB: 1FJN) with the SWISS-MODEL server. The predicted structure of IPDef1 had one α -helix domain at the N-terminus and two β -sheet domains at the C-terminus (Fig. S1B).

As in representative invertebrate defensins, the connectivity of three disulfide bridges (Cys⁴-Cys²⁵, Cys¹¹-Cys³³, and Cys¹⁵-Cys³⁵) formed the core skeleton of IPDef1 (Fig. S1B). In addition, IP-O presented nearly the same secondary structure in different solutions, including water, 0.9% sodium chloride and PBS (Fig. S1C), suggesting that IP-O has a stable structure in different solutions. Because diverse bacteria are present in the habitats of ticks and because IPDef1 belongs to the ancient

invertebrate-type defensin family and contains six disulfide-paired cysteines, we performed minimum inhibitory concentration (MIC) experiments to investigate the antimicrobial activities of IPDef1. IP-O effectively inhibited the growth of five tested standard gram-positive bacteria, exhibiting MIC values of 0.5-2 μ M, whereas its reduced form IP-R showed much weaker bioactivity against these bacteria, showing MIC values of 6-16 μ M (Fig. S2). However, both forms of IPDef1 (IP-R and IP-O) did not seem to exert bioactivity against two tested standard gram-negative bacteria, exhibiting MIC values of more than 30 μ M (Fig. S2). These results suggest that only the oxidized form of IPDef1 (IP-O) exerts excellent antibacterial effects against gram-positive bacteria. Additionally, the conformation of CS $\alpha\beta$ is closely related to the antimicrobial activity of IPDef1.

IPDef1 causes histamine-independent itch in mice.

After preparing IPDef1, we examined whether the peptide can induce itching and scratching responses in mice. The reduced and oxidized forms of IPDef1 (IP-R and IP-O) were intradermally injected into the cheeks of mice, and IP-O but not IP-R elicited significant scratching behavior (vehicle, 11.7 ± 4.0 ; IP-R, 23.17 ± 6.1 ; IP-O, 116.6 ± 9.9 ; $P < 0.0001$; Fig. 1D). Moreover, IP-O induced scratching behavior in a dose-dependent manner (Fig. 1E). The best-characterized type of itch in humans and rodents, histamine-dependent itch, can be triggered by histamine (HIS)¹⁸. Histamine is mainly secreted by skin mast cells and excites nearby sensory fibers by acting on histamine receptors¹⁹. In our study, there was no significant difference in the total number of scratching bouts elicited by IP-O and that induced by histamine (IP-O, 116.6 ± 9.9 ; histamine, 118.4 ± 8.6 ; $P = 0.2$; Fig. 1D). These data suggest that IPDef1 probably has strong pruritogenic potential and that this potential may be dependent on its secondary structure. Considering that histamine induces pruritus in a histamine-dependent manner^{20, 21}, we wanted to know whether IP-O induces scratching responses in mice via a histamine-dependent pathway. Histamine-induced itch can be almost completely blocked by histamine receptor H1/4 antagonists. The H1R antagonist cetirizine (CETY, 10 mg/kg)²¹ and the H4R antagonist JNJ7777120 (JNJ, 40 mg/kg) were administered intraperitoneally 30 min prior to the injection of IP-O or HIS. Compared with vehicle (saline), CETY and JNJ significantly reduced histamine-induced scratching (vehicle, 116.5 ± 8.8 ; CETY, 64.0 ± 6.4 ; $P = 0.0007$; JNJ, 69.5 ± 7.8 ; $P = 0.0017$), but both failed to reduce IP-O-induced scratching

behavior (vehicle, 114.7 ± 13.5 ; CETY, 92.6 ± 12.6 ; $P = 0.2627$; JNJ, 93.5 ± 8.8 ; $P = 0.2332$) (Fig. 1F). These results suggest that IP-O may cause histamine-independent itch in mice. We selected another tick salivary peptide, IRDef2 from *I. ricinus*, to determine the universal ability of tick salivary peptides to cause itch^{22, 23}. Consistently, compared to vehicle and the reduced form of IRDef2 (IR-R), the oxidized product of IRDef2 (IR-O) elicited significant scratching behavior (vehicle, 7.2 ± 2.0 ; IR-R, 15.0 ± 2.0 ; IR-O, 69.7 ± 5.5 ; Fig. S3).

IP-O activates DRG neurons with extracellular Ca^{2+} influx.

To further investigate the neural mechanism underlying IP-O-induced itch, we examined whether the peptide IP-O directly acts on mouse DRG neurons. Consistent with the behavioral data (Fig. 1D), IP-O but not IP-R induced a robust increase in $[\text{Ca}^{2+}]_i$ in DRG neurons, exhibiting an EC_{50} value of $1.47 \pm 0.74 \mu\text{M}$ (Fig. 2A, B and C); this finding indicates that IPDef1 directly acts on DRG neurons to evoke itch through a mechanism dependent on the proper folding of the secondary structure of the peptide. This increase in $[\text{Ca}^{2+}]_i$ in cultured DRG neurons was also seen in representative Fura-2 ratiometric images showing IP-R-evoked ($10 \mu\text{M}$, white arrowheads) and IP-O-evoked ($10 \mu\text{M}$, yellow arrowheads) responses (Fig. 2C). Notably, approximately 7% of the cultured mouse DRG neurons evoked by IP-O ($10 \mu\text{M}$) exhibited a robust increase in $[\text{Ca}^{2+}]_i$ in each experiment, which was similar to the percentage of DRG neurons that exhibited an increase in $[\text{Ca}^{2+}]_i$ following treatment with CQ or BAM8-22 (BAM) (IP-O, $7.5 \pm 1.0\%$; CQ, $6.8 \pm 0.9\%$; BAM, $5.8 \pm 0.9\%$; $P = 0.2413$; Fig. 2D). Previous studies have shown that some pruritogens, including CQ and BAM, mediate itch sensation through activating a highly restricted population of small-diameter neurons in the DRG^{3, 24, 25}. In addition, the specific neurons that selectively detect itch-inducing chemicals and peptides comprise approximately 5% of all DRG neurons. These results suggest that the IP-O-evoked increase in $[\text{Ca}^{2+}]_i$ seen in DRG cultures reflects the activation of a specific subset of DRG neurons, which may be the same population activated by CQ and/or BAM. We performed further experiments to characterize the increase in $[\text{Ca}^{2+}]_i$ by extracellular Ca^{2+} influx or intracellular Ca^{2+} store release in mouse DRG neurons. Extracellular Ca^{2+} was found to be necessary for the increase in $[\text{Ca}^{2+}]_i$ induced by IP-O- and CQ, but not that induced by BAM, because the effects of these two substances were almost completely abolished in Ca^{2+} -free bath solution (Fig. 2E). This result suggests that in

the absence of extracellular Ca^{2+} , IP-O and CQ, unlike BAM, did not mobilize Ca^{2+} release from intracellular stores. However, IP-O, CQ or BAM application in the presence of extracellular Ca^{2+} triggered Ca^{2+} influx across the plasma membrane (Fig. 2E). These data show that IP-O acts on DRG neurons and triggers the influx of Ca^{2+} through transduction channels on the plasma membrane.

Mouse MrgprC11 and human MrgprX1 are the main itch receptors for IP-O on DRG neurons.

IP-O directly acts on primary sensory neurons to evoke itch, and the proportion of IP-O-sensitive neurons among total DRG neurons is similar to the proportions of neurons activated by the two well-known mrgpr-dependent pruritogens CQ and BAM. Thus, it can be inferred that IP-O-induced itch is mediated by an Mrgpr-dependent neural pathway. We cloned each of the 12 mouse Mrgpr gene that have been reported to be itch-related functional receptors into a mammalian expression vector and transfected them individually into human embryonic kidney 293T (HEK293T) cells. By fusing green fluorescent protein (GFP) to the C-terminus of the Mrgpr coding sequences, we were able to visualize the transfected cells and confirm the proper membrane localization of the receptors. Then, we examined the effects of IP-O on the 12 mouse Mrgprs by calcium imaging. The results showed that $22.4 \pm 2.0\%$ MrgprA3-overexpressing HEK293T cells and $76.4 \pm 2.0\%$ MrgprC11-overexpressing HEK293T cells responded to IP-O ($10 \mu\text{M}$). MrgprC11 conferred the strongest responses to the peptide, with an EC_{50} value of $3.61 \pm 0.74 \mu\text{M}$, whereas the other receptors conferred either weak or no responses to the peptide IP-O (MrgprA1, $2.0 \pm 0.7\%$; MrgprA2, $4.0 \pm 0.7\%$; MrgprA4, $6.4 \pm 1.0\%$; MrgprA10, $1.0 \pm 0.3\%$; MrgprA12, $6.0 \pm 0.7\%$; MrgprA14, $1.2 \pm 0.3\%$; MrgprA16, $4.2 \pm 0.5\%$; MrgprA19, $2.2 \pm 0.5\%$; MrgprB4, $5.4 \pm 0.9\%$; MrgprB5, $3.6 \pm 0.6\%$; $P < 0.0001$; Fig. 3A, B, C, D and Fig. S4). In contrast, MrgprC11-overexpressing HEK293T cells exhibited nearly no response to IP-R (Fig. S5), indicating that IPDef1 activates Mrgpr receptors through a mechanism dependent on the folding of its secondary structure. MrgprA3 and MrgprC11 were activated by their agonists CQ and BAM, respectively, confirming that they are functional receptors and are sensitive to IP-O (Fig. 3A and B). The main MrgprXs of the human Mrgpr family (MrgprX1, X2, X3 and X4) are much smaller than those of the murine Mrgpr family. We examined the effects of IP-O on MrgprXs and found that IP-O intensely activated MrgprX1, exhibiting an EC_{50} value

of $4.22 \pm 0.48 \mu\text{M}$. IP-O moderately activated MrgprX2 but did not affect hMrgprX3 or MrgprX4 (MrgprX1, $77.6 \pm 2.8\%$; MrgprX2, $28.2 \pm 2.7\%$; MrgprX3, $6.0 \pm 0.7\%$; MrgprX4, $2.6 \pm 0.5\%$; $P < 0.0001$; Fig. 3C, F, G, H, and Fig. S6). Like MrgprC11-overexpressing HEK293T cells, MrgprX1-overexpressing HEK293T cells did not respond to the reduced form of IPDef1 (IP-R) (Fig. S7). Some studies have shown that MrgprX1/C11 is preferentially activated by peptides that terminate in RYG or RF-amide^{26, 27}. Given that IP-O does not terminate with either motif, these results suggest that IP-O represents a completely new type of ligand for MrgprX1/C11. In addition, like the endogenous ligand BAM, IP-O has the highest affinity for the itch receptors MrgprC11 and MrgprX1 (MrgprC11, $76.4 \pm 2.0\%$; MrgprX1, $77.6 \pm 2.8\%$; Fig. 3C and G), indicating that the molecular mechanism of IP-O-induced itch may be similar to that of BAM.

Coimmunoprecipitation was also used to determine whether the peptide IP-O directly interacts with the tested mouse and human Mrgprs. The peptide His-IP, which contains IPDef1 fused to a six-histidine residue tag at the N-terminus, was chemically synthesized and oxidatively refolded according to the procedure described for the peptide IP-O above (Fig. S8). HEK293T cells were transfected with a plasmid expressing an N-flag-tagged mouse Mrgpr (MrgprC11 or A3) or human Mrgpr (MrgprX1, X2, X3 or X4). The results of coimmunoprecipitation showed that both mouse MrgprC11 and MrgprA3 directly interacted with the peptide IP-O (Fig. S9A). Furthermore, human MrgprX1 and MrgprX2 but not MrgprX3 and MrgprX4 directly interacted with the peptide IP-O (Fig. S9B).

To further investigate the role that MrgprC11 plays in IP-O-induced itch, MrgprC11 knockout C57BL/6 mice were generated by the CRISPR/Cas9 approach (Fig. S10). Then, we compared IP-O-evoked Ca^{2+} signals in DRG neurons isolated from MrgprC11-deficient mice to those in DRG neurons isolated from wild-type (WT) littermates and found that Ca^{2+} signals evoked by IP-O were significantly attenuated in MrgprC11-deficient DRG neurons compared to WT DRG neurons (Fig. 3I). Moreover, BAM-evoked responses were also attenuated in MrgprC11-deficient neurons compared to WT neurons. These results suggest that MrgprC11 is the key neuroreceptor for both IP-O and BAM. Compared to those isolated from WT mice, the cultured DRG neurons isolated from MrgprC11-deficient mice showed a decrease in the proportion of IP-O-sensitive neurons (Fig. 3J). A similar decrease in the percentage of BAM-sensitive neurons was observed in DRG neurons isolated from

MrgprC11-deficient mice compared with those isolated from WT mouse DRG neurons (Fig. 3J). To further investigate the *in vivo* role of MrgprC11 in IP-O-induced itch, we evaluated the itching responses induced by IP-O in MrgprC11-deficient mice. As expected, the scratching response induced by IP-O were dramatically alleviated in MrgprC11-deficient mice compared with those of WT mice (Fig. 3K). Together, these data indicate that MrgprC11 is the main itch receptor for IP-O on mouse DRG neurons and plays a key role in IP-O-induced itch.

TRPV1 is the downstream ion channel coupled to IP-O-activated MrgprC11/X1 on DRG neurons.

Mouse MrgprC11 and human MrgprX1 were identified as the main receptors for the peptide IP-O. However, the signaling pathway and ion channel downstream of IP-O are unknown. It has been observed that many G protein-coupled receptors (GPCRs) on DRG neurons transduce signals via TRP channels^{28, 29}. TRPV1-expressing afferents mediate responses to a variety of pruritogens, and TRPV1-deficient mice display reduced responses to histamine³⁰. CQ and BAM activate a subset of TRPV1-positive neurons³¹. These findings suggest that TRPV1 is a likely candidate transduction channel in Mrgpr pruritic pathways that should not be ignored. Accordingly, we used live-cell calcium imaging to examine the overlap between the sensitivity of WT mouse DRG neurons to IP-O and the TRPV1 agonist capsaicin. AMG9810, an inhibitor of TRPV1, severely attenuated the effect of IP-O (10 μ M) on WT mouse DRG neurons (Fig. 4A). After washout, IP-O induced a relatively normal increase in $[Ca^{2+}]_i$ (Fig. 4A). Subsequent exposure to capsaicin (1 μ M) produced a further increase in $[Ca^{2+}]_i$ in all IP-O-positive cells (Fig. 4A). These data indicate that TRPV1 is likely involved in the signaling pathway associated with IP-O-induced itch. We then compared IP-O-evoked Ca^{2+} signals in DRG neurons isolated from TRPV1-deficient mice to those in DRG neurons isolated from WT littermates and found that Ca^{2+} signals evoked by IP-O were significantly attenuated in TRPV1-deficient DRG neurons compared to WT DRG neurons (Fig. 4B). As expected, capsaicin-evoked responses were also attenuated in TRPV1-deficient neurons compared to WT neurons, but the TRPA1 agonist allyl isothiocyanate (AITC) evoked Ca^{2+} signals in both TRPV1-deficient and WT DRG neurons (Fig. 4B). These results indicate that IP-O-activated DRG neurons express both TRPV1 and TRPA1, whereas TRPV1 but not TRPA1 is required for the IP-O-evoked signaling pathway.

Cultured DRG neurons isolated from TRPV1-deficient mice showed a decrease in the proportion of IP-O-sensitive neurons compared with that exhibited by WT mouse DRG neurons (Fig. 4C). A similar decrease in the proportion of IP-O-sensitive neurons was observed in WT mouse DRG neurons treated with the TRPV1 antagonist AMG9810 (Fig. 4C). Furthermore, there was a significant reduction in the proportion of histamine-sensitive cells in the DRG from TRPV1-deficient mice and AMG9810-treated DRG neurons from WT mice compared with that in WT DRG neurons (Fig. 4C). These findings are consistent with the previous finding that TRPV1 is required for histamine signaling in sensory neurons³². In contrast, the number of CQ-responsive cells was similar in WT mouse DRG neurons, AMG9810-treated WT mouse DRG neurons, and *Trpv1*^{-/-} mouse DRG neurons (Fig. 4C), indicating that TRPV1 is not required for CQ signaling, which is consistent with the findings of a previous study²⁹. In addition, we evaluated the itching responses induced by these three substances in TRPV1-deficient mice. Similar to that evoked by histamine, the scratching response induced by IP-O was significantly alleviated in TRPV1-deficient mice compared with WT mice (Fig. 4D). There was no significant difference in the total number of scratching bouts induced by CQ over a period of 30 min between TRPV1-deficient and WT mice (Fig. 4D). Thus, our results indicate that the functional TRPV1 channel is required for IP-O-evoked DRG activation and IP-O-induced itch in mice.

Consistent with this conclusion, calcium imaging indicated that capsaicin but not IP-O affected the Ca^{2+} response of mouse TRPV1 expressed in heterologous HEK293T cells (Fig. S11A). Whole-cell patch-clamp recordings showed that the peptide IP-O did not affect the currents of TRPV1-overexpressing HEK293T cells (Fig. S11B). Furthermore, co-immunoprecipitation experiments showed that the peptide IP-O did not directly interact with TRPV1 channels (Fig. S11C). All these results suggest that the peptide IP-O is not a direct agonist of TRPV1.

TRPA1 is not required for MrgprC11/X1-mediated DRG neuron activation by IP-O.

Although TRPV1 is required for IP-O-evoked Ca^{2+} signals, it does not mediate all forms of itch. TRPA1, which is highly expressed in a subset of TRPV1-positive neurons, is activated by a number of pain-producing compounds, including isothiocyanates³³. In addition, TRPA1 is activated downstream of some GPCRs. Thus,

we further examined whether there is an overlap between the sensitivity of WT mouse DRG neurons to IP-O and the TRPA1 agonist AITC. HC-030031, an antagonist of the TRPA1 channel³⁴, had little effect on IP-O-evoked Ca^{2+} signals in DRG neurons from WT mice (Fig. 5A). We then compared IP-O-evoked Ca^{2+} signals in DRG neurons isolated from TRPA1-deficient mice to those isolated from WT littermates. The results indicated that IP-O-evoked Ca^{2+} signals were similar in TRPA1-deficient and WT DRG neurons (Fig. 5B). In addition, the TRPV1 agonist capsaicin induced similar Ca^{2+} signals in both TRPA1-deficient and WT DRG neurons, whereas AITC-evoked responses were significantly attenuated in TRPA1-deficient DRG neurons compared to WT DRG neurons (Fig. 5B). These data indicate that TRPA1 is unlikely to be involved in the IP-O signaling pathway. Both DRG neurons isolated from TRPA1-deficient mice and AMG9810-treated DRG neurons isolated from WT mice showed similar responses to IP-O as DRG neurons isolated from WT mice (Fig. 5C). Likewise, no difference was observed in the proportion of histamine-sensitive neurons among these three kinds of DRG neurons (Fig. 5C). In contrast, the proportion of CQ-sensitive neurons among cultured neurons isolated from TRPA1-deficient mice was decreased compared with that among neurons isolated from WT mice (Fig. 5C). Similar results were observed for WT neurons treated with the TRPA1 antagonist HC-030031 (Fig. 5C). These findings show that TRPA1 is required for CQ signaling in sensory neurons. In contrast, the numbers of CQ-responsive cells among WT mouse neurons, AMG9810-treated WT mouse neurons, and mutant neurons were similar (Fig. 5C), indicating that TRPV1 is not required for CQ signaling. These results were completely consistent with the finding of a previous report³⁵. Furthermore, we evaluated the itching responses induced by IP-O, HIS and CQ in TRPA1-deficient mice. No significant difference in the total number of scratching bouts induced by IP-O over a period of 30 min, which was similar to that induced by HIS, was found between TRPA1-deficient and WT mice (Fig. 5D). However, the scratching response induced by CQ in TRPA1-deficient mice was significantly alleviated compared with that in WT mice.

Correspondingly, the results of the calcium imaging experiment indicated that AITC but not IP-O activated mouse TRPA1 expressed on heterologous HEK293T cells (Fig. S12A). Whole-cell patch-clamp recordings showed that the peptide IP-O did not affect the currents of the TRPA1 channel overexpressed in HEK293T cells (Fig. S12B). Moreover, coimmunoprecipitation experiments showed that the peptide

IP-O did not directly interact with the ion channel TRPA1 (Fig. S12C). Taken together, our results indicate that TRPA1 is not required for IP-O-evoked excitation of DRG neurons or subsequent IP-O-induced itch in mice.

IP-O activates mast cells through MrgprB2 and induces acute inflammation in mice.

The above results show that IP-O evokes itch by directly activating MrgprC11/X1 to regulate downstream TRPV1 on pruriceptors and that the MrgprC11/X1-TRPV1 pathway is an important signaling pathway for IP-O-induced itch. However, we found that IP-O moderately activated human MrgprX2 (an ortholog of mouse MrgprB2) (Fig. 3G) selectively expressed on mast cells but not on primary sensory neurons. It is possible that some mast cell-derived mediators, such as proteases and 5-HT, are involved in IP-O-induced itch. Therefore, we examined the effect of IP-O on mouse MrgprB2 by calcium imaging as described above. The results showed that some MrgprB2-overexpressing HEK293T cells responded to IP-O (Fig. 6A). We also found that IP-O directly activated peritoneal mast cells (PMCs) isolated from mice (Fig. 6B). Sodium cromoglicate (cromolyn), a mast cell stabilizer, can effectively inhibit granule release. We evaluated the itching responses induced by IP-O, PAMP9-20 and anti-IgE in cromolyn-treated mice. In contrast to the scratching responses induced by PAMP9-20 and anti-IgE, there was no significant difference in the total number of scratching bouts induced by IP-O over a period of 30 min between cromolyn-treated and vehicle-treated mice (Fig. 6C). It is likely that mast cells activated by IP-O made little contribution to itching and had unknown effects in mice. Evans blue extravasation assays showed that intraplantar injection of IP-O induced acute inflammation in mice (Fig. 6D, E). We measured paw thickness of the mice before and after IP-O treatment, and found that the paw thickness was significantly increased after the injection of IP-O (Fig. S13). In addition, compared with that in vehicle-treated mice, acute inflammation induced by IP-O was reduced in cromolyn-treated mice (Fig. 6F).

The activation of mast cells by intraplantar injection of IP-O caused acute inflammation in mice, but it was unclear which mediators released from mast cells were required for this effect. We detected *in vitro* release of histamine, serotonin, tryptase beta 2, TNF- α and MCP-1 from mouse peritoneal mast cells upon stimulation by IP-O (12, 25, 50 μ M), PAMP (100 μ M), Anti-IgE (25 μ g/mL). Compared with

vehicle, IP-O resulted in the releases of histamine, serotonin and tryptase beta 2 (Fig. 6G-I). In addition, IP-O also induced the releases of TNF- α and MCP-1 with an moderate increasement (Fig. S14). These mediators released from mast cells may have an effect on recruiting immune cells and facilitating the progress of inflammation. Further, we analyzed the mRNA expression of more cytokines and chemokines in IP-O-treated PMCs. Among the test cytokine and chemokine genes, no significant change was observed at the level of mRNA expression after IP-O stimulation in PMCs (Fig. S15), which was consistent with the degranulation release of mast cells. Taken together, these results suggest that IP-O activates mast cells through MrgprB2/X2 and induces acute inflammation but that mast cell activation appears to make little contribution to IP-O-induced itch.

Discussion

As vectors of various pathogens, ticks commonly induce skin pruritus by biting humans and animals. Which class of substance causes scratching and itching following a tick bite: the carried pathogens or endogenous components expressed by ticks? We postulated that endogenous components of ticks are most likely responsible for itch induction for two reasons. First, dogs bitten by the tick *O. brasiliensis* exhibit continuous and intense itching behavior, but typical tick-borne pathogens are not detected in the sera of bitten dogs¹⁰. Tick bites with and without pathogens both lead to skin pruritus in dogs. Second, many arthropod bites and stings can cause itching behavior, but ants, bees, spiders and scorpions have not yet been found to transmit pathogens.

We found that two tick salivary defensin peptides, IPDef1 and IRDef2, significantly induced itching and scratching behavior in mice upon intradermal injection. Defensins in the saliva of ticks were discovered to act as new pruritogenic agents, at least partially explaining the pathological phenomenon of skin pruritus caused by tick bites. Defensins in the saliva of ticks share high homology and structural similarity with ancient invertebrate defensins. Thus, we found a new large class of pruritogenic peptide agents that is completely different from previously reported pruritogenic peptides such as BAM8-22²⁴ and mouse/human beta-defensins^{36, 37}. Our findings provide many new molecular probes and tools for studying itch receptors.

Interestingly, our study revealed that the tick salivary peptide IPDef1 exerts two activities through two different signaling pathways in mice: MrgprC11/X1-mediated DRG neuron activation and MrgprB2/X2-mediated mast cell activation. First, IPDef1 triggers DRG neuron activation by specifically acting on MrgprC11/X1 on DRG neurons and induces cellular calcium influx into DRG neurons through the downstream TRPV1 channel, which causes itching in mice. Second, IPDef1 also activates mast cells through MrgprB2/X2, a recently discovered membrane receptor on mast cells that induces acute inflammation in mice. Besides taking part in the pathology and mortality caused by envenomation, mast cells were previously found to play an important role in detoxification of harmful poisons. Mast cell degranulation releases carboxypeptidase A3 (CPA3) and chymase, and reduces the toxicity of animal venoms (like scorpions, bees and snakes) by degrading their venom peptides^{38, 39}. It is still unclear whether IPDef1 induces mast cells to release CPA3 or chymase for detoxification by activating MrgprB2/X2.

Coimmunoprecipitation, calcium imaging, genetic ablation and behavior experiments revealed the molecular mechanism by which the peptide IPDef1 directly interacts with Mrgprs (mainly MrgprC11/X1) and activates DRG neurons to induce itch. However, this mechanism is not related to the histamine signaling pathway. We not only found a class of new pruritogenic peptide agents responsible for arthropod bite- or sting-induced itch but also revealed the related neural mechanisms, laying the foundation for the development of anti-itch drugs to combat arthropod bite- or sting-induced pruritus. Moreover, an inhibitor of TRPV1 was shown to specifically block calcium influx into DRG neurons activated by the tick peptide IPDef1 and inhibit the pruritus induced by IPDef1 in cell and animal behavior experiments. TRPV1 is a promising target of anti-itch drugs, and its inhibitors are potential candidates for preventing and treating the pruritus induced by tick bites. Coincidentally, Li et al. reported a case of a man bitten by the tick *I. persulcatus* on Yunmeng Mountain in Beijing, China. The patient subsequently developed topical edematous erythema and itching, but oral antihistamine and topical calamine lotion did not improve his itching symptoms. This observation suggests that antihistamine drugs do not have an effect against pruritus induced by tick bites, providing evidence that tick bite-induced pruritus is independent of the histamine-related pathway. In short, our study identifies potential therapeutic targets and drugs for the prevention

and treatment of pruritus induced by the bites or stings of arthropods such as ticks, mites, fleas, mosquitoes, bees, wasps, spiders, and scorpions.

The generation of peptides that induce itch resulted from the interaction between and coevolution of parasites and hosts or prey and predators. We speculate that there are two possible driving forces of this phenomenon. First, the generation of itchiness is a self-alarm and self-defense mechanism in hosts or predators. Itch helps hosts or predators scratch away external threats². It is easily understood that hosts or predators evolve to produce itching signals against arthropod bites or stings. Second, the salivary or venom glands of these arthropods have evolved to produce peptides that induce pruritus, which are their molecular weapons for predation and defense, to attack hosts, protect against enemies or deter competitors⁴⁰. During long-term evolution, multiple peptide and protein families have been recruited to the animal salivary or venom systems⁴¹. Defensins have also been recruited to animal saliva and venoms as chemical weapons for predation and defence. Defensins belong to a class of ancient cationic peptides that are widely distributed in fungi, plants and animals and are effector molecules of the innate immune system, exhibiting broad-spectrum antimicrobial activity against a range of bacteria and viruses⁴². Consistent with our finding that the ancient invertebrate defensin IPDef1 from the tick *I. persalcatatus* evokes itch by directly activating MrgprC11/X1 expressed on DRG neurons, mouse and human beta-defensins have also been identified as pruritogens that activate Mrgprs or Toll-like receptor 4 (TLR4)^{36, 37}. These results suggest that ancient invertebrate defensins were recruited early to the tick salivary systems and innovated a new toxicological function of itch induction. Our study not only reveals a new toxicological effect and mechanism of defensins in saliva or venoms but also brings to light a new link between neurobiology and immunology.

Acknowledgments

We thank Dr. Qin Liu (Washington University School of Medicine, St. Louis) for kindly providing plasmids of mouse MrgprA3 and MrgprC11. We also thank Dr. Qin Liu and Dr. Weishan Yang (Washington University School of Medicine, St. Louis) for their valuable communications with us. This work was supported by grants from National Science Fund of China (Nos. 32070525, 31872239, and 81630091), Hubei

Science Fund for Excellent Scholars (No. 2020CFA015), and the Open Research Program of the State Key Laboratory of Virology of China (No. 2020KF002).

Author Contributions

X. L., H. Y., and Z. C. designed the experiments and analyzed the data. X. L., and H. Y. performed most of the experiments. Y. H. cloned the cDNA sequences of human *MrgprXs* and did itch-related animal experiments. S. Y. completed patch-clamp experiments. B. S. performed the experiment of acute inflammation in mice. Y. W. and W. L. analyzed experimental data and revised the manuscript. X. L., and Z. C. wrote the manuscript.

References

1. Cevikbas F, Lerner EA. Physiology and pathophysiology of itch. *Physiological Reviews* 2020; 100:945-82.
2. Lay M, Dong X. Neural mechanisms of itch. *Annual Review of Neuroscience* 2020.
3. Liu Q, Tang ZX, Surdenikova L, Kim S, Patel KN, Kim A, et al. Sensory neuron-specific GPCR Mrgprs are itch receptors mediating chloroquine-induced pruritus. *Cell* 2009; 139:1353-65.
4. Feng J, Luo JL, Yang P, Du JH, Kim BS, Hu HZ. Piezo2 channel-Merkel cell signaling modulates the conversion of touch to itch. *Science* 2018; 360:530.
5. Hoogstraal H, Gallagher M. Blisters, pruritus, and fever after bites by the Arabian tick *Ornithodoros (Alectorobius) Muesebecki*. *The Lancet* 1982; 320:288-9.
6. Jones AV, Tilley M, Gutteridge A, Hyde C, Nagle M, Ziemek D, et al. GWAS of self-reported mosquito bite size, itch intensity and attractiveness to mosquitoes implicates immune-related predisposition loci. *Hum Mol Genet* 2017; 26:1391-406.
7. Cyranoski D. East Asia braces for surge in deadly tick-borne virus. *Nature* 2018; 556:282-3.
8. Holding M, Dowall S, Hewson R. Detection of tick-borne encephalitis virus in the UK. *The Lancet* 2020; 395:411.
9. Fisher EJ, Mo J, Lucky AW. Multiple pruritic papules from lone star tick larvae bites. *JAMA Dermatology* 2006; 142:491-4.
10. Reck J, Soares JF, Termignoni C, Labruna MB, Martins JR. Tick toxicosis in a dog bitten by *Ornithodoros brasiliensis*. *Veterinary Clinical Pathology* 2011; 40:356-60.
11. Cao Z, Yu Y, Wu Y, Hao P, Di Z, He Y, et al. The genome of *Mesobuthus martensii* reveals a unique adaptation model of arthropods. *Nature Communications* 2013; 4:2602.
12. Luo L, Li B, Wang S, Wu F, Wang X, Liang P, et al. Centipedes subdue giant

- 831 prey by blocking KCNQ channels. *Proceedings of the National Academy of*
 832 *Sciences* 2018; 115:1646.
- 833 13. Yang S, Yang F, Wei N, Hong J, Li B, Luo L, et al. A pain-inducing centipede
 834 toxin targets the heat activation machinery of nociceptor TRPV1. *Nature*
 835 *Communications* 2015; 6:8297.
- 836 14. Lin King JV, Emrick JJ, Kelly MJS, Herzig V, King GF, Medzihradsky KF, et
 837 al. A cell-penetrating scorpion toxin enables mode-specific modulation of
 838 TRPA1 and pain. *Cell* 2019; 178:1362-74.e16.
- 839 15. Osteen JD, Herzig V, Gilchrist J, Emrick JJ, Zhang C, Wang X, et al. Selective
 840 spider toxins reveal a role for the Nav1.1 channel in mechanical pain. *Nature*
 841 2016; 534:494-9.
- 842 16. Meng L, Xie Z, Zhang Q, Li Y, Yang F, Chen Z, et al. Scorpion potassium
 843 channel-blocking defensin highlights a functional link with neurotoxin.
 844 *Journal of Biological Chemistry* 2016; 291:7097-106.
- 845 17. Thangamani S, Wikel SK. Differential expression of *Aedes aegypti* salivary
 846 transcriptome upon blood feeding. *Parasit Vectors* 2009; 2:34.
- 847 18. Ikoma A, Steinhoff M, Ständer S, Yosipovitch G, Schmelz M. The
 848 neurobiology of itch. *Nature Reviews Neuroscience* 2006; 7:535-47.
- 849 19. Thurmond RL, Kazerouni K, Chaplan SR, Greenspan AJ. Antihistamines and
 850 itch. *Handb Exp Pharmacol* 2015; 226:257-90.
- 851 20. MeiXiong J, Dong XZ. Mas-related G protein-coupled receptors and the
 852 biology of itch sensation. *Annual Review of Genetics* 2017; 51:103-21.
- 853 21. Wooten M, Weng HJ, Hartke TV, Borzan J, Klein AH, Turnquist B, et al.
 854 Three functionally distinct classes of C-fibre nociceptors in primates. *Nature*
 855 *Communications* 2014; 5:4122.
- 856 22. Chrudimská T, Slaninová J, Rudenko N, Růžek D, Grubhoffer L. Functional
 857 characterization of two defensin isoforms of the hard tick *Ixodes ricinus*.
 858 *Parasit Vectors* 2011; 4:63.
- 859 23. Rudenko N, Golovchenko M, Grubhoffer L. Gene organization of a novel
 860 defensin of *Ixodes ricinus*: first annotation of an intron/exon structure in a
 861 hard tick defensin gene and first evidence of the occurrence of two isoforms of
 862 one member of the arthropod defensin family. *Insect Mol Biol* 2007; 16:501-7.
- 863 24. Lembo PM, Grazzini E, Groblewski T, O'Donnell D, Roy MO, Zhang J, et al.
 864 Proenkephalin A gene products activate a new family of sensory
 865 neuron-specific GPCRs. *Nature neuroscience* 2002; 5:201-9.
- 866 25. Sikand P, Dong XZ, LaMotte RH. BAM8-22 peptide produces itch and
 867 nociceptive sensations in humans independent of histamine release. *The*
 868 *Journal of Neuroscience* 2011; 31:7563.
- 869 26. Han SK, Dong XZ, Hwang JI, Zylka MJ, Anderson DJ, Simon MI. Orphan G
 870 protein-coupled receptors MrgA1 and MrgC11 are distinctively activated by
 871 RF-amide-related peptides through the $G\alpha_{q/11}$ pathway. *Proceedings of the*
 872 *National Academy of Sciences* 2002; 99:14740.
- 873 27. Grazzini E, Puma C, Roy MO, Yu XH, Donnell D, Schmidt R, et al. Sensory
 874 neuron-specific receptor activation elicits central and peripheral nociceptive
 875 effects in rats. *Proceedings of the National Academy of Sciences of the United*
 876 *States of America* 2004; 101:7175.
- 877 28. Jordt SE, Bautista DM, Chuang HH, McKemy DD, Zygmunt PM, Högestätt
 878 ED, et al. Mustard oils and cannabinoids excite sensory nerve fibres through
 879 the TRP channel ANKTM1. *Nature* 2004; 427:260-5.
- 880 29. Wilson SR, Gerhold KA, Bifulck-Fisher A, Liu Q, Patel KN, Dong XZ, et al.

- 881 TRPA1 is required for histamine-independent, Mas-related G protein-coupled
 882 receptor-mediated itch. *Nature Neuroscience* 2011; 14:595-602.
- 883 30. Valtcheva MV, Davidson S, Zhao CS, Leitges M, Gereau RW. Protein kinase
 884 C δ mediates histamine-evoked itch and responses in pruriceptors. *Molecular*
 885 *Pain* 2015; 11:1.
- 886 31. Imamachi N, Park GH, Lee HS, Anderson DJ, Simon MI, Basbaum AI, et al.
 887 TRPV1-expressing primary afferents generate behavioral responses to
 888 pruritogens via multiple mechanisms. *Proceedings of the National Academy of*
 889 *Sciences* 2009; 106:11330.
- 890 32. Immke DC, Gavva NR. The TRPV1 receptor and nociception. *Seminars in*
 891 *Cell & Developmental Biology* 2006; 17:582-91.
- 892 33. McNamara CR, Mandel-Brehm J, Bautista DM, Siemens J, Deranian KL,
 893 Zhao M, et al. TRPA1 mediates formalin-induced pain. *Proceedings of the*
 894 *National Academy of Sciences* 2007; 104:13525.
- 895 34. Yin SJ, Luo JL, Qian AH, Du JH, Yang Q, Zhou ST, et al. Retinoids activate
 896 the irritant receptor TRPV1 and produce sensory hypersensitivity. *The Journal*
 897 *of Clinical Investigation* 2013; 123:3941-51.
- 898 35. Ru F, Sun H, Jurcakova D, Herbstsomer RA, MeiXong J, Dong X, et al.
 899 Mechanisms of pruritogen-induced activation of itch nerves in isolated mouse
 900 skin. *The Journal of Physiology* 2017; 595:3651-66.
- 901 36. Zhang L, McNeil BD. Beta-defensins are proinflammatory pruritogens that
 902 activate Mrgprs. *Journal of Allergy and Clinical Immunology* 2019;
 903 143:1960-2.e5.
- 904 37. Feng J, Luo JL, Mack MR, Yang P, Zhang F, Wang G, et al. The antimicrobial
 905 peptide human beta-defensin 2 promotes itch through Toll-like receptor 4
 906 signaling in mice. *Journal of Allergy and Clinical Immunology* 2017;
 907 140:885-8.e6.
- 908 38. Akahoshi M, Song CH, Piliponsky AM, Metz M, Guzzetta A, Abrink M, et al.
 909 Mast cell chymase reduces the toxicity of Gila monster venom, scorpion
 910 venom, and vasoactive intestinal polypeptide in mice. *J Clin Invest* 2011;
 911 121:4180-91.
- 912 39. Metz M, Piliponsky AM, Chen CC, Lammel V, Abrink M, Pejler G, et al. Mast
 913 cells can enhance resistance to snake and honeybee venoms. *Science* 2006;
 914 313:526-30.
- 915 40. Casewell NR, Wüster W, Vonk FJ, Harrison RA, Fry BG. Complex cocktails:
 916 the evolutionary novelty of venoms. *Trends Ecol Evol* 2013; 28:219-29.
- 917 41. Undheim EA, Jones A, Clauser KR, Holland JW, Pineda SS, King GF, et al.
 918 Clawing through evolution: toxin diversification and convergence in the
 919 ancient lineage Chilopoda (centipedes). *Mol Biol Evol* 2014; 31:2124-48.
- 920 42. Schneider T, Kruse T, Wimmer R, Wiedemann I, Sass V, Pag U, et al. Plectasin,
 921 a fungal defensin, targets the bacterial cell wall precursor Lipid II. *Science*
 922 2010; 328:1168-72.

923

Figure Legends

Fig. 1. The tick peptide IPDef1 causes histamine-independent itch in mice. (A) Amino acid sequence of the peptide IPDef1 from the tick *Ixodes persulcatus*. IP-R and IP-O are the reduced and oxidized forms of IPDef1, respectively. SH represents the thiol group of cysteine. The connectivity of disulfide bonds is indicated by the solid line with S-S. The cysteine residues are shaded in yellow, and the basic residues are displayed in blue. **(B)** Oxidative refolding of chemically synthetic linear IPDef1. RP-HPLC shows the difference in retention time (T_R) between IP-R and IP-O. **(C)** MALDI-TOF MS analysis of IP-R (small) and IP-O (large). **(D)** Scratching responses induced by intradermal injection of vehicle (saline, $n = 8$), IP-R (50 μg , $n = 6$), IP-O (50 μg , $n = 7$), histamine (HIS, 10 μmol , $n = 7$) and chloroquine (CQ, 200 μg , $n = 9$) in mice. **(E)** Dose-dependent scratching responses induced by intradermal injection of vehicle (saline, $n = 6$), IP-O (12 μg , $n = 6$), IP-O (25 μg , $n = 7$) and IP-O (50 μg , $n = 6$). **(F)** Difference in scratching responses induced by intradermal injection of IP-O (25 μg , $n = 6$), HIS (10 μmol , $n = 6$) and saline in vehicle-treated (white), cetirizine-treated (CETY, light gray) and JNJ7777120-treated mice (JNJ, dark gray). Each dot represents an individual mouse. All data are presented as the means $\pm$ SEMs. n.s, not significant, $P > 0.5$; ** $P < 0.01$; **** $P < 0.0001$.

Fig. 2. IP-O activates mouse DRG neurons with extracellular Ca^{2+} influx. (A) Representative calcium traces of cultured mouse DRG neurons in the presence of IP-R (10 μM) and IP-O (10 μM). **(B)** Representative Fura-2 ratiometric images of IP-R-evoked (10 μM , white arrowheads) and IP-O-evoked (10 μM , yellow arrowheads) responses in cultured mouse DRG neurons. The scale bar represents 20 μm . **(C)** Dose-response curve of cultured mouse DRG neurons to IP-O (1, 2, 4, 8 and 16 μM , respectively). $n = 3$ experiments/group. **(D)** Percentage of cultured mouse DRG neurons that responded to IP-R (10 μM), IP-O (10 μM), BAM8-22 (BAM, 50 μM) and CQ (1 mM). $n = 3$ experiments/group. All data are presented as the means $\pm$ SEMs. n.s, not significant, $P > 0.5$; *** $P < 0.001$. **(E)** Representative calcium traces of cultured mouse DRG neurons that responded to IP-O (10 μM , left), CQ (1 mM, middle) and BAM (50 μM , right) in the absence and presence of extracellular calcium (2 mM Ca^{2+}).

Fig. 3. Mouse MrgprC11 and human MrgprX1 are two itch receptors for IP-O.

(A, B) Representative calcium traces showing the responses of mouse MrgprA3 and MrgprC11 expressed on HEK293T cells to IP-O (10 μ M). (C) Percentage of 12 mouse Mrgprs (MrgprA1, A2, A3, A4, A10, A12, A14, A16, A19, B4, B5 and C11) expressed on HEK293T cells that responded to IP-O. $n = 5$ experiments/group. (D) Dose-response curve for MrgprC11 expressed on HEK293T cells to IP-O (1, 2, 4, 8 and 16 μ M). $n = 3$ experiments/group. (E, F) Representative calcium traces showing the responses of human MrgprX1 and MrgprX2 expressed on HEK293T cells to IP-O (10 μ M). (G) Percentage of human Mrgprs (MrgprX1, X2, X3 and X4) expressed on HEK293T cells that responded to IP-O. $n = 5$ experiments/group. (H) Dose-response curve for MrgprX1 expressed on HEK293T cells to IP-O (1, 2, 4, 8 and 16 μ M, respectively). $n = 3$ experiments/group. (I) Representative calcium response traces of wild-type (left) and *MrgprC11*^{-/-} (right) mouse DRG neurons exposed to IP-O (10 μ M), BAM (50 μ M) and KCl (50 mM), respectively. (J) Prevalence of IP-O and BAM sensitivity in WT (white) and *MrgprC11*^{-/-} DRG neurons (black). $n = 4$ experiments/group. (K) Difference in scratching responses induced by intradermal injection of IP-O (25 μ g) in WT (white, $n = 8$) and *MrgprC11*^{-/-} mice (black, $n = 10$). All data are presented as the means $\pm$ SEMs. * $P < 0.05$; *** $P < 0.001$.

Fig. 4. TRPV1 is the downstream ion channel that mediates IP-O-evoked DRG neuron activation and IP-O-induced itch in mice.

(A) Representative calcium traces of WT DRG neurons exposed to IP-O (10 μ M) following pretreatment (3 min) with or without AMG9810 (50 μ M). (B) Representative calcium traces of WT (left) and *Trpv1*^{-/-} (right) DRG neurons exposed to IP-O (10 μ M) followed by allyl isothiocyanate (AITC, 100 μ M) and capsaicin (Cap, 1 μ M). (C) Prevalence of IP-O, histamine (HIS, 1 μ M) and CQ (1 mM) sensitivity in WT(white), AMG9810-treated WT DRG (50 μ M, gray) and *Trpv1*^{-/-} DRG neurons (black). $n = 4$ experiments/group. (D) Difference in scratching responses induced by intradermal injection of IP-O (25 μ g, $n = 6$), HIS (10 μ mol, $n = 6$) and CQ (200 μ g, $n = 6$) in WT mice (white) and *Trpv1*^{-/-} mice (black). All data are presented as the means $\pm$ SEMs. n.s, not significant, $P > 0.5$; ** $P < 0.01$; *** $P < 0.001$.


Fig. 5. TRPA1 is not required for IP-O-evoked DRG neuron activation or


IP-O-induced itch in mice. (A) Representative calcium traces of WT DRG neurons exposed to IP-O (10 μ M) following pretreatment (3 min) with or without the HC 030031 (500 μ M). (B) Representative calcium traces of WT (left) and *Trpa1*^{-/-} (right) DRG neurons exposed to IP-O (10 μ M) followed by capsaicin (Cap, 1 μ M) and allyl isothiocyanate (AITC, 100 μ M). (C) Prevalence of IP-O, histamine (1 μ M) and CQ (1 mM) sensitivity in WT (white), HC 030031-treated WT (500 μ M, light gray) and *Trpa1*^{-/-} DRG neurons (dark gray). n = 4 experiments/group. (D) Difference in scratching responses induced by intradermal injection of IP-O (25 μ g, n = 6), HIS (10 μ mol, n = 6) and CQ (200 μ g, n = 6) in WT (white) and *Trpa1*^{-/-} mice (dark gray). All data are presented as the means $\pm$ SEMs. n.s, not significant, $P > 0.5$.


Fig. 6. IP-O activates mast cells through MrgprB2 and induces acute inflammation in mice. (A) Representative calcium traces of mouse MrgprB2-overexpressing HEK293T cells to IP-O (10 μ M). (B) Representative calcium traces of isolated mouse peritoneal mast cells (PMCs) to IP-O (10 μ M). (C) Difference in scratching responses induced by intradermal injection of IP-O (25 μ g, n = 7), PAMP (PAMP9-20, 25 μ g, n = 10) and anti-IgE (1 μ g, n = 9) in vehicle-treated (saline, white) or cromolyn-treated mice (light gray). (D) Representative images of Evans blue extravasation 15 min after intraplantar injection of saline (5 μ L, left paw) or IP-O (5 μ L, 2 mg/mL, right paw). n = 6. (E) Quantification of Evans blue content in the paws after injection of saline or IP-O. (F) Quantification of Evans blue content in the paws after injection of saline or IP-O in PBS-treated and cromolyn-treated mice. (G-I) *In vitro* release of histamine (G), serotonin (H) and tryptase beta 2 (I) from mouse PMCs upon stimulation by IP-O (12, 25, 50 μ M) or PAMP (100 μ M) or Anti-IgE (25 μ g/mL) or vehicle alone (IP-O = 0 μ M). Each dot represents an independent biological replicate from PMCs isolated from >4 animals. All concentrations n = 3. All data are presented as the means $\pm$ SEMs. n.s, not significant, $P > 0.5$; * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$.


Table S1. The qRT-PCR primer sequences for the test mouse cytokines and chemokines


Name	Direction	Sequence (5'-3')
TPH1	+	ACGTTCTCTCTTGGCTGAA
	-	TAGCACGTTGCCAGTTTTTG
SERT	+	TCACATATGCGGAGGCAATA
	-	CTATCCAAACCCAGCGTGAT
Mcpt6	+	CATTGATAATGACGAGCCTCTCC
	-	CATCTCCCGTGTAGAGGCCAG
TNF- α	+	TAGCCAGGAGGGAGAACAGA
	-	CCAGTGAGTGAAAGGGACAGA
IL-1 β	+	TACATCAGCACCTCACAAGC
	-	AGAAACAGTCCAGCCCATACT
MCP-1	+	TTAAAAACCTGGATCGGAACCAA
	-	GCATTAGCTTCAGATTACGGGT
VEGF	+	CAACTTCTGGGCTCTTCTCG
	-	CCTCTCCTCTTCCTTCTCTTCC
CXCL1	+	GTCAGTGCCTGCAGACCATG
	-	TGACTTCGGTTTGGGTGCAG
CXCL2	+	GCCAAGGGTTGACTTCAAGA
	-	TTCAGGGTCAAGGCAAACCTT
GAPDH	+	AGGTCGGTGTGAACGGATTG
	-	TGTAGACCATGTAGTTGAGGTCA


























A**B**













A



His-IP : HHHHHHGF^{GC}PFNQG^{ACH}R^{HC}RSIG^{RR}GGY^{CC}AGLF^{KQT}C^{TC}YS^R : 44


B**C**


A**B**



A


Chromosome 7: 48,020,971- 48,027,597


 Exons of *MrgprC11* (1180 bp) Deleted region (1890 bp)**B**





A**B**





Supplementary Informations:

Tick peptides evoke itch by activating MrgprC11/X1 to sensitize TRPV1 in pruriceptors

Xueke Li, MSc^{1,§}, Haifeng Yang, MSc^{1,§}, Yewen Han, MSc¹, Shijin Yin, PhD², Bingzheng Shen, PhD¹, Yingliang Wu, PhD¹, Wenxin Li, PhD¹, Zhijian Cao, PhD^{1,3,4,¶}

¹ State Key Laboratory of Virology, College of Life Sciences, Wuhan University, Wuhan 430072, P. R. China

² School of Pharmaceutical Sciences, South-Central University for Nationalities, Wuhan 430074, P. R. China.

³ Bio-drug Research Center, Wuhan University, Wuhan 430072, P. R. China

⁴ Hubei Province Engineering and Technology Research, Center for Fluorinated Pharmaceuticals, Wuhan University, Wuhan 430072, P. R. China

Disclosure of potential conflict of interest:

The authors declare that they have no relevant conflicts of interest.

[§] These authors contributed equally

[¶] Correspondence: State Key Laboratory of Virology, College of Life Sciences, Wuhan University, Wuhan 430072, P. R. China. Tel: +86-27-68752831, Fax: +86-27-68756746. E-mail: zjcao@whu.edu.cn (Zhijian Cao)

Supplementary Figure Legends

Fig. S1. Structural features of the peptide IPDef1 from the tick *Ixodes persalcatus*.

(A) Secondary structure analysis of the reduced (IP-R) and oxidated (IP-O) forms of IPDef1. CD spectrum shows structure difference between IP-R and IP-O. (B) Secondary structure analysis of IP-O in different solutions. (C) The homologous model of IPDef1. The 3D-structure of IPDef1 is shown as a solid ribbon model and three disulfide bonds are displayed as a line ribbon. Diagram was generated using SWISS-MODEL.

Fig. S2. Antimicrobial activities of the peptide IPDef1 (IP-R and IP-O) against standard bacteria strains.

Y axis shows MIC values of IP-R and IP-O against five gram-positive bacteria and two gram-negative bacteria. n.s, not significant, $P > 0.5$; * $P < 0.05$; ** $P < 0.01$; **** $P < 0.0001$.

Fig. S3. The tick salivary peptide IRDef2 induces itch responses in mice.

(A) Amino acid sequence of the peptide IRDef2 from the tick *I. ricinus* saliva. Def2-R and Def2-O are the reduced and oxidatized forms of IRDef2, respectively. SH represents the thiol group of cysteine. The connection mode of disulfide bond is displayed in a solid line with S-S. The cysteine residues are shaded with brilliant yellow, acidic residues are displayed in pink and basic residues are displayed in blue. (B) Scratching responses induced by intradermal injection of vehicle (saline), Def2-R (25 μ g) and Def2-O (25 μ g) in mice. Each dot represents an individual mouse. All groups $n = 6$. All data are presented as means $\pm$ SEM. n.s, not significant, $P > 0.5$; **** $P < 0.0001$.

Fig. S4. IP-O fails to activate mouse MrgprA1, A2, A4, A10, A12, A14, A16, A19, B4, and B5.

(A-J) Representative calcium traces of mouse Mrgprs (MrgprA1, A2, A4, A10, A12, A14, A16, A19, B4, and B5) expressed on HEK293T cells to IP-O (10 μ M). n = 3 experiments/group.

Fig. S5. IP-R fails to activate mouse MrgprC11.

Representative calcium traces of mouse MrgprC11 expressed on HEK293T cells to IP-R (10 μ M).

Fig. S6. IP-O fails to activate human MrgprX3 and MrgprX4.

(A, B) Representative calcium traces of human MrgprX3 and MrgprX4 expressed on HEK293T cells to IP-O (10 μ M). n = 3 experiments/group.

Fig. S7. IP-R fails to activate human MrgprX1.

Representative calcium traces of human MrgprX1 expressed on HEK293T cells to IP-R (10 μ M).

Fig. S8. Preparation and structural feature of the peptide His-IP-O.

(A) Amino acid sequence of the peptide fused with a six his-tag at the N-terminus of IPDef1 (His-IP). The connection mode of disulfide bond is displayed in a solid line. Cysteine residues are shaded with brilliant yellow, and basic residues are displayed in blue. (B) Oxidative refolding of chemically synthetic His-IP. RP-HPLC shows retention time (T_R) difference between the reduced (His-IP-R) and oxidatized (His-IP-O) peptides. (C) Secondary structure analysis of His-IPDef1. CD spectrum shows structure difference between His-IP-R and His-IP-O.

Fig. S9. IP-O directly interacts with mouse MrgprA3/C11 and human MrgprX1/X2. (A, B) Co-immunoprecipitation analysis of the peptide IP-O with the mouse MrgprA3/C11 and human MrgprX1-X4. HEK293T cells were transfected with the plamid pcDNA3.1 expressing different N-flag-tagged mouse Mrgprs (MrgprA3 and C11) (A) and human Mrgprs (MrgprX1, X2, X3 and X4) (B), respectively.

Fig. S10. Creation of MrgprC11 knockout mice by CRISPR/Cas9.

(A, B) Strategy and genotyping results of MrgprC11 knockout mouse.

Fig. S11. IP-O fails to interact with TRPV1 directly. (A) Representative calcium traces of TRPV1 expressed on HEK293T cells to IP-O (10 μ M). (B) Effect of IP-O on the current of TRPV1-overexpressing HEK293T cells. (C) Co-immunoprecipitation analysis of the peptide IP-O with the mouse TRPV1.

Fig. S12. IP-O fails to interact with TRPA1 directly. (A) Representative calcium traces of TRPA1-overexpressing HEK293T cells to IP-O (10 μ M). (B) Effect of IP-O on the current of TRPA1-overexpressing HEK293T cells. (C) Co-immunoprecipitation analysis of the peptide IP-O with the mouse TRPA1.

Fig. S13. IP-O increases the paw thickness of mice. (A) Change in the paw thickness (%) of mice after the intraplantar injection of saline (5 μ L, left paw) and IP-O (5 μ L, 2 mg/mL, right paw). $n = 6$. (B) Change in the paw thickness (%) of mice after the intraplantar injection of saline (5 μ L) and IP-O (5 μ L, 2 mg/mL) in PBS-treated mice and cromolyn-treated mice. $n = 6$. All data are presented as means $\pm$ SEM. n.s, not significant, $P > 0.5$; * $P < 0.05$; *** $P < 0.001$.

Fig. S14. IP-O induces the release of TNF- α and MCP-1 from mouse PMCs. (A, B) *In vitro* release of TNF- α (A) and MCP-1 (B) from mouse PMCs upon stimulation by IP-O (12, 25, 50 μ M), PAMP (PAMP9-20, 100 μ M), Anti-IgE (25 μ g/mL) or vehicle alone (IP-O = 0 μ M). All concentrations $n = 3$. All data are presented as the means $\pm$ SEMs. n.s, not significant, $P > 0.5$; * $P < 0.05$.

Fig. S15. Effect of IP-O on the mRNA expression of the inflammatory cytokines and chemokines in mouse PMCs. Mouse PMCs (1×10^4 - 5×10^4) were incubated with test substances for 30 minutes before the total intracellular RNA were collected. TPH1, SERT, Mcpt6, TNF- α , IL-1 β , MCP-1, VEGF, CXCL1 and CXCL2 were analyzed by qPCR. All groups $n = 3$. All data are presented as the means $\pm$ SEMs. n.s, not significant, $P > 0.5$.