


Allergologia et immunopathologia

www.elsevier.es/ai



LETTER TO THE EDITOR

Heat immersion therapy—A new approach for contact dermatitis to propolis?

To the Editor,

Recently, I summarised the current evidence on the treatment of bee stings in a German beekeeping magazine. One major aspect was the use of direct heat and heat immersion therapy. I discussed the study by Müller et al. who showed that patients who suffered from wasp, mosquito and bee stings had fast amelioration of symptoms after locally administrated concentrated heat.¹ An absence of symptoms was reported 10 min after administration. It was also discussed that the concept of heat treatment was found to be effective in a number of other scenarios, like in bluebottle (*Physalia* sp.) stings and acute jellyfish (*Carybdea alata*) stings.^{2,3}

After publication of the article I was contacted by a 62-year-old beekeeper who wrote that he has kept bees since 1958 and has suffered from propolis contact allergy since 1985. In spite of this, he continued beekeeping, but reduced the number of attended bee hives and wore cotton gloves as a protective measure. Since contact with propolis could not be avoided completely, he frequently suffered from contact allergy with itchy rashes and ablation of areas of skin. He wrote that a while ago he started to immerse his hands in water with a temperature of about 50 °C, which intensified itching at the beginning but gave relief after a few minutes. He reported that he repeated this five to six times if necessary. Most importantly he wrote that the itching was gone after the procedure and that all skin affections did not occur. He was also able to stop taking corticoid containing creams which he had previously required.

It may be hypothesised that protease-activated receptor-2 plays an important role in contact allergies as it does in atopic dermatitis which can be inhibited/inactivated by heat.⁴⁻⁸ Consequently, there is less histamine liberation and a lesser contact allergic reaction.

I believe that the concept of heat therapy should be tested since there is currently no effective treatment for propolis contact allergy or other contact allergies.

Funding

There was no funding.

Conflict of interest

The author has no conflict of interest to declare.

References

- Müller C, Großjohann B, Fischer L. The use of concentrated heat after insect bites/stings as an alternative to reduce swelling, pain, and pruritus: an open cohort-study at German beaches and bathing-lakes. *Clin Cosmet Investig Dermatol*. 2011;4:191–6.
- Loten C, Stokes B, Worsley D, Seymour JE, Jiang S, Isbister GK. A randomised controlled trial of hot water (45 °C) immersion versus ice packs for pain relief in bluebottle stings. *Med J Aust*. 2006;184:329–33.
- Nomura JT, Sato RL, Ahern RM, Snow JL, Kuwaye TT, Yamamoto LG. A randomised paired comparison trial of cutaneous treatments for acute jellyfish (*Carybdea alata*) stings. *Am J Emerg Med*. 2002;20:624–6.
- Yosipovitch G, Duque MI, Fast K, Dawn AG, Coghill RC. Scratching and noxious heat stimuli inhibit itch in humans: a psychophysical study. *Br J Dermatol*. 2007;156:629–34.
- Yosipovitch G, Fast K, Bernhard JD. Noxious heat and scratching decrease histamine-induced itch and skin blood flow. *J Invest Dermatol*. 2005;125:1268–72.
- Steinhoff M, Neisius U, Ikoma A, Fartasch M, Heyer G, Skov PS, et al. Proteinase-activated receptor-2 mediates human itch: a novel pathway for pruritus in human skin. *J Neurosci*. 2003;23:6176–80.
- Kawagoe J, Takizawa T, Matsumoto J, Tamiya M, Meek SE, Smith AJ, et al. Effect of protease-activated receptor-2 deficiency on allergic dermatitis in the mouse ear. *Jpn J Pharmacol*. 2002;88:77–84.
- Lee SE, Jeong SK, Lee SH. Protease and protease-activated receptor-2 signaling in the pathogenesis of atopic dermatitis. *Yonsei Med J*. 2010;51:808–22.

Karsten Münstedt

Department of Obstetrics and Gynaecology,
Justus-Liebig-University of Giessen, Giessen, Germany
E-mail address: karsten.muenstedt@gyn.med.uni-giessen.de