

Comparison of Acute Inflammatory and Chronic Structural Asthma-Like Responses between C57BL/6 and BALB/c Mice

Chris L. Van Hove^a Tania Maes^a Didier D. Cataldo^c Maud M. Guéders^c
Els Palmans^b Guy F. Joos^a Kurt G. Tournoy^a

^aDepartment of Respiratory Medicine, Ghent University, and ^bHogeschool Gent, Ghent and

^cDepartment of Respiratory Diseases, University of Liège and CHU of Liège, GIGA-Research, Liège, Belgium

Key Words

Airway remodelling · Asthma · Mouse models · TGF- β

Abstract

Background: The interactions between airway responsiveness, structural remodelling and inflammation in allergic asthma remain poorly understood. Prolonged challenge with inhaled allergen is necessary to replicate many of the features of airway wall remodelling in mice. In both mice and humans, genetic differences can have a profound influence on allergy, inflammation, airway responsiveness and structural changes. **Methods:** The aim of this study was to provide a comparative analysis of allergen-induced airway changes in sensitized BALB/c and C57BL/6 mice that were exposed to inhaled allergen for 2 ('acute'), 6 or 9 weeks ('chronic'). Inflammation, remodelling and responsiveness were analyzed. **Results:** Both strains developed a Th-2-driven airway inflammation with allergen-specific IgE, airway eosinophilia and goblet cell hyperplasia upon 2 weeks of allergen inhalation. This was accompanied by a significant increase in airway smooth muscle mass and hyperresponsiveness in BALB/c but not in C57BL/6 mice. However, airway eosinophilia was more pronounced in the C57BL/6 strain. Chronic allergen exposure (6 or 9 weeks) resulted in an increase in airway smooth muscle mass as well as subepithelial collagen

and fibronectin deposition in both strains. The emergence of these structural changes paralleled the disappearance of inflammation in both C57BL/6 and BALB/c mice and loss of hyperresponsiveness in the BALB/c strain. TGF- β_1 was accordingly elevated in both strains. **Conclusion:** Airway inflammation, remodelling and hyperresponsiveness are closely intertwined processes. Genetic background influences several aspects of the acute allergic phenotype. Chronic allergen exposure induces a marked airway remodelling that parallels a decreased inflammation, which was largely comparable between the two strains.

Copyright © 2009 S. Karger AG, Basel

Introduction

In addition to inflammation-related changes such as cellular infiltration and oedema, asthmatic airways display structural alterations referred to as 'airway remodelling' [1, 2]. Historically, animal models have proven useful for the elucidation of the immunological mechanisms of the Th-2-driven airway inflammation that occurs in the asthmatic airway. These models of allergic airway inflammation usually rely on short-term (1 day to 2 weeks) inhaled allergen after a systemic sensitization [3]. These acute models are not adequate to study the features of




Fig. 1. Allergen exposure protocols. Sensitized C57BL/6 and BALB/c mice were exposed to OVA aerosols. **a** Protocol A: mice exposed for 2 weeks. **b** Protocol B: mice exposed for 6 weeks. **c** Protocol C: mice exposed for 9 weeks.

airway remodelling observed in patients with asthma, as longer periods of allergen exposure are required to mimic these changes [4, 5]. The pathophysiology of airway remodelling remains largely unknown, but the structural alterations in the airway are thought to result from (inadequate) attempts at tissue repair in response to inflammation [6]. In addition, it is currently unclear to what extent this process contributes to the deterioration of asthma symptoms [7] or represents the translation of a repair response that protects against further allergen-induced airway inflammation and bronchoconstriction [8]. The pattern of these structural changes differs between mouse strains [9], suggesting the genetic background might be important. Whether remodelling develops uniformly in all asthma patients is equally unclear [10].

Remarkably, prolonged challenge with inhaled allergen in sensitized mice is often associated with disappearance of eosinophilic airway inflammation and the induction of a state of respiratory tolerance [11–14]. While the immunological mechanisms behind this phenomenon are still poorly understood, tolerance appears to develop in most models using chronic allergen challenge [9, 14].

Although a wide variety of mouse asthma models exist [3, 9, 15, 16], it remains puzzling how airway inflammation, remodelling and responsiveness relate to each other in these models. Thus, we here aimed to provide a thorough characterisation encompassing an assessment of inflammation, airway (hyper)responsiveness (AHR) and the remodelling features seen in the 2 mouse strains most commonly used in the study of asthma, C57BL/6 and BALB/c. We provide this comparative analysis in order to gain insight into the relationship between these asthma characteristics in both strains. We found that upon short-

term challenge, the strains developed differential responses in inflammation, AHR and airway remodelling, particularly airway smooth muscle (ASM) proliferation. Upon prolonged exposure, the strains developed more similar structural airway alterations that paralleled a disappearance of eosinophilic inflammation and AHR.

Materials and Methods

Animals

C57BL/6 and BALB/c mice (all males, 6–8 weeks old) were purchased from Harlan (Zeist, The Netherlands). All experimental procedures were approved by the local ethics committee for animal experiments (Faculty of Medicine and Health Sciences, Ghent University).

Allergen Exposure Protocols

All groups of mice (6–12 mice/group) were sensitized with 10 µg intraperitoneal (i.p.) ovalbumin (OVA; Grade III; Sigma, St. Louis, Mo., USA) adsorbed to 1 mg Al(OH)₃ on days 0 and 7. From day 14 onward, the mice were exposed to aerosolized (Ultrasonic nebulizer Sirius Nova, Heyer Medizintechnologie, Bad Ems, Germany) OVA (1% wt/v) or PBS 30 min/day, 3 times a week for 2 ('acute'), 6 or 9 ('chronic') weeks (fig. 1).

Bronchoalveolar Lavage Fluid: Cellular Analysis

Twenty-four hours after the last aerosol exposure, mice were sacrificed with an i.p. pentobarbital injection (60 mg/kg; Sanofi, Libourne, France). Briefly, bronchoalveolar lavage fluid (BALF) was taken by instillation of Hanks' balanced salt solution via a tracheal cannula. Three lavages with 0.3 ml Hanks' solution followed by 3 lavages with 1 ml Hanks' solution were performed. The recovered BALF from the first 3 fractions was centrifuged and the supernatant was used for cytokine detection. The cell pellet was then added to the rest of the lavage fluid, centrifuged, subjected to red blood cell lysis and resuspended for cell counts on cytospins (May-Grünwald/Giemsa).

Histology

After fixation of the left lung with 4% paraformaldehyde, slices from all left lobes were embedded in paraffin for histological analysis. Sections of 3 μm were stained with Congo Red to highlight eosinophils, Periodic-Acid Schiff (PAS) to highlight goblet cells and with Sirius Red for collagen staining.

Immunohistochemistry

Immunohistochemistry (IHC) for fibronectin was done in an analogous manner to that previously described in the rat [17]. Briefly, lung sections of 3 μm were deparaffinized, rehydrated and the nonspecific binding sites were blocked with 1% blocking reagent in PBS (Boehringer, Mannheim, Germany). Excess reagent was removed and the sections were incubated for 1 h with a mouse anti-mouse fibronectin antibody (dilution factor 1/750; Dako A/S, Glostrup, Denmark). Afterwards, sections were rinsed and incubated with a biotinylated secondary antibody for 30 min. Next, the primary antibody-secondary antibody complex was detected by streptavidin-biotinylated horseradish peroxidase complex. The substrate for the peroxidase was 3, 3'-diaminobenzidine (DAB; Dako), resulting in a brown reaction product for fibronectin quantification. IHC for α -actin smooth muscle contractile elements in the airways (ASM) was performed in an analogous manner, using an anti-mouse α -actin SM antibody (dilution factor 1/100; Dako) and DAB substrate, also resulting in a brown reaction product.

Quantitative Measurements in the Airway Wall

Quantitative measurements were performed in the airways of each animal with a perimeter of basement membrane (Pbm) ranging from 800 to 2,000 μm , provided a reasonable cross section was available (ratio of maximal to minimal internal diameter <1.8). Measurements were performed on the digital representation of the airways using a Zeiss KS400 image analyser system (Oberkochen, Germany), as described earlier [17]. Morphometrical parameters were marked manually: the area defined by the basement membrane (Abm) and the area defined by the total adventitial perimeter (Ao). The total bronchial wall area (Wat) was calculated ($\text{Wat} = \text{Ao} - \text{Abm}$) and normalized to the square of the Pbm. For the quantification of collagen deposition, the area in the airway wall covered by the Sirius Red stain (WCt) was determined by the software and normalized to Pbm. To evaluate the subepithelial fibronectin deposition and the ASM content of the airways, the area covered by the DAB stain (WFt and W α A) was determined in an analogous manner and also normalized to Pbm. Goblet cells were quantified on PAS stained sections. Results were expressed as the number of goblet cells per millimeter of basement membrane. Peribronchial infiltration with eosinophils was evaluated in lung sections stained with Congo Red and the total number of eosinophils per mm^2 of bronchial wall was determined. All measurements were performed on 5 airways per mouse.

IgE and Protein Quantification (ELISA)

OVA-IgE in serum was measured with ELISA using coated microtiter plates and biotinylated polyclonal rabbit anti-mouse IgE (S. Florquin, Université Libre de Bruxelles, Brussels, Belgium). OVA-specific IgE are expressed in arbitrary units (U) per ml serum (an OVA-IgE U). One OVA-IgE U was defined as a 1:100 dilution of an internal standard serum pool obtained from

mice sensitized with OVA (10 μg)/alum (1 mg). TGF- β_1 was determined in the bronchoalveolar lavage fluid (BALF) using ELISA kits (R&D Systems, Abingdon, UK).

Assessment of Airway Responsiveness

Twenty-four hours after the last aerosol exposure, AHR to methacholine was assessed in spontaneously breathing animals using a whole body plethysmograph system (Buxco, Buxco Electronics Inc., Troy, N.Y., USA). Before performing readings, the system was calibrated by rapid injection of 1 ml air. Pressure differences between the main chamber containing mice and a reference chamber were recorded using the software BioSystem XA (version 157, Buxco). This pressure signal is caused by flow changes in the main box during the respiratory cycle of the animal. The value of 'enhanced pause' (Penh) is used here to monitor airway function [18], since it shows strong correlation with the airway resistance [19]. $\text{Penh} = [(\text{Te} - \text{Tr})/\text{Tr}] \times (\text{PEP}/\text{PIP})$, where Te is expiratory time (in seconds), Tr is relaxation time or time of the pressure decay to 36% of total box pressure during expiration, PEP is peak expiratory pressure (in $\text{cm H}_2\text{O}$) and PIP is peak inspiratory pressure (in $\text{cm H}_2\text{O}$). For each mouse, data were recorded at baseline, and after exposure to PBS or increasing concentrations of nebulized methacholine (2–81 mg/ml), to assess AHR. The solutions were nebulized through an inlet of the main chamber during 1 min. Readings were taken during 5 min after each nebulization. Between readings, the Penh value returned to baseline. Cumulative dose-response curves were constructed for the changes in Penh after increasing doses of methacholine. The changes in Penh are expressed as percentage increase of maximum Penh value following methacholine challenge compared to the average Penh value after PBS exposure (calculated by Analyst 1.29, Buxco).

Statistical Analysis


Data were analysed with the statistical package SPSS 15.0 (SPSS Inc., Chicago, Ill., USA). Reported values are expressed as mean $\pm$ standard error of the mean. Mean values of parameters (including quantitative measurements in the airway wall as well as inflammatory cells and mediators, but not Penh values) were compared between the groups through the Kruskal-Wallis test for multiple comparisons (nonparametric testing). When significant differences were observed, post-hoc comparisons between groups were made, using the Mann-Whitney U test with Bonferroni corrections. $p < 0.05$ was considered significant. The dose-response curves of Penh were compared using ANOVA. The concentration of methacholine causing a 250% increase in baseline Penh was calculated by log-linear interpolation of the dose-response curve.

Results

Airway Inflammation in the BALF and Bronchial Walls

OVA exposure for 2 weeks (fig. 1a) altered the composition of the BALF leukocytes in both strains (fig. 2) as compared to the PBS-exposed mice. Both strains developed a significant influx of mononuclear cells and eo-

Fig. 2. BALF total cell count and differentiation in OVA-sensitized C57BL/6 and BALB/c mice exposed to OVA or PBS aerosols for 2 weeks (n = 8–12/group; **a**) and 9 weeks (n = 8–12/group; **b**). The BALF total cell counts after 6 weeks of OVA or PBS exposure are not included, as the results are similar to the 9 weeks experiment in both strains. ^a p < 0.05 OVA vs. PBS. ^b p < 0.05 BALB/c vs. C57BL/6.



sinophils. The percentage of eosinophils in BALF was $21.16 \pm 4.41\%$ in C57BL/6 versus $12.66 \pm 4.08\%$ in BALB/c mice ($p < 0.05$ BALB/c vs. C57BL/6). Only C57BL/6 mice had a discrete but significant concomitant BALF neutrophilia and a significant increase in total BALF cellularity and macrophages. Moreover, both strains developed peribronchial and perivascular eosinophilic inflammation upon short-term challenge (fig. 3a, b). C57BL/6 mice, but not BALB/c mice also possessed some degree of interstitial inflammation (fig. 3c, d).

The numbers of eosinophils counted around the airways on Congo Red stained sections were significantly higher in the C57BL/6 strain compared to the BALB/c strain (table 1).

Upon longer OVA exposure (6 or 9 weeks; fig. 1b, c) the BALF cell composition in OVA-challenged mice of both strains returned to the composition discerned in mice exposed to PBS (fig. 2). At those points in time, only a slightly elevated number of eosinophils remained present in the bronchial walls of both mouse strains (table 1).

Serum OVA-IgE

Whereas the PBS-challenged C57BL/6 mice had very low OVA-IgE levels at all points in time, the BALB/c counterparts developed increasing IgE levels attributable uniquely to the i.p. sensitization (table 2). In the OVA-challenged mice of both strains, OVA-IgE had already in-




Fig. 3. Airway histology (Congo Red staining) of OVA-sensitized and challenged BALB/c mice (a) and C57BL/6 mice (b) after 2 weeks of OVA exposure. $\times 200$. c Congo Red stained pictures of the lung parenchyma of OVA-sensitized and challenged (2 weeks) C57BL/6 mice with interstitial inflammation. $\times 400$. d Congo Red stained pictures of the lung parenchyma of OVA-sensitized and challenged (2 weeks) BALB/c mice without interstitial inflammation. $\times 400$.

creased after 2 weeks as compared to their PBS-challenged counterparts. Upon chronic exposure, both strains had elevated IgE; however, the titres were markedly higher in the BALB/c mice as compared to the C57BL/6 mice ($p < 0.05$ BALB/c vs. C57BL/6).

Epithelial Remodelling: Goblet Cells

Exposure to OVA resulted in a significant increase in goblet cells from 2 weeks onward (fig. 4a and 5a), which was comparable in both strains. This increase persisted after 6 and 9 weeks (fig. 4a). However, in both allergen-challenged C57BL/6 and BALB/c mice, the goblet cells did not increase any further upon prolonged challenge.

Airway Smooth Muscle

The amount of ASM did not change over time in the PBS-exposed mice, either in BALB/c or in C57BL/6 mice (fig. 4b and 5b). Exposure to OVA allergen for 2 weeks resulted in a significant increase in ASM staining only in BALB/c mice. However, longer allergen exposure periods induced an increase in ASM in both strains.

Table 1. Peribronchial eosinophils in epithelium in BALB/c and C57BL/6 mice

Exposure time	Treatment	Eosinophils ($\times 10^2$ cells/mm ² airway wall)	
		BALB/c	C57BL/6
2 weeks	PBS	1.49 ± 0.17	1.05 ± 0.17
	OVA	$5.06 \pm 0.30^{a, c}$	6.61 ± 0.46^a
6 weeks	PBS	0.93 ± 0.10	0.57 ± 0.09^b
	OVA	$1.51 \pm 0.13^{a, b}$	$1.11 \pm 0.09^{a, b}$
9 weeks	PBS	0.79 ± 0.11^b	0.69 ± 0.10
	OVA	$1.03 \pm 0.08^{a, b}$	0.79 ± 0.08^b

^a $p < 0.05$, OVA versus PBS.

^b $p < 0.05$, 6 and 9 weeks versus 2 weeks.

^c $p < 0.05$, BALB/c versus C57BL/6.




Fig. 4. Measurement of indicators of airway remodelling with increasing duration of OVA challenge in C57BL/6 and BALB/c mice (n = 6–12 mice/group). **a** Goblet cells (PAS). **b** ASM. * p < 0.05 OVA vs. PBS.

Table 2. OVA-specific IgE levels in the serum after sensitization and exposure of BALB/c and C57BL/6 mice

Exposure time	Treatment	OVA-specific IgE (U/ml)	
		BALB/c	C57BL/6
2 weeks	PBS	7.60 ± 1.02	2.22 ± 0.55
	OVA	41.75 ± 5.28 ^a	33.50 ± 7.19 ^a
6 weeks	PBS	29.11 ± 2.54 ^{b, c}	6.38 ± 1.80
	OVA	96.30 ± 10.67 ^{a-c}	49.18 ± 9.00 ^a
9 weeks	PBS	23.00 ± 1.90 ^{b, c}	3.43 ± 1.14
	OVA	102.20 ± 10.63 ^{a-c}	21.83 ± 4.01 ^a

^a p < 0.05, OVA versus PBS

^b p < 0.05, 6 weeks and 9 weeks versus 2 weeks.

^c p < 0.05, BALB/c versus C57BL/6.

Morphometry and Deposition of Extracellular Matrix Proteins

Morphometry

WAt or airway wall thickness of large-sized airways did not differ in either strain at 2, 6 or 9 weeks of PBS (fig. 6a). Exposure to OVA allergen for 2 weeks resulted in a significant increase in WAt only in the BALB/c mice when compared to sham-exposed mice. However, longer allergen exposures induced a significant and persistent increase in WAt in both strains.

Fibronectin and Collagen

Fibronectin and collagen deposition were similarly increased in both genotypes after 6 and 9 weeks, respectively (fig. 5c, d and 6b, c).




Fig. 5. Histology pictures of the airways of C57BL/6 mice challenged with OVA or PBS aerosols for 2 or 9 weeks. Original magnifications $\times 100$, an inset with higher magnification is included. All stains were performed in the BALB/c strain as well, leading to analogous results at these points in time. **a** PAS staining to highlight goblet cells in C57BL/6 mice challenged with OVA or PBS for 2 weeks. **b** IHC with DAB substrate to highlight the increase in ASM contractile elements (brown colour) in BALB/c mice challenged with OVA or PBS for 9 weeks. **c** IHC with DAB substrate to highlight fibronectin deposition (brown colour) in C57BL/6 mice challenged with OVA or PBS for 9 weeks. **d** Sirius Red staining to highlight collagen deposition in C57BL/6 mice challenged with OVA or PBS for 9 weeks.




Fig. 6. Measurement of indicators of airway remodelling with increasing duration of OVA challenge in C57BL/6 and BALB/c mice (n = 6–12 mice/group). **a** Airway wall thickening (Congo Red). **b** Fibronectin deposition (IHC). **c** Collagen deposition (Sirius Red). * p < 0.05 OVA vs. PBS.

Airway Hyperresponsiveness

BALB/c but not C57BL/6 mice developed a significant leftward shift in the dose-response curve to inhaled methacholine after 2 weeks of OVA exposure (ANOVA; p < 0.005 vs. PBS; figure 7 a, d). After a longer exposure period (6 or 9 weeks), no difference in the dose-response curve was observed for either strain (fig. 7 b–c, e–f). The

C57BL/6 mice were less responsive to methacholine than BALB/c mice. At 2 weeks, the concentration of methacholine that caused a 250% increase in Penh from baseline in allergen-exposed C57BL/6 was 17.40 ± 5.92 mg/ml versus 4.69 ± 1.81 mg/ml in BALB/c mice (p < 0.05 BALB/c vs. C57BL/6).










Fig. 7. AHR to nebulized methacholine in sensitized BALB/c (**a–c**) and C57BL/6 mice (**d–f**) exposed to OVA for 2 weeks (**a, d**), 6 weeks (**b, e**) and 9 weeks (**c, f**). ANOVA OVA vs. PBS at 2 weeks: $p < 0.005$.

Fig. 8. Levels of TGF- β_1 in the BALF of BALB/c mice (a) and C57BL/6 mice (b) sensitized to OVA and exposed to OVA or PBS aerosols for 2 weeks and 9 weeks. * $p < 0.05$ OVA vs. PBS.



BALF Proteins
TGF- β_1 levels in BALF were persistently increased after 2 or 9 weeks of allergen exposure compared to control PBS-exposed mice in both strains (fig. 8).

Discussion

We here document that in the 2 mouse strains most commonly used to model asthma (C57BL/6 and BALB/c), genetic background and the duration of allergen exposure have a profound impact on different aspects of the resulting pathological phenotype. Although the acute response in both strains was predominated by eosinophilic airway inflammation, only the BALB/c strain developed a significant ASM hyperplasia and AHR. In an attempt to mimic more closely the features of chronic asthma, the response to longer periods of allergen exposure was predominated by airway remodelling, while inflammation and AHR decreased in both strains. Although other groups have earlier described strain differences in response to various durations of OVA aerosol exposure [9], the current study gives a more detailed insight into the relationship between inflammation, remodelling and responsiveness in the 2 most commonly used mouse strains.

Inflammation, AHR and airway remodelling are thought to be interdependent processes. A large variety of mouse models focus on a limited number of characteristics of asthma to study the underlying pathogenesis [14, 15]. To study inflammation, these models mainly rely on (hyper-)acute, short-term exposure protocols. The reason for this is that the use of short-term exposure protocols is practical but also avoids the establishment of re-

spiratory tolerance that can develop after prolonged allergen exposure [11–14]. However, long-term exposure protocols are usually applied to study the main features of airway remodelling [4, 5, 14, 16]. In these models, inflammation and lung function tests are often either absent or partially described [4, 5, 20]. We chose to provide here a comprehensive analysis encompassing not only the assessment of inflammation and responsiveness, but also a thorough characterisation of remodelling characteristics.

Here, we found that inflammation and remodelling behave differently in both strains, especially in the acute models, based on short-term allergen exposure (2 weeks; fig. 1a). Both strains developed eosinophilic inflammation, as evident from BALF analysis and quantification of peribronchial eosinophilia. However, airway eosinophilia was more pronounced in the C57BL/6 mice in the acute model. This is in line with one earlier study, that also found that C57BL/6 mice are more susceptible to the development of airway eosinophilia [21]. In contrast, Shinagawa and Kojima [9] found a more pronounced eosinophilia in BALB/c mice after repeated (chronic) intranasal OVA instillation. We presume the method of antigen exposure is critical in this respect, as we administered OVA 1% by aerosol whereas Shinagawa and Kojima administered OVA 0.1% intranasally. Serum OVA-IgE levels were more elevated in the BALB/c mice than in C57BL/6 after prolonged OVA challenge (6 and 9 weeks; fig. 1a, b) confirming earlier findings that showed that BALB/c mice are IgE prone [22]. This is also underscored by the fact that BALB/c mice, but not C57BL/6, already have measurable IgE titres after sensitization only.

In the BALB/c mice we could observe a significant increase in ASM after just 2 weeks of OVA challenge ('acute';

fig. 1a), in contrast to the C57BL/6 strain, where longer durations of allergen exposure were required to obtain such an ASM increase. In addition, AHR was present in BALB/c mice at that point in time. This histological observation now adds up to the earlier descriptions that inbred mouse strains exhibit significant genetic variability in the airway calibre and development of AHR after antigen sensitization and exposure [9, 23]. Although the contraction of ASM is the ultimate translation of asthma pathology into physiologic disturbance in patients, it remains an unresolved issue if a larger smooth muscle mass is really the cause of AHR [24, 25]. Some studies indicate a clear link between ASM and AHR in both animal models and humans, since both AHR and increased ASM can persist after resolution of the airway inflammation [26–28], and mathematical models predict that airways with increased ASM narrow to a much greater extent than airways with a smaller ASM volume [29]. One could speculate that the lack of AHR after 2 weeks of OVA exposure in C57BL/6 mice could be related to the absence of ASM hyperplasia. However, other factors could be involved. Firstly, the C57BL/6 strain developed a certain degree of interstitial inflammation, which was earlier also reported by another group [30]. Secondly, the airway anatomy of both strains is different [31]. Both of these factors could be of potential relevance to explain why AHR was observed only in the BALB/c strain.

Prolonged allergen challenge (fig. 1b, c) results in increased fibronectin and collagen deposition in the airway wall in both strains which is concordant with findings described by other authors [20, 26, 32–35]. We here document that this structural remodelling occurs in parallel with a decrease in inflammatory cells in the BALF compartment to baseline in both strains and loss of AHR in BALB/c mice. We and others previously described that prolonged OVA challenge in these mice results in the establishment of respiratory tolerance, as they are unresponsive to new sensitizations [12]. To date, the precise mechanisms of the development of respiratory tolerance in mice repeatedly challenged with OVA aerosols have not been elucidated, and both regulatory T cell-dependent [36] and regulatory T cell-independent [37] mechanisms have been proposed. One factor that may be important in the down-regulation of allergic airway inflammation and AHR over time could be the aging of mice, because some studies found that the capability of T cells to differentiate into Th-2 cells was reduced in aging mice [38, 39]. In contrast, one recent study found that older mice developed increased allergic inflammation, but less pronounced AHR [40]. Moreover, we earlier documented

that mice over 20 weeks old can easily develop Th-2-mediated airway inflammation [12]. Thus, the precise effect of aging on allergic airway disease remains a matter of debate [39–41].

The exact role of the eosinophils in remodelling, inflammation and AHR is speculative. It is possible that eosinophils could contribute to the initiation of collagen synthesis by fibroblasts during the acute inflammatory phase [42]. In our opinion, the few remaining eosinophils in the bronchial wall upon prolonged allergen exposure represent a small but measurable fraction deemed to vanish shortly thereafter. In BALB/c mice, the decrease of eosinophilic inflammation and increase of fibrosis in the airway wall occurs in parallel with the loss of AHR. It is thus tempting to speculate that the remodelling process could occur in an attempt to protect against allergen-induced AHR and inflammation. Collagen and fibronectin deposition could indeed increase airway wall stiffness and oppose extended narrowing of the airway wall. Moreover, the subepithelial fibrosis could decrease the amount of allergen exposure by shielding the immune system from the allergens ('wash-away' effect).

TGF- β_1 is a key molecule involved in the regulation of both inflammation and remodelling in both humans and mouse models [43–46], and therefore represents a potential therapeutic target of interest. Increases in TGF- β_1 are a constant finding for both mouse strains used here, in the acute and in the chronic exposure models. It could therefore be speculated that TGF- β_1 is activated quite early as an endogenous 'anti-asthma' molecule to control inflammation [45, 46], to promote remodelling [43, 44, 47] and thus to inhibit AHR [48]. However, previous studies using neutralising antibodies for TGF- β_1 produced conflicting results on its role. One study found that anti-TGF- β_1 antibody treatment could antagonize matrix deposition, smooth muscle cell proliferation and mucus production without affecting the airway inflammation [47]. In contrast, in a recent report based on an intranasal HDM administration it was shown that anti-TGF- β_1 treatment did not inhibit remodelling, but rather induced an exacerbation of the allergen-induced airway changes suggesting that this mediator is not per se responsible for airway remodelling [49]. These authors even warned against the use of therapeutic strategies aimed at interfering with TGF- β_1 .

The current study has some limitations. Firstly, for measuring airway responsiveness, we used the whole body plethysmograph, a method that has been criticised. There is evidence that Penh and airway resistance do not correlate under certain conditions, although plethysmography

remains a valuable tool for explorative analysis [19, 50]. When Penh can be used as a surrogate for invasive measurements, it can decrease cost, time and the number of animals required for experiments. Secondly, in this study we could not provide in-depth explorations into the precise mechanisms that regulate the relationship between airway inflammation, remodelling and responsiveness. Although the limitations of this study should be taken into account, the current data provide researchers with clues to choose an appropriate asthma model for the particular aspects of the disease they want to investigate. There is currently no mouse model that completely mimics the pathology and physiology of patients with asthma.

In conclusion, genetic background has an impact on the different aspects of the acute allergic phenotype.

These differences are far less pronounced when studying chronic allergen exposure, albeit the latter models are characterized mainly by remodelling but not inflammation or AHR.

Acknowledgements

The authors gratefully acknowledge Marie-Rose Mouton, Eliane Castrique, Ann Neessen, Indra De Borle, Christelle Snaauwaert and Kathleen De Saedeleer for their technical assistance. Funding was received from the Fund for Scientific Research – Flanders (FWO-Vlaanderen, project G.0052.06) Belgian IAP P6/35. K.G.T. is a clinical investigator supported by FWO-Vlaanderen.

References

- Jeffery PK: Remodeling in asthma and chronic obstructive lung disease. *Am J Respir Crit Care Med* 2001;164:S28–S38.
- Bousquet J, Jeffery PK, Busse WW, Johnson M, Vignola AM: Asthma: from bronchoconstriction to airways inflammation and remodeling. *Am J Respir Crit Care Med* 2000;161:1720–1745.
- Lloyd CM, Gonzalo JA, Coyle AJ, Gutierrez-Ramos JC: Mouse models of allergic airway disease. *Adv Immunol* 2001;77:263–295.
- Kips JC, Anderson GP, Fredberg JJ, Herz U, Inman MD, Jordana M, Kemeny DM, Lotvall J, Pauwels RA, Plummer CG, Schmidt D, Sterk PJ, van Oosterhout AJ, Vargaftig BB, Chung KF: Murine models of asthma. *Eur Respir J* 2003;22:374–382.
- Ramos-Barbon D, Ludwig MS, Martin JG: Airway remodeling: lessons from animal models. *Clin Rev Allergy Immunol* 2004;27:3–21.
- Fixman ED, Stewart A, Martin JG: Basic mechanisms of development of airway structural changes in asthma. *Eur Respir J* 2007;29:379–389.
- Holgate ST, Davies DE, Powell RM, Howarth PH, Haitchi HM, Holloway JW: Local genetic and environmental factors in asthma disease pathogenesis: chronicity and persistence mechanisms. *Eur Respir J* 2007;29:793–803.
- James AL, Wenzel S: Clinical relevance of airway remodelling in airway diseases. *Eur Respir J* 2007;30:134–155.
- Shinagawa K, Kojima M: Mouse model of airway remodeling: strain differences. *Am J Respir Crit Care Med* 2003;168:959–967.
- Amin K, Ludviksdottir D, Janson C, Nettelbladt O, Bjornsson E, Roomans GM, Boman G, Seveus L, Venge P: Inflammation and structural changes in the airways of patients with atopic and nonatopic asthma: BHR Group. *Am J Respir Crit Care Med* 2000;162:2295–2301.
- Schramm CM, Puddington L, Wu C, Guernsey L, Gharaee-Kermani M, Phan SH, Thrall RS: Chronic inhaled ovalbumin exposure induces antigen-dependent but not antigen-specific inhalational tolerance in a murine model of allergic airway disease. *Am J Pathol* 2004;164:295–304.
- Van Hove CL, Maes T, Joos GF, Tournoy KG: Prolonged inhaled allergen exposure can induce persistent tolerance. *Am J Respir Cell Mol Biol* 2007;36:573–584.
- Sakai K, Yokoyama A, Kohno N, Hamada H, Hiwada K: Prolonged antigen exposure ameliorates airway inflammation but not remodeling in a mouse model of bronchial asthma. *Int Arch Allergy Immunol* 2001;126:126–134.
- Kumar RK, Foster PS: Modeling allergic asthma in mice: pitfalls and opportunities. *Am J Respir Cell Mol Biol* 2002;27:267–272.
- Zosky GR, Sly PD: Animal models of asthma. *Clin Exp Allergy* 2007;37:973–988.
- Kariyawasam HH, Robinson DS: Airway remodelling in asthma: models and supermodels? *Clin Exp Allergy* 2005;35:117–121.
- Palmans E, Kips JC, Pauwels RA: Prolonged allergen exposure induces structural airway changes in sensitized rats. *Am J Respir Crit Care Med* 2000;161:627–635.
- Hamelmann E, Schwarze J, Takeda K, Oshiba A, Larsen GL, Irvin CG, Gelfand EW: Noninvasive measurement of airway responsiveness in allergic mice using barometric plethysmography. *Am J Respir Crit Care Med* 1997;156:766–775.
- Finkelman FD: Use of unrestrained, single-chamber barometric plethysmography to evaluate sensitivity to cholinergic stimulation in mouse models of allergic airway disease. *J Allergy Clin Immunol* 2008;121:334–335.
- Tsuchiya T, Nishimura Y, Nishiuma T, Kotani Y, Funada Y, Yoshimura S, Yokoyama M: Airway remodeling of murine chronic antigen exposure model. *J Asthma* 2003;40:935–944.
- Morokata T, Ishikawa J, Ida K, Yamada T: C57BL/6 mice are more susceptible to antigen-induced pulmonary eosinophilia than BALB/c mice, irrespective of systemic T helper 1/T helper 2 responses. *Immunology* 1999;98:345–351.
- Hamelmann E, Takeda K, Oshiba A, Gelfand EW: Role of IgE in the development of allergic airway inflammation and airway hyperresponsiveness: a murine model. *Allergy* 1999;54:297–305.
- Brewer JP, Kisselgof AB, Martin TR: Genetic variability in pulmonary physiological, cellular, and antibody responses to antigen in mice. *Am J Respir Crit Care Med* 1999;160:1150–1156.
- O'Byrne PM, Inman MD: Airway hyperresponsiveness. *Chest* 2003;123:411S–416S.
- Gil FR, Lauzon AM: Smooth muscle molecular mechanics in airway hyperresponsiveness and asthma. *Can J Physiol Pharmacol* 2007;85:133–140.

- 26 Leigh R, Ellis R, Wattie J, Southam DS, De Hoogh M, Gauldie J, O'Byrne PM, Inman MD: Dysfunction and remodeling of the mouse airway persist after resolution of acute allergen-induced airway inflammation. *Am J Respir Cell Mol Biol* 2002;27:526–535.
- 27 Southam DS, Ellis R, Wattie J, Inman MD: Components of airway hyperresponsiveness and their associations with inflammation and remodeling in mice. *J Allergy Clin Immunol* 2007;119:848–854.
- 28 Kariyawasam HH, Aizen M, Barkans J, Robinson DS, Kay AB: Remodeling and airway hyperresponsiveness but not cellular inflammation persist after allergen challenge in asthma. *Am J Respir Crit Care Med* 2007;175:896–904.
- 29 Moreno RH, Hogg JC, Pare PD: Mechanics of airway narrowing. *Am Rev Respir Dis* 1986;133:1171–1180.
- 30 Takeda K, Haczku A, Lee JJ, Irvin CG, Gelfand EW: Strain dependence of airway hyperresponsiveness reflects differences in eosinophil localization in the lung. *Am J Physiol Lung Cell Mol Physiol* 2001;281:L394–L402.
- 31 Thiesse JR, Namati E, de Ryck JC, Reinhardt JM, Hoffman EA, McLennan G: Anatomy of the normal airway tree in three strains of mice using micro-CT and pathology techniques. *Am J Respir Crit Care Med* 2007;175:A528.
- 32 Locke NR, Royce SG, Wainwright JS, Samuel CS, Tang ML: Comparison of airway remodeling in acute, subacute, and chronic models of allergic airways disease. *Am J Respir Cell Mol Biol* 2007;36:625–632.
- 33 Ellis R, Leigh R, Southam D, O'Byrne PM, Inman MD: Morphometric analysis of mouse airways after chronic allergen challenge. *Lab Invest* 2003;83:1285–1291.
- 34 Blyth DI, Pedrick MS, Savage TJ, Hessel EM, Fattah D: Lung inflammation and epithelial changes in a murine model of atopic asthma. *Am J Respir Cell Mol Biol* 1996;14:425–438.
- 35 Temelkovski J, Hogan SP, Shepherd DP, Foster PS, Kumar RK: An improved murine model of asthma: selective airway inflammation, epithelial lesions and increased methacholine responsiveness following chronic exposure to aerosolised allergen. *Thorax* 1998;53:849–856.
- 36 Carson IV WF, Guernsey LA, Singh A, Vella AT, Schramm CM, Thrall RS: Accumulation of regulatory T cells in local draining lymph nodes of the lung correlates with spontaneous resolution of chronic asthma in a murine model. *Int Arch Allergy Immunol* 2007;145:231–243.
- 37 Niu N, Le Goff MK, Li F, Rahman M, Homer RJ, Cohn L: A novel pathway that regulates inflammatory disease in the respiratory tract. *J Immunol* 2007;178:3846–3855.
- 38 Hasegawa A, Miki T, Hosokawa H, Hossain MB, Shimizu C, Hashimoto K, Kimura MY, Yamashita M, Nakayama T: Impaired GATA3-dependent chromatin remodeling and Th2 cell differentiation leading to attenuated allergic airway inflammation in aging mice. *J Immunol* 2006;176:2546–2554.
- 39 Gelfand EW, Joetham A, Cui ZH, Balhorn A, Takeda K, Taube C, Dakhama A: Induction and maintenance of airway responsiveness to allergen challenge are determined at the age of initial sensitization. *J Immunol* 2004;173:1298–1306.
- 40 Busse PJ, Zhang TF, Srivastava K, Schofield B, Li XM: Effect of ageing on pulmonary inflammation, airway hyperresponsiveness and T and B cell responses in antigen-sensitized and -challenged mice. *Clin Exp Allergy* 2007;37:1392–1403.
- 41 Yagi T, Sato A, Hayakawa H, Ide K: Failure of aged rats to accumulate eosinophils in allergic inflammation of the airway. *J Allergy Clin Immunol* 1997;99:38–47.
- 42 Kariyawasam HH, Robinson DS: The role of eosinophils in airway tissue remodelling in asthma. *Curr Opin Immunol* 2007;19:681–686.
- 43 Lloyd CM, Robinson DS: Allergen-induced airway remodelling. *Eur Respir J* 2007;29:1020–1032.
- 44 Boxall C, Holgate ST, Davies DE: The contribution of transforming growth factor-beta and epidermal growth factor signalling to airway remodelling in chronic asthma. *Eur Respir J* 2006;27:208–229.
- 45 Schmidt-Weber CB, Blaser K: Regulation and role of transforming growth factor-beta in immune tolerance induction and inflammation. *Curr Opin Immunol* 2004;16:709–716.
- 46 Wan YY, Flavell RA: 'Yin-Yang' functions of transforming growth factor-beta and T regulatory cells in immune regulation. *Immunol Rev* 2007;220:199–213.
- 47 McMillan SJ, Xanthou G, Lloyd CM: Manipulation of allergen-induced airway remodeling by treatment with anti-TGF-beta antibody: effect on the Smad signaling pathway. *J Immunol* 2005;174:5774–5780.
- 48 Alcorn JF, Rinaldi LM, Jaffe EF, van Loon M, Bates JH, Janssen-Heininger YM, Irvin CG: Transforming growth factor-beta1 suppresses airway hyperresponsiveness in allergic airway disease. *Am J Respir Crit Care Med* 2007;176:974–982.
- 49 Fattouh R, Midence G, Arias K, Johnson JR, Walker TD, Goncharova S, Souza KP, Gregory RC, Lonning S, Gauldie J, Jordana M: TGF-beta regulates house dust mite-induced allergic airway inflammation but not airway remodeling. *Am J Respir Crit Care Med* 2008;177:593–603.
- 50 Glaab T, Ziegert M, Baelder R, Korolewitz R, Braun A, Hohlfeld JM, Mitzner W, Krug N, Hoymann HG: Invasive versus noninvasive measurement of allergic and cholinergic airway responsiveness in mice. *Respir Res* 2005;6:139.