

✓ ISSN: 2455-8141

⚙ Global Journal of Allergy (<https://www.peertechz.com/journals/global-journal-of-allergy>)

📄 Research Article 🔓 Open Access 📝 Peer-Reviewed

The skin prick test results to saffron, sumac and barberry in patients with atopy

Mozhgan Moghtaderi*

+ Author and article information

+ Abstract

Quick Enquiry

Background: Food allergens appear to play a role in the etiology and deteriorating of atopy in some patients, little is known about hypersensitivity to some common food additives in these patients. The purpose of this study was to identify probable sensitization to saffron, sumac and barberry in patients with atopy.

Materials and Methods: This cross-sectional study included 390 patients with atopy and 300 healthy individuals with no history of atopic diseases. Skin tests were performed in both patient and control group with saffron, sumac and barberry.

Results: A positive skin test to saffron was seen in 61 (15.6%), to sumac in 29 (7.4%) and to barberry in 36 (9.2%) patients with atopy. None of individuals in control group showed sensitization to three studied extracts.

Conclusion: The results of the present study revealed sensitization to saffron, sumac and barberry in patients with atopy. A further research with food challenges is required to confirm food allergy in those patients with sensitization to saffron, sumac and barberry.

+ Main article text

Introduction

Allergens are primarily proteins, capable of stimulating IgE synthesis in genetically susceptible people. They enter to human body by inhalation, contact and ingestion. These allergens may induce allergic diseases such as asthma, Allergic Rhinitis (AR) and Atopic Dermatitis (AD). Ingested allergens are thought to play an important role in the development and worsening the clinical symptoms in patients with atopy [1-3].

Saffron (*Crocus sativus*) is a kind of cooking spice that cultivated in some parts of Iran and near the studied area. It's now an essential part of some Eastern, Middle Eastern, and European dishes. It is expensive because only a small amount of each saffron flower is used and all harvesting must be done by hand. Clinical allergic symptoms to saffron are reported in eye, nose and respiratory system through an IgE-dependent mechanism [4,5].

Sumac (*Rhus coriaria*) in form of dried and powdered fruits is widely used in Asian countries. It is essential ingredient in Middle Eastern cooking. Sumac has been served as acidic taste in cooking prior to the introduction of lemons by the Romans. The reactions to sumac present as dermatitis in children and occupational disease [6,7].

Barberry (*Berberis vulgaris*) as a sharp acid flavor is used the most is Iran. *Berberis vulgaris* grows in the wild in much of Europe and West Asia. They are used in jams, a common flavoring for soft drinks and Persian rice with

sugar. This plant is considered mostly allergy free and causes little or no allergy problems in people [8].

Skin prick test (SPT) is a primary test for the diagnosis of IgE-mediated food allergy and it is safe, inexpensive, informative, and easy to perform [9].

A number of patients with atopy describe clinical allergic symptoms after ingestion of some ingredients of foods. This study was designed to investigate how much saffron, sumac, and barberry may elicit positive skin test in patients with atopy in compared with control group.

Methods and Materials

This cross-sectional study was performed on 390 patients with atopy and 300 individuals without atopy from June to November 2016. All the patients were from southwestern Iran who was regularly referred to an allergy clinic at Namazee Hospital affiliated to Shiraz University of Medical Sciences, Iran. The diagnosis of asthma was based on the Expert Panel Report 3, AR according to allergic rhinitis and its impact on asthma and AD based on the criteria of Hanifin and Rajka in order with the supervision of allergist [10-12]. Clear, watery discharge and itching in the eyes were considered allergic conjunctivitis and the timeline of 6 weeks of daily or nearly daily urticaria was considered as chronic urticaria [13,14].

The study protocol was approved by the Ethics Committee of Shiraz University of Medical Sciences (93-8302), and informed consent was obtained from patients and control individuals after the study was described in detail.

Demographic data including sex, age and type of atopy (allergic rhinitis, asthma, chronic urticaria, allergic conjunctivitis) were collected. Preparation of extract of saffron, sumac and barberry was done based on modified method of Kwaasi, et al., [15,16]. SPT were performed on the forearms using manufactured extracts of saffron, sumac and barberry and the results were

measured 15 minutes after application. The tests were considered positive sensitization when wheal diameter was ≥ 3 mm greater in diameter than the negative control in response to the extract after 15 min application. Histamine (10 mg/mL) and saline were used as positive and negative controls, respectively. Antihistamine medications were stopped 7 days before the skin prick tests in all individuals.

Results and Discussion

Three hundred and ninety patients (233 females and 157 males) with atopy ranging in age from 10 to 50 years (mean age 30.63 ± 10.72) were enrolled in the study. Three hundred individuals (205 female and 95 male) ranging in age between 15 and 52 years (mean age 31.27 ± 10.22) were included as control group. There were no significant differences for sex and age between the two groups.

A positive skin test to saffron was seen in 61 (15.6%), to sumac in 29 (7.4%) and to barberry in 36 (9.2%) patients with atopy. None of individuals in control group showed sensitization to three studied extracts. There was no relationship between sensitization to extract of each saffron, sumac and barberry with age and sex in these patients.

Positive sensitization to saffron extract reported 12.5% (21/167) in saffron workers of Khorasan (Iran), this rate is near similar to our patients with atopy [5]. An earlier study showed sensitization to saffron in Spanish saffron workers 6% and in atopic patients 4.2% [17]. Saffron is commonly grown in Spain similar to Iran for commercial purposes; sensitization is higher in Iranian than Spanish patients. There is no sufficient data for comparison of sensitization rate to sumac and barberry in patients with atopy in the literature.

Table 1 shows type of atopy and the number of patients with positive sensitization to saffron, sumac and barberry.

--

Table 1: The rate of positive sensitization to saffron, sumac and barberry in patients with atopy.

Atopic disease	number (%)	Positive to saffron	Positive to sumac	Positive to barberry
			number	
Allergic rhinitis	117 (30.1)	15	8	9
Asthma	153 (39.2)	22	8	13
Chronic urticaria	73(18.7)	12	6	6
Allergic conjunctivitis	4(1)	2	1	0
Asthma& Allergic rhinitis	43(11)	10	6	8
Total	390(100)	61	29	36

Positive sensitization to three studied allergen was 27.3% (32/117) in patients with allergic rhinitis, 28.1% (43/153) in asthmatic patients and 55.8% (24/43) in individuals with asthma& allergic rhinitis. Having both asthma and allergic rhinitis show severity of atopy, therefore, it causes more chance for sensitization to allergens.

From 73 patients with chronic urticaria 32.8% (24/73) showed sensitization to studied food allergens. Contrast our result; Rajan JP, et al., study reported food additives are a rare cause for chronic urticaria by doing double-blind placebo-controlled food challenge. [18].

We had 4 patients with allergic conjunctivitis, 2 had positive reaction to saffron and one to sumac. It is better considering SPT to studied food allergens in those patients with allergic conjunctivitis, although the number of our patients was small.

Positive skin test is not enough to confirm the presence of allergic disease, however; it shows allergic sensitization which may predict the subsequent onset of allergic symptoms. It is noticeable that cross-reactivity with other allergens may elicit positive skin test in individuals with no clinical symptoms.

The results of the present study revealed sensitization to saffron, sumac and barberry in patients with atopy. Food challenges are considered the gold accurate standard method for diagnosing food allergy; however, this procedure requires cooperation of the patient and preparation of the office for the challenge. The major limitation of this study is need to food challenge for establishing allergy to saffron, sumac and barberry.

Authors' contributions

All authors contributed to this project and article equally. All authors read and approved the final manuscript.

+ References

1. Kelso JM (2014) Potential food allergens in medications. *J Allergy Clin Immunol* 133: 1509-1518. Link: <http://bit.ly/2qbGkjlw> (<http://bit.ly/2qbGkjlw>)
2. Moghtaderi M, Hejrati Z, Dehghani Z, Dehghani F, Kolahi N (2016) Sensitization to Food Additives in Patients with Allergy: A Study Based on Skin Test and Open Oral Challenge. *Iran J Allergy Asthma Immunol* 15: 198-203. Link: <http://bit.ly/2WBvEXJ> (<http://bit.ly/2WBvEXJ>)
3. Gonzales-González VA, Díaz AM, Fernández K, Rivera MF (2018) Prevalence of food allergens sensitization and food allergies in a group of allergic Honduran children. *Allergy Asthma Clin Immunol* 14: 23. Link: <http://bit.ly/2Nc3zTL> (<http://bit.ly/2Nc3zTL>)
4. Moshiri M, Vahabzadeh M, Hosseinzadeh H (2015) Clinical Applications of Saffron (*Crocus sativus*) and its Constituents: A Review. *Drug Res* 65: 287-295. Link: <http://bit.ly/2N87ZuD> (<http://bit.ly/2N87ZuD>)
5. Fereydouni M, Sankian M, Varasteh AR (2002) The prevalence of saffron pollen allergy in saffron workers of Khorasan (Iran) 12: 7-13. Link: <http://bit.ly/2N6ONxu> (<http://bit.ly/2N6ONxu>)
6. Wang S, Zhu F (2018) Quality attributes of bread fortified with staghorn sumac extract. *J Texture Stud* 49: 129-134. Link: <http://bit.ly/32agOIA> (<http://bit.ly/32agOIA>)
7. Rayne S, Mazza G (2007) Biological activities of extracts from sumac (*Rhus* spp): A review. *Plant Foods Hum Nutr* 62: 165-175. Link: <http://bit.ly/36B1usa> (<http://bit.ly/36B1usa>)
8. Kukula-Koch W, Aligiannis N, Halabalaki M, Skaltsounis AL, Glowinski K, et al. (2013) Influence of extraction procedures on phenolic content and antioxidant activity of Cretan barberry herb. *Food Chem* 138: 406-413. Link: <http://bit.ly/2NaMdGJ> (<http://bit.ly/2NaMdGJ>)
9. Carr TF, Saltoun CA (2012) Chapter 2: Skin testing in allergy. *Allergy Asthma Proc* 33: S6-S8. Link: <http://bit.ly/33bCGoy> (<http://bit.ly/33bCGoy>)

10. National Asthma Education and Prevention Program (2007) Expert panel report 3 (EPR-3): Guidelines for the diagnosis and management of asthma-summary report 2007. *J Allergy Clin Immunol* 120: S94-S138. Link: <http://bit.ly/2PFEDFy> (<http://bit.ly/2PFEDFy>)
11. Bousquet J, Schunemann H, Fonseca J, Samolinski B, Bachert C, et al. (2015) MACVIA-ARIA Sentinel Network for allergic rhinitis (MASK-rhinitis): The new generation guideline implementation. *Allergy* 70: 1372-1392. Link: <http://bit.ly/2oDRRaR> (<http://bit.ly/2oDRRaR>)
12. Hanifin JM, Rajka G (1980) Diagnostic features of atopic dermatitis. *Acta Derm Venereol* 92: 44-47. Link: <http://bit.ly/36B2sEO> (<http://bit.ly/36B2sEO>)
13. Baab S, Kinzer EE (2018) Allergic Conjunctivitis. StatPearls [Internet]. Treasure Island (FL): Stat Pearls Publishing.
14. Godse KV, Zawar V, Krupashankar D, Girdhar M, Kandhari S, et al. (2011) Consensus statement on the management of urticaria. *Indian J Dermatol* 56: 485-489. Link: <http://bit.ly/2r3PzTf> (<http://bit.ly/2r3PzTf>)
15. Kwaasi AA, Parhar RS, Harfi H, Tipirneni P, al-Sedairy ST (1992) Characterization of antigens and allergens of date palm (*Phoenix dactylifera*) pollen. Immunologic assessment of atopic patients by whole extract and its fractions [corrected]. *Allergy* 47: 535-544. Link: <http://bit.ly/2pyCj8Q> (<http://bit.ly/2pyCj8Q>)
16. Varasteh AR, Vahedi F, Sankian M, Kaghazian H, Tavallaie S, et al. (2007) Specific IgG antibodies (total and subclasses) against Saffron pollen: a study of their correlation with specific IgE and immediate skin reactions. *Iran J Allergy Asthma Immunol* 6: 189-195. Link: <http://bit.ly/2PLHH2W> (<http://bit.ly/2PLHH2W>)
17. Feo F, Martinez J, Martinez A, Galindo PA, Cruz A, et al. (1997) Occupational allergy in saffron workers. *Allergy* 52: 633-641. Link: <http://bit.ly/2WAbA7N> (<http://bit.ly/2WAbA7N>)
18. Rajan JP, Simon RA, Bosso JV (2014) Prevalence of sensitivity to food and drug additives in patients with chronic idiopathic urticaria. *J Allergy Clin Immunol Pract* 2: 168-171. Link: <http://bit.ly/32aiFgw> (<http://bit.ly/32aiFgw>)

+ Copyright

© 2019 Moghtaderi M. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

+ View Similar Articles

We recommend

A Great Discovery Allergy and Asthma are fully Genetic in Children

Arnaldo Cantani*, Journal of Addiction Medicine and Therapeutic Science, 2015

Listeria Monocytogenes as a Foodborne Pathogen: Biocontrol in Foods using Lytic Bacteriophages

Naim Deniz Ayaz et al., Journal of Addiction Medicine and Therapeutic Science, 2016

Nanotechnology and Nutraceuticals

Swarnali Das Paul et al., International Journal of Nanomaterials, Nanotechnology and Nanomedicine, 2015

Impact of Food Allergies on the Allergic Person's Travel Decision, Trip Organization and Stay Abroad

Dano D et al., Global Journal of Allergy, 2015

Coprecipitation procedure for speciation of chromium in some dairy food product and water samples by FAAS

Şerife Saçmacı*, Journal of Addiction Medicine and Therapeutic Science, 2017

Need for Open Science: Study on Effects of Sucrose Hidden for 50 years

Enago Academy, 2017

Progress on the Chemical Constituents Derived from Glucosinolates in Maca (*Lepidium meyenii*)

Yan-Jie Huang et al., Natural Products and Bioprospecting, 2018

Building the Base for Promoting Research in Food Science & Technology — Workshop at KoSFoST International Symposium, Korea

Enago Academy, 2016

Black phosphorus-based field effect transistor devices for Ag ions detection*

Hui-De Wang et al., Chinese Physics B, 2018

Efficacy of nintedanib and docetaxel in patients with advanced lung adenocarcinoma treated with first-line chemotherapy and second-line immunotherapy in the nintedanib NPU program

Powered by **TREND MD**

I consent to the use of Google Analytics and related cookies across the TrendMD network (widget, website, blog).

[Learn more \(https://www.trendmd.com/google-analytics\)](https://www.trendmd.com/google-analytics)

Yes

No

+ Share your thoughts and experiences

peertechz Comment Policy

<https://www.peertechz.com/privacy-policy>

Please read our [Comment Policy](#) before commenting.



What do you think?

0 Responses

Upvote

Funny

Love

Surprised

Angry

Sad

Comments Community

1 Login

Recommend

Tweet

Share

Sort by Best

Start the discussion...

LOG IN WITH

OR SIGN UP WITH DISQUS ?

Name

Be the first to comment.

ALSO ON PEERTECHZ

Kambo: A ritualistic healing substance from

India – A Preferred Destination for IVF

4 Comments

Sort by Top

Add a comment...

**Peertechz Journals**

Cytoprotection in Multiple Sclerosis and Ischemic Stroke with C-Phycocyanin and Phycocyanobilin

Like · Reply · 3w

**Davis Skola**

Two years ago, I learnt about a successful Hepatitis B Virus treatment from Herbal Health Point (www.herbalhealthpoint.com), I was on the treatment for over 7 months and tested negative to the virus after I completed the treatment. Its almost unbelievable how effective the treatment was alleviating the virus. I completed the treatment two years ago and still negative!

Like · Reply · 2w

**Felicity Tina**

Hello i am TINA, I am publicizing this amazing testimonial of my life to you because. i knew of so many people who will still be having this same issues, which you are finding so difficult to solve. I and my Wife, have been suffering from HIV for the past 2 years. we have visited doctors but they couldn't get the permanent solution to my sickness. I almost loosed it until one faithful day when i saw so many amazing testimonials about DOCTOR, RENE how he has cured patients infected with HIV. So i told my HUSBAND and we both decide to give it a try and contact him, here we are now. All thanks to DOCTOR RENE for curing our HIV permanently. Now we are both perfectly... [See more](#)

Like · Reply · 2w

**Skola Davies**

Five years ago I was admitted to the hospital with abdominal pain and fever, and was diagnosed with Hepatitis B. I was discharged after 2 weeks, I was on Hepsera for few months. But the mild pains still persist on the right side of my abdomen. Two years ago, I learnt about a successful Hepatitis B Virus treatment from Herbal Health Point (www.herbalhealthpoint.com), I was on the treatment for over 7 months and tested negative to the virus after I completed the treatment. Its almost unbelievable how effective the treatment was alleviating the virus. I completed the treatment two years ago and still negative!

Like · Reply · 4w



Quick Enquiry



(https://crossmark.crossref.org/dialog/?doi=10.17352/2455-8141.000022&domain=pdf&date_stamp=2019-11-04)

Download as ▼

Order for reprints (<mailto:reprints@peertechz.com>)

Send URL to Your Friend

Article Alerts

Subscribe to our articles alerts and stay tuned.

Subscribe Now!



(<http://creativecommons.org/licenses/by/4.0/>) This work is licensed under a Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>).