



Differential effect of appendectomy and tonsillectomy on anti-*Kudoa* sp. antibodies in patients with MALTectomy

J.C. Andreu-Ballester^a, F. Ballester^b, J. Pérez-Griera^c, V. Amigo^d, C. Peñarroja-Otero^e, E. Colomer-Rubio^a, I. Ortiz-Tarín^f, V. Pelayo^g, P. García-Hernández^g, M. Rodero^g, C. Cuéllar^{g,*}

^a Research Unit, Emergency Department, Arnau de Vilanova Hospital, 46015 Valencia, Spain

^b Unit of Epidemiology and Statistics, Valencian School of Studies for Health, 46017 Valencia, Spain

^c Biopathology Department, Clinic University Hospital, 46010 Valencia, Spain

^d Hematology Department, Arnau de Vilanova Hospital, 46015 Valencia, Spain

^e Internal Medicine Department, Casa de Salud Hospital, 46021 Valencia, Spain

^f Department of Surgery, Arnau de Vilanova Hospital, 46015 Valencia, Spain

^g Department of Parasitology, Faculty of Pharmacy, Complutense University, 28040 Madrid, Spain

ARTICLE INFO

Article history:

Received 8 May 2009

Received in revised form 31 July 2009

Accepted 4 August 2009

Available online 13 August 2009

Keywords:

Kudoa sp. antigen

Myxosporidia

Appendectomy

Tonsillectomy

Mucosa Associated Lymphoid Tissue (MALT)

system

Ulcerative colitis

ABSTRACT

We found an association between tonsillectomized patients and subsequent appendicitis. We also observed that MALTectomy significantly decreased secretory IgA levels in serum of patients, being this decrease more pronounced when both operations (tonsillectomy and appendectomy) had been performed. The elevated humoral responses detected previously by us in BALB/c mice immunized with *Kudoa* sp. pseudocyst extracts and the high IgG1 and IgE levels induced by the oral administration of *Kudoa* sp. pseudocysts to BALB/c mice showed the possible immunopathological effects in man from the ingestion of *Kudoa* sp. infected fish. We use the ELISA method to investigate the possible relationship between MALTectomy (tonsillectomy and appendectomy) and specific antibody levels to *Kudoa* sp. Both anti-*Kudoa* sp. specific antibody levels and the number of patients that recognized *Kudoa* sp. antigens were greater in tonsillectomy patients when compared to the control and the other studied groups (appendectomized and appendectomized + tonsillectomies patients). Tonsillectomy was associated to a switch in the class of immunoglobulins involved in these responses and these responses may be abrogated by appendectomy. Tonsils and appendix may respond in different ways to *Kudoa* sp. antigens and these different reactions may be involved in some immunopathological reactions.

© 2009 Elsevier Ireland Ltd. All rights reserved.

1. Introduction

The MALT system (Mucosa Associated Lymphoid Tissue), in the digestive tract GALT (Gut Associated Lymphoid Tissue), is an important part of secondary lymphoid tissues and is comprised of Peyer's patches of terminal ileum, and the appendix. In the nasopharyngeal area it is denominated NALT system (Nasopharyngeal Associated Lymphoid Tissues) composed of tonsils palatines, adenoids and tongue [1–3]. In adult human beings, 80–85% of immune cells are located in the digestive tract mucosa, hence the importance of the

MALT in human immunology [4]. The appendix is highly vascular, contains a high concentration of lymphoid tissue, and produces immune system cells normally involved with the MALT [5–7].

In previous studies, we found an association between tonsillectomized patients and subsequent appendicitis [8,9]. Likewise [10], we studied the influence of the surgical removal of two important parts of the MALT (tonsils and appendix) on immune parameters. MALTectomy significantly decreased secretory IgA levels in serum of patients, being this decrease more pronounced when both operations (tonsillectomy and appendectomy) had been performed. Recently, we found that appendectomy significantly diminished specific immunoglobulin levels in serum against two food borne parasite antigens (*Anisakis* and *Kudoa* sp.) [11]. This decrease was detectable from three months to three years post-appendectomy. These facts prompted us to study the influence of the surgical removal of other important parts of the MALT on these anti-parasite humoral immune responses.

The majority of *Kudoa* species infect somatic muscle of fish, establishing cysts which contain many spores. There is currently no effective method to detect infected fish without destroying them, for this reason parasitized fish reach the consumer. Although there is no information about the prevalence of *Kudoa* sp. in fish markets in the

Abbreviations: MALT system, Mucosa Associated Lymphoid Tissue; GALT, Gut Associated Lymphoid Tissue; NALT system, Nasopharyngeal Associated Lymphoid Tissues; AP, Appendectomy; T, Tonsillectomy; T + AP, Appendectomy and tonsillectomy; C, Control group without either operation; Ig's, Total immunoglobulins; HRP, Horse radish peroxidase; O.D., Optical density; OR, Odds Ratio; CI 95%, Confidence Interval of 95%.

* Corresponding author. Tel.: +34 913941818; fax: +34 913941815.

E-mail address: cuellarh@farm.ucm.es (C. Cuéllar).

studied region, multivalvulid myxozoan parasites, including *Kudoa* spp. have been described from fish from the Mediterranean coastal waters of Spain [12]. Likewise, in Spain, *Kudoa*-infected fish have lately been detected in both fresh and frozen imported Chilean hake (*Merluccius gayi gayi*) destined for human consumption. The elevated humoral responses detected previously by us in BALB/c mice immunized with *Kudoa* sp. pseudocyst extracts and the high IgG1 and IgE levels induced by the oral administration of *Kudoa* sp. pseudocysts to BALB/c mice showed the possible immunopathological effects in man from the ingestion of *Kudoa* sp. infected fish [13]. Previously [14] we investigated the seroprevalence of anti-*Kudoa* sp. antibodies in a Spanish healthy population and the possible association between the manifestation of allergic reactions after fish consumption and the humoral responses to *Kudoa* sp. antigens. Likewise, we determined specific anti-*Kudoa* sp. specific antibodies in sera of patients diagnosed with several digestive pathologies. Positive reactions to *Kudoa* sp. were observed in patients by means of the skin prick test [15]. For these reasons, this report aims to investigate the possible relationship between MALTectomy (tonsillectomy and appendectomy) and the levels of specific antibodies to *Kudoa* sp.

2. Materials and methods

2.1. Patients and serum samples

A population of adult patients was randomly selected from the database of appendectomized (AP) patients, treated over the last three years in our hospital (Arnau de Vilanova Hospital, Valencia, Spain). Patients who had been operated on in the last three months were excluded. By means of a telephone interview (in chronological order of intervention, starting with the most recently operated patient), information concerning a possible previous tonsillectomy (T) was obtained. So, patients were split into a group of 40 patients who had had both operations (T + AP), and another group of patients who had only undergone appendectomy. These groups were sorted according to sex, age (± 2 years) and date of operation (the patient nearest on the list). Further, another two groups were also randomly selected from patients presenting to the emergency service: one group of patients who had had a tonsillectomy but not appendectomy; and the other group made up of controls that had not had either operation. These groups of patients were paired by sex and age with the two former groups. We excluded patients with any acute infections or inflammatory diseases, as well as those who had received immunosuppressive treatment or any kind of vaccine during the previous year. The age and the date when they had been operated on were also noted. The anatomopathological report on the appendicitis (type and size of appendices) was included. The tonsillectomy was confirmed by otorrinolaringological exploration. We examined a total of 160 patients. In this way, our study population was made up of four groups of 40 patients per group: Group 1 [patients with AP + T operations]; Group 2 [patients with only appendectomy (AP)]; Group 3 [patients with only tonsillectomy (T)]; Group 4 [control group without either operation (C)]. Each participant in the study gave their consent and the study was approved by the Ethics and Investigation Committee of our hospital.

Total immunoglobulins (Ig's), IgG, IgM, IgA and IgE against *Kudoa* sp. antigen were determined in the sera of the selected patients.

2.2. Antigens

"White" *Kudoa* sp. pseudocysts were manually obtained from the skeletal musculature of Chilean hake (*M. gayi gayi*) from local fish markets and destined for human consumption. Pseudocysts were separated from any associated fish tissue and then homogenised. In order to release the spore contents, glass beads were added and

"white" *Kudoa* sp. pseudocysts were shaken and homogenates were extracted in PBS [13].

2.3. Specific antibody levels

Each well of 96 well microtitre plates (Costar, Corning, NY, USA) was coated by the addition of 10 μ g/ml of *Kudoa* sp. antigen. Duplicate dilutions of sera at 1/100 in PBS-Tween, containing 0.1% BSA were added and incubated. Horse radish peroxidase (HRP) conjugate goat anti-human Ig's, IgM, IgG or IgA (Biosource International, Camarillo, CA, USA) was used [16,17].

For the IgE determination, test sera were added in duplicate at a 1/2 dilution. A murine monoclonal antibody against an epsilon human IgE chain (IgG1 κ , E21A11, INGENASA, Madrid, Spain) was added and incubated, followed by a goat anti-mouse IgG1 (gamma) HRP conjugate (CALTAG Laboratories, Burlingame, California) [13,17]. Anti-*Kudoa* sp. positive and negative sera from mice were used as ELISA controls [13].

Values higher than the mean of the optical density (O.D.) of 160 subjects obtained for each immunoglobulin and antigen plus their respective standard deviation were considered as positive.

2.4. Determination of biological indicators of immunological status

Cells were stained for the surface markers CD8, CD4, CD3, CD19, CD2, CD8 and CD56. The data were collected and analyzed by flow cytometry (Coulter Epics XL-MCL; Beckman-Coulter). Nephelometry (BN-II Analyzer, Behring Diagnostics GmbH, Marburg, Germany) was employed for serum IgG and IgM assay. Total IgA, IgA1, IgA2 and secretory IgA were determined by ELISA using monoclonal mouse anti-human antibodies (HyTest Ltd., Turku, Finland) [10].

2.5. Statistical analysis

Statistical analysis was performed using the SPSS statistical package. Descriptive characteristics of the patients as well as variables of the immunoglobulin levels in each group were obtained. Contingency tables were made for the categorical variables. Their frequencies were compared by means of Pearson's Chi-Square test. To compare the mean values of these variables, a repeated measure ANOVA with contrast analysis of types of differences was used. Normality in distribution of the variables of immunoglobulin levels was confirmed by the Kolmogorov-Smirnov test. Sphericity was verified by the Mauchly test. In the case of non-sphericity, a low limit correction was obtained. Friedman's ANOVA non-parametric test was used to study the variables that did not have a normal distribution, and the exact or asymptotic test was chosen depending on the case. *p* values <0.05 were considered to be statistically significant. The Pearson's test was used to study the correlations.

3. Results

We examined a total of 160 subjects [40 appendectomy (AP), 40 tonsillectomy (T), 40 AP + T patients and 40 controls (C)], 76 were females (47.5%) and 84 were males (52.5%). The mean age ($\pm$ SD) was 39.2 ± 14 years (CI 95%: 37.0–41.4) range 20–75. The four groups were matched for sex and age. In each of the four groups there were 21 men and 19 women. The mean age at which the tonsillectomy was performed was 6.87 years (CI 95%: 5.14–8.61). The mean age at which the appendectomy was performed was 36.33 years (CI 95%: 31.83–40.82). The selected population in our study was appendectomized between 3 months and 3 years before blood testing, and tonsillectomized 20 years before our investigation.

Anti-*Kudoa* sp. immunoglobulin levels are shown in the Fig. 1. Total antibody prevalences for *Kudoa* sp. Ig's, IgG, IgM, IgA and IgE




Fig. 1. Specific serum responses to *Kudoa* sp. antigen, expressed as optical density values in the studied subjects. C, control; AP, appendectomy; T, tonsillectomy; AP + T, appendectomy + tonsillectomy. Each point represents a single serum. * $p < 0.001$, # $p = 0.001$. Repeated measures ANOVA (IgG and IgM) was used when the values were distributed according to a normal distribution in all the groups. Friedman's ANOVA (Ig's, IgA and IgE) was used when in some of the groups distribution was not normal. IgE ($p = 0.101$).

were 20.6%, 23.1%, 20.6%, 13.1% and 11.3%, respectively. The obtained values for each serum are shown in Fig. 1.

The highest levels of specific antibodies were obtained in the tonsillectomized patients, followed by the control group, the appendectomized group and the tonsillectomized–appendectomized group. This fact was also observed when we analyzed the percentages of sera with any positive anti-*Kudoa* sp. antibody class (Fig. 2). In all the




Fig. 2. Percentages of sera with positive anti-*Kudoa* sp. antibodies according to GALTectomy groups. C, control [OR = 0.52 (CI 95% = 0.24–1.13)]; AP, appendectomy [OR = 1.07 (CI 95% = 0.52–2.21)]; T, tonsillectomy [OR = 10.12 (CI 95% = 4.24–24.17)]; AP + T, appendectomy + tonsillectomy [OR = 0.14 (CI 95% = 0.05–0.38)]. * $p < 0.0001$.

cases significant differences were observed with the exception of the IgE (Fig. 1). No correlations were observed between the time from tonsillectomy or the patient's age and the levels of anti-*Kudoa* sp. antibodies. High correlations were observed between the levels of anti-*Kudoa* sp. Ig's, IgG, IgM and IgA ($p < 0.01$).

When the prevalence of detectable antibodies against *Kudoa* sp. in all the groups [control (C), appendectomy (AP), tonsillectomy (T) and AP + T] were compared, the following percentages of positive were obtained: total Ig's (C = 17.5%, AP = 12.5%, T = 52.5%, AP + T = 0%); IgG (C = 17.5%, AP = 25.0%, T = 45.0%, AP + T = 5.0%); IgM (C = 15.0%, AP = 20.0%, T = 45.0%, AP + T = 2.5%); IgA (C = 12.5%, AP = 7.5%, T = 30.0%, AP + T = 2.5%) and IgE (C = 12.5%, AP = 17.5%, T = 10.0%, AP + T = 5%).

The results of lymphocytic populations and non-specific immunoglobulins fall within the parameters of normality. In the lymphocyte population, the low values of CD4⁺ T cells in the tonsillectomized patients are noted, although there are not statistical significances [$975.50 \pm 361.28 \times 10^9/l$ versus 1122.75 ± 482.29 (T + AP), 1009.25 ± 377.16 (AP), 1044.25 ± 344.10 (C)]. When the comparisons of means were carried out, there were no significant differences between the groups, except to the secretory IgA levels. Levels of secretory IgA were significantly lower in the tonsillectomy + appendectomy group than in the other three groups ($p < 0.001$). Levels of secretory IgA correlated to anti-*Kudoa* sp. IgE levels (correlation coefficient = 0.300; $p < 0.05$).

Likewise, the groups of patients that had only one of the operations showed lower values than the control group ($p < 0.001$). Likewise, levels of secretory IgA1 were higher in the sera with positive anti-*Kudoa* sp. antibodies ($p = 0.026$). No correlations were observed when the anti-*Kudoa* sp. specific immunoglobulins were compared to the total immunoglobulin levels.

4. Discussion

In this study, we have found that specific Ig's, IgG, IgM and IgA anti-*Kudoa* sp. levels were raised in tonsillectomized patients. As in the general population, tonsillectomized individuals take contact with parasites several times in their lives. It has been shown that some parasites can be isolated from tonsils suggesting that they may play an important role in avoiding parasites travelling along the digestive tract and preventing potential pathogenicity [18–21]. In the absence of tonsils, parasites may contact other MALT system structures, especially appendix, and induce a strong immunological response and hypertrophy. This could be related with the association previously found between tonsillectomy and subsequent acute appendicitis [8,9].

Interestingly, the results obtained by us in the present study supported this idea. First, the tonsillectomy group of patients showed a very significant increase in the number of anti-*Kudoa* sp. positive individuals with the exception of the IgE ($p < 0.0001$) pointing to the filter role of tonsils as the first MALT defence line. Second, the anti-*Kudoa* sp. antibody levels were more intense in the tonsillectomy group than that observed in the control group ($p < 0.001$) supporting our previous clinical observations. Finally, these responses can be abrogated when appendectomy was undertaken after a previous tonsillectomy (32/40 responders in the tonsillectomy group versus 5/40 responders in the tonsillectomy + appendectomy group, $p < 0.0001$). Taken all together, our results pointed to a central role of the tonsil and associated structures as a potent effector system in gut's immunological responses and their removal may alter the immune response of the gut.

Infectious agents, such parasites, can be present in the appendix of predisposed individuals and the appendix can serve as an amplifier of the proinflammatory chain reactions increasing the exposure to normal enteric microflora antigens [19]. Epidemiological studies have also demonstrated that appendectomy is a protective factor against ulcerative colitis in humans [22–30].

So far, it is conceivable that, under certain conditions—immunological repertoire and constitution or environmental factors including bacterial and non-bacterial antigens, prionic proteins [31,32] or parasites—appendix immunological responses may lead, through abnormal responses, to ulcerative colitis and Crohn's disease in humans. Further clinical and immunological studies of the role of appendix in inflammatory bowel disease have to be undertaken.

Experimental studies are necessary to confirm the reproducibility and significance of these clinical findings. Similar immunological investigations with other antigens are in progress to prove the important immunological consequences of MALTectomy.

In summary, both anti-*Kudoa* sp. specific antibody levels and the number of patients that recognized *Kudoa* sp. antigens were greater in tonsillectomy patients (T) when compared to the control (C) and the other studied groups [(AP) and (AP + T)]. Furthermore, a secretory IgA reduction was observed in tonsillectomized and appendectomized patients, probably due to a reduction in MALT system mass. No correlations were observed when anti-*Kudoa* sp. specific immunoglobulins were compared to the total immunoglobulin levels. This finding showed that the decrease in anti-*Kudoa* sp. levels did not simply reflect the reduction of total immunoglobulin levels. Also, tonsillectomy could be associated with a switch in the class of immunoglobulins involved in these responses and these anti-*Kudoa* sp. responses may be abrogated by appendectomy. Interestingly,

levels of secretory IgA1 were higher in the sera with positive anti-*Kudoa* sp. antibodies which could be suggesting the relationship of this IgA subclass to the immune responses against *Kudoa* sp. antigens. It seems that tonsils and appendix may respond in different ways to *Kudoa* sp. antigens and these different reactions may be involved, under appropriated conditions, in some immunopathological reactions. In order to improve the knowledge on the interaction of *Kudoa* parasites with the MALT system, the examination of secretory antibody responses in saliva, intestinal mucus, or stool, could be an interesting study. Works are in progress to check the immunoglobulin production against other common antigens (including antigens from non-food borne pathogens) to make sure that the immunoglobulin response in control and MALTectomy groups is a generalized phenomenon and it is specific to mucosal immune responses.

Acknowledgments

This work was supported by Grant PR27/05-13950-BSCH. We also are grateful to F. Bru, A. Portero, I. Torres, V. Valls (Emergency Department, Arnau de Vilanova Hospital, Valencia, Spain), A.J. Hueso, A. Payá, C. Enguidanos, A. Almiñana (Hematology and Biological Chemistry Department, Arnau de Vilanova Hospital, Valencia, Spain), G. Martínez de Velasco and Y. Laó (Department of Parasitology, Faculty of Pharmacy, Complutense University, Madrid, Spain).

References

- [1] Bienenstock J, McDermott M, Befus D, O'Neill M. A common mucosal immunologic system involving the bronchus, breast and bowel. *Adv Exp Med Biol* 1978;107:53–9.
- [2] Mestecky J, McGhee JR. Immunoglobulin A (IgA): molecular and cellular interactions involved in IgA biosynthesis and immune response. *Adv Immunol* 1987;40:153–245.
- [3] Mestecky J, Blumberg RS, Kiyono H, McGhee JR. The mucosal immune system. In: Paul WE, editor. *Fundamental immunology*. Philadelphia: Lippincott Williams & Wilkins; 2003. p. 965–1020.
- [4] Brandtzaeg P, Halstensen TS, Kett K, Krajci P, Kvale D, Rognum TO, et al. Immunobiology and immunopathology of human gut mucosa: humoral immunity and intraepithelial lymphocytes. *Gastroenterology* 1989;97:1562–84.
- [5] Fisher RE. The primate appendix: a reassessment. *Anat Rec* 2000;261:228–36.
- [6] Somekh E, Serour F, Gorenstein A, Vohl M, Lehman D. Phenotypic pattern of B cells in the appendix: reduced intensity of CD19 expression. *Immunobiology* 2000;201:461–9.
- [7] Nagler-Anderson C. Man the barrier! Strategic defences in the intestinal mucosa. *Nat Rev Immunol* 2001;1:59–67.
- [8] Andreu-Ballester JC, Colomer RE, Millan Scheiding M, Ballester F. Tonsillectomy and risk of acute appendicitis. *Am J Gastroenterol* 2003;98:2806–7.
- [9] Andreu-Ballester JC, Ballester F, Colomer-Rubio E, Millan Scheiding M. Association between tonsillectomy, adenoidectomy, and appendicitis. *Rev Esp Enferm Dig* 2005;97:179–86.
- [10] Andreu-Ballester JC, Pérez-Griera J, Ballester F, Colomer-Rubio E, Ortiz-Tarin I, Peñarroja-Otero C. Secretory immunoglobulin A (sIgA) deficiency in serum of patients with GALTectomy (appendectomy and tonsillectomy). *Clin Immunol* 2007;123:289–97.
- [11] Andreu-Ballester JC, Pérez-Griera J, Ballester F, Colomer-Rubio E, Ortiz-Tarin I, Pelayo V, et al. *Anisakis simplex* and *Kudoa* sp.: evaluation of specific antibodies in appendectomized patients. *Exp Parasitol* 2008;119:433–6.
- [12] Yurakhno VM, Ovcharenko MO, Holzer AS, Sarabiev VL, Balbuena JA. *Kudoa unicapsula* n. sp. (Myxosporea: Kudoidae) a parasite of the Mediterranean mullets *Liza ramada* and *L. aurata* (Teleostei: Mugilidae). *Parasitol Res* 2007;101:1671–80.
- [13] Martínez de Velasco G, Cuéllar C. Humoral immune responses induced by *Kudoa* sp. (Myxosporea: Multivalvulida) in BALB/c mice: oral administration, immunization and cross-reactions with *Myxobolus aeglefini* (Myxosporea: Bivalvulida). *Parasite Immunol* 2003;25:449–56.
- [14] Martínez de Velasco G, Rodero M, Chivato T, Cuéllar C. Seroprevalence of anti-*Kudoa* sp. (Myxosporea: Multivalvulida) antibodies in a Spanish population. *Parasitol Res* 2007;100:1205–11.
- [15] Martínez de Velasco G, Rodero M, Cuéllar C, Chivato T, Mateos JM, Laguna R. Skin prick test of *Kudoa* sp. antigens in patients with gastrointestinal and/or allergic symptoms related to fish ingestion. *Parasitol Res* 2008;103:713–5.
- [16] Daschner A, Cuéllar C, Sánchez-Pastor S, Pascual CY, Martín-Esteban M. Gastro-allergic anisakiasis as a consequence of simultaneous primary and secondary immune response. *Parasite Immunol* 2002;24:243–51.
- [17] Gutiérrez R, Cuéllar C. Immunoglobulins anti-*Anisakis simplex* in patients with gastrointestinal diseases. *J Helminthol* 2002;76:131–6.
- [18] Amendoeira MR, Coutinho SG. Isolation of *Toxoplasma gondii* from the saliva and tonsils of a three-year-old child. *J Infect Dis* 1982;145:587.

- [19] Rosemberg S, Lopes MB, Masuda Z, Campos R, Vieira Bressan MC. Fatal encephalopathy due to *Lagochilascaris minor* infection. *Am J Trop Med Hyg* 1986;35:575–8.
- [20] Bhargava D, Raman R, El Azzouni MZ, Bhargava K, Bhusnurmath B. Anisakiasis of the tonsils. *J Laryngol Otol* 1996;110:387–8.
- [21] Zhao C, Papadopolou B, Tremblay MJ. *Leishmania infantum* promotes replication of HIV type 1 in human lymphoid tissue cultured ex vivo by inducing secretion of the proinflammatory cytokines TNF-alpha and IL-1 alpha. *J Immunol* 2004;172:3086–93.
- [22] Gilat T, Hachohen D, Lilos P, Langman MJ. Childhood factors in ulcerative colitis and Crohn's disease. An international cooperative study. *Scand J Gastroenterol* 1987;22:1009–24.
- [23] Lowenfels AB. Appendectomy and ulcerative colitis. *Gastroenterology* 1994;107:1570.
- [24] Rutgeerts P, D'Haens G, Hiele M, Vantrappen G. Appendectomy protects against ulcerative colitis. *Gastroenterology* 1994;106:1251–3.
- [25] Russel MG, Dorant E, Brummer RJ, Van der Kruijs MA, Muris JW, Bergers JM, et al. Appendectomy and the risk of developing ulcerative colitis or Crohn's disease: results of a large case control study. South Limburg Inflammatory Bowel Disease Study Group. *Gastroenterology* 1997;113:377–82.
- [26] Andersson RE, Olaison G, Tysk C, Ekblom A. Appendectomy and protection against ulcerative colitis. *N Engl J Med* 2001;344:808–14.
- [27] Koutroubakis IE, Vlachonikolis IG. Appendectomy and the development of ulcerative colitis: results of a metaanalysis of published case-control studies. *Am J Gastroenterol* 2000;95:171–6.
- [28] Koutroubakis IE, Vlachonikolis IG, Kouroumalis EA. Role of appendicitis and appendectomy in the pathogenesis of ulcerative colitis: a critical review. *Inflamm Bowel Dis* 2002;8:277–86.
- [29] Firouzi F, Bahari A, Aghazadeh R, Zali MR. Appendectomy, tonsillectomy, and risk of inflammatory bowel disease: a case control study in Iran. *Int J Colorectal Dis* 2006;21:155–9.
- [30] Jiang L, Xia B, Li J, Ye M, Deng C, Ding Y, et al. Risk factors for ulcerative colitis in a Chinese population: an age-matched and sex-matched case-control study. *J Clin Gastroenterol* 2007;41:280–4.
- [31] Hill AF, Zeidler M, Ironside J, Collinge J. Diagnosis of new variant Creutzfeldt–Jakob disease by tonsil biopsy. *Lancet* 1997;349:99.
- [32] Hilton DA, Fathers E, Edwards P, Ironside JW, Zajicek J. Prion immunoreactivity in appendix before clinical onset of variant Creutzfeldt–Jakob disease. *Lancet* 1998;352:703.