



Biologic therapy in the management of asthma

Jennifer L. McCracken*, Julia W. Tripple*, and William J. Calhoun

Purpose of review

Current asthma management relies on inhaled corticosteroids, but some asthma is not well controlled with inhaled steroids alone or in combination with long-acting bronchodilators or leukotriene pathway inhibitors. The field of biologic therapy has grown dramatically in the past two decades, with current availability of three molecules, with two distinct and highly selective approaches to interfering with the allergic and eosinophilic airway inflammation common to most asthma. This review summarizes current and future options for incorporating biologic therapy into the overall management of asthma.

Recent findings

Two new biologic agents have been recently introduced in the United States market, supported by well controlled, randomized clinical trials. These trials have provided insight into the types of patients who are most likely to benefit from these novel agents.

Summary

In asthma patients with frequent exacerbations, the addition of a biologic agent targeting the interleukin-5 pathway, or immunoglobulin E, can significantly reduce exacerbations and improve asthma control. The clinical predictors of utility of specific agents overlap with one another, highlighting the importance of clinical judgment in the overall management of this complex disorder.

Keywords

benralizumab, biologic therapy, IL-13, IL-17, IL-4, mepolizumab, omalizumab

INTRODUCTION

Asthma is a heterogeneous, chronic disease of the airways characterized by reversible airflow obstruction, bronchial hyperresponsiveness, airway inflammation, and recurrent symptoms [1]. It is estimated that 300 million people worldwide have asthma, and the prevalence of disease has been increasing over the last 40 years [2]. The mainstay of asthma therapy is based on severity of disease and control of symptoms and relies on inhaled glucocorticoids (ICS) for patients with persistent disease or worse [1]. For the majority of patients, current treatment options offer good control of their disease, however 10–20% of patients do not achieve control with current gold standards of care [3,4]. This remaining population of severe refractory asthmatic patients is at increased risk of morbidity and mortality related to their disease and make up the majority of economic costs of asthma [5–7].

Over the last decade, a shift towards evaluating specific phenotypes and endotypes of asthma has led to the creation of targeted therapies to fit patient specific disease [4,8–11]. Through a better understanding of the inflammatory modulators involved in asthma, a number of monoclonal antibodies have

emerged with the aim of providing patient tailored asthma treatment.

ASTHMA PATHOGENESIS

Asthma has traditionally been described as a Th2 lymphocyte-mediated condition in which allergen presentation by antigen presenting cells to naïve T cells results in Th2 cell differentiation. The Th2 lymphocytes produce interleukin-4 (IL-4), IL-5, and IL-13 cytokines that then drive B cells to secrete immunoglobulin E (IgE). Mast cell bound IgE is able to bind allergens causing degranulation and release of chemical mediators such as histamine, prostaglandins, and leukotrienes, which induce smooth

Division of Allergy and Clinical Immunology, University of Texas Medical Branch, Galveston, Texas, USA

Correspondence to Dr William J. Calhoun, MD, Professor and Vice Chair for Research, Department of Internal Medicine, JSA 4.116, 301 University Boulevard, Galveston, TX 77555-0568, USA.
E-mail: wjcalhou@utmb.edu

*These authors contributed equally to this work.

Curr Opin Allergy Clin Immunol 2016, 16:375–382

DOI:10.1097/ACI.0000000000000284

KEY POINTS

- Biologic agents for asthma treatment provide targeted therapy for patients with specific asthma phenotypes.
- Ongoing clinical trials are evaluating new biologic agents to help broaden the current arsenal of asthma treatment options.
- Currently omalizumab, mepolizumab, and reslizumab are the only FDA approved monoclonal antibodies available for the treatment of asthma.
- Despite significant costs related to these agents, the benefits of reduced exacerbations, reduced steroid burden, and the potential for decreased healthcare utilization are encouraging.

muscle bronchoconstriction and further stimulate the inflammatory response. IL-5 also acts as a potent regulator for eosinophil proliferation, differentiation, and activation [12]. Patients who have this allergic mediated phenotype generally respond well to inhaled corticosteroids. IgE, IL-4/13, and IL-5 have also been targets recently for the development of biologic agents as discussed below. However, there is a subset of patients with severe disease that do not respond well to ICS. These patients may have a Th-17 predominate disease with high levels of IL-17 leading to more neutrophil predominance and inflammation, less reversible airflow obstruction, and less bronchial hyperresponsiveness with methacholine challenge testing [12]. Another distinct group of asthmatic patients exists with normal levels of both eosinophils and neutrophils. Similarly to neutrophilic asthma, this subset, paucigranulocytic asthma does not respond well to inhaled corticosteroids [13].

IMMUNOGLOBULIN E

IgE is one of the key contributors to the proinflammatory cascade in allergic asthma. Allergens enter the airways and are presented by antigen presenting cells to T lymphocytes, which initiate the cell-mediated immune response [14]. Th2 cells and their associated cytokine milieu stimulate B cells to produce IgE antibodies and proallergic cytokines, such as IL-4, IL-5, IL-9, and IL-13. Free IgE released from B cells binds to the high affinity FCεRI receptor on the surface of mast cells and basophils. The receptor bound IgE is then cross-linked by an allergen and triggers degranulation and release of prostaglandins, leukotrienes, histamine, proteases, and cytokines which all lead to the early allergic response [15].

Omalizumab

Omalizumab was the first US Food and Drug Administration (FDA)-approved biologic for the treatment of severe asthma. It is a humanized anti-IgE antibody that specifically binds the Fc portion of unbound IgE and forms omalizumab:IgE complexes. This reduces free IgE levels and prevents IgE from binding to FCεR1s [16]. Without the binding of IgE to FCεR1s, there can be no degranulation of mast cells and the release of inflammatory mediators is inhibited. IgE also is responsible for regulating its own receptor and by reducing the levels of free IgE, omalizumab also decreases the expression of FCεRI [16].

Several large clinical trials involving adults, adolescents, and children have shown clinical benefits of treatment with omalizumab in patients with severe allergic asthma. Phase III studies [14,17–23] involving these populations have demonstrated a reduction in the rate of asthma exacerbations compared with control or best standard of care. Bousquet *et al.* [20] pooled the data from seven of these studies and found an average reduction in asthma exacerbations of 38% in the omalizumab group compared with placebo. This reduction in asthma exacerbations was also despite reduction in ICS doses. Several studies [14,17,18] have shown the ability to reduce total daily steroid dose by 50% or greater in 72–79% of those receiving omalizumab versus 50–55% in placebo group. Other studies [20,24,25] have shown a reduction in rates of unscheduled outpatient visits, emergency department (ED) visits, and hospitalizations for exacerbations compared with placebo. Because FDA approval has increased availability of omalizumab treatment to more patients, several ‘real-world’ effectiveness studies including the PERSIST study and the eXperience study have been published that confirm the findings of earlier clinical trials. The PERSIST trial showed that after 1 year of treatment there was an 84% improvement in AQLQ (Juniper Asthma Related Quality of Life scale) and 65% were severe exacerbation free [26]. The eXperience study demonstrated similar results showing that the rate of clinically significant asthma exacerbations decreased by 54% at 12 months and 67% at 24 months. There was also a greater than 50% reduction in symptoms and rescue medication use from baseline and maintenance doses of oral corticosteroids decreased as treatment continued [27]. These studies [26,27] also showed that patients receiving omalizumab had fewer hospitalizations, ED visits, and clinic visits.

Omalizumab was designed to be long-term therapy and no specific guidelines have been established for duration of treatment. The INNOVATE trial examined the relationships among free IgE, omalizumab, and clinical outcomes. Once

omalizumab is started, free IgE is very rapidly suppressed, however total symptom score, peak expiratory flow (PEF), and rescue medication use does not reach their new equilibrium until 12–16 weeks. After treatment cessation, free IgE returns to baseline values 18–20 weeks after the last dose in the majority of patients. The return of this value back to baseline correlates with return of symptoms [28]. The XPORT trial was conducted to further evaluate omalizumab persistence after long-term therapy but data have not been published yet. A multicenter, retrospective study [29] conducted in France showed discontinuation of omalizumab was not associated with any rebound effect or exacerbation of the disease and control was sustained at 6 months in nearly half of all patients.

Investigators have tried to identify markers in patients with moderate to severe persistent asthma that would predict the most beneficial response from omalizumab. Several groups have analyzed data from the INNOVATE trial and reported that total serum IgE and antigen specific-IgE are not predictive of response [30,31]. Another group analyzed data from the EXTRA trial evaluating the Th2 inflammatory biomarkers FE_{NO}, peripheral blood eosinophils, and serum periostin as potential targets to predict response. Hanania *et al.* [32] found that higher levels of these biomarkers were associated with greater response to anti-IgE therapy and reduced exacerbations, however the authors cautioned that additional studies are required to explore the value of these biomarkers in clinical practice.

A few studies have looked at the anti-inflammatory effects of omalizumab as well as the pathologic changes on airways. Djukanovic *et al.* [33] showed patients with mild steroid-naïve asthma who received omalizumab had a decrease in eosinophils, IgE+ cells, FcεRI+, IL-4 secreting cells, and CD3+ T lymphocytes within the epithelium and submucosa compared with those who received placebo. This was supported by a study [34] that showed omalizumab reduced the level of circulating T lymphocytes and eosinophils. Hoshino *et al.* [35] showed omalizumab significantly reduced airway wall thickness relative to placebo.

Omalizumab is generally very well tolerated and overall incidence of adverse events with omalizumab is similar to that of the control group. The most common side-effects in both adults and children were nasopharyngitis, headache, upper respiratory infection, and sinusitis [36]. Omalizumab has a black box warning for anaphylaxis but occurs rarely. Anaphylaxis developed in 0.14% of patients compared with 0.07% in the placebo group and can occur anytime in the treatment course. There also

has been increased interest recently in the possibility of increased risk in malignancy with omalizumab. Initial data from phase I to III clinical trials showed a numerical imbalance in the incidence of malignancy between placebo (0.18%) and omalizumab (0.5%), however the rate in the omalizumab group is comparable to that in the general population [36]. The EXCELS (The Epidemiologic Study of Xolair) study was a postmarketing prospective cohort study that was conducted to look at this issue further. Results did not show any increase in the risk of primary malignancy with long-term omalizumab [37]. Analysis of data from the EXCELS trial also noted an increased rate of transient ischemic attacks and ischemic stroke for those treated with omalizumab. This conclusion prompted the FDA to request a thorough analysis of arterial thromboembolic events (ATEs) in all omalizumab studies but the pooled data did not support an increased risk of ATE with omalizumab [38].

INTERLEUKIN-4/INTERLEUKIN-13

IL-4 and IL-13, expressed by Th2 cells and mast cells, are key cytokines in the pathogenesis of atopic asthma. Redundancy in the IL4/IL-13 pathways results from the complex receptor system that shares IL-4 receptor α (IL-4R α). Binding of this receptor mediates downstream signaling through signal transducer and activator of transcription factor 6 (STAT-6) leading to airway inflammation via eosinophil, macrophage and dendritic cell activation, airway remodeling via fibroblast proliferation and activation, B cell activation leading to IgE class switching, stimulation of airway epithelial and goblet cells leading to mucus secretion, and activation of airway smooth muscle cells resulting in airway hyperresponsiveness (AHR) [39].

Lebrikizumab

Lebrikizumab is an immunoglobulin G4 (IgG4) humanized monoclonal antibody that blocks IL-13. In a randomized, double-blind, placebo-controlled study [40] in patients with poorly controlled asthma despite ICS, lebrikizumab treatment was associated with improved lung function. At 12 weeks, the mean forced expired volume in 1 second (FEV1) was 5.5% higher in the lebrikizumab group than in the placebo group. Interestingly, improvements in lung function were more marked in patients receiving lebrikizumab with a high pretreatment serum periostin levels compared with patients with low pretreatment periostin levels. Periostin is a matricellular protein secreted by bronchial

epithelial cells in response to IL-13 [41,42]. These results suggest that serum periostin may be an important serum biomarker for patients with IL-13-induced asthma and may help identify those patients that will be most responsive to anti-IL-13 treatment [43]. However, a more recent dose-ranging study [44], evaluating subcutaneous lebrikizumab at 125, 250, or 500 mg versus placebo in asthmatic patients not receiving ICS, found that although FEV1 increased in all lebrikizumab groups compared with placebo the increase was not statistically or clinically significant. In addition, unlike previous studies there was no meaningful difference in FEV1 in patients with high serum periostin versus patients with low serum periostin. Further studies are needed to evaluate these findings.

Tralokinumab

Tralokinumab is an IgG4 monoclonal antibody to IL-13 that has been shown to inhibit bronchial hyperresponsiveness and airway eosinophilia in preclinical trials [45]. In a recent phase II study [46] in moderate to severe asthma, tralokinumab was associated with improved lung function as measured by FEV1 as well as decreased use of short-acting beta agonists. Interestingly, the increase in FEV1 was higher in the tralokinumab treated group with elevated sputum IL-13 when compared with patients with no sputum IL-13. In addition, the increase in FEV1 remained elevated for 12 weeks following tralokinumab cessation. The study did not however show any improvement in ACQ-6 (6-item Asthma Control Questionnaire) scores in the tralokinumab group compared with placebo. In general, tralokinumab was well tolerated with an acceptable safety profile.

Pitrakinra

Pitrakinra is a recombinant human IL-4 variant that competitively inhibits the IL-4R α complex thus interfering with downstream signaling of both IL-4 and IL-13. In patients with atopic asthma, pitrakinra was shown to lessen the effects of the late asthmatic response after allergen challenge noted by more attenuated decreases in FEV1 in the pitrakinra treated group compared with placebo [47]. In a pharmacogenetic investigation of patients with moderate to severe asthma, patients with specific single nucleotide polymorphisms in the 3' untranslated region of IL-4R α had significantly reduced exacerbation rates when treated with pitrakinra 10 mg compared with placebo suggesting that this subgroup of patients may be more responsive to pitrakinra therapy [48].

Dupilumab

Another IL-4R α target is dupilumab, a fully human monoclonal antibody to the IL-4R α subunit that inhibits the actions of both IL-4 and IL-13. In a randomized, double-blind, placebo-controlled phase IIA study [49], once weekly dupilumab 300 mg subcutaneous versus placebo significantly reduce asthma exacerbation in patients with moderate to severe asthma and elevated eosinophil levels. In addition, dupilumab led to marked improvement in FEV1, improvement in ACQ scores, decreased symptoms, nocturnal awakening, and short-acting beta agonist usage. Moreover, dupilumab reduced biomarkers associated with Th2 inflammation including FE_{NO}, eotaxin-3, TARC (thymus and activation-regulated chemokine), and serum IgE.

INTERLEUKIN-5

As mentioned earlier, eosinophils are believed to play an important role in the pathogenesis and severity of asthma. Several studies [50–52] have shown a reduction in acute exacerbations when sputum eosinophil counts were maintained at low levels by adjustments in inhaled corticosteroids. IL-5 is a proinflammatory cytokine secreted by T lymphocytes, mast cells, and eosinophils. It is heavily involved in the regulation of eosinophil differentiation, proliferation, and activation [53]. For these reasons, targeting IL-5 as a therapeutic option in patients with eosinophilic asthma has been pursued.

Mepolizumab

Mepolizumab is a humanized monoclonal antibody against IL-5 that selectively inhibits eosinophilic inflammation and reduces the number of eosinophils in both sputum and blood [54,55]. Despite showing reduction in blood and sputum eosinophilia, initial trials failed to show improvement in outcomes with mepolizumab in patients with asthma [54,56]. However, more recent targeted studies have shown benefit in patients with a severe eosinophilic asthma phenotype. Nair *et al.* evaluated the steroid sparing effect of mepolizumab in patients with sputum eosinophilia and airway symptoms despite treatment with prednisone and high-dose ICS. In this small study [57], they found that mepolizumab, when compared with placebo, reduced the number of blood and sputum eosinophils and allowed prednisone sparing without the development of asthma exacerbations. Mepolizumab therapy was also shown to reduce the number of exacerbations and improve AQLQ scores in patients with refractory eosinophilic asthma and a

history of recurrent severe exacerbations despite high doses of corticosteroids [58].

The DREAM (Dose Ranging Efficacy and Safety with Mepolizumab in Severe Asthma) study, a large, multicenter, double-blind, placebo-controlled study, investigated the effect of mepolizumab therapy on the rate of asthma exacerbations in patients with a history of recurrent severe asthma exacerbations and signs of eosinophilic inflammation. Over a year of treatment, mepolizumab reduced the rate of clinically significant exacerbations and was well tolerated in this population. Interestingly, the reduction in the rate of clinically significant exacerbations varied only according to baseline peripheral blood eosinophil count and exacerbation frequency in the previous year. In addition, IgE concentration and atopic status at baseline were not associated with response to mepolizumab [59].

More recently, the MENSA (Mepolizumab as Adjunctive Therapy in Patients with Severe Asthma) [60^{***}] and SIRIUS (Steroid Reduction with Mepolizumab Study) [61^{***}] studies have noted benefit with mepolizumab in patients with eosinophilic asthma. The MENSA study compared the rate of exacerbations in patients with recurrent asthma exacerbations and evidence of eosinophilic inflammation despite high doses of ICS in patients on intravenous or subcutaneous mepolizumab to placebo. When compared with placebo, the rate of exacerbations of patients receiving intravenous mepolizumab was reduced by 47 and 53% for subcutaneous mepolizumab, and exacerbations requiring emergency room visits or hospitalization decreased by 32% for intravenous and 61% for subcutaneous mepolizumab. In addition, patients in both the intravenous and subcutaneous mepolizumab groups showed significant improvement in quality of life and asthma control as assessed by SGRQ (St. George's Respiratory Questionnaire) and ACQ-5 (5-item Asthma Control Questionnaire). The SIRIUS study evaluated the steroid sparing effects of mepolizumab compared with placebo in patients with severe asthma and peripheral blood eosinophilia while on maintenance systemic glucocorticoids and found that the likelihood of a reduction in glucocorticoid dose was 2.39 times greater in the mepolizumab group than placebo with the median percentage reduction in glucocorticoid dose being 50%. In addition, despite being on lower doses of glucocorticoids the mepolizumab group had reduced exacerbation rates and improved asthma control. Taken together, these studies further support evidence showing the effectiveness of mepolizumab in patients with a severe eosinophilic asthma phenotype.

It is important to note that cessation of mepolizumab therapy resulted in return of pretreatment blood and sputum eosinophilia levels within 3 months of cessation and was associated with a loss of asthma control as patients were back to pretreatment exacerbation levels within 3–6 months of cessation of therapy [62]. In addition, despite promising data related to decreased exacerbation rates in patients with severe eosinophilic asthma, in the majority of studies mepolizumab fails to show a significant effect on more traditional markers of asthma control, like FEV1, PEF, AHR, AQLQ, and ACQ [63].

Mepolizumab, under the trade name Nucala, has recently received FDA approval for add-on maintenance treatment of patients 12 years and older with severe asthma and an eosinophilic phenotype.

Reslizumab

Reslizumab, an alternative humanized monoclonal IL-5 antibody, neutralizes circulating IL-5 by preventing its binding to eosinophils. Castro *et al.* [64] evaluated the effectiveness of reslizumab in patients with eosinophilic asthma poorly controlled on high-dose ICS over a 15-week protocol. Compared with placebo, patients receiving reslizumab showed significant improvements in airway function and a trend toward greater asthma control as measured by ACQ score. Interestingly, these effects were most pronounced in patients with asthma and nasal polyps – an eosinophil mediated disorder. More recently, the results of two duplicate, multicenter, double-blind placebo-controlled phase III studies showed efficacy for the use of reslizumab for patients with asthma and eosinophilia who are poorly controlled on ICS therapy [65^{***}]. In these studies, reslizumab when compared with placebo significantly reduced the frequency of asthma exacerbation rate. Reslizumab, under the trade name CINQAIR, recently received FDA approval for add-on maintenance treatment of patients with severe asthma 18 years and older with an eosinophilic phenotype.

Benralizumab

Benralizumab is a humanized afucosylated monoclonal antibody that targets the human IL-5 receptor alpha (IL5R α) found on eosinophils and basophils. *In vitro*, the lack of a fucose moiety on the oligosaccharide core improves binding affinity of benralizumab and enhances antibody-dependent cell-mediated cytotoxicity leading to apoptosis of target cells [66]. In an initial study [67] evaluating patients with mild atopic asthma, a single intravenous dose of benralizumab was found to decrease

peripheral blood eosinophil counts. Subsequently, in a phase I trial evaluating the safety of benralizumab and its effects on eosinophil counts in patients with eosinophilic asthma showed benralizumab was safe and reduced eosinophil counts in airway mucosa, sputum, bone marrow, and peripheral blood [68]. More recently, a year-long phase IIb dose-ranging study [69^{***}] found that benralizumab 100 mg tended to reduce exacerbations and improve lung function and quality of life in patients with uncontrolled, eosinophilic asthma. Improvements in exacerbation rate, FEV1, and ACQ-6 scores were noted specifically for patients with increased blood eosinophil counts compared with patients without peripheral eosinophilia suggesting blood eosinophil counts may be helpful in identifying those patients who will respond best to benralizumab therapy. Further, evaluation and confirmation of these initial promising data are underway in ongoing phase III studies.

Taken together, these three drugs targeting IL-5 show the effects of IL-5 inhibition on eosinophilic inflammation and exacerbations in asthma.

INTERLEUKIN-17

Th17 cells are a subset of CD4⁺ T cells associated with a more severe phenotype of asthma that is less responsive to corticosteroids and are responsible for secreting the cytokines IL-17A, IL-17F, IL-22, and IL-21. Studies have demonstrated increased levels of IL-17A, IL-17F, and IL-22 in the bronchoalveolar lavage (BAL) fluid and bronchial biopsies of patients with moderate and severe asthma [70]. IL-17A, IL-17F, and IL-22 have all been implicated in causing increased neutrophilic airway inflammation, mucous cell metaplasia, and smooth muscle proliferation [71,72].

Secukinumab and brodalumab

Biologic agents targeting IL-17A or IL-17RA signaling are currently in clinical trials for asthma. Secukinumab is a monoclonal antibody targeted at IL-17A and has shown a reduction in clinical symptoms in other Th17-mediated diseases such as psoriasis and rheumatoid arthritis. A phase II clinical trial has been completed involving patients with uncontrolled asthma but results are not yet available. Brodalumab, a monoclonal antibody directed against IL-17RA, was recently tested in a phase II clinical trial for patients with moderate to severe asthma using change in Asthma Quality Control (AQC score) as the primary outcome. Results showed no difference in the AQC score between those treated with brodalumab and those who

Table 1. Biologic agents and their targets

Target	Treatment	Stage of development
IgE	Omalizumab	Marketed
IL-5	Mepolizumab and reslizumab	Marketed
IL-5R	Benralizumab	Phase III
IL-4R α (IL-4/IL-13)	Dupilumab and pitakinra	Phase III, phase II
IL-13	Lebrikizumab and tralokinumab	Phase III
IL-17	Secukinumab and brodalumab	Phase II

received placebo however a clinically meaningful response was seen in bronchodilator reversibility [71–73].

CONCLUSION

Asthma is a complex disease afflicting millions worldwide and is associated with a significant financial healthcare burden. As we have developed a better understanding of the inflammatory markers involved in the disease and the various phenotypes and endotypes of asthmatic patients, biologic agents, like the ones described in this review, present new therapeutic options for the treatment of severe disease (Table 1). Biologic agents are promising as they allow the specific inhibition of relevant asthma pathways, have a long duration of action allowing infrequent dosing schedules, have generally good safety profiles with few off-target effects, and provide a means to accomplish personalized treatment. Although promising, there is significant cost associated with these agents. In addition, as they only offer specific inhibition of particular asthma pathways, their spectrum of efficacy is narrower than standard therapy and does not address all the objectives of asthma management possibly necessitating predictive biomarkers for practical implementation. Despite some of these disadvantages, biologic agents have significant promise for the treatment of refractory asthma and further investigation into their therapeutic benefits is warranted.

Acknowledgements

The support of the UTMB CTSA (UL1 TR001439) is gratefully acknowledged.

Financial support and sponsorship

None.

Conflicts of interest

W.J.C. has consulted for Genentech. The remaining authors have no conflicts of interest.

REFERENCES AND RECOMMENDED READING

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Expert Panel Report 3 (EPR-3). Guidelines for the diagnosis and management of asthma – summary report. *J Allergy Clin Immunol* 2007; 120:S94–S138.
 2. WHO. Global surveillance, prevention, and control of chronic respiratory diseases: a comprehensive approach. 2007.
 3. Bateman ED, Boushey HA, Bousquet J, *et al.* Can guideline-defined asthma control be achieved? The Gaining Optimal Asthma Control study. *Am J Respir Crit Care Med* 2004; 170:836–844.
 4. Bousquet J, Mantzouranis E, Cruz AA, *et al.* Uniform definition of asthma severity, control, and exacerbations: document presented for the World Health Organization Consultation on Severe Asthma. *J Allergy Clin Immunol* 2010; 126:926–938.
 5. Antonicelli L, Bucca C, Neri M, *et al.* Asthma severity and medical resource utilisation. *Eur Respir J* 2004; 23:723–729.
 6. Serra-Batllés J, Plaza V, Morejon E, *et al.* Costs of asthma according to the degree of severity. *Eur Respir J* 1998; 12:1322–1326.
 7. Weiss KB, Sullivan SD. The health economics of asthma and rhinitis. I. Assessing the economic impact. *J Allergy Clin Immunol* 2001; 107:3–8.
 8. Wenzel S. Severe asthma: from characteristics to phenotypes to endotypes. *Clin Exp Allergy* 2012; 42:650–658.
 9. Miranda C, Busacker A, Balzar S, *et al.* Distinguishing severe asthma phenotypes: role of age at onset and eosinophilic inflammation. *J Allergy Clin Immunol* 2004; 113:101–108.
 10. Haldar P, Pavord ID, Shaw DE, *et al.* Cluster analysis and clinical asthma phenotypes. *Am J Respir Crit Care Med* 2008; 178:218–224.
 11. Moore WC, Meyers DA, Wenzel SE, *et al.* Identification of asthma phenotypes using cluster analysis in the Severe Asthma Research Program. *Am J Respir Crit Care Med* 2010; 181:315–323.
 12. Trevor JL, Deshane JS. Refractory asthma: mechanisms, targets, and therapy. *Allergy* 2014; 69:817–827.
 13. Simpson JL, Scott R, Boyle MJ, Gibson PG. Inflammatory subtypes in asthma: assessment and identification using induced sputum. *Respirology* 2006; 11:54–61.
 14. Holgate ST, Chuchalin AG, Hebert J, *et al.* Efficacy and safety of a recombinant anti-immunoglobulin E antibody (omalizumab) in severe allergic asthma. *Clin Exp Allergy* 2004; 34:632–638.
 15. Wenzel SE. Asthma phenotypes: the evolution from clinical to molecular approaches. *Nat Med* 2012; 18:716–725.
 16. Kuhl K, Hanania NA. Targeting IgE in asthma. *Curr Opin Pulm Med* 2012; 18:1–5.
 17. Busse W, Corren J, Lanier BQ, *et al.* Omalizumab, anti-IgE recombinant humanized monoclonal antibody, for the treatment of severe allergic asthma. *J Allergy Clin Immunol* 2001; 108:184–190.
 18. Soler M, Matz J, Townley R, *et al.* The anti-IgE antibody omalizumab reduces exacerbations and steroid requirements in allergic asthmatics. *Eur Respir J* 2001; 18:254–261.
 19. Ayres JG, Higgins B, Chilvers ER, *et al.* Efficacy and tolerability of anti-immunoglobulin E therapy with omalizumab in patients with poorly controlled (moderate-to-severe) allergic asthma. *Allergy* 2004; 59:701–708.
 20. Bousquet J, Cabrera P, Berkman N, *et al.* The effect of treatment with omalizumab, an anti-IgE antibody, on asthma exacerbations and emergency medical visits in patients with severe persistent asthma. *Allergy* 2005; 60:302–308.
 21. Humbert M, Beasley R, Ayers J, *et al.* Benefits of omalizumab as add-on therapy in patients with severe persistent asthma who are inadequately controlled despite best available therapy (GINA 2002 step 4 treatment): INNOVATE. *Allergy* 2005; 60:309–316.
 22. Lanier B, Bridges T, Kulus M, *et al.* Omalizumab for the treatment of exacerbations in children with inadequately controlled allergic (IgE-mediated) asthma. *J Allergy Clin Immunol* 2009; 124:1210–1216.
 23. Milgrom H, Berger W, Nayak A, *et al.* Treatment of childhood asthma with anti-immunoglobulin E antibody (omalizumab). *Pediatrics* 2001; 108:E36.
 24. Corren J, Casale T, Deniz Y, Ashby M. Omalizumab, a recombinant humanized anti-IgE antibody, reduces asthma-related emergency room visits and hospitalizations in patients with allergic asthma. *J Allergy Clin Immunol* 2003; 111:87–90.
 25. Normasell R, Walker S, Milan SJ, *et al.* Omalizumab for asthma in adults and children. *Cochrane Database Syst Rev* 2014; CD003559.
 26. Brusselle G, Michils A, Louis R, *et al.* “Real-life” effectiveness of omalizumab in patients with severe persistent allergic asthma: the PERSIST study. *Respir Med* 2009; 103:1633–1642.
 27. Braunstahl GJ, Chen CW, Maykut R, *et al.* The eXpErience registry: the ‘real-world’ effectiveness of omalizumab in allergic asthma. *Respir Med* 2013; 107:1141–1151.
 28. Slavin RG, Ferioli C, Tannenbaum SJ, *et al.* Asthma symptom re-emergence after omalizumab withdrawal correlates well with increasing IgE and decreasing pharmacokinetic concentrations. *J Allergy Clin Immunol* 2009; 123:107–113; e3.
 29. Molimard M, Mala L, Bourdeix I, Le Gros V. Observational study in severe asthmatic patients after discontinuation of omalizumab for good asthma control. *Respir Med* 2014; 108:571–576.
 30. Bousquet J, Rabe K, Humbert M, *et al.* Predicting and evaluating response to omalizumab in patients with severe allergic asthma. *Respir Med* 2007; 101:1483–1492.
 31. Wahn U, Martin C, Freeman P, *et al.* Relationship between pretreatment specific IgE and the response to omalizumab therapy. *Allergy* 2009; 64:1780–1787.
 32. Hanania NA, Wenzel S, Rosen K, *et al.* Exploring the effects of omalizumab in allergic asthma: an analysis of biomarkers in the EXTRA study. *Am J Respir Crit Care Med* 2013; 187:804–811.
 33. Djukanovic R, Wilson SJ, Kraft M, *et al.* Effects of treatment with anti-immunoglobulin E antibody omalizumab on airway inflammation in allergic asthma. *Am J Respir Crit Care Med* 2004; 170:583–593.
 34. Noga O, Hanf G, Brachmann I, *et al.* Effect of omalizumab treatment on peripheral eosinophil and T-lymphocyte function in patients with allergic asthma. *J Allergy Clin Immunol* 2006; 117:1493–1499.
 35. Hoshino M, Ohtawa J. Effects of adding omalizumab, an anti-immunoglobulin E antibody, on airway wall thickening in asthma. *Respiration* 2012; 83:520–528.
 36. Corren J, Casale TB, Lanier B, *et al.* Safety and tolerability of omalizumab. *Clin Exp Allergy* 2009; 39:788–797.
 37. Long A, Rahmaoui A, Rothman KJ, *et al.* Incidence of malignancy in patients with moderate-to-severe asthma treated with or without omalizumab. *J Allergy Clin Immunol* 2014; 134:560–567; e4.
- This is a postmarketing analysis that addressed the concern that omalizumab may increase the risk of malignancy.
38. Humbert M, Busse W, Hanania NA, *et al.* Omalizumab in asthma: an update ■ on recent developments. *J Allergy Clin Immunol Pract* 2014; 2:525–536; e1. This article provides an excellent review of omalizumab, including pivotal studies and clinical safety.
 39. Oh CK, Geba GP, Molfino N. Investigational therapeutics targeting the IL-4/IL-13/STAT-6 pathway for the treatment of asthma. *Eur Respir Rev* 2010; 19:46–54.
 40. Corren J, Lemanske RF, Hanania NA, *et al.* Lebrikizumab treatment in adults with asthma. *N Engl J Med* 2011; 365:1088–1098.
 41. Woodruff PG, Boushey HA, Dolganov GM, *et al.* Genome-wide profiling identifies epithelial cell genes associated with asthma and with treatment response to corticosteroids. *Proc Natl Acad Sci USA* 2007; 104:15858–15863.
 42. Sidhu S, Yuan S, Innes AL, *et al.* Roles of epithelial cell-driven periostin in TGF- β activation, collagen production, and collagen gel elasticity in asthma. *Proc Natl Acad Sci U S A* 2010; 107:14170–14175.
 43. Jia G, Erickson RW, Choy DF, *et al.* Periostin is a systemic biomarker of eosinophilic airway inflammation in asthmatic patients. *J Allergy Clin Immunol* 2012; 130:647–654; e10.
 44. Noonan M, Korenblat P, Mosesova S, *et al.* Dose-ranging study of lebrikizumab in asthmatic patients not receiving inhaled steroids. *J Allergy Clin Immunol* 2013; 132:567–574; e12.
 45. Blanchard C, Mishra A, Soto-Hakel H, *et al.* Inhibition of human interleukin-13-induced respiratory and esophageal inflammation by anti-human interleukin-13 antibody (CAT-354). *Clin Exp Allergy* 2005; 35:1096–1103.
 46. Piper E, Brightling C, Niven R, *et al.* A phase II placebo-controlled study oftralokinumab in moderate-to-severe asthma. *Eur Respir J* 2013; 41:330–338.
 47. Wenzel S, Wilbraham D, Fuller R, *et al.* Effect of an interleukin-4 variant on late phase asthmatic response to allergen challenge in asthmatic patients: results of two phase 2a studies. *Lancet* 2007; 370:1422–1431.
 48. Slager RE, Otulana BA, Hawkins GA, *et al.* IL-4 receptor polymorphisms predict reduction in asthma exacerbations during response to an anti-IL-4 receptor alpha antagonist. *J Allergy Clin Immunol* 2012; 130:516–522; e4.
 49. Wenzel S, Ford L, Pearlman D, *et al.* Dupilumab in persistent asthma with elevated eosinophil levels. *N Engl J Med* 2013; 368:2455–2466.
 50. Green RH, Brightling CE, McKenna S, *et al.* Asthma exacerbations and sputum eosinophil counts: a randomised controlled trial. *Lancet* 2002; 360:1715–1721.
 51. Jayaram L, Pizzichini MM, Cook RJ, *et al.* Determining asthma treatment by monitoring sputum cell counts: effect on exacerbations. *Eur Respir J* 2006; 27:483–494.
 52. Chlumsky J, Striz I, Terl M, *et al.* Strategy aimed at reduction of sputum eosinophils decreases exacerbation rate in patients with asthma. *J Int Med Res* 2006; 34:129–139.
 53. Takatsu K, Nakajima H. IL-5 and eosinophilia. *Curr Opin Immunol* 2008; 20:288–294.
 54. Flood-Page P, Menzies-Gow A, Phipps S, *et al.* Anti-IL-5 treatment reduces deposition of ECM proteins in the bronchial subepithelial basement membrane of mild atopic asthmatics. *J Clin Invest* 2003; 112:1029–1036.

55. Menzies-Gow A, Flood-Page P, Sehmi R, *et al.* Anti-IL-5 (mepolizumab) therapy induces bone marrow eosinophil maturational arrest and decreases eosinophil progenitors in the bronchial mucosa of atopic asthmatics. *J Allergy Clin Immunol* 2003; 111:714–719.
56. Leckie MJ, ten Brinke A, Khan J, *et al.* Effects of an interleukin-5 blocking monoclonal antibody on eosinophils, airway hyper-responsiveness, and the late asthmatic response. *Lancet* 2000; 356:2144–2148.
57. Nair P, Pizzichini M, Kjarsgaard M, *et al.* Mepolizumab for prednisone-dependent asthma with sputum eosinophilia. *N Engl J Med* 2009; 360:985–993.
58. Haldar P, Brightling C, Hargadon B, *et al.* Mepolizumab and exacerbations of refractory eosinophilic asthma. *N Engl J Med* 2009; 360:973–984.
59. Pavord ID, Korn S, Howarth P, *et al.* Mepolizumab for severe eosinophilic asthma (DREAM): a multicentre, double-blind, placebo-controlled trial. *Lancet* 2012; 380:651–659.
60. Ortega HG, Liu MC, Pavord ID, *et al.* Mepolizumab treatment in patients with severe eosinophilic asthma. *N Engl J Med* 2014; 371:1198–1207. This MENSA study showed decreased exacerbation rates and improved quality of life and asthma control scores when compared with placebo. This study helped lead to licensing of mepolizumab in the US.
61. Bel EH, Wenzel SE, Thompson PJ, *et al.* Oral glucocorticoid-sparing effect of mepolizumab in eosinophilic asthma. *N Engl J Med* 2014; 371:1189–1197. The SIRIUS study showed the steroid sparing effects of mepolizumab in eosinophilic asthma and also helped lead to licensing of mepolizumab in the US.
62. Haldar P, Brightling CE, Singapuri A, *et al.* Outcomes after cessation of mepolizumab therapy in severe eosinophilic asthma: a 12-month follow-up analysis. *J Allergy Clin Immunol* 2014; 133:921–923. This study showed that cessation of mepolizumab therapy resulted in return of pretreatment blood and sputum eosinophilia levels within 3 months and was associated with a loss of asthma control.
63. Liu Y, Zhang S, Li DW, Jiang SJ. Efficacy of anti-interleukin-5 therapy with mepolizumab in patients with asthma: a meta-analysis of randomized placebo-controlled trials. *PLoS One* 2013; 8:e59872.
64. Castro M, Mathur S, Hargreave F, *et al.* Reslizumab for poorly controlled, eosinophilic asthma: a randomized, placebo-controlled study. *Am J Respir Crit Care Med* 2011; 184:1125–1132.
65. Castro M, Zangrilli J, Wechsler ME, *et al.* Reslizumab for inadequately controlled asthma with elevated blood eosinophil counts: results from two multicentre, parallel, double-blind, randomised, placebo-controlled, phase 3 trials. *Lancet Respir Med* 2015; 3:355–366. These phase III trials showed that reslizumab for inadequately controlled eosinophilic asthma improved exacerbation rates compared with placebo and lead to licensing of reslizumab in the US.
66. Kolbeck R, Kozhich A, Koike M, *et al.* MEDI-563, a humanized anti-IL-5 receptor alpha mAb with enhanced antibody-dependent cell-mediated cytotoxicity function. *J Allergy Clin Immunol* 2010; 125:1344–1353; e2.
67. Busse WW, Katial R, Gossage D, *et al.* Safety profile, pharmacokinetics, and biologic activity of MEDI-563, an anti-IL-5 receptor alpha antibody, in a phase I study of subjects with mild asthma. *J Allergy Clin Immunol* 2010; 125:1237–1244; e2.
68. Lavolette M, Gossage DL, Gaureau G, *et al.* Effects of benralizumab on airway eosinophils in asthmatic patients with sputum eosinophilia. *J Allergy Clin Immunol* 2013; 132:1086–1096; e5.
69. Castro M, Wenzel SE, Bleeker ER, *et al.* Benralizumab, an anti-interleukin 5 receptor α monoclonal antibody, versus placebo for uncontrolled eosinophilic asthma: a phase 2b randomised dose-ranging study. *Lancet Respir Med* 2014; 2:879–890. This phase IIb dose-ranging study found that benralizumab reduced exacerbations and improved lung function and quality of life in patients with uncontrolled, eosinophilic asthma.
70. Molet S, Hamid Q, Davoine F, *et al.* IL-17 is increased in asthmatic airways and induces human bronchial fibroblasts to produce cytokines. *J Allergy Clin Immunol* 2001; 108:430–438.
71. Newcomb DC, Peebles RS Jr. Th17-mediated inflammation in asthma. *Curr Opin Immunol* 2013; 25:755–760.
72. Morishima Y, Ano S, Ishii Y, *et al.* Th17-associated cytokines as a therapeutic target for steroid-insensitive asthma. *Clin Dev Immunol* 2013; 2013:609395.
73. Busse WW, Holgate S, Kerwin E, *et al.* Randomized, double-blind, placebo-controlled study of brodalumab, a human anti-IL-17 receptor monoclonal antibody, in moderate to severe asthma. *Am J Respir Crit Care Med* 2013; 188:1294–1302.